

AI MENTAL HEALTH ANA – 2

Report on “Demographic breakdown analysis”

Dataset- Adult Psychiatric Morbidity Survey 2014.

**Submitted By-
Deepak Baranwal
-Data Analyst Intern**

Date – 05 May ,2026

1. Introduction:

Mental health conditions can vary across different groups of people. Studying these variations helps in understanding patterns and trends in the population.

The aim of this analysis is to compare the prevalence of mental health severity levels across different demographic groups, mainly focusing on gender. The analysis is based on data from the Adult Psychiatric Morbidity Survey (APMS) 2014.

2. Dataset Description:

The dataset used in this analysis is taken from APMS 2014. It contains percentage values representing different levels of mental health severity.

Instead of specific disorder names, the dataset uses **CIS-R score categories** to represent severity levels:

- 0–5 → No or minimal disorder
- 6–11 → Mild disorder
- 12–17 → Moderate disorder
- 18 or more → Severe disorder
- 12 or more → Clinically significant disorder

For this analysis, three groups were considered:

- Men
- Women
- All Adults

3. Data Preparation:

The dataset was cleaned and prepared before analysis.

The following steps were performed:

- Removed unnecessary rows and labels
- Extracted separate datasets for Men, Women, and All Adults
- Converted data into numeric format
- Calculated average values for each category

This made the dataset ready for comparison and visualization.

Men:

```
i) df_men = df.iloc[0:6]
df_men
```

	Category	Unnamed: 1	Unnamed: 2	Unnamed: 3	Unnamed: 4	Unnamed: 5	Unnamed: 6	Unnamed: 7	Unnamed: 8
0	0-5	76.5	72.3	69.7	72.1	69.9	81.3	81.9	74.0
1	6-11	14.4	12.5	15.2	14.7	15.2	11.5	12.8	13.9
2	Under 12	90.9	84.7	84.9	86.8	85.1	92.7	94.7	87.8
3	12-17	4.9	7.4	6.8	6.2	5.8	3.7	4.2	5.8
4	18 or more	4.2	7.8	8.3	7.0	9.1	3.6	1.1	6.4
5	12 or more	9.1	15.3	15.1	13.2	14.9	7.3	5.3	12.2

Women:

```
[?] df_women = df.iloc[7:13]
df_women
```

	Category	Unnamed: 1	Unnamed: 2	Unnamed: 3	Unnamed: 4	Unnamed: 5	Unnamed: 6	Unnamed: 7	Unnamed: 8
7	0-5	57.3	60.4	63.2	58.6	62.8	72.3	74.6	63.5
8	6-11	16.7	20.5	16.2	18.7	18.1	14.9	15.4	17.4
9	Under 12	74.0	80.9	79.4	77.3	80.9	87.1	90.0	80.9
10	12-17	10.9	10.5	9.4	10.2	9.8	8.1	5.1	9.3
11	18 or more	15.1	8.6	11.2	12.5	9.3	4.8	4.9	9.8
12	12 or more	26.0	19.1	20.6	22.7	19.1	12.9	10.0	19.1

All Adults:

```
df_adults = df.iloc[14:20]
df_adults
```

	Category	Unnamed: 1	Unnamed: 2	Unnamed: 3	Unnamed: 4	Unnamed: 5	Unnamed: 6	Unnamed: 7	Unnamed: 8
14	0-5	67.1	66.3	66.4	65.3	66.3	76.6	77.6	68.6
15	6-11	15.5	16.5	15.7	16.7	16.7	13.2	14.3	15.7
16	Under 12	82.7	82.8	82.1	82.0	82.9	89.8	91.9	84.3
17	12-17	7.8	9.0	8.1	8.2	7.8	5.9	4.7	7.6
18	18 or more	9.5	8.2	9.8	9.8	9.2	4.2	3.3	8.1
19	12 or more	17.3	17.2	17.9	18.0	17.1	10.2	8.1	15.7

4. Methodology:

To perform the demographic analysis:

- The same severity categories were selected for all groups
- Average percentage values were calculated
- Data was combined into a single comparison table
- Bar charts were used to visualize the differences

This approach makes it easier to compare trends across demographic groups.


Combined data :


▼

	Men	Women	All Adults
Category			
0-5	74.7125	64.0875	69.2750
6-11	13.7750	17.2375	15.5375
Under 12	88.4500	81.3125	84.8125
12-17	5.6000	9.1625	7.3875
18 or more	5.9375	9.5250	7.7625
12 or more	11.5500	18.6875	15.1875

5.Data Visualization:

A grouped bar chart was created to compare mental health severity across Men, Women, and All Adults.





6. Observations:

From the analysis, the following observations can be made:

- The “**Under 12**” category has the highest values across all groups.
- Women show slightly higher values in **moderate and severe categories**.
- Men have higher values in **lower severity categories**.
- The “All Adults” group lies between Men and Women, representing overall population trends.
- Severe mental health conditions are less common compared to mild conditions.

7. Ethnicity:

Ethnicity-based analysis could not be performed using this dataset because ethnicity-related variables were not available in the selected table.

8. Conclusion:

This analysis shows that mental health severity varies slightly across gender groups.

Women tend to have slightly higher percentages in moderate and severe categories, while men show higher values in lower severity levels. The “All Adults” group provides an overall view of the population.

Overall, the dataset shows that most individuals fall under lower severity categories, and severe conditions are relatively less common.