Introductory plant breeding: A practical manual

Deependra Dhakal November, 2019

Contents

1	Introduction	5
2	Study of floral parts of field crops	7
	2.1 Objectives	7
	2.2 Theory	7
	2.3 Materials required	10
	2.4 Procedure	10
3	Conclusion	15
4	Hybridization of crops available in the field	17
5	Plant breeding data recording	19
6	Determining genetic purity	21
7	Maintaining genetic purity in the field	23
8	Disease scoring and determining resistance and susceptibility to pests	25
	8.1 Resistance and susceptibility	25
	8.2 Disease scoring	25
9	Describing the traits for release of a new variety	27
10	Study of activities at National Maize Research Programme (NMRP)	29

4 CONTENTS

11 Study of activities at National Grain Legumes Research Programme (NGLRP) 31

Introduction

Introductory plant breeding practical manual is intended for B.Sc.Ag, 4th semester's course in Introductory plant breeding, although a wider readership (field researchers, lecturers, lab researcher and commercial practitioners) may benefit from it. Meanwhile the manually is continually evolving.

Study of floral parts of field crops

2.1 Objectives

- To be familiar with different parts of flower and their functions
- To know the floral morphology and mode of pollination in crops

2.2 Theory

Flowers are the reproductive organs of a plant which lead to the development of fruit and seed. ? How flowers are important part of evolutionary history in crops ?

The floral morphology is the study of forms and features of flowers. ? Add a brief history to this ?

The genetic constitution of a crop depends on the mode of pollination of the crop?. Floral morphology dictates the mode of pollination in flowers. Therefore, it becomes an essential for a plant breeder to know the structural organization and basic nomenclature associated with commonly cultivated crops' flowers.

 $\label{linear_convert} \begin{tabular}{ll} \# pdftools::pdf_convert("/home/deependra/Desktop/BSc_Ag_lectures/literatures/Plant\ Science\ Strause, and the strain of the property of the prope$

knitr::include_graphics("./images/rice_panicle.png")




Figure 2.1: Flower and fruit morphology of poales. Wheat, *Triticum aestivum* with (c) spelt and (d, e) wheat. (k) Wheat grain (caryopsis). (i, j) Rice, *Oryza sativa*. a point of emergence, c coleoptile, cr coleoptile, d lemma, f fruit furrow, h glume, l vascular bundle, r radicle, s scutellum, v palea, vk shoot apex, w root cap, z cylindrical epithelium

2.2. THEORY 9




Figure 2.2: Flower and fruit morphology of poales. Wheat, *Triticum aestivum* with (c) spelt and (d, e) wheat. (k) Wheat grain (caryopsis). (i, j) Rice, *Oryza sativa*. a point of emergence, c coleoptile, cr coleoptile, d lemma, f fruit furrow, h glume, l vascular bundle, r radicle, s scutellum, v palea, vk shoot apex, w root cap, z cylindrical epithelium

```
knitr::include_graphics("./images/wheat_kernel.png")
```

```
knitr::include_graphics("./images/wheat_spikelet.png")
```

```
knitr::include_graphics("./images/wheat_spike.png")
```

The type of habitat that a species inhabits in determines the reproductive biology of that adapted crop to a large extent. A range of vegetation types (based on habitat) are identified for order poales, including: - Woodland - Grassland - Heathland - Wetland - Desert - Polymorphous

Flower morphology in turn affects the choice of tools and techniques to be used for hybridization. Prominance of reproductive organs – Stigma and anther – which is largely conserved in a species (cleistogamous and ?), is among a number of factors determining the mode of reproduction and complexity of undertaking a manual crossing. More on hybridization techniques in Practical 4.

2.3 Materials required

- Flower,
- Scissor,
- Petri dish,
- · Magnifying glass,
- Notebook,
- Pencil

2.4 Procedure

- A fresh sample of flowers are obtained of crops.
- Flowers are dissected in lab, their nature and morphology looked closely
 with the help of magnifying glass, and the observed impression is drawan
 in notebook.

2.4.1 Following structures can be observed

- 1. Wheat/rice
- Bracteate type flower
- Inflorescence: Spike of spikelets or compound spikelets




Figure 2.3: Flower and fruit morphology of poales. Wheat, $Triticum\ aestivum$ with (c) spelt and (d, e) wheat. (k) Wheat grain (caryopsis). (i, j) Rice, $Oryza\ sativa$. a point of emergence, c coleoptile, cr coleoptiza, d lemma, f fruit furrow, h glume, l vascular bundle, r radicle, s scutellum, v palea, vk shoot apex, w root cap, z cylindrical epithelium



Figure 2.4: Flower and fruit morphology of poales. Wheat, $Triticum\ aestivum\ with\ (c)$ spelt and (d, e) wheat. (k) Wheat grain (caryopsis). (i, j) Rice, $Oryza\ sativa$. a point of emergence, c coleoptile, cr coleoptile, d lemma, f fruit furrow, h glume, l vascular bundle, r radicle, s scutellum, v palea, vk shoot apex, w root cap, z cylindrical epithelium




Figure 2.5: Flower and fruit morphology of poales. Wheat, *Triticum aestivum* with (c) spelt and (d, e) wheat. (k) Wheat grain (caryopsis). (i, j) Rice, *Oryza sativa*. a point of emergence, c coleoptile, cr coleoptile, d lemma, f fruit furrow, h glume, l vascular bundle, r radicle, s scutellum, v palea, vk shoot apex, w root cap, z cylindrical epithelium

14 CHAPTER 2. STUDY OF FLORAL PARTS OF FIELD CROPS

- Mode of pollination: Self pollination
- Flowers of wheat and rice are complete and bisexual.
- ? refer to wheat valentine blog post

2. Pea

- Flowers are pedicilate, zygomorphic and hermaphrodite
- Sepals are gamosepalous, pentamerous
- Petals have axillary aestivation
- Androecium: Superior ovary (epigynous)
- Pollination: Self pollinated

3. Potato:

- Bisexual, complete, regular flower, yellow colored and self pollinated
- ?

4. Maize

- In maize, male and female inflorescence are located on different parts, namely tassel and ear, respectively.
- ? details
- 95% of the times, pollination is cross and 5% of the time only self-pollination takes place.

Conclusion

Hence, study of floral morphology done for some of the common crop species has inspired students about the importance of floral biology in overall reproductive habit of a crop and established that it's knowledge can aid in crop improvement activities, mainly through hybridization.

Hybridization of crops available in the field

Here is a review of existing methods.

18CHAPTER 4. HYBRIDIZATION OF CROPS AVAILABLE IN THE FIELD

Plant breeding data recording

We describe our methods in this chapter.

Determining genetic purity

Genetic purity is determined by purity testing of a variety or cultivar.

Maintaining genetic purity in the field

We have finished a nice book.

Disease scoring and determining resistance and susceptibility to pests

Diseases are bad!

- 8.1 Resistance and susceptibility
- 8.2 Disease scoring

26CHAPTER 8. DISEASE SCORING AND DETERMINING RESISTANCE AND SUSCEPTIBILITY

Describing the traits for release of a new variety

We have finished a nice book.

Study of activities at National Maize Research Programme (NMRP)

We have finished a nice book.

 $30 CHAPTER\ 10.\ STUDY\ OF\ ACTIVITIES\ AT\ NATIONAL\ MAIZE\ RESEARCH\ PROGRAMME\ (NNORMAL MAIZE\ RESEARCH\ PROGRAMME\ ($

Study of activities at National Grain Legumes Research Programme (NGLRP)

Is the NGLRP still in Rampur, Chitwan?