Long table mixing with portrait and landscape

Contents

1	Pag	e display modes and table formatting hacks	1
	1.1	Page display modes: portrait and landscape	1
	1.2	xtable: default and sidewaystable environments	3
	1.3	xtable workaround for landscape and portrait tables in same page	3
	1.4	Long table with footnotes using knitr::kable()	3
	1.5	Long table with footnotes using panodoc (default) syntax	4

1 Page display modes and table formatting hacks

1.1 Page display modes: portrait and landscape

- to get every pages in landscape mode, add: classoption: landscape to the yaml header.
- \blandscape and \elandscape commands come from modifications of command in pdflscape package.
- use LATEX pdflscape package instead of lscape package to get automatically rotated view of landscape pages in pdf viewer.

Portrait page

```
##
       speed
                      dist
                 Min. : 2
   Min. : 4.0
                 1st Qu.: 26
  1st Qu.:12.0
## Median :15.0
                 Median: 36
        :15.4
## Mean
                 Mean: 43
## 3rd Qu.:19.0
                 3rd Qu.: 56
                 Max. :120
## Max.
          :25.0
```

Landscape page

##	speed	dist
##	Min. : 4.0	Min. : 2
##	1st Qu.:12.0	1st Qu.: 26
##	Median :15.0	Median : 36
##	Mean :15.4	Mean : 43
##	3rd Qu.:19.0	3rd Qu.: 56
##	Max. :25.0	Max. :120

1.2 xtable: default and sidewaystable environments

• using sidewaystable environment requires rotating package to be used in latex preamble.

Portrait table using xtable without sideway stable environment is shown in Table 1.

	speed	dist
1	Min.: 4.0	Min. : 2
2	1st Qu.:12.0	1st Qu.: 26
3	Median $:15.0$	Median: 36
4	Mean $:15.4$	Mean: 43
5	3rd Qu.:19.0	3rd Qu.: 56
6	Max. $:25.0$	Max. :120

Table 1: Portrait table

Landscape table using xtable with sidewaystable environment is shown in Table 2.

1.3 xtable workaround for landscape and portrait tables in same page

Portrait table is shown in 3.

Landscape table is shown in 4.

1.4 Long table with footnotes using knitr::kable()

Long table formatted with footnote marks in the table using knitr::kable() is shown in Table 5.

Table 5: A summary of check varieties used in the study

Characteristics	${ m Tilottama}^1$	Bhrikuti ²	$Gautam^3$	Aditya ⁴
Year	2014	1994	2002	2010
Breeding program	Nepal Agriculture Research Council	Nepal Agriculture Research Council	Nepal Agriculture Research Council	Nepal Agriculture Research Council
Synonym	NL 1073; NL1073; NL-1073; FRAN- COLIN; FRANKOLIN		BL 1887; BL- 1887, BL1887	
Growth Habit	Spring	Spring	Spring	Spring
Variety Type	Bread Wheat	Bread Wheat	Bread Wheat	Bread Wheat
Semidwarf (Rht gene)	Yes	Yes	Yes	
Grain color				White grain
Grain hardiness				
Pedigree	WAXWING*- 2/VIVITSI	CONTESTAD- O/COCORAQUE F 75/3/POLLO//F- URY/ANAHUAC F 75	SIDDHARTHA/NG 8319//NL 297 or SIDHHARTHA/NG 8319//NL 297	GS348/NL746//NL748

¹http://wheatatlas.org/varieties/detail/26413

²http://wheatatlas.org/varieties/detail/22378

 $^{^3}$ http://wheatatlas.org/varieties/detail/22383

⁴http://wheatatlas.org/varieties/detail/22373

Selection history		NL 623 FR 2308-2F-1F-0F	NC 1838-4B-020B- 020B-2B-0B	
CID (Cross ID)		251774		
SID (Selection ID)		0		
Origin	Advanced line from CIMMYT	Advanced line from CIMMYT	Cross made in the country, one CIMMYT parent	Cross made in the country, no CIMMYT parents
CIMMYT breeding line Genes	FRANCOLIN			
Selection Code		Line from CIMMYT, further local selection within country		
Ug99	Yes	No	No	No
Stem Rust	Yes			
Yellow Rust	Yes			
Leaf Rust	Yes			
ME1		Spring Wheat	Spring Wheat	Spring Wheat
$\overline{\mathrm{ME2}}$		Spring Wheat	Spring Wheat	SF-1138 (
ME3		~F · · ·	S-1-1-9	
Production Conditions	Irrigated and semi-irrigated areas of Terai and inner Terai		Irrigated, both normal and late sown condition of whole Terai, Taar and foot-hills (<500 m); maturity 105-115; potential yield - 5t/ha	Terai, Tars and Lower Valleys up to 500 500 masl; maturity 118 days, yield potential 4.8 t/ha
Suitable megaenviron- ments	ME1	$\mathrm{ME1}$, $\mathrm{ME5}$	$\mathrm{ME1}$, $\mathrm{ME5}$	ME1
Notes	Resistant to spot blotch. High tillering. Early maturing, stay green, excellent for bread and chapati.			
Cultivated area		153,055 (2011/12)		38263 (2011/12)

1.5 Long table with footnotes using panodoc (default) syntax

Four check varieties (Described in Table 6), released at different periods, adapted to a range of environmental conditions in Nepal were included alongside in the study.

Table 6: A summary of check varieties used in the study

	Aditya ⁵	Bhrikuti ⁶	Gautam ⁷	Tilottama ⁸
Year	2010	1994*	2002	2014
released				

	Aditya ⁹	Bhrikuti ¹⁰	Gautam ¹¹	Tilottama ¹²
Synonyr	HD- 2781*	NL-623*	BL- 1887; NC- 1838-4B- 020B- 020B- 2B-0B*	NL 1073; NL1073; NL-1073; FRAN- COLIN; FRANKOLIN
Wheat type	Bread	Bread	Bread	Bread
Semidwa (Rht	No Info	Yes	Yes	Yes
gene) Grain color	White grain	Unknown	Unknown	Unknown
Grain hardines	Unknown	Unknown	Unknown	Unknown
Pedigree	GS348/NL BOBWHIT 306/C- 591/HW- 2004*	CONTESTAL F 75/3/POLLO F 75; CLEMENT/C 75/3/POLLO 75*	8319//NL 297 or SID- HHARTH	WAXWING*2/VIVITS
Selection history	HD- 2781*	NL 623 FR 2308-2F-1F- 0F; NL-623*	NC 1838-4B- 020B- 020B- 2B-0B	CGSS-01-B- 00056T- 099Y-099M- 099M-099Y- 099M-14Y- 0B*
Breeding program	Nepal Agricul- ture Re- search Council(N.	NARC	NARC	NARC
CID (Cross ID)		251774		
Accessic number		K-63729*		
SID (Selection ID)		0		
Origin	Cross made in the country, no CIM- MYT parents	Advanced line from CIMMYT	Cross made in the country, one CIM- MYT parent	Advanced line from CIMMYT

	$Aditya^9$	${ m Bhrikuti^{10}}$	$Gautam^{11}$	${ m Tilottama}^{12}$
CIMMY breed- ing line				FRANCOLIN
Genes	Sr2*; no- Lr34*; Glu- A1b*,Glu- B1i*,Glu- D1a*	Lr34*; Glu- A1b*,Glu- B1i*,Glu- D1a*		Sr2+
Suitable	ME1	ME1, ME5	ME1,	
megaen Notes	Terai, Tars and Lower Valleys up to 500 500 masl; matu- rity 118 days, yield po- tential 4.8 t/ha		ME5 Irrigated, both normal and late sown condition of whole Terai, Taar and foot- hills (<500 m); ma- turity 105-115; poten- tial yield	Resistant to spot blotch. High tillering. Early maturing, stay green, excellent for bread and chapati.
Resistar to Ug99 rust	No	No	- 5t/ha No	Yes(including to Stem, Yellow and Leaf)
Approxi acer- age vari- ety grown	38,263(201 12)	153,055(2011- 12)		Doary

1 Min.: 4.0 Min.: 2 2 1st Qu.:12.0 1st Qu.: 2 3 Median: 15.0 Median: 3 4 Mean: 15.4 Mean: 43 5 3rd Qu.:19.0 3rd Qu.: 5 6 Max.: 25.0 Max.: 120		$_{ m sbeed}$	dist
1st Qu.: Median : Mean : 4 3rd Qu.: Max. :12	\vdash	Min. : 4.0	Min.: 2
Median : Mean : 4 3rd Qu.: Max. :12	2	1st Qu.:12.0	1st Qu.: 26
Mean: 4 0 3rd Qu.: Max.:12	က	Median :15.0	Median: 36
.0 3rd Qu.: Max. :12	4	Mean :15.4	Mean: 43
	ಬ	3rd Qu.:19.0	3rd Qu.: 56
	9	Max. :25.0	Max. :120

Table 2: Landscape table

	speed	dist
1	Min.: 4.0	Min. : 2
2	1st Qu.:12.0	1st Qu.: 26
3	Median $:15.0$	Median: 36
4	Mean $:15.4$	Mean: 43
5	3rd Qu.:19.0	3rd Qu.: 56
6	Max. $:25.0$	Max. :120

Table 3: Portrait table

	beed	dist
	Min. : 4.0	Min.: 2
2	1st Qu.:12.0	1st Qu.: 26
33	Median:15.0	Median: 36
4	Mean :15.4	Mean: 43
2	3rd Qu.:19.0	3rd Qu.: 56
9	Max. :25.0	Max. :120

Table 4: Landscape table