# 1 Conditions for the Existence of Left and Right Inverses

To establish the conditions for the existence of left and right inverses, we introduce the following notation:

Let $A \in M_{m\times n}$

**Definition 1.**

$$\begin{aligned} \text{Col}(A) = \{Ax \in \mathbb{R}^m : x \in \mathbb{R}^n\} \quad \text{Col}(A) \subseteq \mathbb{R}^m \quad \textit{then} \quad \text{rank}(A) \leq m \\ N(A) = \{x \in \mathbb{R}^n : Ax = 0 \in \mathbb{R}^m\} \quad N(A) \subseteq \mathbb{R}^n \quad \textit{then} \quad \textit{nulity}(A) \leq n \end{aligned}$$

**Theorem 1.** $n = \text{rank}(A) + dim(N(A))$

which implies that

$$n = \text{rank}(A) \leq m \iff \textit{nulity}(A) = 0$$

$$\text{rank}(A) = m \iff m \leq n$$

**Proposition 1.** *A has left inverse* $\implies N(A) = \{0\}$

*Proof.* Suppose that $A_l^{-1} \in M_{n\times m}$ is a left inverse of $A$. Then

$$A_l^{-1}A = I_n.$$

Let $x \in N(A)$. Since $Ax = 0$, it follows that

$$\begin{aligned} x &= I_n x \\ &= A_l^{-1}Ax \\ &= A_l^{-1}(0) \\ &= 0 \end{aligned}$$

Hence,

$$N(A) = \{0\}.$$

Furthermore, by the rank-nullity theorem,

$$n = \text{rank}(A) \leq m$$

□

**Proposition 2.** *If $N(A) = \{0\}$, then A has left inverse*

*Proof.* Suppose that $A^TAx = \vec{0}$, then,

$$\begin{aligned} x^TA^TAx &= (Ax)^TAx \\ &= \|Ax\|^2 \\ &= Ax \\ &= \vec{0} \end{aligned}$$

Therefore, $N(A^TA) = N(A) = \{0\}$.

Since $A^TA$ is an $n \times n$ square matrix, the rank-nullity theorem implies that $\text{rank}(A^TA) = n$.

$$(A^TA)^{-1}A^TA = ((A^TA)^{-1}A^T)A = I_n$$

Therefore,

$$(A^TA)^{-1}A^T \text{ is a left inverse of A}$$

□

**Proposition 3.** *If* $\text{rank}(A) = m$*, then A has a right inverse.*

*Proof.* Let $y$ be an arbitrary vector in $\mathbb{R}^m$.

Since $\text{rank}(A) = m$, there exists $x \in \mathbb{R}^m$ such that $Ax = y$, which means $\text{Col}(A) = \mathbb{R}^m$.

Since $\text{rank}(A) = \text{rank}(A^T) = m$ and $m = \text{rank}(A^T) + nulity A^T$.

$$N(A^T) = \{0\}$$

Suppose that $AA^Tx = 0$. Then

$$x^TAA^Tx = \|A^Tx\|^2 = 0$$

$$N(AA^T) \subseteq N(A^T)$$

Since $N(A^T) = \{0\}$ , it follows that $N(AA^T) = \{0\}$.

By using the rank-nulity theorem,

$$\text{rank}(AA^T) = m$$

Because $AA^T$ is a square matrix, it is therefore invertible.

$$AA^T(AA^T)^{-1} = A(A^T(AA^T)^{-1}) = I_m$$

Hence, $A^T(AA^T)^{-1}$ is right inverse of A $\square$