

VESTIBULAR REHABILITATION SYSTEM USING IMMERSIVE VIRTUAL REALITY

Bonafide record of work done by

ARYAA S A	(21Z210)
DEEPIGA S	(21Z214)
NIROSHA V	(21Z234)
THIRIKSHA S	(21Z263)
BRINDHA L	(22Z435)

19Z620 – INNOVATION PRACTICES LAB

MENTOR: Dr. L. S. Jayashree

Dissertation submitted in the partial fulfillment
of the requirements for the degree of

BACHELOR OF ENGINEERING

BRANCH: COMPUTER SCIENCE AND ENGINEERING



APRIL 2024

DEPARTMENT OF COMPUTER SCIENCE AND ENGINEERING

PSG COLLEGE OF TECHNOLOGY

(Autonomous Institution)

COIMBATORE – 641 004

VESTIBULAR REHABILITATION SYSTEM USING IMMERSIVE VIRTUAL REALITY

Bonafide record of work done by

ARYAA S A	(21Z210)
DEEPIGA S	(21Z214)
NIROSHA V	(21Z234)
THIRIKSHA S	(21Z263)
BRINDHA L	(22Z435)

Dissertation submitted in the partial fulfillment
of the requirements for the degree of

BACHELOR OF ENGINEERING

**BRANCH: COMPUTER SCIENCE AND ENGINEERING
of Anna University**

APRIL 2024

.....
Dr. L. S. Jayashree
Faculty guide

.....
Dr.G.Sudha Sadasivam
Head of the Department

Certified that the candidate was examined in the viva-voce examination held on
04.04.2024

.....
(Internal Examiner)

.....
(External Examiner)

TABLE OF CONTENTS	PAGE
ACKNOWLEDGEMENT	6
SYNOPSIS	7
1. INTRODUCTION	8
1.1 VIRTUAL REALITY	9
1.2 MOTIVATION	10
1.3 SCOPE	10
1.4 OBJECTIVE	11
1.5 PROBLEM STATEMENT	12
2. LITERATURE SURVEY	14
2.1 BIBLIOGRAPHY	21
3. IMPLEMENTATION AND SYSTEM DESIGN	22
3.1 IMPLEMENTATION	22
3.1.1.VR ENVIRONMENT	22
3.1.2.GAZE STABILIZATION EXERCISE	23
3.1.3.EXPERIMENTAL RESULTS	26
3.2 DESIGN	33
3.3 HARDWARE REQUIREMENTS	33
3.4 SOFTWARE REQUIREMENTS	34
3.5 FUNCTIONAL REQUIREMENTS	34
3.6 NON-FUNCTIONAL REQUIREMENTS	35
3.7 FEASIBILITY	36
4. CONCLUSION	38
4.1. SUMMARY	38
4.2. FUTURE DEVELOPMENTS	42
4.3. EVALUATION METRICS	43
4.4. REFERENCE	44

ACKNOWLEDGEMENT:

We would like to extend our sincere thanks to our Principal In-charge, **Dr. K. Prakasan**, for providing us with this opportunity to develop and complete our project in our field of study.

We express our sincere thanks to our Head of the Department, **Dr. G.Sudha Sadasivam**, for encouraging and allowing us to present the project at our department premises for the partial fulfillment of the requirements leading to the award of BE degree.

We take immense pleasure in conveying our thanks and a deep sense of gratitude to the Programme Coordinator, **Dr. N. Arul Anand**, Department of Computer Science and Engineering, Coimbatore, for her relentless support and motivation to constantly up skill ourself throughout this project. We would like to express our gratitude to our faculty guide **Dr. L. S. Jayashree**, Professor, Department of Computer Science and Engineering, PSG College of Technology, Coimbatore, for her valuable suggestions, encouragement, whole-hearted cooperation, constructive criticism, and guidance throughout this project work

SYNOPSIS:

This project aims to develop an innovative Vestibular Rehabilitation System that utilizes immersive Virtual Reality (VR) technology to improve the precision and efficacy of vestibular rehabilitation exercises. The system provides real-time feedback to optimize exercises for each patient, resulting in a more engaging and personalized rehabilitation experience. Through immersive VR environments, This data is then used to tailor rehabilitation exercises in real-time, adjusting intensity and complexity based on individual progress and needs.

CHAPTER 1

INTRODUCTION

Balance disorders and vestibular impairments affect a significant portion of the global population, resulting in debilitating symptoms such as vertigo, dizziness, and an increased risk of falls. Traditional vestibular rehabilitation methods often lack precision and real-time feedback, making it challenging for both patients and healthcare professionals to effectively address these issues. Moreover, existing rehabilitation systems may not fully engage patients in their exercises, hindering treatment effectiveness. This project proposes an innovative approach to vestibular rehabilitation by integrating immersive Virtual Reality (VR) technology. With this technology, we improve patient outcomes and quality of life.

Many vestibular rehabilitation systems using immersive VR are game-based, incorporating gaming elements to enhance engagement while some focus primarily on simulating real-world environments and activities relevant to vestibular rehabilitation that include virtual environments. Some patients may experience motion sickness or discomfort when using VR headsets, which can limit the duration and effectiveness of rehabilitation sessions. This system transitions manual exercises and instructions, typically performed in a physical setting, into a virtual reality environment. It bridges the gap between traditional physical rehabilitation methods and immersive VR technology, that combines the tangible benefits of physical objects with the immersive advantages of VR. It provides gradual exposure to VR environments and smooth movements within the environment to reduce motion sickness and discomfort.

1.1. VIRTUAL REALITY:

Virtual reality (VR) is an immersive technology that creates a simulated environment, often through the use of computer-generated imagery, audio, and other sensory stimuli. Users can interact with and explore these virtual environments in a seemingly real or physical way, typically through specialized equipment such as VR headsets or goggles.

VR technology has applications across various fields, including entertainment, gaming, education, training, healthcare, and therapy. It offers users the ability to experience situations and scenarios that may be difficult, dangerous, or impossible to encounter in the real world.

In healthcare, VR is increasingly being used for medical training, surgical simulation, pain management, exposure therapy, and rehabilitation. By providing immersive and interactive experiences, VR has the potential to enhance learning, improve patient outcomes, and increase engagement in therapeutic interventions.

Overall, virtual reality represents a powerful tool for creating immersive experiences that can entertain, educate, and even transform how we interact with and perceive the world around us.

1.2. MOTIVATION:

The motivation behind developing a Vestibular Rehabilitation System using Immersive Virtual Reality (VR) stems from the significant impact of balance disorders and vestibular impairments on individuals' quality of life worldwide. These conditions often lead to debilitating symptoms such as vertigo, dizziness, and an increased risk of falls, which can significantly impair daily activities and overall well-being.

Traditional rehabilitation methods for vestibular disorders may lack precision, real-time feedback, and patient engagement, hindering effective treatment outcomes. By leveraging immersive VR technology, this project seeks to address these limitations and provide a more effective, engaging, and personalized approach to vestibular rehabilitation.

1.3. SCOPE:

The scope of the Vestibular Rehabilitation System using Immersive Virtual Reality (VR) encompasses several key aspects:

Development of VR environments: This involves creating immersive virtual environments that simulate various scenarios to challenge and trigger vestibular responses in patients.

Customization and personalization: The system will be designed to tailor rehabilitation exercises according to each patient's progress, symptoms, and specific rehabilitation goals.

Evaluation and validation: The effectiveness of the VR-based rehabilitation system will be evaluated through clinical trials and user studies to assess its impact on balance improvement, symptom reduction, patient engagement, and overall treatment outcomes.

Scalability and usability: The system will be designed with scalability and usability in

mind to facilitate its integration into clinical practice and ensure accessibility to a wide range of patients and healthcare settings.

1.4. OBJECTIVE:

The primary objective of this project is to develop an innovative Vestibular Rehabilitation System using Immersive Virtual Reality (VR) that enhances the precision, efficacy, and engagement of vestibular rehabilitation exercises. Specifically, the objectives include:

Designing and developing immersive VR environments that simulate challenging scenarios to trigger vestibular responses. Providing real-time feedback based to customize exercise intensity and complexity for individual patients.

Evaluating the effectiveness of the VR-based rehabilitation system in improving balance, reducing symptoms, and enhancing patient outcomes compared to traditional rehabilitation methods. Enhancing patient engagement and adherence to rehabilitation exercises through immersive and interactive VR experiences.

1.5. PROBLEM STATEMENT:

Current vestibular rehabilitation methods face several challenges, including a lack of precision, real-time feedback, and patient engagement. Traditional approaches often rely on subjective assessments and generic exercises that may not adequately address individual patient needs. Additionally, existing rehabilitation systems may not fully engage patients in their exercises, leading to suboptimal outcomes. These limitations underscore the need for innovative solutions that can provide personalized, precise, and engaging vestibular rehabilitation experiences. This project addresses these challenges by proposing the development of a Vestibular Rehabilitation System that integrates immersive Virtual Reality (VR) technology. With this technology, the system aims to improve the precision and efficacy of rehabilitation exercises while enhancing patient engagement through immersive and personalized experiences.

PROCESS MODEL:




Fig 1. Process Model

CHAPTER 2

LITERATURE SURVEY

Song, J. J.[1] This paper provides an in-depth examination that VR Vestibular Rehabilitation Improves DHI Scores Significantly and also the potential Long-Term Benefits of VR.It provides us with the Side Effects of VR Decrease Over Time.

It examines that With population aging and interest in improving the quality of life, focus on dizziness in the field of ear nose and throat has in-creased. A local study found that 21.7% of adults had experi-enced dizziness in the past year and that dizziness is a major risk factor for falls. Diagnosis and treatment of dizziness would become more important with population aging in the future.

Vestibular rehabilitation is a treatment that, through exercise, promotes vestibular compensation and provides replacement of sensory input and habituation of the motion or situation that causes dizziness, which enhances postural and movement stabil-ity, reduces fall risk, and improves associated anxiety and depression. The recently developed customized vestibular rehabilitation therapy provides treatment by combining various exercis-es depending on the symptoms and conditions of the patient.

Virtual reality (VR) is an interface between humans and com-puters, which includes real-time simulation and interaction through various sensory channels, such as vision, hearing, touch, smell, and taste. The extent of the use of VR varies widely, and recent studies have applied VR technologies in the medical field, including anatomy, three-dimensional patient modeling, virtual surgery, and rehabilitation. Rehabilitation systems using VR technologies have been studied and developed recently. In com-bination with wearable computers, VR can have a wide variety of applications. Further, by applying content such as games to continuously motivate patients to undergo rehabilitation and by making the treatment enjoyable and engaging, the effectiveness can be increased. Currently, studies combining rehabilitation with VR are underway

Studies have been conducted on the use of VR to enhance the efficiency of vestibular rehabilitation. Previously, a balance reha-bilitation unit, force platform, and Nintendo Wii Fit were used, but recently, immersive equipment using the head-mount dis-play (HMD) has been developed.

Rosiak et al. [2] showed that motion trackers and force-plate platforms, which are referred to as hybrid VR units, were effective for subjective reduction of symptoms in patients with peripheral vestibular dysfunction.

Micarelli et al. [3] showed that the home-based VR protocol us-ing HMD can be a safe option to improve vestibular ocular re-flex, postural control, and quality of life in patients with vestibular impairment in whom cognitive decline could hinder rehabili-tation.

Furthermore, Park et al. [4] showed that the VR device combined with eye trackers and HMD can improve goal-orient-ed attention and activation of brain networks while providing vestibular rehabilitation. These VR devices employ games, unlike conventional equip-ment, which are enjoyable, increase concentration, and result in continuing treatment by the patient. Moreover, stimulating mul-tiple senses simultaneously could increase the effectiveness of rehabilitation and help treat patients with visual vertigo. Early models of VR devices could not be commercialized because of expensive equipment and large locations, but recently, training using relatively inexpensive and simple equipment, such as HMD, seemed effective, which is likely to be more practical in the future and provide great help to full-scale rehabilitation.

In Virtual and augmented reality in the vestibular rehabilitation of peripheral vestibular disorders: systematic review and meta-analysis by Austin Heffernan, Mohammed Abdelmalek & Desmond A. Nunez investigates that Vestibular rehabilitation therapy is an established treatment for patients with vestibular dysfunction.

Virtual reality (VR) and augmented reality (AR) can be utilised in vestibular rehabilitation. Evidence of the efficacy of VR and AR delivered rehabilitation in patients with peripheral vestibular disorders is reviewed. MEDLINE, EMBASE, CENTRAL, CINAHL, PsychInfo, PsychBITE, OTSeeker, Ei Compendex, IEE, Clinical trials.gov and WebofScience databases were searched. Reduction in vestibular dysfunction symptoms 0–3 months post-intervention was the primary outcome. Secondary outcomes included long-term symptom improvement and side effects. Risk of bias assessment and meta analyses were planned.

Five studies meeting eligibility criteria were included. Dizziness Handicap Inventory (DHI) scores 0–3 months post-intervention were reported by four studies. Meta-analysis identified a 1.13 (95% CI, – 1.74, – 0.52) standardized mean difference reduction in DHI in VR and AR treated patients compared to controls. Side effects reported by two studies were reduced by week four of VR intervention. Bias assessment identified DHI scores and side effects to be at high risk or of some concern.

Adjunct VR interventions reduced patient DHI significantly more than vestibular rehabilitation alone 0–3 months post-intervention in adult patients diagnosed with unilateral vestibular disease. High quality studies are needed.

It also examines that Dizziness is a common complaint affecting up to 23% of the population at any time in a first world setting¹. In publicly funded healthcare systems 0.8–1.7% of general practitioner attendances are for symptoms of dizziness or vertigo ^{2,3}, 9–13% of whom are referred to other specialists such as neurologists, cardiologists and otolaryngologists ^{3,4}.

Disorders of the vestibular system are identified in 50–65% of patients seen in specialist dizziness clinics ^{5,6}. The vestibular system consists of sensory

organs, cortical and subcortical structures that contribute to balance alongside proprioception and vision. The vestibular apparatus of the inner ear is the primary input for the vestibular system and relays information on head position and motion to the midbrain.

This sensory information leads to adjustments in body movements and posture to maintain balance⁷. The vestibular system can be affected by a variety of peripheral vestibular disorders (PVD), including benign paroxysmal positional vertigo (BPV), Menière's disease (MD), vestibular neuritis (VN) and post-traumatic vestibular dysfunction. Chronic dizziness symptoms can lead to symptoms of anxiety and depression^{8,9}. Dizziness and its sequelae can create a dizziness handicap for symptomatic patients and carries a \$64,929 lifetime burden for affected older adults¹⁰.

Vestibular rehabilitation is the main therapeutic option for many patients with dizziness. McDonnell and Hillier's¹¹ Cochrane review concluded that there was moderate to strong evidence of its effectiveness in individuals with Unilateral Vestibular Hypofunction (UVH).

Vestibular rehabilitation is an umbrella term that covers a range of exercise regimens from the generic such as Cooksey Cawthorne exercises to the customised¹¹. The American Physical Therapy Association Clinical Practice Guidelines indicate that vestibular exercises are effective when compared to no or placebo exercises and that customized exercises are more effective than generic¹².

Vestibular rehabilitation can be done at home, in the clinic or in a combination of these settings. However, home exercises require the patient to be motivated to participate, which is an area where the playful activities utilized in virtual reality (VR) and augmented reality (AR) may add benefit. Seventy three percent of patients suffering from vestibular disease report more enjoyment and motivation with VR and AR interventions than vestibular rehabilitation¹³.

Campbell et al.¹⁴ highlighted the importance of adherence to treatment in determining the benefit achieved through home based interventions aimed at

preventing falls in the visually impaired. Interactive video gaming systems linked to an aerobic training regimen increase adherence and physical fitness attainment more than an aerobic training regimen alone¹⁵. Therefore, the use of VR/AR could improve vestibular rehabilitation compliance and overall outcomes ^{16,17}.

Virtual reality and AR are two ways of delivering potential reality type vestibular rehabilitative interventions. Virtual reality is defined as the immersion of the user in an interactive environment that mimics reality¹⁸. However, much of the current literature considers non-immersive commercially available video gaming systems (non-IGS) (ex. Wii Fit) to be VR.

In an effort to focus on interventions that meet the true definition of virtual reality, non-IGS will be excluded from this systematic review. Immersive VR treats vestibular dysfunction by placing the subject in a simulated real world through two different strategies. One utilizes outpatient systems that use a head mounted display VR device and the second uses total body immersion inpatient systems such as the Immersive Rehabilitation Exercise System (Gesturetek Health)¹⁹.

In contrast to VR, AR augments the real-world environment instead of replacing it. Augmented reality adds to the subject's real world sensory input through computer-generated sound, text and graphics that are projected onto the user's natural visual and auditory fields²⁰. Augmented reality platforms have been used to treat vestibular disorders using AR eyewear²¹.

Virtual reality's utility in the treatment of vestibular dysfunction lies in the possibility of it achieving improved habituation, substitution, and adaptation through a more motivated vestibular rehabilitation ^{22,23}. Virtual reality has been studied as a rehabilitation intervention and has been found to elicit improvements in a variety of PVDs^{19,24}.

VR delivered vestibular rehabilitation has to the best of our knowledge not been assessed in current guidelines on the management of patients with vestibular disorders possibly due to its relative novelty. Furthermore, the previously published systematic review on the efficacy of VR in vestibular rehabilitation was

inconclusive²⁵.

We aim to synthesize the evidence to test the hypothesis that patients suffering from a PVD who have received VR or AR vestibular rehabilitation interventions experience the same improvement in dizziness as patients who received vestibular rehabilitation alone or a comparable control treatment.

The primary objective was to assess the benefits of VR and AR interventions for vestibular dysfunction symptoms (dizziness, vertigo, imbalance or other) in patients suffering from PVD within 3 months of treatment.

Secondary objectives include the long term (> 3 months) benefits, post-intervention change in patient quality of life, determining the adverse events (harms) associated with use of VR or AR interventions and determining the number of patients who failed to complete treatment.

Thus from this article we can infer that Vestibular rehabilitation therapy is an established treatment for patients with vestibular dysfunction. Virtual reality (VR) and augmented reality (AR) can be utilised in vestibular rehabilitation. Evidence of the efficacy of VR and AR delivered rehabilitation in patients with peripheral vestibular disorders is reviewed. MEDLINE, EMBASE, CENTRAL, CINAHL, PsychInfo, PsychBITE, OTSeeker, Ei Compendex, IEE, Clinical trials.gov and Web of Science databases were searched.

Reduction in vestibular dysfunction symptoms 0–3 months post-intervention was the primary outcome. Secondary outcomes included long-term symptom improvement and side effects. Risk of bias assessment and meta analyses were planned. Five studies meeting eligibility criteria were included.

Dizziness Handicap Inventory (DHI) scores 0–3 months post-intervention were reported by four studies. Meta-analysis identified a 1.13 (95% CI, – 1.74, – 0.52) standardized mean difference reduction in DHI in VR and AR treated patients

compared to controls. Side effects reported by two studies were reduced by week four of VR intervention.

Bias assessment identified DHI scores and side effects to be at high risk or of some concern. Adjunct VR interventions reduced patient DHI significantly more than vestibular rehabilitation alone 0–3 months post-intervention in adult patients diagnosed with unilateral vestibular disease. High quality studies are needed.

2.1.BIBLIOGRAPHY

[1]SongJ.J (2019) in “Virtual Reality for Vestibular Rehabilitation”.

[2]Heffernan, A.,Abdelmalek, M.& Nunez, D.A. (2021) in “Virtual and augmented reality in the vestibular rehabilitation of peripheral vestibular disorders”.

CHAPTER 3

IMPLEMENTATION AND SYSTEM DESIGN

3.1. IMPLEMENTATION:

3.1.1 VR ENVIRONMENT:

Incorporating a ball into vestibular rehabilitation exercises enhances the effectiveness and versatility of the therapy, particularly when augmented with Virtual Reality (VR) technology. The ball serves as a multifunctional tool, facilitating a range of dynamic movements that target different aspects of vestibular function.

Patients may engage in activities such as tossing and catching the ball while standing on a balance board or while their visual field is manipulated in the VR environment. These exercises challenge proprioception, coordination, and balance control, fostering adaptive responses within the vestibular system.

Furthermore, the use of a ball introduces an element of unpredictability and variability into the rehabilitation regimen, simulating real-world scenarios where individuals must respond to dynamic stimuli. This variability promotes neuroplasticity and adaptation, crucial components of vestibular rehabilitation that facilitate recovery and symptom management.

Additionally, the ball can be utilized for gaze stabilization exercises, where patients track the ball's motion with their eyes while keeping their head stationary or moving it in a controlled manner. These exercises aim to improve visual tracking and reduce symptoms of dizziness or vertigo.

When integrated with VR technology, the ball becomes an interactive tool within the virtual environment, offering patients engaging and immersive experiences.


For example, patients may engage in virtual games or activities where they

interact with virtual balls while receiving real-time feedback on their performance. The combination of VR and ball-based exercises not only enhances therapeutic outcomes but also increases patient motivation and compliance by making rehabilitation sessions enjoyable and rewarding.

In conclusion, the use of a ball in conjunction with VR technology amplifies the effectiveness and engagement of vestibular rehabilitation exercises. By leveraging the dynamic properties of the ball and the immersive capabilities of VR, healthcare providers can deliver personalized and impactful therapy that promotes recovery, enhances function, and improves overall quality of life for individuals with vestibular disorders.

3.1.2 GAZE STABILIZATION EXERCISE:

- Central Focal Point: At the heart of the virtual scene lies a prominent object.
- Spatial Transitions: The virtual environment transitions the focal object across different spatial directions.
- Directions : Horizontal (right and left), Vertical (up and down), and Diagonal trajectories. In each direction it would be done 5 times.





Gaze stabilization exercises are a crucial component of vestibular rehabilitation therapy, particularly for individuals experiencing vestibular impairments or balance disorders. These exercises aim to improve the ability to maintain a steady visual focus while the head is in motion, which is essential for tasks such as walking, driving, or even simply scanning one's environment without experiencing symptoms like dizziness or vertigo.



During gaze stabilization exercises, patients are typically instructed to fix their gaze on a specific target, such as a stationary object or a visual pattern, while performing controlled head movements. These movements may include turning the head from side to side, nodding up and down, or tilting the head at various angles. The goal is to gradually increase the speed and range of head movements while maintaining a clear and steady focus on the target.



By repeatedly exposing the vestibular system to controlled head movements while simultaneously requiring visual fixation, gaze stabilization exercises help to promote adaptation and compensation within the vestibular system. This process can lead to improvements in sensorimotor integration, reduced sensitivity to motion-related stimuli, and enhanced overall balance and stability.

Gaze stabilization exercises are often tailored to each individual's specific needs and may be prescribed as part of a comprehensive vestibular rehabilitation program overseen by a healthcare professional, such as a physical therapist or an otolaryngologist. Consistency and adherence to prescribed exercise regimens are key to maximizing the benefits of gaze stabilization exercises and achieving optimal outcomes in the management of vestibular disorders.



3.1.3 EXPERIMENTAL RESULTS:











INNOVATION PRACTICES LAB




INNOVATION PRACTICES LAB




INNOVATION PRACTICES LAB



INNOVATION PRACTICES LAB





3.2. DESIGN

The process behind the functioning of the system is depicted in the flow diagram This chapter describes the hardware and software requirements of the system including the libraries and environment used, dataset and pre-processing, and functional and non-functional requirements.

3.3. HARDWARE REQUIREMENTS:

The following are the hardware requirements for the proposed system:

- Oculus Quest VR Headset with hand controllers
- A Console
- Graphics Processing Unit

3.4. SOFTWARE REQUIREMENTS:

The following are the software requirements for the proposed system:

- Unity(version 2021)

Tech Stack:

Language: C#

Plugins : XR Plugin

3.5. FUNCTIONAL REQUIREMENTS:

Virtual Environment Creation:

The system should allow the creation of immersive virtual environments that simulate various scenarios relevant to vestibular rehabilitation exercises. These environments should include scenarios such as walking on uneven terrain, navigating obstacles, and experiencing changes in altitude or visual stimuli.

Real-Time Feedback Mechanism:

The system should provide real-time feedback to users based on their performance during rehabilitation exercises. Feedback should include information on exercise completion, accuracy, and adherence to prescribed protocols.

Customization of Exercises:

Exercises should be customizable based on individual patient needs, progress, and symptoms. The system should adapt exercise intensity, duration, and complexity in real-time according to user performance and feedback.

User Interaction:

The system should provide intuitive user interfaces for patients and healthcare professionals to interact with the VR environments and monitor progress.

It should allow users to navigate through different exercises, adjust settings, and view performance metrics.

Data Logging and Analysis:

The system should log and store exercise performance metrics for each user session.

It should allow for the analysis of collected data to track progress over time and adjust rehabilitation plans accordingly.

3.6. NON-FUNCTIONAL REQUIREMENTS:

Performance:

The system must have low latency and high responsiveness to ensure a seamless and immersive user experience. It should be able to handle complex VR environments and exercises without significant lag or delay.

Reliability:

The system should be reliable and robust, capable of handling extended use without frequent crashes or failures. It should have backup mechanisms to prevent data loss in case of unexpected shutdowns or errors.

Security:

Patient data should be stored securely and comply with relevant healthcare privacy regulations (e.g., HIPAA). The system should implement user authentication and access control measures to prevent unauthorized access to sensitive information.

Usability:

The user interface should be user-friendly and accessible to individuals with varying levels of technical proficiency. Instructions and guidance should be clear and easy to understand, facilitating smooth user interactions.

Scalability:

The system should be scalable to accommodate a growing number of users and support additional features or enhancements in the future. It should be adaptable to different hardware configurations and VR platforms.

Compatibility:

The system should be compatible with a range of VR headsets and accessories to maximize accessibility and usability. It should support integration with existing healthcare IT systems and protocols for seamless interoperability.

3.7. FEASIBILITY:

Technical Feasibility:

The technical feasibility of the Vestibular Rehabilitation System using Immersive Virtual Reality (VR) relies on available VR technology and software development tools. VR hardware such as headsets and controllers are widely accessible, and VR development platforms offer robust tools for creating immersive environments.

The system's technical feasibility also depends on the availability of skilled developers proficient in VR software development.

Financial Feasibility:

The financial feasibility of the project involves assessing the costs associated with hardware, software, development, and maintenance. VR hardware costs have decreased over time, making it more affordable for both healthcare facilities and individual users. However, software development costs, including VR environment creation and system design, may vary based on project complexity and developer expertise.

Operational Feasibility:

Operational feasibility assesses whether the proposed system aligns with the operational needs and workflows of healthcare facilities.

The system should be compatible with existing rehabilitation protocols and integrate seamlessly into clinical practice.

Training and support should be provided to healthcare professionals to ensure effective implementation and utilization of the system.

Legal and Regulatory Feasibility:

Legal and regulatory feasibility involves compliance with healthcare regulations, patient privacy laws, and industry standards. The system must adhere to data protection regulations (e.g., HIPAA) to ensure patient confidentiality and privacy. Compliance with medical device regulations may also be necessary, depending on the intended use and classification of the system.

CHAPTER 4

CONCLUSION

4.1 SUMMARY

In conclusion, the integration of immersive Virtual Reality (VR) technology into vestibular rehabilitation represents a groundbreaking approach to tackling the challenges posed by balance disorders and vestibular impairments globally. By harnessing the power of VR, this innovative system offers a multifaceted solution aimed at addressing the complex array of symptoms, including vertigo, dizziness, and an elevated risk of falls.


Through the simulation of diverse and challenging scenarios, the Vestibular Rehabilitation System Using Immersive Virtual Reality not only enhances the precision and effectiveness of rehabilitation exercises but also fosters greater engagement and motivation among patients. By immersing individuals in realistic virtual environments, the system provides a dynamic platform for tailored therapy interventions, catering to the unique needs and limitations of each patient.

Moreover, the immersive nature of VR therapy holds the potential to expedite recovery processes by facilitating neuroplasticity and sensory reintegration. As a result, this technology-driven approach not only promises to optimize therapy outcomes but also represents a significant step forward in improving the quality of life for those affected by vestibular impairments, offering hope and empowerment to individuals worldwide.

4.2 FUTURE DEVELOPMENTS

Brandt Daroff Exercise:

- Central Focal Point: At the heart of the virtual scene lies a prominent object.
- Spatial Transitions:
 - ◆ The object is moved to the right. (Instruction appears: “Lie down to your left”)
 - ◆ After 30 seconds, the object lies at the focal point.
 - ◆ The object is moved to the left. (Instruction appears: “Lie down to your right”)
 - ◆ After 30 seconds, the object lies at the focal point.




Benign paroxysmal positional vertigo (BPPV) is indeed characterized by brief episodes of vertigo, typically brought on by specific head movements such as tilting the head back, turning over in bed, or looking up. These movements cause the dislodged calcium crystals within the inner ear to stimulate the sensitive hair-like sensors in the semicircular canals, leading to a sensation of spinning or dizziness.

The Brandt-Daroff exercise is one of the methods used to alleviate the symptoms of BPPV. By performing specific movements that involve positioning the head and body in certain ways, the exercises aim to facilitate the relocation of the dislodged calcium crystals to a part of the inner ear where they do not trigger vertigo symptoms. This can help reduce the frequency and severity of vertigo episodes in individuals with BPPV.

It's worth noting that while the Brandt-Daroff exercise can be effective for many people with BPPV, it may not work for everyone, and individual responses can vary. Additionally, proper diagnosis and guidance from a healthcare professional are important to ensure that the exercises are appropriate and performed correctly.

Head Neck Exercise:

- Central Focal Point: At the heart of the virtual scene lies a prominent object.
- Spatial Transitions: The virtual environment transitions the focal object across different spatial directions.
- Directions : Horizontal (right and left), Vertical (up and down), Semi-circular.



Head and neck exercises can indeed be beneficial for improving flexibility, strength, and mobility in the neck area, as well as relieving symptoms such as neck pain, stiffness, and tension headaches. These exercises aim to target the muscles in the neck, shoulders, and upper back, helping to alleviate discomfort and improve overall function.

Regular practice of head and neck exercises can also help prevent future injuries by strengthening the muscles that support the neck and promoting better posture. Additionally, incorporating stretching and relaxation techniques into your routine can help reduce muscle tension and stress in the neck and shoulders, leading to increased comfort and well-being.

It's important to perform head and neck exercises mindfully, paying attention to proper form and avoiding any movements that exacerbate pain or discomfort. If you have any underlying medical conditions or concerns, it's advisable to consult with a healthcare professional or a physical therapist before starting a new exercise program to ensure it's safe and appropriate for your individual needs.

Object Manipulation Exercise:

- Object: A ball is employed as the central object for manipulation within the virtual environment.
- Visual Elements: A virtual wall is prominently displayed in front of the user, serving as the designated target surface for ball interaction.
- Automatic Ball Shooter: The VR system incorporates an automatic ball shooter mechanism, which autonomously propels the ball towards the virtual wall.
- Spatial Transition:
 - The ball is moved towards the wall. It hits the wall.
 - It bounces back to the shooter.
- Object manipulation exercises involve activities that require the coordinated use of hands, fingers, and sometimes other body parts to manipulate objects. These exercises are often used in rehabilitation, therapy, skill development, and entertainment. They can be beneficial for improving fine motor skills, hand-eye coordination, dexterity, concentration, and cognitive abilities.

4.3 EVALUATION METRICS:

Gaze Stabilization Exercise:

1. At what speed does the object move?
2. What is the size of the object?
3. What is the pause duration between movements?
4. What is the horizontal amplitude of the movement?
5. What is the vertical amplitude of the movement?
6. What is the diagonal amplitude of the movement?
7. How many times should it be repeated?

4.4 REFERENCES:

[1] Song, J. J. (2019). Virtual Reality for Vestibular Rehabilitation. Clin Exp Otorhinolaryngol. 2019;12 (4): 329-330. Publication Date (Web): 2019 October 02 (Editorial) doi: <https://doi.org/10.21053/ceo.2019.00983>

[2] Heffernan, A., Abdelmalek, M. & Nunez, D.A. Virtual and augmented reality in the vestibular rehabilitation of peripheral vestibular disorders: systematic review and meta-analysis. Sci Rep 11, 17843 (2021). <https://doi.org/10.1038/s41598-021-97370-9>

[3] Kim M, Kaneko F. Virtual reality-based gait rehabilitation intervention for stroke individuals: a scoping review. J Exerc Rehabil. 2023 Apr 27;19(2):95-104. doi: 10.12965/jer.2346114.057.

[4] Hazzaa NM, Manzour AF, Yahia E, Mohamed Galal E. Effectiveness of virtual reality-based programs as vestibular rehabilitation therapy in peripheral vestibular dysfunction: a meta-analysis. Eur Arch Otorhinolaryngol. 2023 Jul;280(7):3075-3086. doi: 10.1007/s00405-023-07911-3. Epub 2023 Mar 22

[5]Xie M, Zhou K, Patro N, Chan T, Levin M, Gupta MK, Archibald J. Virtual Reality for Vestibular Rehabilitation: A Systematic Review. Otol Neurotol. 2021 Aug 1;42(7):967-977. doi: 10.1097/MAO.0000000000003155.

[6]Chen X, Ai C, Yang Z, Ni Y, Yin X, You J, Li G. Mitigation of Thermal Instability for Electrical Properties in CaZrO₃-Modified (Na, K, Li) NbO₃ Lead-Free Piezoceramics. Materials (Basel). 2023 May 14;16(10):3720. doi: 10.3390/ma16103720.

[7]<https://www.mayoclinic.org/symptoms/dizziness/basics/definition/sym-20050886>

[8]<https://www.apta.org/apta-magazine/2021/11/01/vestibular--rehabilitation-physical-therapy>

[9]<https://m.youtube.com/watch?v=AyBz8QiE2ZY><https://m.youtube.com/watch?v=AyBz8QiE2ZY>

[10]<http://www.youtube.com/watch?v=gGYtahQjmWQ>

[11]<https://m.youtube.com/watch?v=NWBcOy950bc>

[12]<https://docs.unity3d.com/550/Documentation/Manual/VirtualReality.html>