

- Vectors in underline, matrices in capital. Assume all matrices have real entries unless stated otherwise.
- For a vector  $\underline{x}$ , we denote by  $x_i$  the  $i^{\text{th}}$  entry of  $\underline{x}$ .
- The inequality  $A \geq 0$  means that the matrix  $A$  is positive semi definite. The inequality  $\underline{x} \geq 0$  means that all the entries of the vector  $\underline{x}$  are non negative.

1. (5 pts) Given an (arbitrary, non empty) set  $S \subset \mathbb{R}^n$ , define the set of all vectors that make an acute angle with every point in  $S$

$$C = \{\underline{x} : \underline{x}^\top \underline{y} \geq 0 \quad \text{for every } \underline{y} \in S\}.$$

Identify whether  $C$  is

A. a subspace   B. an affine set   C. a convex set   D. a cone .

Do the answers to any of the above depend on the specific structure of  $S$  ?

2. (4 pts) Suppose you are given  $n \times n$  matrix  $A$  and a vector  $\underline{y} \in \mathbb{R}^n$ . Note that  $A$  may have both positive and negative eigen values. Is the function  $f_1(\underline{x}) = \|\underline{y} - A\underline{x}\|_2$  convex ? Answer the same for the function  $f_2(\underline{x}) = \|\underline{y} - A\underline{x}\|_2^2$ .

3. (4 pts) Given a finite set of points  $S = \{\underline{x}_1, \underline{x}_2, \dots, \underline{x}_n\}$ , consider the convex hull  $\text{conv}(S)$  of  $S$ .

(a) For  $S = \{(0, 0), (1, 1), (1, 0)\}$  sketch the convex hull  $\text{conv}(S)$ .

Given a **convex** function  $f : \mathbb{R}^n \mapsto \mathbb{R}$ , we are interested in maximising  $f(\underline{x})$  with the variable  $\underline{x}$  constrained to be in  $\text{conv}(S)$ .

(b) Prove that

$$f(\underline{x}) \leq \max \{f(\underline{x}_1), f(\underline{x}_2), \dots, f(\underline{x}_m)\}, \text{ for } \underline{x} \in \text{conv}(S).$$

This means that the maximum value of  $f$  over  $\text{conv}(S)$  occurs at one of the *vertices* of  $\text{conv}(S)$ .

4. (3 pts) You are given a vector  $\underline{b} \in \mathbb{R}^n$ . Consider the following sets consisting of the pair  $(A, z)$  for  $n \times n$  symmetric matrices  $A$  and scalars  $z$

$$C_1 = \left\{ (A, z) : A \in \mathbb{S}^n, z \in \mathbb{R}, \begin{pmatrix} A & \underline{b} \\ \underline{b}^\top & z \end{pmatrix} \geq 0 \right\}, \quad C_2 = \left\{ (A, z) : A \in \mathbb{S}_{++}^n, z \in \mathbb{R}, z \geq \underline{b}^\top A^{-1} \underline{b} \right\}.$$

Are  $C_1$  and  $C_2$  convex ? (Hint: Argue that  $C_1 = C_2$ )

5. (5 pts) (a) Suppose  $f : \mathbb{R}^2 \mapsto \mathbb{R}$  is a convex symmetric function, i.e.  $f$  is convex and  $f(x_1, x_2) = f(x_2, x_1)$  for all  $x_1, x_2$ . Consider the problem  $\min_{\underline{x} \in C} f(\underline{x})$  where  $C$  is a symmetric convex set. Argue why there exists an optimum  $\underline{x}$  with both the co-ordinates equal.

(b) Find the optimal value of the following problem

$$\begin{aligned} \max_{\underline{x}} \quad & x_1 x_2 x_3 \dots x_n \\ \text{s.t.} \quad & \sum x_i = 1, \quad x_i \geq 0 \text{ for } i = 1, 2, \dots, n. \end{aligned} \tag{1}$$

6. (6 pts) Show the following

- (a) The function  $f(\underline{x}) = \underline{x}^\top \underline{x} / t$ , with  $\text{dom } f = \{(\underline{x}, t) | t > 0\}$  is convex.  
 (b) The function  $f(\underline{x}) = \underline{x}^\top \underline{x} / t^2$ , with  $\text{dom } f = \{(\underline{x}, t) | t > 0\}$  is quasi-convex.  
 (c) The function  $f$  defined below is convex.

$$f(\underline{x}) = \begin{cases} \left\| \underline{x} - \frac{\underline{x}}{\|\underline{x}\|_2} \right\|_2 & \text{if } \|\underline{x}\|_2 \geq 1 \\ 0 & \text{otherwise.} \end{cases}$$