

# Computational study of the glycolytic degradation of poly(ethylene terephthalate) catalized by $N^1, N^2$ -bis(2-aminobenzyl)-1,2-diaminoethane zinc(II)

Pablo E. Alanis González,<sup>\*,†,¶</sup> Isabel del Carmen Saenz Tavera,<sup>‡,¶</sup> and Victor Manuel Rosas García<sup>‡,¶</sup>

<sup>†</sup>*Universidad Autónoma de Nuevo León, San Nicolás de los Garza, N.L.*

<sup>‡</sup>*Facultad de Ciencias Químicas, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, N.L.*

<sup>¶</sup>*Facultad de Ciencias Químicas, UANL*

E-mail: pabloalanis1998@gmail.com

Phone: +52 (81) 1177 3600

## Abstract

A possible reaction mechanism of the glycolytic degradation of poly-ethyleneterephthalate catalyzed with  $N^1, N^2$ -bis(2-aminobenzyl)-1,2-diaminoethane zinc(II) (ABEN) was determined using KS-DFT using the meta-NGA hybrid functional, MN15-L up to def2-SVP level of theory making use of an energy-weighted climbing image nudged elastic band (EW-CI-NEB) algorithm to determine the minimum energy path (MEP) and then optimizing the converged climbing image (CI) using eigenvector-following partitioned rational function optimization (EF P-RFO) to obtain the transition state (TS)

## Methods

The starting configurations for EG and ABEN were proposed and optimized using XTB<sup>1</sup> with the force field GFN2-xTB.<sup>2</sup> The starting configuration for DBHET was proposed using crystallographic data.<sup>3</sup>

The software ORCA (5.0.1)<sup>4</sup> was used to perform the geometrical optimization of the species involved in the depolymerization; PET dimer (DBHET), ABEN and EG in KS-DFT using a meta-NGA functional, MN15-L<sup>5</sup> with the def2-TZVP BS and employing the option to use automatic auxiliary basis set generation.<sup>6</sup>

The optimized structures were then merged onto the same Cartesian coordinates and then re-optimized in MN15-L/def2-SVP. Then, a multidimensional relaxed surface scan was performed keeping constrained, and varying the bond distances in 5 steps to obtain the products of this reaction. The optimized product and the starting geometry of the relaxed surface scan were the input for an Energy-Weighted Nudged Elastic Band (EW-NEB)<sup>7</sup> algorithm to find the path of minimum energy connecting both ends, followed by a P-RFO optimization to find a TS was performed onto the image with the highest energy alongside the MEP.

In order to determine interaction regions between the molecules, an IRI algorithm within multiwfn was utilized<sup>8</sup> with the wavefunctions generated in the previous step.

Using JANPA,<sup>9</sup> CLPOs and the bond orders for the reactants, products and transition states were obtained from the wavefunction generated by ORCA.

## References

- (1) Bannwarth, C.; Caldeweyher, E.; Ehlert, S.; Hansen, A.; Pracht, P.; Seibert, J.; Spicher, S.; Grimme, S. Extended TIGHT-BINDING Quantum Chemistry Methods. *WIREs Comput Mol Sci* **2021**, *11*.
- (2) Bannwarth, C.; Ehlert, S.; Grimme, S. GFN2-xTB—An Accurate and Broadly Parametrized Self-Consistent Tight-Binding Quantum Chemical Method with

Multipole Electrostatics and Density-Dependent Dispersion Contributions. *J. Chem. Theory Comput.* **2019**, *15*, 1652–1671.

- (3) Daubeney, R. D. P.; Bunn Charles William,; Brown, C. J. The Crystal Structure of Polyethylene Terephthalate. *Proc. R. Soc. Lond. A* **1954**, *226*, 531–542.
- (4) Neese, F.; Wennmohs, F.; Becker, U.; Riplinger, C. The ORCA Quantum Chemistry Program Package. *J. Chem. Phys.* **2020**, *152*, 224108.
- (5) Yu, H. S.; He, X.; Truhlar, D. G. MN15-L: A New Local Exchange-Correlation Functional for Kohn–Sham Density Functional Theory with Broad Accuracy for Atoms, Molecules, and Solids. *J. Chem. Theory Comput.* **2016**, *12*, 1280–1293.
- (6) Stoychev, G. L.; Auer, A. A.; Neese, F. Automatic Generation of Auxiliary Basis Sets. *J. Chem. Theory Comput.* **2017**, *13*, 554–562.
- (7) Ásgeirsson, V.; Birgisson, B. O.; Bjornsson, R.; Becker, U.; Neese, F.; Riplinger, C.; Jónsson, H. Nudged Elastic Band Method for Molecular Reactions Using Energy-Weighted Springs Combined with Eigenvector Following. *J. Chem. Theory Comput.* **2021**, *17*, 4929–4945.
- (8) Lu, T.; Chen, Q. Interaction Region Indicator: A Simple Real Space Function Clearly Revealing Both Chemical Bonds and Weak Interactions. *Chem. Methods* **2021**, *1*, 231–239.
- (9) Nikolaienko, T. Y.; Bulavin, L. A.; Hovorun, D. M. JANPA: An Open Source Cross-Platform Implementation of the Natural Population Analysis on the Java Platform. *Comput Theor Chem* **2014**, *1050*, 15–22.