Compact connected abelian groups with no small subgroups are tori

An informal account of Oliver Nash's Lean formalization

Jack McCarthy Claude Opus 5.5 (Anthropic)

19 October 2026

Abstract

We explain Oliver Nash's Lean proof that a topological group is isomorphic to a torus if and only if it is compact, connected and abelian and has no small subgroups.

1 Statement

A topological group G has *no small subgroups* if some neighbourhood U of 1 contains no subgroup other than $\{1\}$.

Theorem 1. *A topological group G is isomorphic, as a topological group, to $\mathbb{T}^m = (S^1)^m$ for some m if and only if G is compact, connected and abelian and has no small subgroups.*

The easy direction. Since $\mathbb{T}^m$ is clearly compact, connected and abelian, it suffices to show that it has no small subgroups. Regard $\mathbb{T}^m$ as a subset of $\mathbb{C}^m$, and let $U = \{z \in \mathbb{T}^m : \operatorname{Re} z_j > 0 \text{ for all } j\}$. Let $H \subseteq U$ be a subgroup and $z \in H$, and suppose some coordinate $w = z_j$ is not 1. Write $w = e^{i\theta}$ with $0 < |\theta| \leq \pi$, and let k be the smallest positive integer with $k|\theta| \geq \pi/2$. Then $(k-1)|\theta| < \pi/2$, so $\pi/2 \leq k|\theta| < \pi/2 + |\theta| \leq 3\pi/2$ and $\operatorname{Re} w^k = \cos k\theta \leq 0$, contradicting $z^k \in U$. Hence $H = \{1\}$, and $\mathbb{T}^m$ has no small subgroups.

A *character* of a topological group G is a continuous homomorphism $G \to S^1$. Let $\widehat{G}$ denote the group of characters. For the hard direction, we use the following lemmas.

Lemma 2 (Characters separate points)**.** *If G is a compact Hausdorff abelian group and $g \neq 1$, there is a character χ with $\chi(g) \neq 1$.*

Lemma 3. *The characters of $\mathbb{T}^m$ are exactly the maps $z \mapsto \prod_{i=1}^m z_i^{n_i}$ with $n_1, \ldots, n_m \in \mathbb{Z}$. In particular, the characters of S^1 are exactly the maps $z \mapsto z^n$ with $n \in \mathbb{Z}$.*

Lemma 4. *A topological group G with no small subgroups is Hausdorff.*

2 Proof of the hard direction

Let G be compact, connected and abelian with no small subgroups, and fix a neighbourhood of 1 containing no subgroup other than $\{1\}$. Its interior U is an open neighbourhood of 1 with the same property. By Lemma 4, G is Hausdorff. The set $K = G \setminus U$ is compact and misses 1. By Lemma 2, the open sets $\{\chi \neq 1\}$, $\chi \in \widehat{G}$, cover K, so finitely many $\chi_1, \ldots, \chi_k$ do.

If χ is a character with $\chi^n = 1$, then χ maps the connected space G into the n-th roots of unity, so $\chi = 1$. Thus $\widehat{G}$ is torsion-free, and the subgroup $M = \langle \chi_1, \dots, \chi_k \rangle$ is free abelian. Let $\psi_1, \dots, \psi_m$ be a basis of M, so that $L\colon \mathbb{Z}^m \to M$, $(n_1, \dots, n_m) \mapsto \prod_i \psi_i^{n_i}$, is an isomorphism, and let $\Psi = (\psi_1, \dots, \psi_m)\colon G \to \mathbb{T}^m$. We show that Ψ is an isomorphism.

Ψ is injective: By the choice of $\chi_1, \dots, \chi_k$,

$$K \subseteq \bigcup_{j=1}^{k} \{\chi_j \neq 1\} \implies \bigcap_{j=1}^{k} \ker \chi_j \subseteq G \setminus K = U \implies \bigcap_{j=1}^{k} \ker \chi_j = \{1\},$$

since U contains no nontrivial subgroup. Each χ_j is a product of powers of $\psi_1, \dots, \psi_m$, so $\chi_j(g) = 1$ whenever $\Psi(g) = 1$. Thus $\ker \Psi \subseteq \bigcap_j \ker \chi_j = \{1\}$.

Ψ is surjective: The image $H = \Psi(G)$ is compact, hence a closed subgroup of $\mathbb{T}^m$. Suppose, for the sake of contradiction, that $t \notin H$ for some $t \in \mathbb{T}^m$. Since H is closed, $\mathbb{T}^m/H$ is a compact Hausdorff abelian group, and $tH \neq 1$. By Lemma 2, there is a character ρ of $\mathbb{T}^m/H$ with $\rho(tH) \neq 1$, so $\tilde{\rho}(z) = \rho(zH)$ is a character of $\mathbb{T}^m$ with $\tilde{\rho}(H) = \{1\}$ and $\tilde{\rho}(t) \neq 1$. By Lemma 3, $\tilde{\rho}(z) = \prod_i z_i^{n_i}$ for some $n_i \in \mathbb{Z}$. Then

$$\begin{aligned} \tilde{\rho}(\Psi(G)) = \tilde{\rho}(H) = \{1\} &\implies \textstyle\prod_i \psi_i(g)^{n_i} = 1 \text{ for all } g \in G \\ &\implies \textstyle\prod_i \psi_i^{n_i} = 1 \text{ in } M \\ &\implies (n_1, \dots, n_m) = L^{-1}\textstyle(\prod_i \psi_i^{n_i}) = L^{-1}(1) = 0, \end{aligned}$$

so $\tilde{\rho}$ is trivial, contradicting $\tilde{\rho}(t) \neq 1$.

Thus Ψ is a continuous bijection from a compact space to a Hausdorff space, hence a homeomorphism, and $G \cong \mathbb{T}^m$. □

3 Proof of Lemma 2

3.1 From Peter–Weyl

For a continuous unitary representation $\pi\colon G \to \mathrm{U}(\mathcal{H})$ on a finite-dimensional complex Hilbert space $\mathcal{H}$ and vectors $v, w \in \mathcal{H}$, the *matrix coefficient* $\pi_{v,w}\colon G \to \mathbb{C}$ is $\pi_{v,w}(x) = \langle \pi(x)v, w \rangle$. We use the Peter–Weyl theorem in the following form (see e.g. Folland, *A Course in Abstract Harmonic Analysis*, §5.2).

Theorem 5 (Peter–Weyl)**.** *If G is a compact Hausdorff group, the linear span of the matrix coefficients of its finite-dimensional continuous unitary representations is dense in $C(G)$ for the uniform norm.*

Proof of Lemma 2. Let $g \neq 1$.

A representation with $\pi(g) \neq 1$: Since G is compact Hausdorff, Urysohn's lemma gives $f \in C(G)$ with $f(1) = 1$ and $f(g) = 0$. By Theorem 5 there are finite-dimensional continuous unitary representations $\pi_1, \dots, \pi_r$ and vectors v_j, w_j such that $q = \sum_{j=1}^{r} (\pi_j)_{v_j, w_j}$ satisfies $\|f - q\|_\infty < 1/2$. Since $|f(1) - f(g)| = 1$, this forces $q(1) \neq q(g)$. If $\pi_j(g) = 1$ for every j, then $(\pi_j)_{v_j,w_j}(g) = \langle v_j, w_j \rangle = (\pi_j)_{v_j,w_j}(1)$ for every j, so $q(g) = q(1)$. Hence $\pi_j(g) \neq 1$ for some j. Put $\pi = \pi_j\colon G \to \mathrm{U}(\mathcal{H})$.

A character with $\chi(g) \neq 1$: Since G is abelian, the operators $\pi(x)$, $x \in G$, are commuting unitary operators on $\mathcal{H}$, so they are simultaneously diagonalisable. That is, $\mathcal{H}$ has an orthonormal basis $e_1, \dots, e_d$ with $\pi(x)e_i = \chi_i(x)e_i$ for all x and i. Each χ_i is multiplicative because π is a homomorphism, takes values in S^1 because $\pi(x)$ is unitary, and is continuous because $\chi_i(x) = \langle \pi(x)e_i, e_i \rangle$. So each χ_i is a character, and since $\pi(g) \neq 1$, some $\chi_i(g) \neq 1$. □

Peter–Weyl rests on the spectral theorem for compact self-adjoint operators. For abelian G the convolution operators of §3.2 commute, so the formalization can use Gelfand theory for commutative C*-algebras instead.

3.2 The formalized proof, via Gelfand theory

Setup. Let G be a compact Hausdorff abelian group and μ the unique regular Haar measure on G with $\mu(G) = 1$. The formalization allows any finite measure that is left- and inversion-invariant, weakly regular and positive on nonempty open sets; the normalisation only simplifies constants. For $f \in C(G)$, $h \in L^2(G, \mu)$ and $a \in G$ put

$$(f * h)(x) = \int_G f(xy^{-1})\, h(y)\, d\mu(y), \qquad f^\sharp(y) = \overline{f(y^{-1})}, \qquad (\tau_a f)(y) = f(ya^{-1}),$$

and let C_f be the convolution operator $h \mapsto f*h$ on L^2. Let $B(L^2)$ denote the C*-algebra of bounded linear operators on $L^2(G, \mu)$, with the operator norm and adjoint. Each $f \in C(G)$ is uniformly continuous, in the sense that for each $\varepsilon > 0$ there is a neighbourhood V of 1 with $|f(u) - f(v)| < \varepsilon$ whenever $uv^{-1} \in V$.

Lemma 6. *Let $f, f' \in C(G)$ and $a, b \in G$.*

(i) $C_f \in B(L^2)$, *and* $F \mapsto C_F$ *is a bounded linear map* $C(G) \to B(L^2)$.

(ii) $f * f' \in C(G)$, *and* $C_f C_{f'} = C_{f*f'} = C_{f'} C_f$.

(iii) $C_f^* = C_{f^\sharp}$.

(iv) $C_f \neq 0$ *whenever* $f \neq 0$.

(v) $\tau_a f * \tau_b f' = \tau_{ab}(f * f')$.

Let A_0 be the unital subalgebra of $B(L^2)$ generated by the C_f. By Lemma 6(ii) the C_f commute, so A_0 is commutative, and by Lemma 6(iii) it is closed under adjoints. Let A be the closure of A_0 in the operator norm. Since multiplication and adjoint are continuous, A is again a commutative unital $*$-subalgebra. Being closed in the complete space $B(L^2)$, it is complete, and it inherits the C*-identity $\|X^*X\| = \|X\|^2$ from $B(L^2)$. So A is a commutative C*-algebra. Let Δ be the set of nonzero multiplicative linear functionals $\varphi\colon A \to \mathbb{C}$, with the weak-$*$ topology, in which it is a compact Hausdorff space. By the commutative Gelfand–Naimark theorem, the Gelfand transform $A \to C(\Delta)$, $X \mapsto (\varphi \mapsto \varphi(X))$, is isometric, hence injective. So for $X \neq Y$ in A there is $\varphi \in \Delta$ with $\varphi(X) \neq \varphi(Y)$.

Proof of Lemma 2 via Gelfand theory. Let $g \neq 1$. By Urysohn choose a real $f \in C(G)$ with $f(g) = 0$ and $f(1) = 1$, and let $T(a) = C_{\tau_a f} \in A$. Then:

- T is continuous: Given $\varepsilon > 0$, choose V for f by uniform continuity. For $a, b \in G$ we have $(ya^{-1})(yb^{-1})^{-1} = a^{-1}b$, as G is abelian, so $\|\tau_a f - \tau_b f\|_\infty = \sup_y |f(ya^{-1}) - f(yb^{-1})| \leq \varepsilon$ whenever $b \in aV$. Thus $a \mapsto \tau_a f$ is continuous from G to $C(G)$ with the sup norm, and composing with the bounded linear map $F \mapsto C_F$ of Lemma 6(i) shows that T is continuous.
- $T(a)T(b) = T(ab)T(1)$: By Lemma 6(ii) and (v), both sides equal $C_{\tau_{ab}(f*f)}$.
- $T(g) \neq T(1)$: We have $T(g) - T(1) = C_{\tau_g f - f}$ and $(\tau_g f - f)(g) = 1$, so this is nonzero by Lemma 6(iv).

Pick $\varphi \in \Delta$ with $\varphi(T(g)) \neq \varphi(T(1))$ and set $c(a) = \varphi(T(a))$, so that $c(a)c(b) = c(ab)c(1)$. If $c(1) = 0$, then $c(a)^2 = c(a^2)c(1) = 0$ for all a, contradicting $c(g) \neq c(1)$. So $\chi = c/c(1)$ is a homomorphism $G \to \mathbb{C}^\times$, and it is continuous because T and φ are. Since G is compact, χ is bounded, say $|\chi| \leq R$. For $a \in G$ and $n \in \mathbb{Z}$ this gives $|\chi(a)|^n = |\chi(a^n)| \leq R$, which forces $|\chi(a)| = 1$. So χ takes values in S^1, it is a character, and $\chi(g) \neq 1$. □

A Proof of Lemma 3

Proof. Each map $z \mapsto \prod_j z_j^{n_j}$ is a continuous homomorphism, hence a character. We prove the converse, first for $m = 1$.

Let ψ be a character of S^1. Since $t \mapsto e^{it}$ is a covering map $\mathbb{R} \to S^1$ and $\psi(1) = 1$, the path lifting lemma gives a continuous $F\colon \mathbb{R} \to \mathbb{R}$ with $F(0) = 0$ and $\psi(e^{it}) = e^{iF(t)}$ for all t. Because ψ is a homomorphism,

$$e^{iF(s+t)} = \psi(e^{is}e^{it}) = \psi(e^{is})\,\psi(e^{it}) = e^{i(F(s)+F(t))},$$

so $F(s+t) = F(s) + F(t) + 2\pi k(s,t)$ with $k(s,t) \in \mathbb{Z}$. By continuity k is constant, and setting $s = t = 0$ gives $k = 0$. That is, F is additive. A continuous additive map is linear, so $F(t) = \lambda t$. Taking $t = 2\pi$ gives $e^{2\pi i\lambda} = \psi(1) = 1$, so $\lambda = n \in \mathbb{Z}$. Hence $\psi(e^{it}) = (e^{it})^n$ for all t, that is, $\psi(z) = z^n$.

Now let ψ be a character of $\mathbb{T}^m$, and let $\iota_j\colon S^1 \to \mathbb{T}^m$ be the inclusion in the j-th coordinate. Each $\psi \circ \iota_j$ is a character of S^1, so $\psi(\iota_j(w)) = w^{n_j}$ for some $n_j \in \mathbb{Z}$. Since $z = \iota_1(z_1) \cdots \iota_m(z_m)$ and ψ is a homomorphism,

$$\psi(z) = \prod_{j=1}^{m} \psi(\iota_j(z_j)) = \prod_{j=1}^{m} z_j^{n_j}.$$ □

B Proof of Lemma 4

Proof. Let U be a neighbourhood of 1 containing no subgroup other than $\{1\}$. Let y lie in the closure of $\{1\}$, so that every neighbourhood of y contains 1. Since left translation by y is a homeomorphism, yU^{-1} is a neighbourhood of y, so $1 \in yU^{-1}$, that is, $y \in U$. Thus the closure of $\{1\}$, which is a subgroup, lies in U. Since U contains no subgroup other than $\{1\}$, the closure is trivial, and $\{1\}$ is closed. Now let $x \neq x'$ in G. Then xx'^{-1} lies in the open set $G \setminus \{1\}$, so by continuity of $(a,b) \mapsto ab^{-1}$ there are neighbourhoods V of x and W of x' with $ab^{-1} \neq 1$ for all $a \in V$ and $b \in W$. Hence $V \cap W = \emptyset$, so G is Hausdorff. □

C Proof of Lemma 6

We use the notation of §3.2, and write $\int$ for $\int_G$ and dy for $d\mu(y)$, and similarly for other variables. Let $f, f' \in C(G)$ and $h \in L^2(G,\mu)$.

Proof. (F1) $\underline{f * h \text{ is continuous:}}$ For $x, x' \in G$, Cauchy–Schwarz and $\mu(G) = 1$ give

$$|(f*h)(x) - (f*h)(x')| \leq \sup_y |f(xy^{-1}) - f(x'y^{-1})|\,\|h\|_2.$$

Given $\varepsilon > 0$, choose V for f by uniform continuity. Taking $u = xy^{-1}$ and $v = x'y^{-1}$ in that condition gives $uv^{-1} = xx'^{-1}$, independent of y. So whenever $xx'^{-1} \in V$, we get $\sup_y |f(xy^{-1}) - f(x'y^{-1})| \leq \varepsilon$, and hence $|(f*h)(x) - (f*h)(x')| \leq \varepsilon\|h\|_2$. Since $\{x' : xx'^{-1} \in V\} = V^{-1}x$ is a neighbourhood of x and ε is arbitrary, $f * h$ is continuous at x.

(F2) $\underline{\|f*h\|_2 \le \|f*h\|_\infty \le \|f\|_\infty \|h\|_2}$: For $x \in G$, Cauchy–Schwarz applied to $y \mapsto f(xy^{-1})$ and h, together with invariance of μ, gives $|(f*h)(x)| \le \|f\|_2 \|h\|_2$. Since $\mu(G) = 1$, we have $\|\cdot\|_2 \le \|\cdot\|_\infty$ for bounded measurable functions, and applying this to $f*h$ and to f gives

$$\|f*h\|_2 \le \|f*h\|_\infty \le \|f\|_2 \|h\|_2 \le \|f\|_\infty \|h\|_2.$$

In particular $F \mapsto C_F$ is a bounded linear map $C(G) \to B(L^2)$.

(F3) $\underline{C_f C_{f'} = C_{f*f'}}$: Take h continuous. Then the integrand below is continuous on the compact space $G \times G$, hence a uniform limit of finite sums of products $\alpha(y)\,\beta(z)$ with $\alpha, \beta \in C(G)$ by Stone–Weierstrass, so the two iterated integrals agree. Thus

$$\begin{aligned}
(f*(f'*h))(x) &= \int f(xy^{-1}) \int f'(yz^{-1})\, h(z)\, dz\, dy \\
&= \int \Big(\int f(xy^{-1})\, f'(yz^{-1})\, dy \Big) h(z)\, dz && \text{(Fubini)} \\
&= \int \Big(\int f(xz^{-1}u^{-1})\, f'(u)\, du \Big) h(z)\, dz && (y = uz) \\
&= \int (f*f')(xz^{-1})\, h(z)\, dz = ((f*f')*h)(x),
\end{aligned}$$

where the substitution $y = uz$ preserves μ by translation invariance. By (F2) both sides are bounded in h, and $C(G)$ is dense in $L^2(G,\mu)$ because μ is regular, so this holds for all $h \in L^2$.

(F4) $\underline{f*f' = f'*f}$: The map $y \mapsto xy^{-1}$ is an inversion followed by a translation, so it preserves μ, and since G is abelian, $x(xy^{-1})^{-1} = y$. Substituting,

$$(f*f')(x) = \int f(xy^{-1})\, f'(y)\, dy = \int f(y)\, f'(xy^{-1})\, dy = (f'*f)(x).$$

(F5) $\underline{C_f^* = C_{f^\sharp}}$: For continuous h and k, Fubini (justified as in (F3)) and $\overline{f(xy^{-1})} = f^\sharp(yx^{-1})$ give

$$\langle f*h, k\rangle = \iint f(xy^{-1})\, h(y)\, \overline{k(x)}\, dy\, dx = \int h(y) \overline{\int f^\sharp(yx^{-1})\, k(x)\, dx}\, dy = \langle h, f^\sharp * k\rangle.$$

By (F2) both sides are continuous in (h,k), so by density this holds for all $h, k \in L^2$.

(F6) $\underline{C_f \neq 0 \text{ whenever } f \neq 0}$: Since μ gives positive mass to every nonempty open set, a continuous function that vanishes almost everywhere vanishes everywhere. Suppose $C_f = 0$. Then $f * f^\sharp = C_f f^\sharp$ vanishes almost everywhere, and it is continuous by (F1), so it vanishes everywhere. In particular

$$0 = (f*f^\sharp)(1) = \int f(y^{-1})\, \overline{f(y^{-1})}\, dy = \int |f(y^{-1})|^2\, dy,$$

so the continuous nonnegative function $y \mapsto |f(y^{-1})|^2$ vanishes almost everywhere, hence everywhere, and $f = 0$.

(F7) $\underline{\tau_a f * \tau_b f' = \tau_{ab}(f*f')}$: Substituting $y = ub$ and using that G is abelian,

$$(\tau_a f * \tau_b f')(x) = \int f(xy^{-1}a^{-1})\, f'(yb^{-1})\, dy = \int f(x(ab)^{-1}u^{-1})\, f'(u)\, du = (f*f')(x(ab)^{-1}).$$

Together these give Lemma 6. Part (i) is (F2), part (ii) follows from (F1), (F3) and (F4), and parts (iii), (iv) and (v) are (F5), (F6) and (F7). □