

# Traits predict forest phenological responses to photoperiod more than temperature

Deirdre Loughnan<sup>1</sup>, Faith A M Jones<sup>1,2</sup>, and E M Wolkovich<sup>1,3,4</sup>

November 28, 2025

<sup>1</sup> Department of Forest and Conservation, Faculty of Forestry, University of British Columbia, 2424 Main Mall Vancouver, BC Canada V6T 1Z4.

<sup>2</sup> Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sciences, 901 83 Umeå, Sweden.

<sup>3</sup> Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

<sup>4</sup> Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA;

Corresponding Author: Deirdre Loughnan [deirdre.loughnan@ubc.ca](mailto:deirdre.loughnan@ubc.ca)

Running title: Traits drive photoperiod cues in budburst

## Summary

Climate change has advanced the timing of major life cycle events globally, with considerable variation across species and communities. In plants, differences in species growth strategies correlate with variation in leaf and structural traits, creating the basis of frameworks like the leaf economic spectrum. But whether traits related to the timing of growth fit within these frameworks is unclear. We combined trait data from six commonly measured traits with experimental data of woody plant budburst to identify shared relationships between traits, environmental cues—like temperature and photoperiod—and the timing of spring growth. Working across eight forest communities, we collected *in situ* trait measurements from 1428 individuals of 48 species. We paired trait data with budburst data from 4211 cuttings taken from the same individuals in a controlled environment study in which temperature and photoperiod cues varied. We found differences in traits between communities and across latitude, but only four of our six traits related to budburst and only in relation to photoperiod. Despite photoperiod being the weakest cue of woody species budburst, our results suggest that its association with other traits influences the timing of species growth, potentially impacting community composition and species interactions under future climates.

## Introduction

Climate change is causing species phenologies—the timing of life history events—to shift, with widespread advances being observed across the tree of life (Parmesan and Yohe, 2003; Hoegh-Guldberg et al., 2018).

This common phenological fingerprint, however, averages over high variability across species (Thackeray et al., 2016; Cohen et al., 2018; Kharouba et al., 2018), posing a challenge to accurate forecasts.

In plants, species variation can be explained, in part, by differences in growth strategies, which are generally inferred from traits (Violle et al., 2007). Decades of research on plant traits have worked to build predictive models of species responses to their environment (Green et al., 2022), which could explain species-level variability in phenological responses. Phenology, however, has generally been excluded from plant trait research due to its high inter- and intra-specific variability, making it difficult to leverage existing frameworks to explain phenological variation and predict future changes.

Previous studies have found high variation in phenology for the same species when observed over different years or sites (Primack et al., 2009; Chuine et al., 2010). But many studies of phenology are observational and conducted at the landscape scale (Fitter and Fitter, 2002; Menzel et al., 2006). In contrast, findings from controlled environment experiments suggest that phenological variation can be consistently decomposed into its cues (Chuine and Cour, 1999; Harrington and Gould, 2015; Flynn and Wolkovich, 2018). Given the importance of phenology in defining when a species grows and the environmental conditions during growth periods, we predict that these phenology-cue relationships may also correlate with other functional traits and vary along an axis of acquisitive to conservative growth strategies.

Correlations between plant traits, growth strategies, and responses to environments have been synthesized into several global frameworks, including the leaf economic spectrum (Wright et al., 2004) and wood economic spectrum (Chave et al., 2009). These frameworks have identified key traits that exhibit distinct gradients, ranging from acquisitive strategies—fast growing plants that produce cheaper tissue—to conservative strategies—with plants that invest in long-lived tissue but slower growth rates (Wright et al., 2004; Díaz et al., 2016). In temperate systems, changes in temperature and frost risk in spring can produce gradients in abiotic stress, including frost risk, soil nutrients, and light availability (Sakai and Larcher, 1987; Gotelli and Graves, 1996; Augspurger, 2009), in addition to differences in biotic interactions from herbivory or competition later in the season (Lopez et al., 2008; Wolkovich and Ettinger, 2014). Species that vary in their timing of leafout, should therefore exhibit traits and growth strategies that allow them to tolerate or avoid these abiotic and biotic factors. Leveraging insights from predictive models of phenology with these well established trait frameworks, we can begin to disentangle the environmental cues that shape phenology from those shaped by other trait differences in plant growth strategies.

To determine whether phenology fits within major functional trait frameworks will require a better understanding of the causes of variation within species. Phenological variation is generally observed in natural conditions where considerable differences in the environmental cues that trigger many phenological events—primarily temperature and photoperiod (Chuine, 2000; Körner and Basler, 2010)—vary across space and time. Within-species variation may also occur across other plant traits (e.g., leaf and wood structure traits), and across latitudinal (Wiemann and Bruce, 2002) or environmental gradients (Pollock et al., 2012), though generally to a smaller scale compared to phenology. While our previous growth chamber study found no evidence that phenological cues varied spatially (Loughnan and Wolkovich, in prep), previous work in functional traits has found some traits to exhibit greater spatial variation in response to environmental gradients (Gross et al., 2000; Wright et al., 2003). These results suggest that to better understand how phenology and other traits correlate across species will require methods that incorporate spatial variation within species.

Here, we tested whether phenological variation was aligned with existing trait frameworks using data on spring budburst paired and a suite of traits that capture acquisitive to conservative growth strategies. We decompose the high phenological variation in budburst date, using experiments to estimate three major phenological cues for woody plant budburst: chilling (cool winter temperatures), forcing (warm

spring temperatures), and photoperiod. We predict that early spring species will budburst before canopy closure—when chilling is lower, forcing temperatures are cooler, and photoperiods shorter—exhibited as smaller responses to temperature and photoperiod. These species should have traits associated with acquisitive growth, particularly shorter heights, smaller trunk or stem diameters, with lower investment in wood structure and leaf tissue, resulting in low wood specific density, diffuse-porous wood anatomy, and low leaf mass area, but high leaf nitrogen content for a greater photosynthetic potential. In contrast, we predict species with later budburst to predominately include canopy species that express more conservative growth strategies and require more chilling, warmer forcing, and longer photoperiods. These species should incur greater investments in long-lived tissue, with ring-porous wood anatomy, taller heights and greater diameter, denser wood and high leaf mass area, but low leaf nitrogen content. We used a powerful joint-modeling approach to estimate the relationships between these plant traits and phenological responses to cues, while partitioning the variance from species- and population-level differences.

## Materials and Methods

### Field sampling

We combined *in situ* trait data with budburst data from two growth chamber cutting experiments conducted across eastern and western temperate deciduous forests in North America. We collected both suites of data from populations that span a latitudinal gradient of 4-6° for the eastern and western communities respectively. We took trait measurements from across eight populations, of which there were four eastern populations—Harvard Forest, Massachusetts, USA (42.55°N, 72.20°W), White Mountains, New Hampshire, USA (44.11°N, 52.14°W), Second College Grant, New Hampshire, USA (44.79°N, 50.66°W), and St. Hippolyte, Quebec, Canada (45.98°N, 74.01°W), and four western population—E.C. Manning Park (49.06°N, 120.78°W), Sun Peaks (50.88°N, 119.89°W), Alex Fraser Research Forest (52.14°N, 122.14°W), and Smithers (54.78°N, 127.17°W), British Columbia (BC), Canada (Fig. 1). For the two growth chamber studies on budburst phenology, we collected cuttings from the most southern and northern populations in each transect ( $n_{pop}=4$ ).

### Functional traits

We measured all traits in the summer prior to each growth chamber study (eastern transect: 8-25 June 2015, western transect: 29 May to 30 July 2019), following full leafout but before budset. At each population and for each species, we measured a total of five traits from 1-10 healthy adult individuals: height, diameter of the main trunk or stem (hereafter referred to as diameter), wood specific density, leaf mass area, and the percent leaf nitrogen content. We also obtained xylem structure data from the WSL xylem database (Schweingruber and Landolt, 2010) for 72.3% of our species.

We measured traits in accordance to the methods discussed by Pérez-Harguindeguy et al. (2013). We calculated tree height using trigonometric methods and used a base height of 1.37 m to measure diameter (Magarik et al., 2020). For shrub heights, we measured the distance from the ground to the height of the top foliage and measured stem diameters at approximately 1 cm above ground-level. All stem and leaf samples were kept cool during transport and measurements of leaf area and stem volume taken within 3 and 12 hours of sample collection respectively. To measure wood specific density, we collected a 10 cm sample of branch wood, taken close to the base of the branch at the stem and calculated stem volume using the water displacement method. For our leaf traits, we haphazardly selected and sampled five, fully expanded, and hardened leaves, with no to minimal herbivore damage. We took a high resolution scan of each leaf using a flatbed scanner and estimated leaf area using the ImageJ software (version 2.0.0).

## Growth chamber study

For our growth chamber studies, we collected branch cuttings from our highest and lowest latitude populations in each transect, with sampling in our eastern study occurring from 20-28 January 2015 and sampling for our western study from 19-28 October 2019. In both studies, we included a total of eight distinct treatments consisting of two levels of chilling, forcing, and photoperiods (Fig. 1). We recorded budburst stages of each sample every 1-3 days for up to four months, defining the day of budburst as the day of budbreak or shoot elongation (denoted as code 07 by Finn et al. (2007)). For a more detailed discussion of study sample collection and methods see Flynn and Wolkovich (2018) for details on our eastern study and Loughnan and Wolkovich (in prep) for details on our western study.

## Statistical Analysis

Our analysis combined our *in situ* trait data with budburst data from the controlled environment. For each trait, we developed a joint Bayesian model, in which the relationship between traits and cues is used to estimate budburst. This statistical approach improves upon previous analyses of multiple traits, as it allows us to carry through uncertainty between trait and phenology data—and better partitions the drivers of variation in species phenologies

Our joint model consists of two parts. The first is a hierarchical linear model, which partitions the variation of individual observations ( $i$ ) of a given trait value ( $Y_{\text{trait}}$ ) to account for the effects of species ( $j$ ), population-level differences arising from transects, latitude, as well as the interaction between transects and latitude ( $\text{transect} \cdot \text{latitude}$ ), and finally, residual variation or ‘measurement error’ ( $\sigma_m^2$ ).

$$Y_{\text{trait}_{i,j}} \sim \text{Normal}(\mu_{i,j}, \sigma_m^2) \quad (1)$$

$$\mu_{i,j} = \alpha_{\text{grand trait}} + \alpha_{\text{trait}_j} + \beta_{\text{transect}} \times \text{transect} + \quad (2)$$

$$\beta_{\text{latitude}} \times \text{latitude} + \beta_{\text{transect} \cdot \text{latitude}} \times (\text{transect} \cdot \text{latitude}) \quad (3)$$

$$(4)$$

$$\alpha_{\text{trait}} \begin{bmatrix} \alpha_{\text{trait}_1} \\ \alpha_{\text{trait}_2} \\ \dots \\ \alpha_{\text{trait}_n} \end{bmatrix} \text{ such that } \alpha_{\text{trait}} \sim \text{Normal}(0, \sigma_{\text{trait}}^2) \quad (5)$$

$$(6)$$

We include transect as a dummy variable (0/1) and latitude as a continuous variable in our model. We modeled traits using natural units, with the exception of leaf mass area, which was rescaled by 100 for numeric stability in the model. Our model also includes partial pooling for species—which controls for variation in the number of trait estimates per species and trait variability—using these species-level estimates as predictors for each cue ( $\beta_{\text{chilling},j}$ ,  $\beta_{\text{forcing},j}$ ,  $\beta_{\text{photoperiod},j}$ ).

$$\beta_{\text{chilling}_j} = \alpha_{\text{chilling},j} + \beta_{\text{trait.chilling}} \times \alpha_{\text{trait},j} \quad (7)$$

$$\beta_{\text{forcing}_j} = \alpha_{\text{forcing},j} + \beta_{\text{trait.forcing}} \times \alpha_{\text{trait},j}$$

$$\beta_{\text{photoperiod}_j} = \alpha_{\text{photoperiod},j} + \beta_{\text{trait.photoperiod}} \times \alpha_{\text{trait},j}$$

164 In addition to the species-level estimates, the second part of our model estimates the overall effect of  
 165 each trait on each cue ( $\beta_{\text{trait.chilling}}$ ,  $\beta_{\text{trait.forcing}}$ ,  $\beta_{\text{trait.photoperiod}}$ ). From this we can estimate how well  
 166 traits explain species-level differences—by estimating the the species-level cue variation not explained  
 167 by traits ( $\alpha_{\text{chilling},j}$ ,  $\alpha_{\text{forcing},j}$ ,  $\alpha_{\text{photoperiod},j}$ ) and individual species responses to cues (*chilling*, *forcing*,  
 168 *photoperiod*, respectively). Finally, our model estimates the residual budburst variation across species  
 169 ( $Y_{\text{pheno},j}$ ), observations ( $\sigma_d^2$ ), as well as the variation in cues not attributed to the trait (using partial  
 170 pooling).

$$Y_{\text{pheno}_{i,j}} \sim \mathcal{N}(\mu_{i,j}, \sigma_d^2) \quad (8)$$

171 with

$$\mu_{i,j} = \alpha_{\text{pheno}_j} + \beta_{\text{chilling}_j} \cdot \textit{chilling} + \beta_{\text{forcing}_j} \cdot \textit{forcing} + \beta_{\text{photoperiod}_j} \cdot \textit{photoperiod} \quad (9)$$

172 where  $\alpha_{\text{pheno}_j}$ ,  $\alpha_{\text{chilling}_j}$ ,  $\alpha_{\text{forcing}_j}$ , and  $\alpha_{\text{photoperiod}_j}$  are elements of the normal random vectors:

$$\begin{aligned} \boldsymbol{\alpha}_{\text{pheno}} &= \{\alpha_{\text{pheno}_1}, \dots, \alpha_{\text{pheno}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{pheno}} \sim \mathcal{N}(\mu_{\text{pheno}}, \sigma_{\text{pheno}}^2) \\ \boldsymbol{\alpha}_{\text{chilling}} &= \{\alpha_{\text{chilling}_1}, \dots, \alpha_{\text{chilling}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{chilling}} \sim \mathcal{N}(\mu_{\text{chilling}}, \sigma_{\text{chilling}}^2) \\ \boldsymbol{\alpha}_{\text{forcing}} &= \{\alpha_{\text{forcing}_1}, \dots, \alpha_{\text{forcing}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{forcing}} \sim \mathcal{N}(\mu_{\text{forcing}}, \sigma_{\text{forcing}}^2) \\ \boldsymbol{\alpha}_{\text{photo}} &= \{\alpha_{\text{photo}_1}, \dots, \alpha_{\text{photo}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{photo}} \sim \mathcal{N}(\mu_{\text{photo}}, \sigma_{\text{photo}}^2) \end{aligned} \quad (10)$$

173 We modeled each trait individually, with the exception of ring-porosity, which we compared across  
 174 species using the posterior estimates of our wood stem density model, allowing us to account for inher-  
 175 ent differences in wood anatomy across species and growth form. We included all three cues (chilling,  
 176 forcing, and photoperiod) as continuous variables in our model, as well as all two-way interactions  
 177 between cues and between cues and sites. We converting chilling temperatures to total chill portions,  
 178 including both the chilling experienced in the field prior to sampling and during the experiment. For  
 179 this we used local weather station data and the chillR package (v. 0.73.1, Luedeling, 2020). To account  
 180 for differences in thermoperiodicity between the two studies (Buonaiuto et al., 2023), we also converted  
 181 forcing temperatures to mean daily temperatures for each treatment. Finally, we *z*-scored each cue  
 182 and site using two standard deviations to allow direct comparisons between results across parameters  
 183 (Gelman, 2008).

184  
 185 For each model we used trait specific priors that were weakly informative. We validated our choice  
 186 of priors using prior predictive checks and confirmed model stability under wider priors. All models  
 187 were coded in the Stan programming language for Bayesian models using the rstan package (Stan  
 188 Development Team, 2018) in R version 4.3.1 (R Development Core Team, 2017). All models met basic  
 189 diagnostic checks, including no divergences, high effective sample sizes ( $n_{\text{eff}}$ ) that exceeded 10% of  
 190 the number of iterations, and  $\hat{R}$  values close to 1. We report our model estimates as the mean values  
 191 with 90% uncertainty intervals (UI), interpreting parameter estimates with intervals that overlap to  
 192 be statistically similar to each other and those that include zero to have small effects.  
 193

## 194 Results

195 Across our eight populations, we measured 47 species of which 28 were in our eastern transect and  
 196 22 in our western transect. These include species dominant in both the understory and canopy layer,  
 197 with our eastern community consisting of 13 shrubs and 15 trees, our western community consisting of  
 198 18 shrubs and 4 trees, and three species that occurred in both transects. In total we measured traits

of 1428 unique individuals between the two transects across our five *in situ* traits: height ( $n = 1317$ ), diameter ( $n = 1220$ ), wood stem density ( $n = 1359$ ), leaf mass area ( $n = 1345$ ), leaf nitrogen content ( $n = 1351$ ). Across our two growth chamber studies, we made observations of 4211 samples, with our observations of budburst spanning 82 and 113 days for our eastern and western studies respectfully.

Most of our traits showed some variation by latitude within each transect, with a strong interactive effect between transect and latitude (Fig. 2). Leaf nitrogen content was the only trait to vary with latitude alone, with low latitude communities on both our eastern and western transects having greater values of leaf nitrogen content than communities at higher latitudes (-0.1, UI: -0.2, 0.0, Table S6). The strongest negative interaction was observed for height (-0.2, UI: -0.4, 0.0), while the strongest positive interaction was observed for leaf mass area (0.5, UI: 0.4, 0.6). Overall, woody species in our eastern community exhibited greater heights, diameter, and wood stem density but decreases in leaf mass area with increasing latitude. But in our western communities height and wood stem density decreased with latitude, while diameter and leaf mass area both increased (Fig. 2). In addition to the differences we found across populations, we also observed considerable differences between individual species, which varied considerably and up to 7 fold for some traits (Fig. 3).

We found that three of our four traits had a strong relationship with photoperiod, but not always in the direction we predicted. Taller species with larger trunk diameters and leaves with high nitrogen content had stronger responses with longer photoperiods (Fig. 3 c, i, o; Tables S2, S3, S6). But, contrary to our expectation, species with denser, high leaf mass area leaves had weaker photoperiod responses, allowing them to potentially budburst under shorter photoperiods (Fig. 3f).

Temperature cues ( $\beta_{\text{trait.chilling}}$  and  $\beta_{\text{trait.forcing}}$ ) exhibited no relationships with individual traits, but by accounting for the effects of leaf or wood traits, we found the importance of our three cues the vary by trait. Our models of height (-13.4, UI: -17.2, -9.9), diameter (-12.5, UI: -16.2, -8.6), wood stem density (-20.9, UI: -33.2, -9.8), and leaf nitrogen content (-35.1, UI: -68.1, -4.1) all estimated chilling as the strongest cue of budburst ( $\beta_{\text{chilling}}$ ), with more chilling advancing budburst. Our model of leaf mass area, however, estimated photoperiod as the strongest cue ( $\beta_{\text{photoperiod}}$ , -14.0, UI: -23.1, -3.5). After accounting for the effects of traits, only our height and diameter model found all three environmental cues to drive budburst timing (Tables S2, S3). Our models of wood stem density and leaf nitrogen content in turn found temperature cues alone to shape budburst (Tables S4, S6), while our model of leaf mass area found a large response to only photoperiod (Table S5).

In synthesizing the effects of multiple traits across species, our results can be used to make generalizations across ecologically important groups of species. But only some of our models estimated clear gradients in species timing between trees and shrubs. In particular, we found height to have large correlations between budburst timing and trait values, with earlier estimates of budburst for shrubs (with a mean day of budburst of 10)—especially under greater cues—and later budburst estimates for trees (with a mean day of budburst of 17.3, Fig. S1). Diameter at breast height showed similar trends as estimates from our height model (results not shown). But this was not the case for our two leaf traits. Leaf nitrogen content, for example, showed no distinct separation between shrub and tree functional groups (Fig. S1).

## Discussion

Using our joint modeling approach, we estimated how leaf and wood traits interact with temperature and photoperiod cues to shape species budburst. We found that photoperiod—often the weakest cue of budburst (Laube et al., 2014; Zohner et al., 2016; Flynn and Wolkovich, 2018)—was the most important cue in trait-phenology relationships. In general, we also found trait patterns varied between

our eastern and western transects and with latitude. These spatial differences in trait variation may be due to differences in the community assemblages, as our western community is more shrub dominated, with shorter plants with less dense branch wood. This more acquisitive growth strategy suggests these species are more likely to utilize resources early in the season prior to canopy closure. Collectively our results provide new insights into the complexity of the mechanisms that underlie the relationships between traits and environmental cues, while also challenging our existing understanding of these processes.

## Cues and functional traits

We found only partial support for our prediction that species with acquisitive traits—particularly small trees with low wood density, low leaf mass area, and high leaf nitrogen content—would have early budburst via small temperature and photoperiod responses. We did find species with smaller heights and diameters to have smaller photoperiod responses (associated with earlier budburst). But contrary to our prediction, species with less dense leaves showed larger responses to photoperiod, while leaves with high nitrogen content had stronger photoperiod responses. We also did not find any relationships between any of our focal traits and temperature. These unexpected trends in our results may be due to selection on other physiological processes, as many of our traits are associated with one or more ecological function (Wright et al., 2004; Pérez-Harguindeguy et al., 2013; Reich, 2014). In particular, leaf mass area is known to correlate with other traits like leaf lifespan or decomposition rates in addition to light capture (De La Riva et al., 2016). So while our results highlight the ways in which phenology partially aligns with gradients found in established trait frameworks, they offer new insight into potential tradeoffs in how varying physiological processes shape species growth strategies.

Decades of previous research that only considered phenology found budburst timing to be primarily driven by temperature and weakly by photoperiod (Chuine et al., 2010; Basler and Körner, 2014; Laube et al., 2014). We, however, found no traits that correlate with responses to temperature, suggesting other cues are impacting leaf and structural traits in temperate forests. Leaf mass area also varies with soil moisture, with variation in leaf area allowing plants to reduce evaporation under dry conditions, and thus selecting for high values of leaf mass area (De La Riva et al., 2016). Soil moisture also shapes other phenological events in woody plants, including radial growth phenology and shoot elongation (Cabon et al., 2020; Peters et al., 2021). If selection by soil moisture is shaping phenological responses, it may be contributing to the unexpected trends we observed in leaf traits and the absence of relationships with temperature. Thus, fully understanding how species growth strategies correlate with phenology may require the consideration of cues known to shape plant traits more broadly.

The absence of trait-cue relationships between budburst and wood structure and wood stem density also contrasts the findings of previous work linking these traits. Previous studies have found some evidence that trees with diffuse-porous wood structure leafout earlier than species with ring-porous structures (Lechowicz, 1984; Panchen et al., 2014; Yin et al., 2016; Osada, 2017; Savage et al., 2022). But using wood density as a proxy for wood structure, with wood density positively correlating with xylem resistance to embolism (Hacke et al., 2001), we did not find clear association between our three cues and xylem structure. But the positive relationship between wood density and latitude in our eastern community could be caused by the differences in winter conditions experienced by canopy versus understory species. The canopy tree species that dominate our eastern communities may experience greater horizontal stress from wind and downward pressure from snow, explaining the greater wood densities they exhibit at higher latitudes (MacFarlane and Kane, 2017; MacFarlane, 2020), while species in the shrub dominated western communities experience greater protection from being in the understory.

In comparing our results with a global meta-analysis of tree trait relationships with budburst cues

(Loughnan et al., 2025), we found similar trait-cue relationships for height and leaf mass area. At both the global and continental scales, we found taller tree heights to leafout with longer photoperiods. We also found species with high specific leaf area—which is the inverse of leaf mass area and thus equivalent to low values—exhibited large responses to photoperiod (Loughnan et al., 2025). The consistency of these results, despite the differences in the two spatial scales of these datasets, provides further evidence that alternate underlying mechanisms are shaping how woody species respond to photoperiod cues.

## Functional traits predict climate change responses

Our results offer novel insights into how broader correlations between growth strategies and phenological cues can help predict responses in woody plant communities with climate change. As temperatures rise, particularly at higher latitudes (Hoegh-Guldberg et al., 2018), warmer winter and spring temperatures may select for earlier budburst in some species. But since photoperiod will remain fixed, our observed relationships between photoperiod and other traits has the potential to limit species abilities to track temperatures and constrain the extent to which some species phenologies will advance with climate change. Our results suggest that these effects will likely be greater for taller species or canopy trees and species with relatively low leaf mass area. These constraints could have cascading effects on forest communities, as variable species responses to increasing temperatures further alter species growth strategies and their interactions with competitors or herbivores within their communities.

Our findings of correlations between phenology and other commonly measured traits highlight how accurate forecasts of future changes in phenology can benefit from accounting for the response of other traits to climate change. Across temperature and precipitation gradients, leaf size and shape also change, as species shift to conserve water and mitigate effects of transpiration under higher temperatures (De La Riva et al., 2016). These changes could impact species photosynthetic potential and ultimately ecosystem services, such as carbon sequestration. While phenological research has focused on forecasting responses to temperature, the correlation of other traits with photoperiod suggests it is also an important cue. It is therefore advantageous for additional traits that are likely to respond to photoperiod to be further studied in relation to phenology when forecasting the future impacts of climate change on communities.

## References

- Augspurger, C. K. 2009. Spring 2007 warmth and frost: phenology, damage and refoliation in a temperate deciduous forest. *Functional Ecology* 23:1031–1039.
- Basler, D., and C. Körner. 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* 34:377–388.
- Buonaiuto, D. M., E. M. Wolkovich, and M. J. Donahue. 2023. Experimental designs for testing the interactive effects of temperature and light in ecology : The problem of periodicity. *Functional Ecology* 37:1747–1756.
- Cabon, A., L. Fernández-de-Uña, G. Gea-Izquierdo, F. C. Meinzer, D. R. Woodruff, J. Martínez-Vilalta, and M. De Cáceres. 2020. Water potential control of turgor-driven tracheid enlargement in Scots pine at its xeric distribution edge. *New Phytologist* 225:209–221.
- Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a worldwide wood economics spectrum. *Ecology Letters* 12:351–366.



342 Chuine, I. 2000. A unified model for budburst of trees. *Journal of Theoretical Biology* 207:337–347.

343 Chuine, I., and P. Cour. 1999. Climatic determinants of budburst seasonality in four temperate-zone  
344 tree species. *New Phytologist* 143:339–349.

345 Chuine, I., X. Morin, and H. Bugmann. 2010. Warming, photoperiods, and tree phenology. *Science*  
346 329:277–278.

347 Cohen, J. M., M. J. Lajeunesse, and J. R. Rohr. 2018. A global synthesis of animal phenological  
348 responses to climate change. *Nature Climate Change* 8:224–228.

349 De La Riva, E. G., M. Olmo, H. Poorter, J. L. Uberta, and R. Villar. 2016. Leaf Mass per Area (LMA)  
350 and Its Relationship with Leaf Structure and Anatomy in 34 Mediterranean Woody Species along a  
351 Water Availability Gradient. *PLOS ONE* 11:e0148788.

352 Díaz, S., J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer, C. Wirth,  
353 I. Colin Prentice, E. Garnier, G. Bönsch, M. Westoby, H. Poorter, P. B. Reich, A. T. Moles, J. Dickie,  
354 A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright, S. N. Sheremet'ev, H. Jactel, C. Baraloto,  
355 B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig, A. Günther, V. Falczuk,  
356 N. Rüger, M. D. Mahecha, and L. D. Gorné. 2016. The global spectrum of plant form and function.  
357 *Nature* 529:167–171.

358 Finn, G. A., A. E. Straszewski, and V. Peterson. 2007. A general growth stage key for describing trees  
359 and woody plants. *Annals of Applied Biology* 151:127–131.

360 Fitter, A. H., and R. S. Fitter. 2002. Rapid changes in flowering time in British plants. *Science*  
361 296:1689–1691.

362 Flynn, D. F. B., and E. M. Wolkovich. 2018. Temperature and photoperiod drive spring phenology  
363 across all species in a temperate forest community. *New Phytologist* 219:1353–1362.

364 Gelman, A. 2008. Scaling regression inputs by dividing by two standard deviations. *Statistics in*  
365 *Medicine* 27:2865–2873.

366 Gotelli, N. J., and G. R. Graves. 1996. The temporal niche. Pages 95–112 *in* *Null Models In Ecology*.  
367 Smithsonian Institution Press, Washington, D. C.

368 Green, S. J., C. B. Brookson, N. A. Hardy, and L. B. Crowder. 2022. Trait-based approaches to  
369 global change ecology: moving from description to prediction. *Proceedings of the Royal Society B:*  
370 *Biological Sciences* 289:1–10.

371 Gross, K. L., M. R. Willig, L. Gough, R. Inouye, and S. B. Cox. 2000. Patterns of species density and  
372 productivity at different spatial scales in herbaceous plant communities. *Oikos* 89:417–427.

373 Hacke, U. G., J. S. Sperry, W. T. Pockman, S. D. Davis, and K. A. McCulloh. 2001. Trends in wood  
374 density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia*  
375 126:457–461.

376 Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying dormancy  
377 requirements for Pacific Northwest tree species. *Frontiers in Plant Science* 6:1–12.

378 Hoegh-Guldberg, O., D. Jacob, M. Taylor, M. Bindi, S. Brown, I. Camilloni, A. Diedhiou, R. Djalante,  
379 K. Ebi, F. Engelbrecht, J. Guiot, Y. Hijioka, S. Mehrotra, A. Payne, S. Seneviratne, A. Thomas,  
380 R. Warren, and G. Zhou. 2018. Impacts of 1.5 °C Global Warming on Natural and Human Systems.  
381 In: *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C*  
382 *above pre-industrial levels and related global greenhouse gas emission pathways, in the context of* .  
383 *Tech. rep.*, Cambridge University Press, Cambridge, UK and New York, NY, USA.

384 Kharoubá, H. M., J. Ehrlén, A. Gelman, K. Bolmgren, J. M. Allen, S. E. Travers, and E. M. Wolkovich.  
385 2018. Global shifts in the phenological synchrony of species interactions over recent decades. *Pro-*  
386 *ceedings of the National Academy of Sciences* 115:5211–5216.

387 Körner, C., and D. Basler. 2010. Phenology Under Global Warming. *Science* 327:1461–1463.

388 Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling outweighs  
389 photoperiod in preventing precocious spring development. *Global Change Biology* 20:170–182.

390 Lechowicz, M. J. 1984. Why Do Temperate Deciduous Trees Leaf Out at Different Times? *Adaptation*  
391 *and Ecology of Forest Communities*. *The American Naturalist* 124:821–842.

392 Lopez, O. R., K. Farris-Lopez, R. A. Montgomery, and T. J. Givnish. 2008. Leaf phenology in relation  
393 to canopy closure in southern Appalachian trees. *American Journal of Botany* 95:1395–1407.

394 Loughnan, D., F. A. Jones, G. Legault, C. J. Chamberlain, D. M. Buonaiuto, A. K. Ettinger, M. Gar-  
395 ner, D. S. Sodhi, and E. M. Wolkovich. 2025. Budburst timing within a functional trait framework.  
396 *Journal of Ecology* 00:1–12.

397 Loughnan, D., and E. M. Wolkovich. in prep. Temporal assembly of woody plant communities shaped  
398 equally by evolutionary history as by current environments .

399 Luedeling, E. 2020. chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees.  
400 <https://CRAN.R-project.org/package=chillR>.

401 MacFarlane, D. W. 2020. Functional Relationships Between Branch and Stem Wood Density for  
402 Temperate Tree Species in North America. *Frontiers in Forests and Global Change* 3.

403 MacFarlane, D. W., and B. Kane. 2017. Neighbour effects on tree architecture: functional trade-offs  
404 balancing crown competitiveness with wind resistance. *Functional Ecology* 31:1624–1636.

405 Magarik, Y. A., L. A. Roman, and J. G. Henning. 2020. How should we measure the dbh of multi-  
406 stemmed urban trees? *Urban Forestry & Urban Greening* 47:1–11.

407 Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aaasa, R. Ahas, K. Alm-Kübler, P. Bissolli,  
408 O. Braslavská, A. Briede, F. M. Chmielewski, Z. Crepinsek, Y. Curnel, Å. Dahl, C. Defila, A. Don-  
409 nelly, Y. Filella, K. Jatzak, F. Måge, A. Mestre, Ø. Nordli, J. Peñuelas, P. Pirinen, V. Remišová,  
410 H. Scheffinger, M. Striz, A. Susnik, A. J. Van Vliet, F. E. Wielgolaski, S. Zach, and A. Zust. 2006.  
411 European phenological response to climate change matches the warming pattern. *Global Change*  
412 *Biology* 12:1969–1976.

413 Osada, N. 2017. Relationships between the timing of budburst, plant traits, and distribution of 24  
414 coexisting woody species in a warm-temperate forest in Japan. *American Journal of Botany* 104:550–  
415 558.

416 Panchen, Z. A., R. B. Primack, B. Nordt, E. R. Ellwood, A. Stevens, S. S. Renner, C. G. Willis,  
417 R. Fahey, A. Whitemore, Y. Du, and C. C. Davis. 2014. Leaf out times of temperate woody  
418 plants are related to phylogeny, deciduousness, growth habit and wood anatomy. *New Phytologist*  
419 203:1208–1219.

420 Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across  
421 natural systems. *Nature* 421:37–42.

422 Pérez-Harguindeguy, N., S. Díaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S. Bret-  
423 Harte, W. K. Cornwell, J. M. Craine, D. E. Gurvich, C. Urcelay, E. J. Veneklaas, P. B. Reich,  
424 L. Poorter, I. J. Wright, P. Ray, L. Enrico, J. G. Pausas, A. C. de Vos, N. Buchmann, G. Funes,  
425 F. Quétier, J. G. Hodgson, K. Thompson, H. D. Morgan, H. ter Steege, M. G. A. van der Heijden,

426 L. Sack, B. Blonder, P. Poschlod, M. V. Vaieretti, G. Conti, A. C. Staver, S. Aquino, and J. H. C.  
 427 Cornelissen. 2013. New handbook for standardized measurement of plant functional traits worldwide.  
 428 Australian Journal of Botany 61:167–234.

429 Peters, R. L., K. Steppe, H. E. Cuny, D. J. De Pauw, D. C. Frank, M. Schaub, C. B. Rathgeber,  
 430 A. Cabon, and P. Fonti. 2021. Turgor – a limiting factor for radial growth in mature conifers along  
 431 an elevational gradient. New Phytologist 229:213–229.

432 Pollock, L. J., W. K. Morris, and P. A. Vesk. 2012. The role of functional traits in species distributions  
 433 revealed through a hierarchical model. Ecography 35:716–725.

434 Primack, R. B., I. Ibáñez, H. Higuchi, S. D. Lee, A. J. Miller-Rushing, A. M. Wilson, and J. A. Silander.  
 435 2009. Spatial and interspecific variability in phenological responses to warming temperatures.  
 436 Biological Conservation 142:2569–2577.

437 R Development Core Team. 2017. R: A language and environment for statistical computing.

438 Reich, P. B. 2014. The world-wide ‘fast–slow’ plant economics spectrum: a traits manifesto. Journal  
 439 of Ecology 102:275–301.

440 Sakai, A., and W. Larcher. 1987. Frost Survival of Plants: Responses and adaptation to freezing stress.  
 441 Springer-Verlag, Berlin, Heidelberg.

442 Savage, J. A., T. Kiecker, N. McMann, D. Park, M. Rothendler, and K. Mosher. 2022. Leaf out time  
 443 correlates with wood anatomy across large geographic scales and within local communities. New  
 444 Phytologist 235:953–964.

445 Schweingruber, F., and W. Landolt. 2010. The xylem database.

446 Stan Development Team. 2018. RStan: the R interface to Stan. R package version 2.17.3.

447 Thackeray, S. J., P. A. Henrys, D. Hemming, J. R. Bell, M. S. Botham, S. Burthe, P. Helaouet,  
 448 D. G. Johns, I. D. Jones, D. I. Leech, E. B. MacKay, D. Massimino, S. Atkinson, P. J. Bacon,  
 449 T. M. Brereton, L. Carvalho, T. H. Clutton-Brock, C. Duck, M. Edwards, J. M. Elliott, S. J. Hall,  
 450 R. Harrington, J. W. Pearce-Higgins, T. T. Høye, L. E. Kruuk, J. M. Pemberton, T. H. Sparks,  
 451 P. M. Thompson, I. White, I. J. Winfield, and S. Wanless. 2016. Phenological sensitivity to climate  
 452 across taxa and trophic levels. Nature 535:241–245.

453 Violle, C., M. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let the  
 454 concept of trait be functional! Oikos 116:882–892.

455 Wiemann, M. C., and W. G. Bruce. 2002. Geographic variation in wood specific gravity: effects of  
 456 latitude, temperature, and precipitation. Wood and Fiber Science 34:96–107.

457 Wolkovich, E. M., and A. K. Ettinger. 2014. Back to the future for plant phenology research. New  
 458 Phytologist 203:1021–1024.

459 Wright, I. J., M. Westoby, P. B. Reich, J. Oleksyn, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-  
 460 Bares, T. Chapin, J. H. C. Cornelissen, M. Diemer, J. Flexas, J. Gulias, E. Garnier, M. L. Navas,  
 461 C. Roumet, P. K. Groom, B. B. Lamont, K. Hikosaka, T. Lee, W. Lee, C. Lusk, J. J. Midgley,  
 462 Ü. Niinemets, H. Osada, H. Poorter, P. Pool, E. J. Veneklaas, L. Prior, V. I. Pyankov, S. C.  
 463 Thomas, M. G. Tjoelker, and R. Villar. 2004. The worldwide leaf economics spectrum. Nature  
 464 428:821–827.

465 Wright, J. P., A. S. Flecker, and C. G. Jones. 2003. Local vs. landscape controls on plant species  
 466 richness in Beaver meadows. Ecology 84:3162–3173.

- 467 Yin, J., J. D. Fridley, M. S. Smith, and T. L. Bauerle. 2016. Xylem vessel traits predict the leaf  
468 phenology of native and non-native understorey species of temperate deciduous forests. *Functional*  
469 *Ecology* 30:206–214.
- 470 Zohner, C. M., B. M. Benito, J.-C. Svenning, and S. S. Renner. 2016. Day length unlikely to constrain  
471 climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate Change* 6:1120–  
472 1123.




Figure 1: We collected traits data and branch cuttings from eight sites, across two transects in eastern and western North America. Cuttings were used in two controlled environment studies in which we varied chilling, forcing, and photoperiod treatments and recorded the day of budburst of each individual. Pairing our *in situ* trait data with the results of these experiments, we tested whether earlier budbursting species exhibited traits associated with more acquisitive growth strategies and smaller responses to cues and later budbursting species a more conservative growth strategy and larger responses to cues.




Figure 2: We found a. height, b. diameter, c. branch stem density, and d. leaf mass area to all experience a strong interaction between latitude and transect, e. while leaf nitrogen content showed a strong effect of latitude alone.




Figure 3: Relationships between species traits and cue responses showed considerable variation across a-c. height, d-f. leaf mass area, g-i. diameter, j-l. wood specific density, and m-o. the leaf nitrogen content. Point colours representing different species groups, with tree species shown in red and shrub species in blue. Crosses depict the 50% uncertainty interval of the model estimates of species trait values and estimated responses to cues. The grey band depicts the 90% uncertainty interval and the black line the mean response.




Figure 4: Despite previous studies finding relationships between leaf out timing and species wood xylem structures, we did not find clear differences in species-level estimates of cue responses with wood structure or relative to their wood specific densities. Each cross represents the 50% uncertainty interval of **a.** forcing, **b.** chilling, and **c.** photoperiod responses and WSD, with colors depicting different types of wood structure.