

Traits predict forest phenological responses to photoperiod more than temperature

Deirdre Loughnan¹, Faith A M Jones^{1,2}, and E M Wolkovich^{1,3,4}

November 28, 2025

¹ Department of Forest and Conservation, Faculty of Forestry, University of British Columbia, 2424 Main Mall Vancouver, BC Canada V6T 1Z4.

² Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sciences, 901 83 Umeå, Sweden.

³ Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

⁴ Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA;

Corresponding Author: Deirdre Loughnan deirdre.loughnan@ubc.ca

Running title: Traits drive photoperiod cues in budburst

Summary

Climate change has advanced the timing of major life cycle events globally, with considerable variation across species and communities. In plants, differences in species growth strategies correlate with variation in leaf and structural traits, creating the basis of frameworks like the leaf economic spectrum. But whether traits related to the timing of growth fit within these frameworks is unclear. We combined trait data from six commonly measured traits with experimental data of woody plant budburst to identify shared relationships between traits, environmental cues—like temperature and photoperiod—and the timing of spring growth. Working across eight forest communities, we collected *in situ* trait measurements from 1428 individuals of 48 species. We paired trait data with budburst data from 4211 cuttings taken from the same individuals in a controlled environment study in which temperature and photoperiod cues varied. We found differences in traits between communities and across latitude, but only four of our six traits related to budburst and only in relation to photoperiod. Despite photoperiod being the weakest cue of woody species budburst, our results suggest that its association with other traits influences the timing of species growth, potentially impacting community composition and species interactions under future climates.

Introduction

Climate change is causing species phenologies—the timing of life history events—to shift, with widespread advances being observed across the tree of life (Parmesan and Yohe, 2003; Hoegh-Guldberg et al., 2018).

This common phenological fingerprint, however, averages over high variability across species (Thackeray et al., 2016; Cohen et al., 2018; Kharouba et al., 2018), posing a challenge to accurate forecasts.

In plants, species variation can be explained, in part, by differences in growth strategies, which are generally inferred from traits (Violle et al., 2007). Decades of research on plant traits have worked to build predictive models of species responses to their environment (Green et al., 2022), which could explain species-level variability in phenological responses. Phenology, however, has generally been excluded from plant trait research due to its high inter- and intra-specific variability, making it difficult to leverage existing frameworks to explain phenological variation and predict future changes.

Previous studies have found high variation in phenology for the same species when observed over different years or sites (Primack et al., 2009; Chuine et al., 2010). But many studies of phenology are observational and conducted at the landscape scale (Fitter and Fitter, 2002; Menzel et al., 2006). In contrast, findings from controlled environment experiments suggest that phenological variation can be consistently decomposed into its cues (Chuine and Cour, 1999; Harrington and Gould, 2015; Flynn and Wolkovich, 2018). Given the importance of phenology in defining when a species grows and the environmental conditions during growth periods, we predict that these phenology-cue relationships may also correlate with other functional traits and vary along an axis of acquisitive to conservative growth strategies.

Correlations between plant traits, growth strategies, and responses to environments have been synthesized into several global frameworks, including the leaf economic spectrum (Wright et al., 2004) and wood economic spectrum (Chave et al., 2009). These frameworks have identified key traits that exhibit distinct gradients, ranging from acquisitive strategies—fast growing plants that produce cheaper tissue—to conservative strategies—with plants that invest in long-lived tissue but slower growth rates (Wright et al., 2004; Díaz et al., 2016). In temperate systems, changes in temperature and frost risk in spring can produce gradients in abiotic stress, including frost risk, soil nutrients, and light availability (Sakai and Larcher, 1987; Gotelli and Graves, 1996; Augspurger, 2009), in addition to differences in biotic interactions from herbivory or competition later in the season (Lopez et al., 2008; Wolkovich and Ettinger, 2014). Species that vary in their timing of leafout, should therefore exhibit traits and growth strategies that allow them to tolerate or avoid these abiotic and biotic factors. Leveraging insights from predictive models of phenology with these well established trait frameworks, we can begin to disentangle the environmental cues that shape phenology from those shaped by other trait differences in plant growth strategies.

To determine whether phenology fits within major functional trait frameworks will require a better understanding of the causes of variation within species. Phenological variation is generally observed in natural conditions where considerable differences in the environmental cues that trigger many phenological events—primarily temperature and photoperiod (Chuine, 2000; Körner and Basler, 2010)—vary across space and time. Within-species variation may also occur across other plant traits (e.g., leaf and wood structure traits), and across latitudinal (Wiemann and Bruce, 2002) or environmental gradients (Pollock et al., 2012), though generally to a smaller scale compared to phenology. While our previous growth chamber study found no evidence that phenological cues varied spatially (Loughnan and Wolkovich, in prep), previous work in functional traits has found some traits to exhibit greater spatial variation in response to environmental gradients (Gross et al., 2000; Wright et al., 2003). These results suggest that to better understand how phenology and other traits correlate across species will require methods that incorporate spatial variation within species.

Here, we tested whether phenological variation was aligned with existing trait frameworks using data on spring budburst paired and a suite of traits that capture acquisitive to conservative growth strategies. We decompose the high phenological variation in budburst date, using experiments to estimate three major phenological cues for woody plant budburst: chilling (cool winter temperatures), forcing (warm

spring temperatures), and photoperiod. We predict that early spring species will budburst before canopy closure—when chilling is lower, forcing temperatures are cooler, and photoperiods shorter—exhibited as smaller responses to temperature and photoperiod. These species should have traits associated with acquisitive growth, particularly shorter heights, smaller trunk or stem diameters, with lower investment in wood structure and leaf tissue, resulting in low wood specific density, diffuse-porous wood anatomy, and low leaf mass area, but high leaf nitrogen content for a greater photosynthetic potential. In contrast, we predict species with later budburst to predominately include canopy species that express more conservative growth strategies and require more chilling, warmer forcing, and longer photoperiods. These species should incur greater investments in long-lived tissue, with ring-porous wood anatomy, taller heights and greater diameter, denser wood and high leaf mass area, but low leaf nitrogen content. We used a powerful joint-modeling approach to estimate the relationships between these plant traits and phenological responses to cues, while partitioning the variance from species- and population-level differences.

Materials and Methods

Field sampling

We combined *in situ* trait data with budburst data from two growth chamber cutting experiments conducted across eastern and western temperate deciduous forests in North America. We collected both suites of data from populations that span a latitudinal gradient of 4-6° for the eastern and western communities respectively. We took trait measurements from across eight populations, of which there were four eastern populations—Harvard Forest, Massachusetts, USA (42.55°N, 72.20°W), White Mountains, New Hampshire, USA (44.11°N, 52.14°W), Second College Grant, New Hampshire, USA (44.79°N, 50.66°W), and St. Hippolyte, Quebec, Canada (45.98°N, 74.01°W), and four western population—E.C. Manning Park (49.06°N, 120.78°W), Sun Peaks (50.88°N, 119.89°W), Alex Fraser Research Forest (52.14°N, 122.14°W), and Smithers (54.78°N, 127.17°W), British Columbia (BC), Canada (Fig. 1). For the two growth chamber studies on budburst phenology, we collected cuttings from the most southern and northern populations in each transect ($n_{pop}=4$).

Functional traits

We measured an assemblage of 47 species from the understory and canopy layers, with 28 species at our eastern transect and 22 species at our western transect, with three species occurring in both transects. We measured all traits in the summer prior to each growth chamber study (eastern transect: 8-25 June 2015, western transect: 29 May to 30 July 2019), following full leafout but before budset. At each population and for each species, we measured a total of five traits from 1-10 healthy adult individuals: height, diameter of the main trunk or stem (hereafter referred to as diameter), wood specific density, leaf mass area, and the percent leaf nitrogen content. We also obtained xylem structure data from the WSL xylem database (Schweingruber and Landolt, 2010) for 72.3% of our species.

We measured traits in accordance to the methods discussed by Pérez-Harguindeguy et al. (2013). We calculated tree height using trigonometric methods and used a base height of 1.37 m to measure diameter (Magarik et al., 2020). For shrub heights, we measured the distance from the ground to the height of the top foliage and measured stem diameters at approximately 1 cm above ground-level. All stem and leaf samples were kept cool during transport and measurements of leaf area and stem volume taken within 3 and 12 hours of sample collection respectively. To measure wood specific density, we collected a 10 cm sample of branch wood, taken close to the base of the branch at the stem and calculated stem volume using the water displacement method. For our leaf traits, we haphazardly selected and sampled five, fully expanded, and hardened leaves, with no to minimal herbivore damage.

135 We took a high resolution scan of each leaf using a flatbed scanner and estimated leaf area using the
 136 ImageJ software (version 2.0.0).

137 Growth chamber study

138 For our growth chamber studies, we collected branch cuttings from our highest and lowest latitude
 139 populations in each transect, with sampling in our eastern study occurring from 20-28 January 2015
 140 and sampling for our western study from 19-28 October 2019. In both studies, we included a total of
 141 eight distinct treatments consisting of two levels of chilling, forcing, and photoperiods (Fig. 1). We
 142 recorded budburst stages of each sample every 1-3 days for up to four months, defining the day of
 143 budburst as the day of budbreak or shoot elongation (denoted as code 07 by Finn et al. (2007)). For
 144 a more detailed discussion of study sample collection and methods see Flynn and Wolkovich (2018)
 145 for details on our eastern study and Loughnan and Wolkovich (in prep) for details on our western study.
 146

147 Statistical Analysis

148 Our analysis combined our *in situ* trait data with budburst data from the controlled environment. For
 149 each trait, we developed a joint Bayesian model, in which the relationship between traits and cues
 150 is used to estimate budburst. This statistical approach improves upon previous analyses of multiple
 151 traits, as it allows us to carry through uncertainty between trait and phenology data—and better
 152 partitions the drivers of variation in species phenologies
 153

154 Our joint model consists of two parts. The first is a hierarchical linear model, which partitions the
 155 variation of individual observations (i) of a given trait value (Y_{trait}) to account for the effects of species
 156 (j), population-level differences arising from transects, latitude, as well as the interaction between tran-
 157 sects and latitude ($\text{transect} \times \text{latitude}$), and finally, residual variation or ‘measurement error’ (σ_m^2).
 158

$$Y_{\text{trait}_{i,j}} \sim \text{Normal}(\mu_{i,j}, \sigma_m^2) \quad (1)$$

$$\mu_{i,j} = \alpha_{\text{grand trait}} + \alpha_{\text{trait}_j} + \beta_{\text{transect}} \times \text{transect} + \quad (2)$$

$$\beta_{\text{latitude}} \times \text{latitude} + \beta_{\text{transect} \cdot \text{latitude}} \times (\text{transect} \cdot \text{latitude}) \quad (3)$$

$$(4)$$

159

$$\alpha_{\text{trait}} \begin{bmatrix} \alpha_{\text{trait}_1} \\ \alpha_{\text{trait}_2} \\ \dots \\ \alpha_{\text{trait}_n} \end{bmatrix} \text{ such that } \alpha_{\text{trait}} \sim \text{Normal}(0, \sigma_{\text{trait}}^2) \quad (5)$$

$$(6)$$

160 We include transect as a dummy variable (0/1) and latitude as a continuous variable in our model.
 161 We modeled traits using natural units, with the exception of LMA, which was rescaled by 100 for
 162 numeric stability in the model. Our model also includes partial pooling for species—which controls for
 163 variation in the number of trait estimates per species and trait variability—using these species-level
 164 estimates as predictors for each cue ($\beta_{\text{chilling},j}$, $\beta_{\text{forcing},j}$, $\beta_{\text{photoperiod},j}$).
 165

$$\beta_{\text{chilling}_j} = \alpha_{\text{chilling},j} + \beta_{\text{trait.chilling}} \cdot \alpha_{\text{trait},j} \quad (7)$$

$$\beta_{\text{forcing}_j} = \alpha_{\text{forcing},j} + \beta_{\text{trait.forcing}} \cdot \alpha_{\text{trait},j}$$

$$\beta_{\text{photoperiod}_j} = \alpha_{\text{photoperiod},j} + \beta_{\text{trait.photoperiod}} \cdot \alpha_{\text{trait},j}$$

166 In addition to the species-level estimates, the second part of our model estimates the overall effect of
 167 each trait on each cue ($\beta_{\text{trait.chilling}}$, $\beta_{\text{trait.forcing}}$, $\beta_{\text{trait.photoperiod}}$). From this we can estimate how well
 168 traits explain species-level differences—by estimating the the species-level cue variation not explained
 169 by traits ($\alpha_{\text{chilling},j}$, $\alpha_{\text{forcing},j}$, $\alpha_{\text{photoperiod},j}$) and individual species responses to cues (*chilling*, *forcing*,
 170 *photoperiod*, respectively). Finally, our model estimates the residual budburst variation across species
 171 ($Y_{\text{pheno},j}$), observations (σ_d^2), as well as the variation in cues not attributed to the trait (using partial
 172 pooling).

$$Y_{\text{pheno}_{i,j}} \sim \mathcal{N}(\mu_{i,j}, \sigma_d^2) \quad (8)$$

173 with

$$\mu_{i,j} = \alpha_{\text{pheno}_j} + \beta_{\text{chilling}_j} \cdot \textit{chilling} + \beta_{\text{forcing}_j} \cdot \textit{forcing} + \beta_{\text{photoperiod}_j} \cdot \textit{photoperiod} \quad (9)$$

174 where α_{pheno_j} , $\alpha_{\text{chilling}_j}$, $\alpha_{\text{forcing}_j}$, and $\alpha_{\text{photoperiod}_j}$ are elements of the normal random vectors:

$$\begin{aligned} \boldsymbol{\alpha}_{\text{pheno}} &= \{\alpha_{\text{pheno}_1}, \dots, \alpha_{\text{pheno}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{pheno}} \sim \mathcal{N}(\mu_{\text{pheno}}, \sigma_{\text{pheno}}^2) \\ \boldsymbol{\alpha}_{\text{chilling}} &= \{\alpha_{\text{chilling}_1}, \dots, \alpha_{\text{chilling}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{chilling}} \sim \mathcal{N}(\mu_{\text{chilling}}, \sigma_{\text{chilling}}^2) \\ \boldsymbol{\alpha}_{\text{forcing}} &= \{\alpha_{\text{forcing}_1}, \dots, \alpha_{\text{forcing}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{forcing}} \sim \mathcal{N}(\mu_{\text{forcing}}, \sigma_{\text{forcing}}^2) \\ \boldsymbol{\alpha}_{\text{photo}} &= \{\alpha_{\text{photo}_1}, \dots, \alpha_{\text{photo}_n}\}^T \text{ such that } \boldsymbol{\alpha}_{\text{photo}} \sim \mathcal{N}(\mu_{\text{photo}}, \sigma_{\text{photo}}^2) \end{aligned} \quad (10)$$

175 We modeled each trait individually, with the exception of ring-porosity, which we compared across
 176 species using the posterior estimates of our WSD model, in order to account for inherent differences
 177 in wood anatomy across species and growth form. We included all three cues (chilling, forcing, and
 178 photoperiod) as continuous variables in our model, as well as all two-way interactions between cues
 179 and between cues and sites. We converting chilling temperatures to total chill portions, including both
 180 the chilling experienced in the field prior to sampling and during the experiment in our 4°C chilling
 181 chambers, using local weather station data and used the chillR package to calculate total chill portions
 182 (v. 0.73.1, Luedeling, 2020). To account for differences in thermoperiodicity between the two studies
 183 (Buonaiuto et al., 2023), we also converted forcing temperatures to mean daily temperatures for each
 184 treatment. Finally, we *z*-scored each cue and site using two standard deviations to allow direct com-
 185 parisons between results across parameters (Gelman, 2008).

186
 187 We used weakly informative priors unique to each trait model. We validated our choice of priors us-
 188 ing prior predictive checks and confirmed model stability under wider priors. All models were coded
 189 in the Stan programming language for Bayesian models using the rstan package (Stan Development
 190 Team, 2018) in R version 4.3.1 (R Development Core Team, 2017). All models met basic diagnostic
 191 checks, including no divergences, high effective sample sizes (n_{eff}) that exceeded 10% of the number
 192 of iterations, and \hat{R} values close to 1. We report our model estimates as the mean values with 90%
 193 uncertainty intervals, interpreting parameter estimates with UI that overlap to be statistically similar
 194 to each other and intervals that include zero to have small effects.

195

Results

Across our eight populations, we measured 47 species of which 28 were in our eastern transect and 22 in our western transect. These include species dominant in both the understory and canopy layer, with our eastern community consisting of 13 shrubs and 15 trees, our western community consisting of 18 shrubs and 4 trees, and three species that occurred in both transects. In total we measured traits of 1428 unique individuals between the two transects across our five traits: height ($n = 1317$), diameter ($n = 1220$), wood stem density ($n = 1359$), leaf mass area (LMA, $n = 1345$), leaf nitrogen content (LNC, $n = 1351$). Across our two growth chamber studies, we made observations of 4211 samples, with our observations of budburst spanning 82 and 113 days for our eastern and western studies respectfully.

Most of our traits showed some variation by latitude within each transect (with a strong interactive effect between transect and latitude, Fig. 2). Leaf nitrogen content was the only trait to vary with latitude alone, with low latitude communities on both our eastern and western transects having greater values of LNC than communities at higher latitudes (-0.1, UI: -0.2, -0.06, Table S6). The strongest negative interaction was observed for height (-0.2, UI: -0.4, 0), while the strongest positive interaction was observed for LMA (0.5, UI: 0.4, 0.6). Overall, woody species in our eastern community exhibited greater heights, diameter, and WSD but decreases in LMA with increasing latitude. But in our western communities height and WSD decreased with latitude, while diameter and leaf mass area both increased (Fig. 2). In addition to the differences we found across populations, we also observed considerable differences between individual species, which varied considerably and up to 7 fold for some traits (Fig. 3).

We found that three of our four traits had a strong relationship with photoperiod, but not always in the direction we predicted. Taller species with larger trunk diameters and leaves with high nitrogen content had stronger responses to photoperiod (Table S6), Fig. 3 c, i, o; Tables S2, S3, S5). Contrary to our expectation, species with denser, high LMA leaves, however, had weaker photoperiod responses, allowing them to potentially budburst under shorter photoperiods (Fig. 3 f).

Temperature cues exhibited no relationships with individual traits ($\beta_{\text{trait.chilling}}$), but by accounting for the effects of leaf or wood traits, we found the importance of our three cues the vary by trait. Our models of height (-13.4, UI: -17.2, -9.9), diameter (-12.5, UI: -16.2, -8.6), WSD (-20.9, UI: -33.2, -9.8), and LNC (-35.1, UI: -68.1, -4.1) all estimated chilling as the strongest cue of budburst (β_{chilling}), with more chilling advancing budburst. Our model of LMA, however, estimated photoperiod as the strongest cue ($\beta_{\text{photoperiod}}$, -2.7, UI: -30.3, 26). After accounting for the effects of traits, only our height and diameter model found all three environmental cues to drive budburst timing (Tables S2, S3). Our models of WSD and LNC in turn found temperature cues alone to shape budburst (Tables S4, S6), while our model of LMA found a large response to only photoperiod (Table S5).

In synthesizing the effects of multiple traits across species, our results can be used to make generalizations across ecologically important groups of species. But only some of our models estimated clear gradients in species timing between trees and shrubs. In particular, we found height to have large correlations between budburst timing and trait values, with earlier estimates of budburst for shrubs (with a mean day of budburst of 10)—especially under greater cues—and later budburst estimates for trees (with a mean day of budburst of 17.3). Diameter at breast height showed similar trends as estimates from our height model (results not shown). But this was not the case for our two leaf traits. Leaf nitrogen content, for example, showed no distinct separation between shrub and tree functional groups

Discussion

Using our joint modeling approach, we estimated how leaf and wood traits interact with temperature and photoperiod cues to shape species budburst. We found that photoperiod—often the weakest cue of budburst (Laube et al., 2014; Zohner et al., 2016; Flynn and Wolkovich, 2018)—was the most important cue in trait-phenology relationships. In general, we also found trait patterns varied between our eastern and western transects and with latitude. These spatial differences in trait variation may be due to differences in the community assemblages found in our eastern versus western forests. Plants at our western sites were shorter with less dense branch wood, suggesting a more acquisitive growth strategies on average. Given that our western community is shrub dominated, this suggests these species are more likely to utilize resources early in the season prior to canopy closure. Collectively our results provide new insights into the complexity of the mechanisms that underlie the relationships between traits and environmental cues, while also challenging our existing understanding of these processes.

Cues and functional traits

We predicted species with acquisitive traits—particularly small trees with low wood densities, and less dense leaves with low leaf mass area, and high leaf nitrogen content—to have early budburst via small temperature and photoperiod responses. But we found only partial support for this prediction. In line with our predictions, acquisitive species with smaller heights and diameter did have smaller photoperiod responses (associated with later budburst). But contrary to our prediction, species with less dense leaves showed larger responses to photoperiod, while leaves with high nitrogen content had stronger photoperiod responses. We also did not find any relationships between any of our focal traits and temperature. These unexpected trends in our results may be due to selection on other physiological processes, as many of our traits are associated with one or more ecological function (Wright et al., 2004; Pérez-Harguindeguy et al., 2013; Reich, 2014). Our results highlight the ways in which phenology partially aligns with gradients found in established trait frameworks, while offering new insight into potential tradeoffs in how varying physiological processes shape species growth strategies.

Decades of previous research that only considered phenology found budburst timing to be primarily driven by temperature and weakly by photoperiod (Chuine et al., 2010; Basler and Körner, 2014; Laube et al., 2014). We, however, found no traits that correlate with responses to temperature, suggesting other cues are impacting leaf and structural traits in temperate forests. In particular, leaf mass area is known to correlate with other traits like leaf lifespan or decomposition rates in addition to light capture (De La Riva et al., 2016), traits which may respond more strongly to other cues. Leaf mass area also varies with soil moisture, with variation in the area of a leaf allowing plants to reduce evaporation under dry conditions, thus selecting for high values of leaf mass area (De La Riva et al., 2016). Soil moisture is known to shape other phenological events in woody plants, including radial growth phenology and shoot elongation (Cabon et al., 2020; Peters et al., 2021). If selection by soil moisture is shaping phenological responses, it may be contributing to the unexpected trends we observed in leaf mass area and the absence of relationships with temperature. Thus, fully understanding how species growth strategies correlate with phenology may require the consideration of cues known to shape other plant traits.

The absence of trait-cue relationships between budburst and wood structure and WSD also contrasts the findings of previous work linking these traits. Previous studies have found some evidence that trees with diffuse-porous wood structure leaf out earlier than species with ring-porous structures (Lechowicz, 1984; Panchen et al., 2014; Yin et al., 2016; Osada, 2017; Savage et al., 2022). But using wood density as a proxy for wood structure, with wood density positively correlating with xylem resistance to embolism (Hacke et al., 2001), we did not find clear association between our three cues and xylem structure. However, the positive relationship between wood density and latitude in our east-

ern community may in part be caused by the differences in winter conditions experienced by canopy versus understory species. The canopy tree species that dominate our eastern communities may experience greater horizontal stress from wind and downward pressure from snow, explaining the greater wood densities they exhibit at higher latitudes (MacFarlane and Kane, 2017; MacFarlane, 2020), while species in the shrub dominated western communities experience greater protection from being in the understory.

In comparing our results with a global meta-analysis of tree trait relationships with budburst cues (Loughnan et al., 2025), we found similar trait-cue relationships for several of our traits. At both the global and continental scales, we found taller tree heights to leaf out with longer photoperiods, while in both studies, species with high specific leaf area (which is the inverse and thus equivalent to low leaf mass area) exhibited large responses to photoperiod (Loughnan et al., 2025). The consistency of these results, despite the differences in their two spatial scales of study, provides further evidence that alternate underlying mechanisms are shaping how species respond to photoperiod cues.

Functional traits predict climate change responses

Our results offer novel insights into how broader correlations between growth strategies and phenological cues can help predict responses in plant communities with climate change. As temperatures rise, particularly at higher latitudes (Hoegh-Guldberg et al., 2018), warmer winter and spring temperatures, may select for earlier budburst in some species, but constrain others as photoperiod will remain fixed. The relationships between photoperiod and other traits we observed have the potential to limit species abilities to track temperatures and constrain the extent to which woody plant phenologies will advance with climate change. Our results suggest that these effects will likely be greater for taller species or canopy trees and species with relatively low leaf mass area. These constraints could have cascading effects on forest communities, as variable species responses to increasing temperatures further alter species growth strategies and their interactions with competitors or herbivores within their communities.

Our findings of correlations between phenology and other commonly measured traits highlight how accurate forecasts of future changes in phenology can benefit from accounting for the response of other traits to climate change. Across temperature and precipitation gradients, leaf size and shape also change, as species shift to conserve water and mitigate effects of transpiration under higher temperatures (De La Riva et al., 2016). These changes could impact species photosynthetic potential and ultimately ecosystem services, such as carbon sequestration. While phenological research has focused on forecasting responses to temperature, the correlation of other traits with photoperiod suggests its importance as a cue. It is therefore advantageous for additional traits that are likely to respond to photoperiod to further studied in relation to phenology when forecasting the future impacts of climate change on communities.

References

- Augspurger, C. K. 2009. Spring 2007 warmth and frost: phenology, damage and refoliation in a temperate deciduous forest. *Functional Ecology* 23:1031–1039.
- Basler, D., and C. Körner. 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* 34:377–388.
- Buonaiuto, D. M., E. M. Wolkovich, and M. J. Donahue. 2023. Experimental designs for testing the

interactive effects of temperature and light in ecology : The problem of periodicity. *Functional Ecology* 37:1747–1756.

Cabon, A., L. Fernández-de-Uña, G. Gea-Izquierdo, F. C. Meinzer, D. R. Woodruff, J. Martínez-Vilalta, and M. De Cáceres. 2020. Water potential control of turgor-driven tracheid enlargement in Scots pine at its xeric distribution edge. *New Phytologist* 225:209–221.

Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a worldwide wood economics spectrum. *Ecology Letters* 12:351–366.

Chuine, I. 2000. A unified model for budburst of trees. *Journal of Theoretical Biology* 207:337–347.

Chuine, I., and P. Cour. 1999. Climatic determinants of budburst seasonality in four temperate-zone tree species. *New Phytologist* 143:339–349.

Chuine, I., X. Morin, and H. Bugmann. 2010. Warming, photoperiods, and tree phenology. *Science* 329:277–278.

Cohen, J. M., M. J. Lajeunesse, and J. R. Rohr. 2018. A global synthesis of animal phenological responses to climate change. *Nature Climate Change* 8:224–228.

De La Riva, E. G., M. Olmo, H. Poorter, J. L. Uberta, and R. Villar. 2016. Leaf Mass per Area (LMA) and Its Relationship with Leaf Structure and Anatomy in 34 Mediterranean Woody Species along a Water Availability Gradient. *PLOS ONE* 11:e0148788.

Díaz, S., J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer, C. Wirth, I. Colin Prentice, E. Garnier, G. Bönsch, M. Westoby, H. Poorter, P. B. Reich, A. T. Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright, S. N. Sheremet’ev, H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig, A. Günther, V. Falczuk, N. Rüger, M. D. Mahecha, and L. D. Gorné. 2016. The global spectrum of plant form and function. *Nature* 529:167–171.

Finn, G. A., A. E. Straszewski, and V. Peterson. 2007. A general growth stage key for describing trees and woody plants. *Annals of Applied Biology* 151:127–131.

Fitter, A. H., and R. S. Fitter. 2002. Rapid changes in flowering time in British plants. *Science* 296:1689–1691.

Flynn, D. F. B., and E. M. Wolkovich. 2018. Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytologist* 219:1353–1362.

Gelman, A. 2008. Scaling regression inputs by dividing by two standard deviations. *Statistics in Medicine* 27:2865–2873.

Gotelli, N. J., and G. R. Graves. 1996. The temporal niche. Pages 95–112 *in* Null Models In Ecology. Smithsonian Institution Press, Washington, D. C.

Green, S. J., C. B. Brookson, N. A. Hardy, and L. B. Crowder. 2022. Trait-based approaches to global change ecology: moving from description to prediction. *Proceedings of the Royal Society B: Biological Sciences* 289:1–10.

Gross, K. L., M. R. Willig, L. Gough, R. Inouye, and S. B. Cox. 2000. Patterns of species density and productivity at different spatial scales in herbaceous plant communities. *Oikos* 89:417–427.

Hacke, U. G., J. S. Sperry, W. T. Pockman, S. D. Davis, and K. A. McCulloh. 2001. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia* 126:457–461.

377 Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying dormancy
378 requirements for Pacific Northwest tree species. *Frontiers in Plant Science* 6:1–12.

379 Hoegh-Guldberg, O., D. Jacob, M. Taylor, M. Bindi, S. Brown, I. Camilloni, A. Diedhiou, R. Djalante,
380 K. Ebi, F. Engelbrecht, J. Guiot, Y. Hijioka, S. Mehrotra, A. Payne, S. Seneviratne, A. Thomas,
381 R. Warren, and G. Zhou. 2018. Impacts of 1.5 °C Global Warming on Natural and Human Systems.
382 In: *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C*
383 *above pre-industrial levels and related global greenhouse gas emission pathways, in the context of .*
384 *Tech. rep.*, Cambridge University Press, Cambridge, UK and New York, NY, USA.

385 Kharouba, H. M., J. Ehrlén, A. Gelman, K. Bolmgren, J. M. Allen, S. E. Travers, and E. M. Wolkovich.
386 2018. Global shifts in the phenological synchrony of species interactions over recent decades. *Pro-*
387 *ceedings of the National Academy of Sciences* 115:5211–5216.

388 Körner, C., and D. Basler. 2010. Phenology Under Global Warming. *Science* 327:1461–1463.

389 Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling outweighs
390 photoperiod in preventing precocious spring development. *Global Change Biology* 20:170–182.

391 Lechowicz, M. J. 1984. Why Do Temperate Deciduous Trees Leaf Out at Different Times? *Adaptation*
392 *and Ecology of Forest Communities. The American Naturalist* 124:821–842.

393 Lopez, O. R., K. Farris-Lopez, R. A. Montgomery, and T. J. Givnish. 2008. Leaf phenology in relation
394 to canopy closure in southern Appalachian trees. *American Journal of Botany* 95:1395–1407.

395 Loughnan, D., F. A. Jones, G. Legault, C. J. Chamberlain, D. M. Buonaiuto, A. K. Ettinger, M. Gar-
396 ner, D. S. Sodhi, and E. M. Wolkovich. 2025. Budburst timing within a functional trait framework.
397 *Journal of Ecology* 00:1–12.

398 Loughnan, D., and E. M. Wolkovich. in prep. Temporal assembly of woody plant communities shaped
399 equally by evolutionary history as by current environments .

400 Luedeling, E. 2020. chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees.
401 <https://CRAN.R-project.org/package=chillR>.

402 MacFarlane, D. W. 2020. Functional Relationships Between Branch and Stem Wood Density for
403 Temperate Tree Species in North America. *Frontiers in Forests and Global Change* 3.

404 MacFarlane, D. W., and B. Kane. 2017. Neighbour effects on tree architecture: functional trade-offs
405 balancing crown competitiveness with wind resistance. *Functional Ecology* 31:1624–1636.

406 Magarik, Y. A., L. A. Roman, and J. G. Henning. 2020. How should we measure the dbh of multi-
407 stemmed urban trees? *Urban Forestry & Urban Greening* 47:1–11.

408 Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aaasa, R. Ahas, K. Alm-Kübler, P. Bissolli,
409 O. Braslavská, A. Briede, F. M. Chmielewski, Z. Crepinsek, Y. Curnel, Å. Dahl, C. Defila, A. Don-
410 nelly, Y. Filella, K. Jactzak, F. Måge, A. Mestre, Ø. Nordli, J. Peñuelas, P. Pirinen, V. Remišová,
411 H. Scheffinger, M. Striz, A. Susnik, A. J. Van Vliet, F. E. Wielgolaski, S. Zach, and A. Zust. 2006.
412 European phenological response to climate change matches the warming pattern. *Global Change*
413 *Biology* 12:1969–1976.

414 Osada, N. 2017. Relationships between the timing of budburst, plant traits, and distribution of 24
415 coexisting woody species in a warm-temperate forest in Japan. *American Journal of Botany* 104:550–
416 558.

417 Panchen, Z. A., R. B. Primack, B. Nordt, E. R. Ellwood, A. Stevens, S. S. Renner, C. G. Willis,
418 R. Fahey, A. Whitemore, Y. Du, and C. C. Davis. 2014. Leaf out times of temperate woody
419 plants are related to phylogeny, deciduousness, growth habit and wood anatomy. *New Phytologist*
420 203:1208–1219.

421 Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across
422 natural systems. *Nature* 421:37–42.

423 Pérez-Harguindeguy, N., S. Díaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S. Bret-
424 Harte, W. K. Cornwell, J. M. Craine, D. E. Gurvich, C. Urcelay, E. J. Veneklaas, P. B. Reich,
425 L. Poorter, I. J. Wright, P. Ray, L. Enrico, J. G. Pausas, A. C. de Vos, N. Buchmann, G. Funes,
426 F. Quétier, J. G. Hodgson, K. Thompson, H. D. Morgan, H. ter Steege, M. G. A. van der Heijden,
427 L. Sack, B. Blonder, P. Poschlod, M. V. Vaieretti, G. Conti, A. C. Staver, S. Aquino, and J. H. C.
428 Cornelissen. 2013. New handbook for standardized measurement of plant functional traits worldwide.
429 *Australian Journal of Botany* 61:167–234.

430 Peters, R. L., K. Steppe, H. E. Cuny, D. J. De Pauw, D. C. Frank, M. Schaub, C. B. Rathgeber,
431 A. Cabon, and P. Fonti. 2021. Turgor – a limiting factor for radial growth in mature conifers along
432 an elevational gradient. *New Phytologist* 229:213–229.

433 Pollock, L. J., W. K. Morris, and P. A. Vesk. 2012. The role of functional traits in species distributions
434 revealed through a hierarchical model. *Ecography* 35:716–725.

435 Primack, R. B., I. Ibáñez, H. Higuchi, S. D. Lee, A. J. Miller-Rushing, A. M. Wilson, and J. A. Silander.
436 2009. Spatial and interspecific variability in phenological responses to warming temperatures.
437 *Biological Conservation* 142:2569–2577.

438 R Development Core Team. 2017. R: A language and environment for statistical computing.

439 Reich, P. B. 2014. The world-wide ‘fast–slow’ plant economics spectrum: a traits manifesto. *Journal*
440 *of Ecology* 102:275–301.

441 Sakai, A., and W. Larcher. 1987. Frost Survival of Plants: Responses and adaptation to freezing stress.
442 Springer-Verlag, Berlin, Heidelberg.

443 Savage, J. A., T. Kiecker, N. McMann, D. Park, M. Rothendler, and K. Mosher. 2022. Leaf out time
444 correlates with wood anatomy across large geographic scales and within local communities. *New*
445 *Phytologist* 235:953–964.

446 Schweingruber, F., and W. Landolt. 2010. The xylem database.

447 Stan Development Team. 2018. RStan: the R interface to Stan. R package version 2.17.3.

448 Thackeray, S. J., P. A. Henrys, D. Hemming, J. R. Bell, M. S. Botham, S. Burthe, P. Helaouet,
449 D. G. Johns, I. D. Jones, D. I. Leech, E. B. MacKay, D. Massimino, S. Atkinson, P. J. Bacon,
450 T. M. Brereton, L. Carvalho, T. H. Clutton-Brock, C. Duck, M. Edwards, J. M. Elliott, S. J. Hall,
451 R. Harrington, J. W. Pearce-Higgins, T. T. Høye, L. E. Kruuk, J. M. Pemberton, T. H. Sparks,
452 P. M. Thompson, I. White, I. J. Winfield, and S. Wanless. 2016. Phenological sensitivity to climate
453 across taxa and trophic levels. *Nature* 535:241–245.

454 Violle, C., M. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let the
455 concept of trait be functional! *Oikos* 116:882–892.

456 Wiemann, M. C., and W. G. Bruce. 2002. Geographic variation in wood specific gravity: effects of
457 latitude, temperature, and precipitation. *Wood and Fiber Science* 34:96–107.

458 Wolkovich, E. M., and A. K. Ettinger. 2014. Back to the future for plant phenology research. *New*
459 *Phytologist* 203:1021–1024.

- 460 Wright, I. J., M. Westoby, P. B. Reich, J. Oleksyn, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-
461 Bares, T. Chapin, J. H. C. Cornellissen, M. Diemer, J. Flexas, J. Gulias, E. Garnier, M. L. Navas,
462 C. Roumet, P. K. Groom, B. B. Lamont, K. Hikosaka, T. Lee, W. Lee, C. Lusk, J. J. Midgley,
463 Ü. Niinemets, H. Osada, H. Poorter, P. Pool, E. J. Veneklaas, L. Prior, V. I. Pyankov, S. C.
464 Thomas, M. G. Tjoelker, and R. Villar. 2004. The worldwide leaf economics spectrum. *Nature*
465 428:821–827.
- 466 Wright, J. P., A. S. Flecker, and C. G. Jones. 2003. Local vs. landscape controls on plant species
467 richness in Beaver meadows. *Ecology* 84:3162–3173.
- 468 Yin, J., J. D. Fridley, M. S. Smith, and T. L. Bauerle. 2016. Xylem vessel traits predict the leaf
469 phenology of native and non-native understorey species of temperate deciduous forests. *Functional*
470 *Ecology* 30:206–214.
- 471 Zohner, C. M., B. M. Benito, J.-C. Svenning, and S. S. Renner. 2016. Day length unlikely to constrain
472 climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate Change* 6:1120–
473 1123.




Figure 1: We found a. height, b. diameter, c. branch stem density, and d. leaf mass area to all experience a strong interaction between latitude and transect, e. while leaf nitrogen content showed a strong effect of latitude alone.




Figure 2: We found a. height, b. diameter, c. branch stem density, and d. leaf mass area to all experience a strong interaction between latitude and transect, e. while leaf nitrogen content showed a strong effect of latitude alone.




Figure 3: Relationships between species traits and cue responses showed considerable variation across a-c. height, d-f. leaf mass area, g-i. diameter, j-l. wood specific density, and m-o. the leaf nitrogen content. Point colours representing different species groups, with tree species shown in red and shrub species in blue. Crosses depict the 50% uncertainty interval of the model estimates of species trait values and estimated responses to cues. The grey band depicts the 90% uncertainty interval and the black line the mean response.




Figure 4: Despite previous studies finding relationships between leaf out timing and species wood xylem structures, we did not find clear differences in species-level estimates of cue responses with wood structure or relative to their wood specific densities. Each cross represents the 50% uncertainty interval of **a.** forcing, **b.** chilling, and **c.** photoperiod responses and WSD, with colors depicting different types of wood structure.