

1 Traits predict forest phenological responses to photoperiod
2 more than temperature

3 Deirdre Loughnan¹, Faith A M Jones^{1,2}, and E M Wolkovich^{1,3,4}

4 ¹ Department of Forest and Conservation, Faculty of Forestry, University of British Columbia, 2424
5 Main Mall Vancouver, BC Canada V6T 1Z4.

6
7 ² Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sci-
8 ences, 901 83 Umeå, Sweden.

9
10 ³ Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

11
12 ⁴ Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts,
13 USA;

14
15 Corresponding Author: Deirdre Loughnan

16 deirdre.loughnan@ubc.ca

17 604-802-6917

18 6270 University Blvd #4200,

19 Vancouver, BC V6T 1Z4

20 Canada

21
22 Faith A M Jone: faith.jones@slu.se

23 E M Wolkovich: e.wolkovich@ubc.ca

24
25 Running title: Traits mediate budburst responses to photoperiod

26 Keywords: Phenology, Budburst, Plant traits, Forest communities, Chilling, Forcing, Photoperiod

27 Article type: letter

28
29 Abstract length: 149

30 Main text length: XXXX

31 Number of references: 65

32 Number of figures: 4

33

Summary

Shifts in the timing of spring phenological events, like leafout, can have major ecosystem consequences. Understanding how species phenology fits within broader frameworks of plant growth strategies could help predict these impacts. Theory predicts that species with early spring phenology should have a more acquisitive strategy while later species should be more conservative. Yet testing these predictions has been slow given the high natural variability of phenology across space and time. Using controlled environment studies, we reduce phenological variation into its component responses to winter and spring temperatures and photoperiod, then test for growth strategy relationships via five traits measured across 47 species sampled from eight forests across North America. We find phenology connects to plant traits via photoperiod responses, but not temperature. Our results suggest photoperiod responses may be a critical component of how spring phenology fits within plant growth strategies, which could help predict shifts in forest ecosystems with future climate change.

Keywords

Phenology, budburst, leafout, traits, forest communities, woody plants.

Introduction

Climate change is causing species phenologies—the timing of life history events—to shift, with widespread advances observed across the tree of life (Parmesan & Yohe, 2003; Hoegh-Guldberg *et al.*, 2018). This common phenological fingerprint, however, averages over high variability across species (Thackeray *et al.*, 2016; Cohen *et al.*, 2018; Kharouba *et al.*, 2018), posing a challenge to accurate forecasts.

In plants, species variation can be explained, in part, by differences in growth strategies, which are generally inferred from traits (Violle *et al.*, 2007). Decades of research on plant traits have worked to build predictive models of species responses to their environment (Green *et al.*, 2022), which could explain species-level variability in phenological responses. Phenology, however, has generally been excluded from plant trait research due to its high variability, even for the same species when observed over different years or sites (Primack *et al.*, 2009; Chuine *et al.*, 2010). Within-species variation also occurs across other plant traits (e.g. leaf and wood structure traits), but generally to a much smaller scale compared to phenology (Violle *et al.*, 2007). Yet increasing evidence that spring phenology is much less variable when carefully measured (Ettinger *et al.*, 2020) holds promise to leverage existing frameworks to explain phenological variation and predict future changes.

Correlations between plant traits, growth strategies, and responses to environments have been synthesized into several global frameworks, including the leaf economic spectrum (Wright *et al.*, 2004) and wood economic spectrum (Chave *et al.*, 2009). These frameworks have identified key traits that exhibit distinct gradients, ranging from acquisitive strategies—fast growing plants that produce cheaper tissue—to conservative strategies—with plants that invest in long-lived tissue but slower growth rates

71 (Wright *et al.*, 2004; Díaz *et al.*, 2016). In temperate systems, changes in temperature in the spring
72 can produce gradients in abiotic stress, including frost risk, soil nutrients, and light availability (Sakai
73 & Larcher, 1987; Gotelli & Graves, 1996; Augspurger, 2009), in addition to differences in biotic inter-
74 actions from herbivory or competition later in the season (Lopez *et al.*, 2008; Wolkovich & Ettinger,
75 2014). Species that vary in their timing of leafout should therefore exhibit traits and growth strategies
76 that allow them to tolerate or avoid these abiotic and biotic factors, but testing how phenology fits
77 within major functional trait frameworks requires working across within- and between-species variation.

78
79 Phenology is generally observed in natural conditions where variation in the environment across space
80 and time generates high variation in leafout, flowering and other phenological events (Chaine, 2000;
81 Körner & Basler, 2010), which has stymied previous efforts to understand the role of phenology in
82 existing trait frameworks. Increasing evidence, however, shows that phenology—especially spring
83 phenology—is often far less variable when measured in experiments (Basler & Körner, 2014; Vitasse
84 *et al.*, 2009). Experiments in controlled environments (growth chambers) in particular can consistently
85 decompose spring phenology for woody plants into its individual environmental cues (Chaine & Cour,
86 1999; Harrington & Gould, 2015; Flynn & Wolkovich, 2018): chilling (cool winter temperatures), forc-
87 ing (warm spring temperatures), and photoperiod. Decades of experiments now show how these cues
88 combine to determine early versus late spring phenology across species. While all species show strong
89 responses to temperature cues, with chilling generally the strongest cue, later species tend to have
90 larger responses to forcing, chilling and photoperiod (Ettinger *et al.*, 2020). Within species, evidence
91 further suggests spatial gradients in climate may drive variation in responses to cues (Vitasse *et al.*,
92 2018; Wang *et al.*, 2025).

93
94 Here, we tested whether phenological variation was aligned with existing trait frameworks using data on
95 spring budburst paired with a suite of traits that capture acquisitive to conservative growth strategies
96 measured across eight sites across North America. We decomposed the high phenological variation in
97 budburst date by using experiments to estimate chilling, forcing, and photoperiod cues for each species.
98 We predicted that shrub and early spring species, which generally budburst before canopy closure
99 (Augspurger, 2008; Vitasse *et al.*, 2013), will have smaller responses to temperature and photoperiod
100 and have traits associated with acquisitive growth. These individuals would thus be shorter, with
101 smaller trunks or stem diameters, and a lower investment in wood structure and leaf tissue, resulting
102 in low branch wood density, diffuse-porous wood anatomy, and low leaf mass per area, but high leaf
103 nitrogen content for a greater photosynthetic potential (Fig. 1). In contrast, we predicted species with
104 later budburst (requiring more chilling, forcing, and longer photoperiods) to predominately include tree
105 species that have more conservative growth strategies. These species should incur greater investments
106 in long-lived tissue, with ring-porous wood anatomy, taller heights and greater diameter, denser wood
107 and high leaf mass per area, but low leaf nitrogen content (Fig. 1). We further expected traits to vary
108 spatially, with individuals at more northern sites being relatively smaller, with a lower investment in leaf
109 tissue relative to our southern sites. We used a joint-modeling approach to estimate the relationships
110 between these plant traits and phenological responses to cues, while partitioning the variance from
111 species- and population-level differences.

Materials and Methods

Field sampling

We combined *in situ* trait data with budburst data from two growth chamber cutting experiments conducted across eastern and western temperate deciduous forests in North America. We collected both suites of data from populations that span a latitudinal gradient of 4-6° for the eastern and western sites respectively. We took trait measurements from across eight populations, of which there were four eastern populations—Harvard Forest, Massachusetts, USA (42.55°N, 72.20°W), White Mountains, New Hampshire, USA (44.11°N, 52.14°W), Second College Grant, New Hampshire, USA (44.79°N, 50.66°W), and St. Hippolyte, Quebec, Canada (45.98°N, 74.01°W), and four western populations—E.C. Manning Park (49.06°N, 120.78°W), Sun Peaks (50.88°N, 119.89°W), Alex Fraser Research Forest (52.14°N, 122.14°W), and Smithers (54.78°N, 127.17°W), British Columbia (BC), Canada (Fig. 1). For the two growth chamber studies on budburst phenology, we collected cuttings from the most southern and northern populations in both our western and eastern transect ($n_{pop}=4$).

Across our eight populations, we measured 47 unique species of which 28 were in our eastern transect and 22 in our western transect (Table S1). At each site, we sampled the dominant deciduous species in both the understory and canopy layer. At our eastern sites, we sampled 13 shrub species and 15 species of tree, while at our western sites we sampled 18 shrub and 4 tree species. Across the two transects, we sampled one shrub and two tree species that occurred in both transects.

Functional traits

We measured all traits in the summer prior to each growth chamber study (eastern transect: 8-25 June 2015, western transect: 29 May to 30 July 2019), following full leafout but before budset. At each population and for each species, we measured a total of five traits from 1-10 healthy adult individuals: height, diameter of the main trunk or stem (hereafter referred to as diameter), wood density, leaf mass per area, and the percent leaf nitrogen content. We also obtained xylem structure data from the WSL xylem database (Schweingruber & Landolt, 2010) for 72.3% of our species.

We measured traits in accordance to the methods discussed by Pérez-Harguindeguy *et al.* (2013). We calculated tree height using trigonometric methods and used a base height of 1.37 m to measure tree diameter (Magarik *et al.*, 2020). For shrub heights, we measured the distance from the ground to the height of the top foliage and measured stem diameters at approximately 1 cm above ground-level. All stem and leaf samples were kept cool during transport and measurements of leaf area and stem volume taken within 3 and 12 hours of sample collection respectively. To measure wood density, we collected a 10 cm sample of branch wood, taken close to the base of the branch at the stem and calculated stem volume using the water displacement method. For our leaf traits, we haphazardly selected and sampled five, fully expanded, and hardened leaves, with no to minimal herbivore damage. We took a high resolution scan of each leaf using a flatbed scanner and estimated leaf area using the ImageJ software (version 2.0.0).

151 Growth chamber study

152 For our growth chamber studies, we collected dormant branch cuttings from our highest and lowest
 153 latitude populations in each transect (total of four sites), with sampling in our eastern study occurring
 154 from 20-28 January 2015 and sampling for our western study from 19-28 October 2019. Dormant
 155 branch cuttings have been repeatedly shown to approximate whole plant responses in budburst (Vi-
 156 tasse & Basler, 2014), allowing us to estimate responses to environmental cues. In both our eastern
 157 and western controlled environment study, we included two temperature treatments and photoperiod,
 158 fully crossed, for a total of eight distinct treatments (Fig. 1). Our treatments included two levels
 159 of chilling—with our eastern study having no additional chilling or 30 days at 4°C, and our western
 160 study 21 days or 56 days of chilling at 4°C, with all non-field chilling occurring under dark conditions.
 161 Our forcing treatments included either a cool regime of 15:5°C or a warm regime of 20:10°C, and our
 162 photoperiod treatments consisted of either a 8 or 12 hour photoperiod. These treatment levels, while
 163 greater than the conditions individuals are likely to experience in the field, are highly comparable to
 164 treatments commonly used in controlled environment experiments and allow us to make similar infer-
 165 ences as studies that apply site-specific conditions (Ettinger *et al.*, 2020, 2021). We recorded budburst
 166 stages of each sample every 1-3 days for up to four months, defining the day of budburst as the day of
 167 budbreak or shoot elongation (denoted as code 07 by Finn *et al.*, 2007). For a more detailed discussion
 168 of study sample collection and methods see Flynn & Wolkovich (2018) for details on our eastern study
 169 and Loughnan & Wolkovich (in review) for details on our western study.

171 Statistical Analysis

172 We combined our *in situ* trait data with budburst data from the controlled environment experiments
 173 through a joint Bayesian model for each trait. This approach improves upon previous analyses of trait
 174 correlations, as it allows us to carry through uncertainty between trait and phenology estimates—and
 175 better partitions the drivers of variation in species phenologies.

176
 177 Our joint model consists of two parts. The first is a hierarchical linear model, which partitions the vari-
 178 ation of individual observations (i) of a given trait value (Y_{trait}) to account for the effects of species
 179 (j), population-level differences arising from transects, latitude, as well as the interaction between
 180 transects and latitude ($\text{transect} \cdot \text{latitude}$), and finally, residual variation or ‘measurement error’ (σ_m^2).

$$Y_{\text{trait}_{i,j}} \sim \text{Normal}(\mu_{i,j}, \sigma_m^2) \quad (1)$$

$$\mu_{i,j} = \alpha_{\text{grand trait}} + \alpha_{\text{trait}_j} + \beta_{\text{transect}} \times \text{transect} + \beta_{\text{latitude}} \times \text{latitude} + \beta_{\text{transect} \cdot \text{latitude}} \times (\text{transect} \cdot \text{latitude}) \quad (2)$$

182 Where α_{trait} is an element of the normal random vector:

$$\alpha_{\text{trait}} \begin{bmatrix} \alpha_{\text{trait}_1} \\ \alpha_{\text{trait}_2} \\ \dots \\ \alpha_{\text{trait}_n} \end{bmatrix} \text{ such that } \alpha_{\text{trait}} \sim \text{Normal}(0, \sigma_{\text{trait}}^2)$$

183 We included transect as a dummy variable (0/1) and latitude as a continuous variable and modeled
 184 traits in their original units, with the exception of leaf mass per area which was rescaled by 100 and
 185 wood density which was rescaled by 10 for numeric stability. Our model also included partial pooling
 186 for species—which controls for uneven sampling and trait variability. These species-level estimates are
 187 then predictors for each cue ($\beta_{\text{chilling},j}$, $\beta_{\text{forcing},j}$, $\beta_{\text{photoperiod},j}$).

188

$$\begin{aligned} \beta_{\text{chilling}_j} &= \alpha_{\text{chilling},j} + \beta_{\text{trait.chilling}} \times \alpha_{\text{trait}_j} \\ \beta_{\text{forcing}_j} &= \alpha_{\text{forcing},j} + \beta_{\text{trait.forcing}} \times \alpha_{\text{trait}_j} \\ \beta_{\text{photoperiod}_j} &= \alpha_{\text{photoperiod},j} + \beta_{\text{trait.photoperiod}} \times \alpha_{\text{trait}_j} \end{aligned} \quad (3)$$

189 In addition to the species-level estimates, this part of our model estimates the overall effect of each
 190 trait on each cue ($\beta_{\text{trait.chilling}}$, $\beta_{\text{trait.forcing}}$, $\beta_{\text{trait.photoperiod}}$). From this we could estimate how well
 191 traits explain species-level differences—by estimating the the species-level cue variation not explained
 192 by traits ($\alpha_{\text{chilling},j}$, $\alpha_{\text{forcing},j}$, $\alpha_{\text{photoperiod},j}$) and individual species responses to cues (*chilling*, *forcing*,
 193 *photoperiod*, respectively). Finally, our model estimates the residual budburst variation across species
 194 (α_{pheno_j}), observations (σ_d^2), and the variation in cues not attributed to the trait using partial pooling,
 195 *with the day of observed budburst ($Y_{\text{pheno}_{i,j}}$) reflecting the additive effects of these sources of variation*
 196 *and the differences in species responses to the experimental treatments.*

$$Y_{\text{pheno}_{i,j}} \sim \text{Normal}(\mu_{i,j}, \sigma_d^2) \quad (4)$$

197 with

$$\mu_{i,j} = \alpha_{\text{pheno}_j} + \beta_{\text{chilling}_j} \cdot \textit{chilling} + \beta_{\text{forcing}_j} \cdot \textit{forcing} + \beta_{\text{photoperiod}_j} \cdot \textit{photoperiod} \quad (5)$$

198 where α_{pheno_j} , $\alpha_{\text{chilling}_j}$, $\alpha_{\text{forcing}_j}$, and $\alpha_{\text{photoperiod}_j}$ are elements of the normal random vectors:

$$\boldsymbol{\alpha}_{\text{pheno}} = \begin{bmatrix} \alpha_{\text{pheno}_1} \\ \alpha_{\text{pheno}_2} \\ \dots \\ \alpha_{\text{pheno}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{pheno}} \sim \text{Normal}(\mu_{\text{pheno}}, \sigma_{\text{pheno}}^2) \quad (6)$$

$$\boldsymbol{\alpha}_{\text{chilling}} = \begin{bmatrix} \alpha_{\text{chilling}_1} \\ \alpha_{\text{chilling}_2} \\ \dots \\ \alpha_{\text{chilling}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{chilling}} \sim \text{Normal}(\mu_{\text{chilling}}, \sigma_{\text{chilling}}^2) \quad (7)$$

$$\boldsymbol{\alpha}_{\text{forcing}} = \begin{bmatrix} \alpha_{\text{forcing}_1} \\ \alpha_{\text{forcing}_2} \\ \dots \\ \alpha_{\text{forcing}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{forcing}} \sim \text{Normal}(\mu_{\text{forcing}}, \sigma_{\text{forcing}}^2) \quad (8)$$

$$\boldsymbol{\alpha}_{\text{photoperiod}} = \begin{bmatrix} \alpha_{\text{photoperiod}_1} \\ \alpha_{\text{photoperiod}_2} \\ \dots \\ \alpha_{\text{photoperiod}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{photoperiod}} \sim \text{Normal}(\mu_{\text{photoperiod}}, \sigma_{\text{photoperiod}}^2) \quad (9)$$

199 We modeled each trait individually, with the exception of ring-porosity, which we compared across
 200 species using the posterior estimates of our wood density model, allowing us to account for inherent
 201 differences in wood anatomy across species and growth form. We included all three cues (chilling,
 202 forcing, and photoperiod) as continuous variables in our model. We converted chilling temperatures
 203 to total chill portions, including both the chilling experienced in the field prior to sampling and during
 204 the experiment. For this we used local weather station data and the chillR package (v. 0.73.1, Luedel-
 205 ing, 2020). To account for differences in thermoperiodicity between our eastern and western growth
 206 chamber studies (Buonaiuto *et al.*, 2023), we also converted forcing temperatures to mean daily tem-
 207 peratures for each treatment. Finally, we *z*-scored each cue and site using two standard deviations to
 208 allow direct comparisons between results across parameters (Gelman, 2008).

209

210 We used trait-specific priors that were weakly informative. We validated our choice of priors using
 211 prior predictive checks and confirmed model stability under wider priors. All models were coded
 212 in the Stan programming language for Bayesian models using the rstan package (Stan Development
 213 Team, 2018) in R version 4.3.1 (R Development Core Team, 2017). All models met basic diagnostic
 214 checks, including no divergences, high effective sample sizes (n_{eff}) that exceeded 10% of the number of
 215 iterations, and $\hat{R}$ values close to 1. We report our model estimates as the mean values with 90% uncer-
 216 tainty intervals (UI), interpreting parameter estimates with intervals that include zero to have no effect.

217

Results

Across our eight populations, we measured traits of 1428 unique individuals between the two transects across our five *in situ* traits: height ($n = 1317$), diameter ($n = 1220$), wood density ($n = 1359$), leaf mass per area ($n = 1345$), leaf nitrogen content ($n = 1351$). Between our two growth chamber studies, we made observations of 4211 branch cuttings, with our observations of budburst spanning 82 and 113 days for our eastern and western studies respectfully.

We found that four of our five measured traits had a strong relationship with photoperiod, but not always in the direction we predicted. Taller species with larger trunk diameters and leaves with high nitrogen content did have larger responses with longer photoperiods (Fig. 3 c, i, o; Tables S3, S4, S7), as we predicted. But, contrary to our expectation, species with denser, high leaf mass per area leaves had smaller photoperiod responses, allowing them to potentially budburst earlier, under shorter photoperiods (Fig. 3f, Table S6).

Temperature cues ($\beta_{\text{trait.chilling}}$ and $\beta_{\text{trait.forcing}}$) exhibited no relationships with individual traits, but we found the individual importance of our three budburst cues (chilling, forcing and photoperiod) differed once the effects of leaf or wood traits had been accounted for. Chilling (β_{chilling}) had the strongest effect in our models of height (-13.4 days per standardized chill portions, UI: -17.2, -9.9), diameter (-12.5 days per standardized chill portions, UI: -16.2, -8.6), wood density (-20.9 days per standardized chill portions, UI: -33.2, -9.8), and leaf nitrogen content (-35.1 days per standardized chill portions, UI: -68.1, -4.1), with more chilling advancing budburst. Our model of leaf mass per area estimated photoperiod ($\beta_{\text{photoperiod}}$) as the strongest cue (-14.0 days per standardized photoperiod, UI: -23.1, -3.5), but only two of our trait models—accounting for the effects of height and diameter—estimated strong responses in budburst timing to all three cues (Tables S3, S4). But in our models accounting for the effects of wood density and leaf nitrogen content, only temperature cues (chilling and forcing) shaped budburst timing (Tables S5, S7), and in our model of leaf mass per area, budburst was estimated to respond only to photoperiod (Table S6).

Of our five trait models, we found clear gradients in species timing between trees and shrubs in our models of height and diameter. Species height exhibited large correlations between budburst timing and trait values, with earlier estimates of budburst for shrubs (with a mean day of budburst of 10)—especially under greater cues—and later budburst estimates for trees (with a mean day of budburst of 17.3, Fig. S1). But this was not the case for the leaf traits. Leaf nitrogen content, in particular, showed no distinct separation between shrub and tree functional groups (Fig. S1). We also found no differences between xylem structure and their responses to cues (Fig. 4).

Most of our traits varied across latitude within each transect and had strong interactive effects between transect and latitude (Fig. 2). Leaf nitrogen content varied with latitude alone, with low latitude sites on both our eastern and western transects having more nitrogen rich leaves than sites at higher latitudes (-0.1 percent per degree latitude, UI: -0.2, 0.0, Table S7). Plants also had wider trunk or stem diameters at higher latitudes in both eastern and western sites (0.5 cm per degree latitude, UI: 0.1,

0.9), with a larger effect in eastern sites (Fig. 2d). Relationships with height, woody density and leaf mass per area were, however, more complex. Plants heights and wood density increased with increasing latitude in western sites, but decreased with latitude in eastern sites (-0.2 m per degree latitude, UI: -0.4, 0.0 for our height model and -0.01 g/cm³ per degree latitude, UI: -0.02, 0.0 for our wood density model; Fig. 2 a and c). For leaf mass per area we found the opposite relationship (decreasing leaf mass per area with increasing latitudes in western sites, while it increased with latitude in our eastern sites (0.005 g/cm² per degree latitude, UI: 0.004, 0.006; Fig. 2d). In addition to the differences we found across sites, we also observed large differences between individual species, which varied up to 7 fold for some traits (Fig. 3).

Discussion

Using a joint modeling approach to understand how phenology may fit within acquisitive to conservative plant strategies, we found that photoperiod was the only cue related to our suite of leaf and wood traits. While budburst responses to photoperiod are often smaller than temperature responses (chilling and forcing; Laube *et al.*, 2014; Zohner *et al.*, 2016; Flynn & Wolkovich, 2018), our results suggest photoperiod may be the most important cue in linking spring phenology to functional traits. To accurately predict these responses likely requires considering additional traits and variation across space, as we observed differences in the relative importance of cues across our trait models and between our eastern and western transects with latitude. The observed spatial differences in trait variation may be due to differences in the community assemblages, as our western sites were more shrub dominated, with shorter plants with less dense branch wood, suggesting a more acquisitive growth strategy to allow species to utilize resources early in the season before canopy closure.

Cues and functional traits

We found only partial support for our prediction that species with acquisitive traits—particularly small trees with low wood density, low leaf mass per area, and high leaf nitrogen content—would have smaller temperature and photoperiod responses (associated with early budburst). Species with smaller heights and diameters did have smaller photoperiod responses, but—contrary to our prediction—species with less dense leaves showed larger responses to photoperiod, while leaves with high nitrogen content had stronger photoperiod responses.

None of our focal traits showed a relationship with temperature (chilling or forcing), which may be due to selection on other physiological processes. Many of our traits are associated with one or more ecological function (Wright *et al.*, 2004; Pérez-Harguindeguy *et al.*, 2013; Reich, 2014). In particular, leaf mass per area is known to correlate with traits like leaf lifespan or decomposition rates in addition to light capture (De La Riva *et al.*, 2016). In accounting for the relationships between other traits and budburst cues, we also found the relative importance of temperature and photoperiod on budburst timing varied, suggesting other traits may play a role in shaping phenologies.

Decades of previous phenology research have found budburst timing to be primarily driven by temperature (chilling and forcing) and weakly by photoperiod (Chuine *et al.*, 2010; Basler & Körner, 2014; Laube *et al.*, 2014). Yet we found no traits that correlated with responses to temperature, suggesting other drivers may impact leaf and structural traits in temperate forests. One potential abiotic driver we did not consider is soil moisture, which covaries with a number of traits, including leaf mass per area, as higher leaf area allows plants to reduce evaporation under dry conditions (De La Riva *et al.*, 2016). Soil moisture also shapes other phenological events in woody plants, including radial growth phenology and shoot elongation (Cabon *et al.*, 2020; Peters *et al.*, 2021). Though temperate forests are generally moist compared to other systems, previous studies suggests soil moisture can shape spring phenology across xeric systems as well (Crimmins *et al.*, 2011; Park, 2014; Ettinger *et al.*, 2019), thus including soil moisture in future studies could help understand how species growth strategies correlate with phenology.

Our finding that budburst was not related to wood structure or wood density contrasts the findings of previous work linking these traits. Previous studies have found some evidence that trees with diffuse-porous wood structure leafout earlier than species with ring-porous structures (Lechowicz, 1984; Panchen *et al.*, 2014; Yin *et al.*, 2016; Osada, 2017; Savage *et al.*, 2022). Using wood density as an alternative measure of wood structure (wood density positively correlates with xylem resistance to embolism, Hacke *et al.*, 2001), we did not find an association between our three phenological cues and xylem structure, despite our data including similar temperate forest species to previous studies. This could be because we captured additional variation by sampling many individuals for each species across a large spatial gradient. In particular, we found larger wood densities at higher latitudes in our western transect, which could be caused by the differences in winter conditions experienced across this transect. Higher wood densities are especially favorable in communities that experience greater horizontal stress from wind and downward pressure from snow (MacFarlane & Kane, 2017; MacFarlane, 2020). Relationships with wood density may also be more apparent over a larger span of values, as our observations varied from 0.2 to 0.6 g/cm³, which is smaller than seen in studies that span a more global distribution (Galvão *et al.*, 2021; Savage *et al.*, 2022; Mo *et al.*, 2024).

In addition to our study providing insight into how trait-budburst relationships vary with latitude, our sites also spanned North America, encompassing a gradient of 55° in longitude. At this continental scale we found correlations between phenology and traits for woody species in temperate forests, but these may shift with data that expands across biomes and more plant functional groups. Indeed, both the leaf and wood economic spectra find that large variation in traits across species arises from large global datasets and include both temperate and tropical species (Wright *et al.*, 2004; Díaz *et al.*, 2016; Chave *et al.*, 2009) and functional groups that span from trees to grasses (Wright *et al.*, 2004). Global patterns in trait variation may also be absent at smaller spatial scales (Wright & Sutton-Grier, 2012; Messier *et al.*, 2017a,b), suggesting that comparisons of trait-phenology relationships at larger spatial scales or across more diverse pools of species may better align with the patterns predicted by existing economic spectra.

In comparing spring phenology of mainly temperate woody species, our results support previous work through which a global meta-analysis found that spring phenology may correlate with height and leaf mass per area (Loughnan *et al.*, 2025). At both the global and continental scales, trees with taller heights leafed out with longer photoperiods. We also found species with high specific leaf area—which is the inverse of leaf mass per area and thus equivalent to low values—exhibited large responses to photoperiod (Loughnan *et al.*, 2025). The consistency of these results, despite the differences in the two spatial scales of these datasets, provides further support that woody species responses to photoperiod may be part of a larger plant strategy.

Functional traits predict climate change responses

Our results offer novel insights into how broader correlations between growth strategies and phenological cues can help predict responses in woody plant communities with climate change. As temperatures rise, particularly at higher latitudes (Hoegh-Guldberg *et al.*, 2018), warmer winter and spring temperatures may select for earlier budburst in some species. But, since photoperiod will remain fixed, our observed relationships between photoperiod and other traits has the potential to limit how well species track temperatures. Photoperiod cues could further constrain the extent to which some species growth will advance with climate change (Guy, 2014). Our results suggest that these effects will likely be greater for taller species or canopy trees and species with relatively low leaf mass per area. These constraints could have cascading effects on forest communities, as variable species responses to increasing temperatures further alter species growth strategies and their interactions with competitors or herbivores within their communities.

Our findings of correlations between phenology and other commonly measured traits highlight how accurate forecasts of future changes in phenology could benefit from accounting for the response of other traits to climate change. Across temperature and precipitation gradients, leaf size and shape also change, as species shift to conserve water and mitigate the effects of transpiration under higher temperatures (De La Riva *et al.*, 2016). These changes could impact species photosynthetic potential and ultimately ecosystem services, such as carbon sequestration. While phenological research has focused on forecasting responses to temperature, the correlation of other traits with photoperiod suggests it is also an important cue. Considering the tradeoffs and differences in cues that together shape plants growth strategies may yield more accurate forecasts of species phenology and community dynamics under future climates.

Acknowledgements

We thank M. Yasutake, K. Slimon, S. Larter, P. Autio, and S. Collins for their help running the Western part of this experiment; D. Flynn and T. Savas for contributions to the Eastern experimental data; and to J. Hopps and K. Wilson for their support during our work at E. C. Manning Park and P. Crawford and D. Belford for their support while working in Babine Provincial Park. Funding was provided by a

376 Natural Sciences and Engineering Research Council (NSERC) Canada Graduate Scholarship-Doctoral
377 award to Deirdre Loughnan and Canada Research Chair award in Temporal Ecology and NSERC
378 Discovery awards to E.M. Wolkovich.
379

380 Competing interests

381 The authors have no competing interests.
382

383 Author contributions

384 Both D.L. and E.M.W. conceived the study, D.L., F.A.J. and E.M.W. all contributed to the analysis
385 and code and contributed to the writing and revision of the manuscript. DL led field and experimental
386 data collection.
387

388 Data availability

389 The phenology data used in this study is published on the Knowledge Network of Biocomplexity
390 for both the eastern transect (urn:uuid:efeccd2b-b850-4b48-b721-9bed8040562e) and western transect
391 (doi:10.5063/F1W66J82). Trait data from the eastern transect is available through the Harvard Forest
392 Data Archive (doi:10.6073/pasta/dc68c63764f0c5edf6f09e7a60a0468a). Trait data from the western
393 transect will be published on the Knowledge Network of Biocomplexity.

394 References

- 395 Augspurger, C.K. (2008). Early spring leaf out enhances growth and survival of saplings in a temperate
396 deciduous forest. *Oecologia*, 156, 281–286.
- 397 Augspurger, C.K. (2009). Spring 2007 warmth and frost: phenology, damage and refoliation in a
398 temperate deciduous forest. *Functional Ecology*, 23, 1031–1039.
- 399 Basler, D. & Körner, C. (2014). Photoperiod and temperature responses of bud swelling and bud burst
400 in four temperate forest tree species. *Tree Physiology*, 34, 377–388.
- 401 Buonaiuto, D.M., Wolkovich, E.M. & Donahue, M.J. (2023). Experimental designs for testing the
402 interactive effects of temperature and light in ecology: The problem of periodicity. *Functional*
403 *Ecology*, 37, 1747–1756.
- 404 Cabon, A., Fernández-de-Uña, L., Gea-Izquierdo, G., Meinzer, F.C., Woodruff, D.R., Martínez-Vilalta,
405 J. & De Cáceres, M. (2020). Water potential control of turgor-driven tracheid enlargement in Scots
406 pine at its xeric distribution edge. *New Phytologist*, 225, 209–221.

Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A.E. (2009). Towards a worldwide wood economics spectrum. *Ecology Letters*, 12, 351–366.

Chuine, I. (2000). A unified model for budburst of trees. *Journal of Theoretical Biology*, 207, 337–347.

Chuine, I. & Cour, P. (1999). Climatic determinants of budburst seasonality in four temperate-zone tree species. *New Phytologist*, 143, 339–349.

Chuine, I., Morin, X. & Bugmann, H. (2010). Warming, photoperiods, and tree phenology. *Science*, 329, 277–278.

Cohen, J.M., Lajeunesse, M.J. & Rohr, J.R. (2018). A global synthesis of animal phenological responses to climate change. *Nature Climate Change*, 8, 224–228.

Crimmins, T.M., Crimmins, M.A. & Bertelsen, C.D. (2011). Onset of summer flowering in a ‘Sky Island’ is driven by monsoon moisture. *New Phytologist*, 191, 468–479.

De La Riva, E.G., Olmo, M., Poorter, H., Ubera, J.L. & Villar, R. (2016). Leaf Mass per Area (LMA) and Its Relationship with Leaf Structure and Anatomy in 34 Mediterranean Woody Species along a Water Availability Gradient. *PLOS ONE*, 11, e0148788.

Díaz, S., Kattge, J., Cornelissen, J.H., Wright, I.J., Lavorel, S., Dray, S., Reu, B., Kleyer, M., Wirth, C., Colin Prentice, I., Garnier, E., Bönsch, G., Westoby, M., Poorter, H., Reich, P.B., Moles, A.T., Dickie, J., Gillison, A.N., Zanne, A.E., Chave, J., Joseph Wright, S., Sheremet Ev, S.N., Jactel, H., Baraloto, C., Cerabolini, B., Pierce, S., Shipley, B., Kirkup, D., Casanoves, F., Joswig, J.S., Günther, A., Falczuk, V., Rüger, N., Mahecha, M.D. & Gorné, L.D. (2016). The global spectrum of plant form and function. *Nature*, 529, 167–171.

Ettinger, A.K., Buonaiuto, D.M., Chamberlain, C.J., Morales-Castilla, I. & Wolkovich, E.M. (2021). Spatial and temporal shifts in photoperiod with climate change. *New Phytologist*, 230, 462–474.

Ettinger, A.K., Chamberlain, C.J., Morales-Castilla, I., Buonaiuto, D.M., Flynn, D.F.B., Savas, T., Samaha, J.A. & Wolkovich, E.M. (2020). Winter temperatures predominate in spring phenological responses to warming. *Nature Climate Change*, 10, 1137–1142.

Ettinger, A.K., Chuine, I., Cook, B.I., Dukes, J.S., Ellison, A.M., Johnston, M.R., Panetta, A.M., Rollinson, C.R., Vitasse, Y. & Wolkovich, E.M. (2019). How do climate change experiments alter plot-scale climate? *Ecology Letters*, 22, 748–763.

Finn, G.A., Straszewski, A.E. & Peterson, V. (2007). A general growth stage key for describing trees and woody plants. *Annals of Applied Biology*, 151, 127–131.

Flynn, D.F.B. & Wolkovich, E.M. (2018). Temperature and photoperiod drive spring phenology across all species in a temperate forest community. *New Phytologist*, 219, 1353–1362.

Galvão, F.G., Alves De Lima, A.L., Candeia De Oliveira, C., Da Silva, V.F. & Rodal, M.J.N. (2021). The importance of wood density in determining the phenology of tree species in a coastal rain forest. *Biotropica*, 53, 1134–1141.

442 Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics in*
443 *Medicine*, 27, 2865–2873.

444 Gotelli, N.J. & Graves, G.R. (1996). The temporal niche. In: *Null Models In Ecology*. Smithsonian
445 Institution Press, Washington, D. C., pp. 95–112.

446 Green, S.J., Brookson, C.B., Hardy, N.A. & Crowder, L.B. (2022). Trait-based approaches to global
447 change ecology: moving from description to prediction. *Proceedings of the Royal Society B: Biological*
448 *Sciences*, 289, 1–10.

449 Guy, R.D. (2014). The early bud gets to warm. *New Phytologist*, 202, 7–9.

450 Hacke, U.G., Sperry, J.S., Pockman, W.T., Davis, S.D. & McCulloh, K.A. (2001). Trends in wood
451 density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia*,
452 126, 457–461.

453 Harrington, C.A. & Gould, P.J. (2015). Tradeoffs between chilling and forcing in satisfying dormancy
454 requirements for Pacific Northwest tree species. *Frontiers in Plant Science*, 6, 1–12.

455 Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante,
456 R., Ebi, K., Engelbrecht, F., Guiot, J., Hijioka, Y., Mehrotra, S., Payne, A., Seneviratne, S., Thomas,
457 A., Warren, R. & Zhou, G. (2018). Impacts of 1.5 °C Global Warming on Natural and Human
458 Systems. In: *Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming*
459 *of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the*
460 *context of . Tech. rep., Cambridge University Press, Cambridge, UK and New York, NY, USA.*

461 Kharouba, H.M., Ehrlén, J., Gelman, A., Bolmgren, K., Allen, J.M., Travers, S.E. & Wolkovich,
462 E.M. (2018). Global shifts in the phenological synchrony of species interactions over recent decades.
463 *Proceedings of the National Academy of Sciences*, 115, 5211–5216.

464 Körner, C. & Basler, D. (2010). Phenology Under Global Warming. *Science*, 327, 1461–1463.

465 Laube, J., Sparks, T.H., Estrella, N., Höfler, J., Ankerst, D.P. & Menzel, A. (2014). Chilling outweighs
466 photoperiod in preventing precocious spring development. *Global Change Biology*, 20, 170–182.

467 Lechowicz, M.J. (1984). Why Do Temperate Deciduous Trees Leaf Out at Different Times? Adaptation
468 and Ecology of Forest Communities. *The American Naturalist*, 124, 821–842.

469 Lopez, O.R., Farris-Lopez, K., Montgomery, R.A. & Givnish, T.J. (2008). Leaf phenology in relation
470 to canopy closure in southern Appalachian trees. *American Journal of Botany*, 95, 1395–1407.

471 Loughnan, D., Jones, F.A., Legault, G., Chamberlain, C.J., Buonaiuto, D.M., Ettinger, A.K., Garner,
472 M., Sodhi, D.S. & Wolkovich, E.M. (2025). Budburst timing within a functional trait framework.
473 *Journal of Ecology*, 113, 3787–3798.

474 Loughnan, D. & Wolkovich, E.M. (in review). Temporal assembly of woody plant communities shaped
475 equally by evolutionary history as by current environments.

- 476 Luedeling, E. (2020). *chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees*.
477 <https://CRAN.R-project.org/package=chillR>.
- 478 MacFarlane, D.W. (2020). Functional Relationships Between Branch and Stem Wood Density for
479 Temperate Tree Species in North America. *Frontiers in Forests and Global Change*, 3, 1–16.
- 480 MacFarlane, D.W. & Kane, B. (2017). Neighbour effects on tree architecture: functional trade-offs
481 balancing crown competitiveness with wind resistance. *Functional Ecology*, 31, 1624–1636.
- 482 Magarik, Y.A., Roman, L.A. & Henning, J.G. (2020). How should we measure the dbh of multi-
483 stemmed urban trees? *Urban Forestry & Urban Greening*, 47, 1–11.
- 484 Messier, J., Lechowicz, M.J., McGill, B.J., Violle, C. & Enquist, B.J. (2017a). Interspecific integration
485 of trait dimensions at local scales: the plant phenotype as an integrated network. *Journal of Ecology*,
486 105, 1775–1790.
- 487 Messier, J., McGill, B.J., Enquist, B.J. & Lechowicz, M.J. (2017b). Trait variation and integration
488 across scales: is the leaf economic spectrum present at local scales? *Ecography*, 40, 685–697.
- 489 Mo, L., Crowther, T.W., Maynard, D.S. & Van Den Hoogen, e.a. (2024). The global distribution and
490 drivers of wood density and their impact on forest carbon stocks. *Nature Ecology & Evolution*, 8,
491 2195–2212.
- 492 Osada, N. (2017). Relationships between the timing of budburst, plant traits, and distribution of 24
493 coexisting woody species in a warm-temperate forest in Japan. *American Journal of Botany*, 104,
494 550–558.
- 495 Panchen, Z.A., Primack, R.B., Nordt, B., Ellwood, E.R., Stevens, A., Renner, S.S., Willis, C.G.,
496 Fahey, R., Whittemore, A., Du, Y. & Davis, C.C. (2014). Leaf out times of temperate woody plants
497 are related to phylogeny, deciduousness, growth habit and wood anatomy. *New Phytologist*, 203,
498 1208–1219.
- 499 Park, I.W. (2014). Impacts of differing community composition on flowering phenology throughout
500 warm temperate, cool temperate and xeric environments. *Global Ecology and Biogeography*, 23,
501 789–801.
- 502 Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across
503 natural systems. *Nature*, 421, 37–42.
- 504 Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte,
505 M.S., Cornwell, W.K., Craine, J.M., Gurvich, D.E., Urcelay, C., Veneklaas, E.J., Reich, P.B.,
506 Poorter, L., Wright, I.J., Ray, P., Enrico, L., Pausas, J.G., de Vos, A.C., Buchmann, N., Funes,
507 G., Quétier, F., Hodgson, J.G., Thompson, K., Morgan, H.D., ter Steege, H., van der Heijden,
508 M.G.A., Sack, L., Blonder, B., Poschlod, P., Vaieretti, M.V., Conti, G., Staver, A.C., Aquino, S. &
509 Cornelissen, J.H.C. (2013). New handbook for standardized measurement of plant functional traits
510 worldwide. *Australian Journal of Botany*, 61, 167–234.

511 Peters, R.L., Steppe, K., Cuny, H.E., De Pauw, D.J., Frank, D.C., Schaub, M., Rathgeber, C.B.,
512 Cabon, A. & Fonti, P. (2021). Turgor – a limiting factor for radial growth in mature conifers along
513 an elevational gradient. *New Phytologist*, 229, 213–229.

514 Primack, R.B., Ibáñez, I., Higuchi, H., Lee, S.D., Miller-Rushing, A.J., Wilson, A.M. & Silander,
515 J.A. (2009). Spatial and interspecific variability in phenological responses to warming temperatures.
516 *Biological Conservation*, 142, 2569–2577.

517 R Development Core Team (2017). R: A language and environment for statistical computing.

518 Reich, P.B. (2014). The world-wide ‘fast–slow’ plant economics spectrum: a traits manifesto. *Journal*
519 *of Ecology*, 102, 275–301.

520 Sakai, A. & Larcher, W. (1987). *Frost Survival of Plants: Responses and adaptation to freezing stress*.
521 Springer-Verlag, Berlin, Heidelberg.

522 Savage, J.A., Kiecker, T., McMann, N., Park, D., Rothendler, M. & Mosher, K. (2022). Leaf out time
523 correlates with wood anatomy across large geographic scales and within local communities. *New*
524 *Phytologist*, 235, 953–964.

525 Schweingruber, F. & Landolt, W. (2010). The xylem database.

526 Stan Development Team (2018). RStan: the R interface to Stan. R package version 2.17.3.

527 Thackeray, S.J., Henrys, P.A., Hemming, D., Bell, J.R., Botham, M.S., Burthe, S., Helaouet, P.,
528 Johns, D.G., Jones, I.D., Leech, D.I., MacKay, E.B., Massimino, D., Atkinson, S., Bacon, P.J.,
529 Brereton, T.M., Carvalho, L., Clutton-Brock, T.H., Duck, C., Edwards, M., Elliott, J.M., Hall, S.J.,
530 Harrington, R., Pearce-Higgins, J.W., Høye, T.T., Kruuk, L.E., Pemberton, J.M., Sparks, T.H.,
531 Thompson, P.M., White, I., Winfield, I.J. & Wanless, S. (2016). Phenological sensitivity to climate
532 across taxa and trophic levels. *Nature*, 535, 241–245.

533 Violle, C., Navas, M., Vile, D., Kazakou, E., Fortunel, C., Hummel, I. & Garnier, E. (2007). Let the
534 concept of trait be functional! *Oikos*, 116, 882–892.

535 Vitasse, Y. & Basler, D. (2014). Is the use of cuttings a good proxy to explore phenological responses
536 of temperate forests in warming and photoperiod experiments? *Tree Physiology*, 34, 174–183.

537 Vitasse, Y., Hoch, G., Randin, C.F., Lenz, A., Kollas, C., Scheepens, J.F. & Körner, C. (2013).
538 Elevational adaptation and plasticity in seedling phenology of temperate deciduous tree species.
539 *Oecologia*, 171, 663–678.

540 Vitasse, Y., Josée, A., Kremer, A., Michalet, R. & Delzon, S. (2009). Responses of canopy duration
541 to temperature changes in four temperate tree species : relative contributions of spring and autumn
542 leaf phenology. *Oecologia*, 161, 187–198.

543 Vitasse, Y., Signarbieux, C. & Fu, Y.H. (2018). Global warming leads to more uniform spring phenology
544 across elevations. *PNAS*, 115, 1004–1008.

- 545 Wang, N., Wu, Z., Gong, Y., Cui, Y. & Fu, Y.H. (2025). Background climate determines the response
546 of spring leaf-out to climate change—Results from a national-scale twig-cutting experiment. *Journal*
547 *of Ecology*, 113, 2981–2991.
- 548 Wolkovich, E.M. & Ettinger, A.K. (2014). Back to the future for plant phenology research. *New*
549 *Phytologist*, 203, 1021–1024.
- 550 Wright, I.J., Westoby, M., Reich, P.B., Oleksyn, J., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-
551 Bares, J., Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Gullas, J., Garnier, E., Navas,
552 M.L., Roumet, C., Groom, P.K., Lamont, B.B., Hikosaka, K., Lee, T., Lee, W., Lusk, C., Midgley,
553 J.J., Niinemets, Ü., Osada, H., Poorter, H., Pool, P., Veneklaas, E.J., Prior, L., Pyankov, V.I.,
554 Thomas, S.C., Tjoelker, M.G. & Villar, R. (2004). The worldwide leaf economics spectrum. *Nature*,
555 428, 821–827.
- 556 Wright, J.P. & Sutton-Grier, A. (2012). Does the leaf economic spectrum hold within local species
557 pools across varying environmental conditions? *Functional Ecology*, 26, 1390–1398.
- 558 Yin, J., Fridley, J.D., Smith, M.S. & Bauerle, T.L. (2016). Xylem vessel traits predict the leaf phenology
559 of native and non-native understorey species of temperate deciduous forests. *Functional Ecology*, 30,
560 206–214.
- 561 Zohner, C.M., Benito, B.M., Svenning, J.C. & Renner, S.S. (2016). Day length unlikely to constrain
562 climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate Change*, 6, 1120–
563 1123.




Figure 1: We collected trait data and branch cuttings from woody plants growing within eight sites, across two transects in eastern and western North America (map at left). Cuttings were used in two controlled environment (growth chamber) studies in which we applied high and low chilling, forcing, and photoperiod treatments and recorded the day of budburst. Using our paired *in situ* trait and experimental budburst data, we tested whether earlier budbursting species exhibited traits associated with more acquisitive growth strategies and smaller responses to cues and later budbursting species a more conservative growth strategy and larger responses to cues (see ‘Predictions’ at right).




Figure 2: We found (a) height, (b) diameter, (c) branch wood density, and (d) leaf mass per area to all experience a strong interaction between latitude and transect, (e) while leaf nitrogen content showed a strong effect of latitude alone.




Figure 3: Relationships between species traits (rows) and cue responses (columns) showed considerable variation across: (a-c) height, (d-f) leaf mass per area, (g-i) diameter, (j-l) wood density, and (m-o) leaf nitrogen content. Point colours represent different species groups, with tree species shown in red and shrub species in blue. Crosses depict the 50% uncertainty interval of the model estimates of species trait values and estimated responses to cues. Grey bands depict large relationships between a trait and cue, representing the 90% uncertainty interval, and black lines the mean response.




Figure 4: Despite previous studies finding relationships between leaf out timing and wood xylem structures, we did not find clear differences in species-level estimates of cue responses with wood structure or relative to their wood densities. Each cross represents the 50% uncertainty interval of (a). forcing, (b) chilling, and (c) photoperiod responses and wood density for each species, with colors depicting different types of wood structure. The lowest wood density was estimated for *Sambucus racemosa* and the highest wood density for *Viburnum lantanoides*.