Traits predict forest phenological responses to photoperiod more than temperature

Deirdre Loughnan[1], Faith A M Jones[1,2], and E M Wolkovich[1,3,4]

April 20, 2026

[1] Department of Forest and Conservation, Faculty of Forestry, University of British Columbia, 2424 Main Mall Vancouver, BC Canada V6T 1Z4.

[2] Department of Wildlife, Fish and Environmental Studies, Swedish University of Agricultural Sciences, 901 83 Umeå, Sweden.

[3] Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

[4] Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA;

Corresponding Author: Deirdre Loughnan deirdre.loughnan@ubc.ca
Running title: Traits and budburst both related photoperiod cues

Summary

Shifts in the timing of spring phenology, such as budburst and leafout, can have major ecosystem consequences. Understanding how species phenological responses fit within broader frameworks of plant strategies could help predict these consequences and other impacts on plant communities. Theory suggests that species with early spring phenology would have a more acquisitive growth strategy while later species would be more conservative. Yet testing these predictions has been slow given the high variability of spring phenology when measured in natural settings across space and time. Using controlled environment studies, we reduce phenological variation into its component responses to chilling (cool winter temperatures), forcing (spring warming temperatures) and photoperiod, and test for trait relationships across 47 species sampled from eight forest communities (1428 individuals) across North America. We find phenology connects to four major plant functional traits—height, diameter, leaf mass per area and nitrogen content—via responses to photoperiod, but not temperature. Our results suggest photoperiod responses may be a critical component of how spring phenology fits within plant strategies, which could help predict future shifts in forest growth with continued climate change.

Introduction

Climate change is causing species phenologies—the timing of life history events—to shift, with widespread advances being observed across the tree of life (Parmesan and Yohe, 2003; Hoegh-Guldberg et al., 2018). This common phenological fingerprint, however, averages over high variability across species (Thackeray et al., 2016; Cohen et al., 2018; Kharouba et al., 2018), posing a challenge to accurate forecasts.

In plants, species variation can be explained, in part, by differences in growth strategies, which are generally inferred from traits (Violle et al., 2007). Decades of research on plant traits have worked to build predictive models of species responses to their environment (Green et al., 2022), which could explain species-level variability in phenological responses. Phenology, however, has generally be excluded from plant trait research due to its high variability, even for the same species when observed over different years or sites (Primack et al., 2009; Chuine et al., 2010). Within species variation also occurs across other plant traits (e.g. leaf and wood structure traits), but generally to a much smaller scale compared to phenology (Violle et al., 2007). Yet increasing evidence that spring phenology is much less variable when carefully measured (Zeng and Wolkovich, 2024; Ettinger et al., 2020) holds promise to leverage existing frameworks to explain phenological variation and predict future changes.

Correlations between plant traits, growth strategies, and responses to environments have been synthesized into several global frameworks, including the leaf economic spectrum (Wright et al., 2004) and wood economic spectrum (Chave et al., 2009). These frameworks have identified key traits that exhibit distinct gradients, ranging from acquisitive strategies—fast growing plants that produce cheaper tissue —to conservative strategies—with plants that invest in long-lived tissue but slower growth rates (Wright et al., 2004; Díaz et al., 2016). In temperate systems, changes in temperature in spring can produce gradients in abiotic stress, including frost risk, soil nutrients, and light availability (Sakai and Larcher, 1987; Gotelli and Graves, 1996; Augspurger, 2009), in addition to differences in biotic interactions from herbivory or competition later in the season (Lopez et al., 2008; Wolkovich and Ettinger, 2014). Species that vary in their timing of leafout should therefore exhibit traits and growth strategies that allow them to tolerate or avoid these abiotic and biotic factors, but testing how phenology fits within major functional trait frameworks requires working across within- and between-species variation.

Phenological variation is generally observed in natural conditions where the environmental cues that trigger many phenological events—primarily temperature and photoperiod (Chuine, 2000; Körner and Basler, 2010)—vary across space and time, generating high variation that has stymied previous efforts to understand the role of phenology in existing trait frameworks. Increasing evidence, however, shows that phenology—especially spring phenology—is often far less variable when measured in experiments (Zeng and Wolkovich, 2024; Basler and Körner, 2014; Vitasse et al., 2009). Experiments in controlled environments (growth chambers) in particular can consistently decompose spring phenology for woody plants into its individual environmental cues (Chuine and Cour, 1999; Harrington and Gould, 2015; Flynn and Wolkovich, 2018): chilling (cool winter temperatures), forcing (warm spring temperatures), and photoperiod. Decades of experiments now show how these cues combine to determine early versus late spring phenology across species. While all species show strong responses to temperature cues, with chilling generally the strongest cue, later species tend to have larger responses to forcing, chilling and photoperiod (Ettinger et al., 2020). Within species, some evidence suggests spatial gradients in climate may drive variation in response to cues (Menzel et al., 2006; Vitasse et al., 2018; Wang et al., 2025).

Here, we tested whether phenological variation was aligned with existing trait frameworks using data on spring budburst paired with a suite of traits that capture acquisitive to conservative growth strategies. We decomposed the high phenological variation in budburst date by using experiments to estimate chilling, forcing, and photoperiod cues. We predicted that early spring species, which generally budburst before canopy closure (Augspurger, 2008; Vitasse et al., 2013), will have smaller responses to temperature and photoperiod and have traits associated with acquisitive growth. They would thus be shorter, with smaller trunks or stem diameters, and a lower investment in wood structure and leaf tissue, resulting in low wood specific density, diffuse-porous wood anatomy, and low leaf mass per area, but high leaf nitrogen content for a greater photosynthetic potential. In contrast, we predicted species with later budburst (requiring more chilling, forcing, and longer photoperiods) to predominately include canopy species that have more conservative growth strategies. These species should incur greater investments in long-lived tissue, with ring-porous wood anatomy, taller heights and greater diameter,

denser wood and high leaf mass per area, but low leaf nitrogen content. We further expected traits to vary spatially, with individuals at more northern sites being relatively smaller, with lower leaf mass per area relative to our southern sites, but follow similar responses in both our eastern and western transect. We used a joint-modeling approach to estimate the relationships between these plant traits and phenological responses to cues, while partitioning the variance from species- and population-level differences.

Materials and Methods

Field sampling

We combined *in situ* trait data with budburst data from two growth chamber cutting experiments conducted across eastern and western temperate deciduous forests in North America. We collected both suites of data from populations that span a latitudinal gradient of 4-6° for the eastern and western communities respectively. We took trait measurements from across eight populations, of which there were four eastern populations—Harvard Forest, Massachusetts, USA (42.55°N, 72.20°W), White Mountains, New Hampshire, USA (44.11°N, 52.14°W), Second College Grant, New Hampshire, USA (44.79°N, 50.66°W), and St. Hippolyte, Quebec, Canada (45.98°N, 74.01°W), and four western populations—E.C. Manning Park (49.06°N, 120.78°W), Sun Peaks (50.88°N, 119.89°W), Alex Fraser Research Forest (52.14°N, 122.14°W), and Smithers (54.78°N, 127.17°W), British Columbia (BC), Canada (Fig. 1). For the two growth chamber studies on budburst phenology, we collected cuttings from the most southern and northern populations in both our western and eastern transect (n_{pop}=4).

Functional traits

We measured all traits in the summer prior to each growth chamber study (eastern transect: 8-25 June 2015, western transect: 29 May to 30 July 2019), following full leafout but before budset. At each population and for each species, we measured a total of five traits from 1-10 healthy adult individuals: height, diameter of the main trunk or stem (hereafter referred to as diameter), wood specific density, leaf mass per area, and the percent leaf nitrogen content. We also obtained xylem structure data from the WSL xylem database (Schweingruber and Landolt, 2010) for 72.3% of our species.

We measured traits in accordance to the methods discussed by Pérez-Harguindeguy et al. (2013). We calculated tree height using trigonometric methods and used a base height of 1.37 m to measure tree diameter (Magarik et al., 2020). For shrub heights, we measured the distance from the ground to the height of the top foliage and measured stem diameters at approximately 1 cm above ground-level. All stem and leaf samples were kept cool during transport and measurements of leaf area and stem volume taken within 3 and 12 hours of sample collection respectively. To measure wood specific density, we collected a 10 cm sample of branch wood, taken close to the base of the branch at the stem and calculated stem volume using the water displacement method. For our leaf traits, we haphazardly selected and sampled five, fully expanded, and hardened leaves, with no to minimal herbivore damage. We took a high resolution scan of each leaf using a flatbed scanner and estimated leaf area using the ImageJ software (version 2.0.0).

Growth chamber study

For our growth chamber studies, we collected dormant branch cuttings from our highest and lowest latitude populations in each transect (total of four sites), with sampling in our eastern study occurring from 20-28 January 2015 and sampling for our western study from 19-28 October 2019. Dormant branch cuttings have been repeatedly shown to approximate whole plant responses in budburst (Vitasse and Basler, 2014), allowing us to estimate responses to environmental cues. In both studies, we

included a total of eight distinct treatments consisting of two levels of chilling, forcing, and photoperiod (Fig. 1). We recorded budburst stages of each sample every 1-3 days for up to four months, defining the day of budburst as the day of budbreak or shoot elongation (denoted as code 07 by Finn et al., 2007). For a more detailed discussion of study sample collection and methods see Flynn and Wolkovich (2018) for details on our eastern study and Loughnan and Wolkovich (in prep) for details on our western study.

Statistical Analysis

We combined our *in situ* trait data with budburst data from the controlled environment experiments through a joint Bayesian model for each trait. This approach improves upon previous analyses of trait correlations, as it allows us to carry through uncertainty between trait and phenology estimates—and better partitions the drivers of variation in species phenologies.

Our joint model consists of two parts. The first is a hierarchical linear model, which partitions the variation of individual observations (i) of a given trait value (Y_{trait}) to account for the effects of species (j), population-level differences arising from transects, latitude, as well as the interaction between transects and latitude ($transect \cdot latitude$), and finally, residual variation or 'measurement error' (σ^2_{m}).

$$Y_{\text{trait}_{i,j}} \sim Normal(\mu_{i,j}, \sigma^2_{\text{m}}) \tag{1}$$

$$\begin{aligned} \mu_{i,j} = \alpha_{\text{grand trait}} + \alpha_{\text{trait}_j} + \beta_{\text{transect}} \times transect + \\ \beta_{\text{latitude}} \times latitude + \beta_{\text{transect}\cdot\text{latitude}} \times (transect \cdot latitude) \end{aligned} \tag{2}$$

Where α_{trait} is an element of the normal random vector:

$$\boldsymbol{\alpha}_{\text{trait}} \begin{bmatrix} \alpha_{\text{trait}_1} \\ \alpha_{\text{trait}_2} \\ \cdots \\ \alpha_{\text{trait}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{trait}} \sim Normal(0, \sigma^2_{\text{trait}})$$

We included transect as a dummy variable (0/1) and latitude as a continuous variable and modeled traits in their original units, with the exception of leaf mass per area which was rescaled by 100 and wood specific density which was rescaled by 10 for numeric stability. Our model also included partial pooling for species—which controls for uneven sampling and trait variability. These species-level estimates are then predictors for each cue ($\beta_{\text{chilling,j}}$, $\beta_{\text{forcing,j}}$, $\beta_{\text{photoperiod,j}}$).

$$\begin{aligned} \beta_{\text{chilling}_j} &= \alpha_{\text{chilling},j} + \beta_{\text{trait.chilling}} \times \alpha_{\text{trait},j} \\ \beta_{\text{forcing}_j} &= \alpha_{\text{forcing},j} + \beta_{\text{trait.forcing}} \times \alpha_{\text{trait},j} \\ \beta_{\text{photoperiod}_j} &= \alpha_{\text{photoperiod},j} + \beta_{\text{trait.photoperiod}} \times \alpha_{\text{trait},j} \end{aligned} \tag{3}$$

In addition to the species-level estimates, this part of our model estimates the overall effect of each trait on each cue ($\beta_{\text{trait.chilling}}$, $\beta_{\text{trait.forcing}}$, $\beta_{\text{trait.photoperiod}}$). From this we could estimate how well traits explain species-level differences—by estimating the the species-level cue variation not explained by traits ($\alpha_{\text{chilling,j}}$, $\alpha_{\text{forcing,j}}$, $\alpha_{\text{photoperiod,j}}$) and individual species responses to cues (*chilling*, *forcing*, *photoperiod*, respectively). Finally, our model estimates the residual budburst variation across species

($\alpha_{\text{pheno,j}}$), observations (σ^2_{d}), as well as the variation in cues not attributed to the trait (using partial pooling).

$$Y_{\text{pheno}_{i,j}} \sim Normal(\mu_{i,j}, \sigma^2_{\text{d}}) \tag{4}$$

with

$$\mu_{i,j} = \alpha_{\text{pheno}_j} + \beta_{\text{chilling}_j} \cdot chilling + \beta_{\text{forcing}_j} \cdot forcing + \beta_{\text{photoperiod}_j} \cdot photoperiod \tag{5}$$

where α_{pheno_j}, $\alpha_{\text{chilling}_j}$, $\alpha_{\text{forcing}_j}$, and $\alpha_{\text{photoperiod}_j}$ are elements of the normal random vectors:

$$\boldsymbol{\alpha}_{\text{pheno}} = \begin{bmatrix} \alpha_{\text{pheno}_1} \\ \alpha_{\text{pheno}_2} \\ \cdots \\ \alpha_{\text{pheno}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{pheno}} \sim Normal(\mu_{\text{pheno}}, \sigma^2_{\text{pheno}}) \tag{6}$$

$$\boldsymbol{\alpha}_{\text{chilling}} = \begin{bmatrix} \alpha_{\text{chilling}_1} \\ \alpha_{\text{chilling}_2} \\ \cdots \\ \alpha_{\text{chilling}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{chilling}} \sim Normal(\mu_{\text{chilling}}, \sigma^2_{\text{chilling}}) \tag{7}$$

$$\boldsymbol{\alpha}_{\text{forcing}} = \begin{bmatrix} \alpha_{\text{forcing}_1} \\ \alpha_{\text{forcing}_2} \\ \cdots \\ \alpha_{\text{forcing}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{forcing}} \sim Normal(\mu_{\text{forcing}}, \sigma^2_{\text{forcing}}) \tag{8}$$

$$\boldsymbol{\alpha}_{\text{photoperiod}} = \begin{bmatrix} \alpha_{\text{photoperiod}_1} \\ \alpha_{\text{photoperiod}_2} \\ \cdots \\ \alpha_{\text{photoperiod}_n} \end{bmatrix} \text{ such that } \boldsymbol{\alpha}_{\text{photoperiod}} \sim Normal(\mu_{\text{photoperiod}}, \sigma^2_{\text{photoperiod}})$$

We modeled each trait individually, with the exception of ring-porosity, which we compared across species using the posterior estimates of our wood stem density model, allowing us to account for inherent differences in wood anatomy across species and growth form. We included all three cues (chilling, forcing, and photoperiod) as continuous variables in our model. We converted chilling temperatures to total chill portions, including both the chilling experienced in the field prior to sampling and during the experiment. For this we used local weather station data and the chillR package (v. 0.73.1, Luedeling, 2020). To account for differences in thermoperiodicity between our eastern and western growth chamber studies (Buonaiuto et al., 2023), we also converted forcing temperatures to mean daily temperatures for each treatment. Finally, we z-scored each cue and site using two standard deviations to allow direct comparisons between results across parameters (Gelman, 2008).

We used trait-specific priors that were weakly informative. We validated our choice of priors using prior predictive checks and confirmed model stability under wider priors. All models were coded in the Stan programming language for Bayesian models using the rstan package (Stan Development Team, 2018) in R version 4.3.1 (R Development Core Team, 2017). All models met basic diagnostic checks, including no divergences, high effective sample sizes (n_{eff}) that exceeded 10% of the number of iterations, and $\hat{R}$ values close to 1. We report our model estimates as the mean values with 90% uncertainty intervals (UI), interpreting parameter estimates with intervals that include zero to have no effect.

Results

Across our eight populations, we measured 47 unique species of which 28 were in our eastern transect and 22 in our western transect. These include species dominant in both the understory and canopy

layer, with our eastern community consisting of 13 shrubs and 15 trees, our western community consisting of 18 shrubs and 4 trees, of which three species occurred in both transects. In total we measured traits of 1428 unique individuals between the two transects across our five *in situ* traits: height ($n = 1317$), diameter ($n = 1220$), wood stem density ($n = 1359$), leaf mass per area ($n = 1345$), leaf nitrogen content ($n = 1351$). Across our two growth chamber studies, we made observations of 4211 branch cuttings, with our observations of budburst spanning 82 and 113 days for our eastern and western studies respectfully.

We found that three of our four traits had a strong relationship with photoperiod, but not always in the direction we predicted. As predicted, taller species with larger trunk diameters and leaves with high nitrogen content had larger responses with longer photoperiods (Fig. 2 c, i, o; Tables S2, S3, S6). But, contrary to our expectation, species with denser, high leaf mass per area leaves had smaller photoperiod responses, allowing them to potentially budburst earlier, under shorter photoperiods (Fig. 2f).

Temperature cues ($\beta_{\text{trait.chilling}}$ and $\beta_{\text{trait.forcing}}$) exhibited no relationships with individual traits, but by accounting for the effects of leaf or wood traits, we found the individual importance of our three cues on budburst varied by trait. Of the three cues, chilling (β_{chilling}) was the strongest in our models of height (-13.4 days per standardized chill portions, UI: -17.2, -9.9), diameter (-12.5 days per standardized chill portions, UI: -16.2, -8.6), wood stem density (-2.1 days per standardized chill portions, UI: -3.3, -9.8), and leaf nitrogen content (-35.1 days per standardized chill portions, UI: -68.1, -4.1), with more chilling advancing budburst. Our model of leaf mass per area, however, estimated photoperiod as the strongest cue ($\beta_{\text{photoperiod}}$, -14.0 days per standardized photoperiod, UI: -23.1, -3.5). After accounting for the effects of traits, only our height and diameter model found all three environmental cues to drive budburst timing (Tables S2, S3). Our models of wood stem density and leaf nitrogen content in turn found temperature cues alone to shape budburst (Tables S4, S6), while our model of leaf mass per area found a large response to only photoperiod (Table S5).

Only some of our trait models estimated clear gradients in species timing between trees and shrubs. In particular, we found height had large correlations between budburst timing and trait values, with earlier estimates of budburst for shrubs (with a mean day of budburst of 10)—especially under greater cues—and later budburst estimates for trees (with a mean day of budburst of 17.3, Fig. S1). But this was not the case for the leaf traits. Leaf nitrogen content, in particular, showed no distinct separation between shrub and tree functional groups (Fig. S1). We also found no differences between xylem structure and their responses to cues (Fig. 4).

Most of our traits showed some spatial variation, differing across latitude within each transect and with a strong interactive effect between transect and latitude (Fig. 3). Leaf nitrogen content was the only trait to vary with latitude alone, with low latitude communities on both our eastern and western transects having greater values of leaf nitrogen content than communities at higher latitudes (-0.1 percent per degree latitude, UI: -0.2, 0.0, Table S6). Plant diameter increased with increasing (higher) latitudes in both eastern and western communities (0.5 cm per degree latitude, UI: 0.1, 0.9), with a larger effect in eastern communities (Fig. 3d). Relationships with height, woody stem density and leaf mass per area were, however, more complex. Heights and wood stem density increased with increasing latitude in western communities, but decreased with latitude in eastern communities (-0.2 m per degree latitude, UI: -0.4, 0.0 for our height model and -0.01 g/cm^3 per degree latitude, UI: -0.02, 0.0 for our wood stem density model; Fig. 3 a and c), while for leaf mass per area we found the reverse (decreasing leaf mass per area with increasing latitudes in western communities, while increasing with latitude in eastern communities (0.005 g/cm^2 per degree latitude, UI: 0.004, 0.006; Fig. 3d). In addition to the differences we found across populations, we also observed large differences between individual species, which varied up to 7 fold for some traits (Fig. 2).

Discussion

Using a joint modeling approach to understand how phenology may fit within acquisitive to conservative plant strategies, we found that photoperiod was the only cue related to our suite of leaf and wood traits. While budburst responses to photoperiod are often smaller than temperature responses (chilling and forcing; Laube et al., 2014; Zohner et al., 2016; Flynn and Wolkovich, 2018), our results suggest photoperiod may be the most important cue in linking spring phenology to functional traits.

Our results suggest that fully predicting these responses also requires considering additional traits and variation across space given differences in the relative importance of cues across our trait models and between our eastern and western transects with latitude. The observed spatial differences in trait variation may be due to differences in the community assemblages, as our western community was more shrub dominated, with shorter plants with less dense branch wood, suggesting a more acquisitive growth strategy to allow species to utilize resources early in the season before canopy closure.

Cues and functional traits

We found only partial support for our prediction that species with acquisitive traits—particularly small trees with low wood density, low leaf mass per area, and high leaf nitrogen content—would have smaller temperature and photoperiod responses (associated with early budburst). Species with smaller heights and diameters did have smaller photoperiod responses, but—contrary to our prediction—species with less dense leaves showed larger responses to photoperiod, while leaves with high nitrogen content had stronger photoperiod responses.
None of our focal traits, however, showed a relationship with temperature (chilling or forcing), which may be due to selection on other physiological processes. Many of our traits are associated with one or more ecological function (Wright et al., 2004; Pérez-Harguindeguy et al., 2013; Reich, 2014). In particular, leaf mass per area is know to correlate with traits like leaf lifespan or decomposition rates in addition to light capture (De La Riva et al., 2016). In accounting for the relationships between other traits and budburst cues, we also found the relative importance of temperature and photoperiod on budburst timing varied. Despite strong evidence that budburst is driven by temperature and photoperiod cues, we found the strength of cue responses to differ between our trait models.

Decades of previous phenology research have found budburst timing to be primarily driven by temperature (chilling and forcing) and weakly by photoperiod (Chuine et al., 2010; Basler and Körner, 2014; Laube et al., 2014). Yet we found no traits that correlated with responses to temperature, suggesting other drivers may impact leaf and structural traits in temperate forests. One potential abiotic driver we did not consider is soil moisture, which covaries with a number of traits, including leaf mass per area, as higher leaf area allows plants to reduce evaporation under dry conditions (De La Riva et al., 2016). Soil moisture also shapes other phenological events in woody plants, including radial growth phenology and shoot elongation (Cabon et al., 2020; Peters et al., 2021). Though temperate forests are generally moist compared to other systems, previous studies suggests soil moisture can shape spring phenology across xeric systems as well (Crimmins et al., 2011; Park, 2014; Ettinger et al., 2019), thus including it in future studies could help understand how species growth strategies correlate with phenology.

Our finding that budburst was not related to wood structure or wood stem density contrasts the findings of previous work linking these traits. Previous studies have found some evidence that trees with diffuse-porous wood structure leafout earlier than species with ring-porous structures (Lechowicz, 1984; Panchen et al., 2014; Yin et al., 2016; Osada, 2017; Savage et al., 2022). Using wood density as an alternative measure of wood structure (wood density positively correlates with xylem resistance to embolism, Hacke et al., 2001), we did not find an association between our three phenological cues and xylem structure, despite our data including similar temperate forest species to previous studies.

This could be because we captured additional variation by sampling many individuals for each species across a large spatial gradient. In particular, we found larger wood densities at higher latitudes in our western transect, which could be caused by the differences in winter conditions experienced across this transect. Higher wood densities are especially favorable in communities that experience greater horizontal stress from wind and downward pressure from snow (MacFarlane and Kane, 2017; MacFarlane, 2020). Relationships with wood density may also be more apparent over a larger span of wood specific densities (our values varied from 0.2 to 0.6 g/cm^3, which is smaller than seen in studies that span a more global distribution, Galvão et al., 2021; Savage et al., 2022; Mo et al., 2024).

In addition to our study providing insight into how trait-budburst relationships vary with latitude, our sites also spanned North America, encompassing a gradient of 55° in longitude. At this continental scale we found correlations between phenology and traits for woody species in temperate forests, but these may shift with data that expands across biomes and more plant functional groups. Indeed, both the leaf and wood economic spectra find that large variation in traits across species arises from large global datasets and include both temperate and tropical species (Wright et al., 2004; Díaz et al., 2016; Chave et al., 2009) and functional groups that span from trees to grasses (Wright et al., 2004). Global patterns in trait variation may also be absent at smaller spatial scales (Wright and Sutton-Grier, 2012; Messier et al., 2017*a*,*b*), suggesting that comparisons of trait-phenology relationships at larger spatial scales or across more diverse pools of species may better align with the patterns predicted by existing economic spectra.

In comparing spring phenology of mainly temperate woody species, our results support previous work through which a global meta-analysis found that spring phenology may correlate with height and leaf mass per area (Loughnan et al., 2025). At both the global and continental scales, trees with taller heights leafed out with longer photoperiods. We also found species with high specific leaf area—which is the inverse of leaf mass per area and thus equivalent to low values—exhibited large responses to photoperiod (Loughnan et al., 2025). The consistency of these results, despite the differences in the two spatial scales of these datasets, provides further support that woody species responses to photoperiod may be part of a larger plant strategy.

Functional traits predict climate change responses

Our results offer novel insights into how broader correlations between growth strategies and phenological cues can help predict responses in woody plant communities with climate change. As temperatures rise, particularly at higher latitudes (Hoegh-Guldberg et al., 2018), warmer winter and spring temperatures may select for earlier budburst in some species. But, since photoperiod will remain fixed, our observed relationships between photoperiod and other traits has the potential to limit species abilities to track temperatures. This could constrain the extent to which some species growth will advance with climate change. Our results suggest that these effects will likely be greater for taller species or canopy trees and species with relatively low leaf mass per area. These constraints could have cascading effects on forest communities, as variable species responses to increasing temperatures further alter species growth strategies and their interactions with competitors or herbivores within their communities.

Our findings of correlations between phenology and other commonly measured traits highlight how accurate forecasts of future changes in phenology could benefit from accounting for the response of other traits to climate change. Across temperature and precipitation gradients, leaf size and shape also change, as species shift to conserve water and mitigate effects of transpiration under higher temperatures (De La Riva et al., 2016). These changes could impact species photosynthetic potential and ultimately ecosystem services, such as carbon sequestration. While phenological research has focused on forecasting responses to temperature, the correlation of other traits with photoperiod suggests it is also an important cue. Considering the tradeoffs and differences in cues that together shape plants

growth strategies may yield more accurate forecasts of species phenology and community dynamics under future climates.

References

Augspurger, C. K. 2008. Early spring leaf out enhances growth and survival of saplings in a temperate deciduous forest. Oecologia 156:281–286.

———. 2009. Spring 2007 warmth and frost: phenology, damage and refoliation in a temperate deciduous forest. Functional Ecology 23:1031–1039.

Basler, D., and C. Körner. 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. Tree Physiology 34:377–388.

Buonaiuto, D. M., E. M. Wolkovich, and M. J. Donahue. 2023. Experimental designs for testing the interactive effects of temperature and light in ecology : The problem of periodicity. Functional Ecology 37:1747–1756.

Cabon, A., L. Fernández-de-Uña, G. Gea-Izquierdo, F. C. Meinzer, D. R. Woodruff, J. Martínez-Vilalta, and M. De Cáceres. 2020. Water potential control of turgor-driven tracheid enlargement in Scots pine at its xeric distribution edge. New Phytologist 225:209–221.

Chave, J., D. Coomes, S. Jansen, S. L. Lewis, N. G. Swenson, and A. E. Zanne. 2009. Towards a worldwide wood economics spectrum. Ecology Letters 12:351–366.

Chuine, I. 2000. A unified model for budburst of trees. Journal of Theoretical Biology 207:337–347.

Chuine, I., and P. Cour. 1999. Climatic determinants of budburst seasonality in four temperate-zone tree species. New Phytologist 143:339–349.

Chuine, I., X. Morin, and H. Bugmann. 2010. Warming, photoperiods, and tree phenology. Science 329:277–278.

Cohen, J. M., M. J. Lajeunesse, and J. R. Rohr. 2018. A global synthesis of animal phenological responses to climate change. Nature Climate Change 8:224–228.

Crimmins, T. M., M. A. Crimmins, and C. D. Bertelsen. 2011. Onset of summer flowering in a 'Sky Island' is driven by monsoon moisture. New Phytologist 191:468–479.

De La Riva, E. G., M. Olmo, H. Poorter, J. L. Ubera, and R. Villar. 2016. Leaf Mass per Area (LMA) and Its Relationship with Leaf Structure and Anatomy in 34 Mediterranean Woody Species along a Water Availability Gradient. PLOS ONE 11:e0148788.

Díaz, S., J. Kattge, J. H. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer, C. Wirth, I. Colin Prentice, E. Garnier, G. Bönisch, M. Westoby, H. Poorter, P. B. Reich, A. T. Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright, S. N. Sheremet Ev, H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig, A. Günther, V. Falczuk, N. Rüger, M. D. Mahecha, and L. D. Gorné. 2016. The global spectrum of plant form and function. Nature 529:167–171.

Ettinger, A. K., C. J. Chamberlain, I. Morales-Castilla, D. M. Buonaiuto, D. F. B. Flynn, T. Savas, J. A. Samaha, and E. M. Wolkovich. 2020. Winter temperatures predominate in spring phenological responses to warming. Nature Climate Change 10:1137–1142.

Ettinger, A. K., I. Chuine, B. I. Cook, J. S. Dukes, A. M. Ellison, M. R. Johnston, A. M. Panetta, C. R. Rollinson, Y. Vitasse, and E. M. Wolkovich. 2019. How do climate change experiments alter plot-scale climate? Ecology Letters 22:748–763.

Finn, G. A., A. E. Straszewski, and V. Peterson. 2007. A general growth stage key for describing trees and woody plants. Annals of Applied Biology 151:127–131.

Flynn, D. F. B., and E. M. Wolkovich. 2018. Temperature and photoperiod drive spring phenology across all species in a temperate forest community. New Phytologist 219:1353–1362.

Galvão, F. G., A. L. Alves De Lima, C. Candeia De Oliveira, V. F. Da Silva, and M. J. N. Rodal. 2021. The importance of wood density in determining the phenology of tree species in a coastal rain forest. Biotropica 53:1134–1141.

Gelman, A. 2008. Scaling regression inputs by dividing by two standard deviations. Statistics in Medicine 27:2865–2873.

Gotelli, N. J., and G. R. Graves. 1996. The temporal niche. Pages 95–112 *in* Null Models In Ecology. Smithsonian Institution Press, Washington, D. C.

Green, S. J., C. B. Brookson, N. A. Hardy, and L. B. Crowder. 2022. Trait-based approaches to global change ecology: moving from description to prediction. Proceedings of the Royal Society B: Biological Sciences 289:1–10.

Hacke, U. G., J. S. Sperry, W. T. Pockman, S. D. Davis, and K. A. McCulloh. 2001. Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. Oecologia 126:457–461.

Harrington, C. A., and P. J. Gould. 2015. Tradeoffs between chilling and forcing in satisfying dormancy requirements for Pacific Northwest tree species. Frontiers in Plant Science 6:1–12.

Hoegh-Guldberg, O., D. Jacob, M. Taylor, M. Bindi, S. Brown, I. Camilloni, A. Diedhiou, R. Djalante, K. Ebi, F. Engelbrecht, J. Guiot, Y. Hijioka, S. Mehrotra, A. Payne, S. Seneviratne, A. Thomas, R. Warren, and G. Zhou. 2018. Impacts of 1.5 °C Global Warming on Natural and Human Systems. In: Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of . Tech. rep., Cambridge University Press, Cambridge, UK and New York, NY, USA.

Kharouba, H. M., J. Ehrlén, A. Gelman, K. Bolmgren, J. M. Allen, S. E. Travers, and E. M. Wolkovich. 2018. Global shifts in the phenological synchrony of species interactions over recent decades. Proceedings of the National Academy of Sciences 115:5211–5216.

Körner, C., and D. Basler. 2010. Phenology Under Global Warming. Science 327:1461–1463.

Laube, J., T. H. Sparks, N. Estrella, J. Höfler, D. P. Ankerst, and A. Menzel. 2014. Chilling outweighs photoperiod in preventing precocious spring development. Global Change Biology 20:170–182.

Lechowicz, M. J. 1984. Why Do Temperate Deciduous Trees Leaf Out at Different Times? Adaptation and Ecology of Forest Communities. The American Naturalist 124:821–842.

Lopez, O. R., K. Farris-Lopez, R. A. Montgomery, and T. J. Givnish. 2008. Leaf phenology in relation to canopy closure in southern Appalachian trees. American Journal of Botany 95:1395–1407.

Loughnan, D., F. A. Jones, G. Legault, C. J. Chamberlain, D. M. Buonaiuto, A. K. Ettinger, M. Garner, D. S. Sodhi, and E. M. Wolkovich. 2025. Budburst timing within a functional trait framework. Journal of Ecology 00:1–12.

Loughnan, D., and E. M. Wolkovich. in prep. Temporal assembly of woody plant communities shaped equally by evolutionary history as by current environments .

Luedeling, E. 2020. chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees. https://CRAN.R-project.org/package=chillR.

MacFarlane, D. W. 2020. Functional Relationships Between Branch and Stem Wood Density for Temperate Tree Species in North America. Frontiers in Forests and Global Change 3.

MacFarlane, D. W., and B. Kane. 2017. Neighbour effects on tree architecture: functional trade-offs balancing crown competitiveness with wind resistance. Functional Ecology 31:1624–1636.

Magarik, Y. A., L. A. Roman, and J. G. Henning. 2020. How should we measure the dbh of multi-stemmed urban trees? Urban Forestry & Urban Greening 47:1–11.

Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aaasa, R. Ahas, K. Alm-Kübler, P. Bissolli, O. Braslavská, A. Briede, F. M. Chmielewski, Z. Crepinsek, Y. Curnel, Å. Dahl, C. Defila, A. Donnelly, Y. Filella, K. Jatczak, F. Måge, A. Mestre, Ø. Nordli, J. Peñuelas, P. Pirinen, V. Remišová, H. Scheifinger, M. Striz, A. Susnik, A. J. Van Vliet, F. E. Wielgolaski, S. Zach, and A. Zust. 2006. European phenological response to climate change matches the warming pattern. Global Change Biology 12:1969–1976.

Messier, J., M. J. Lechowicz, B. J. McGill, C. Violle, and B. J. Enquist. 2017*a*. Interspecific integration of trait dimensions at local scales: the plant phenotype as an integrated network. Journal of Ecology 105:1775–1790.

Messier, J., B. J. McGill, B. J. Enquist, and M. J. Lechowicz. 2017*b*. Trait variation and integration across scales: is the leaf economic spectrum present at local scales? Ecography 40:685–697.

Mo, L., T. W. Crowther, D. S. Maynard, and e. a. Van Den Hoogen. 2024. The global distribution and drivers of wood density and their impact on forest carbon stocks. Nature Ecology & Evolution 8:2195–2212.

Osada, N. 2017. Relationships between the timing of budburst, plant traits, and distribution of 24 coexisting woody species in a warm-temperate forest in Japan. American Journal of Botany 104:550–558.

Panchen, Z. A., R. B. Primack, B. Nordt, E. R. Ellwood, A. Stevens, S. S. Renner, C. G. Willis, R. Fahey, A. Whittemore, Y. Du, and C. C. Davis. 2014. Leaf out times of temperate woody plants are related to phylogeny, deciduousness, growth habit and wood anatomy. New Phytologist 203:1208–1219.

Park, I. W. 2014. Impacts of differing community composition on flowering phenology throughout warm temperate, cool temperate and xeric environments. Global Ecology and Biogeography 23:789–801.

Parmesan, C., and G. Yohe. 2003. A globally coherent fingerprint of climate change impacts across natural systems. Nature 421:37–42.

Pérez-Harguindeguy, N., S. Díaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S. Bret-Harte, W. K. Cornwell, J. M. Craine, D. E. Gurvich, C. Urcelay, E. J. Veneklaas, P. B. Reich, L. Poorter, I. J. Wright, P. Ray, L. Enrico, J. G. Pausas, A. C. de Vos, N. Buchmann, G. Funes, F. Quétier, J. G. Hodgson, K. Thompson, H. D. Morgan, H. ter Steege, M. G. A. van der Heijden, L. Sack, B. Blonder, P. Poschlod, M. V. Vaieretti, G. Conti, A. C. Staver, S. Aquino, and J. H. C. Cornelissen. 2013. New handbook for standardized measurement of plant functional traits worldwide. Australian Journal of Botany 61:167–234.

Peters, R. L., K. Steppe, H. E. Cuny, D. J. De Pauw, D. C. Frank, M. Schaub, C. B. Rathgeber, A. Cabon, and P. Fonti. 2021. Turgor – a limiting factor for radial growth in mature conifers along an elevational gradient. New Phytologist 229:213–229.

Primack, R. B., I. Ibáñez, H. Higuchi, S. D. Lee, A. J. Miller-Rushing, A. M. Wilson, and J. A. Silander. 2009. Spatial and interspecific variability in phenological responses to warming temperatures. Biological Conservation 142:2569–2577.

R Development Core Team. 2017. R: A language and environment for statistical computing.

Reich, P. B. 2014. The world-wide 'fast–slow' plant economics spectrum: a traits manifesto. Journal of Ecology 102:275–301.

Sakai, A., and W. Larcher. 1987. Frost Survival of Plants: Responses and adaptation to freezing stress. Springer-Verlag, Berlin, Heidelberg.

Savage, J. A., T. Kiecker, N. McMann, D. Park, M. Rothendler, and K. Mosher. 2022. Leaf out time correlates with wood anatomy across large geographic scales and within local communities. New Phytologist 235:953–964.

Schweingruber, F., and W. Landolt. 2010. The xylem database.

Stan Development Team. 2018. RStan: the R interface to Stan. R package version 2.17.3.

Thackeray, S. J., P. A. Henrys, D. Hemming, J. R. Bell, M. S. Botham, S. Burthe, P. Helaouet, D. G. Johns, I. D. Jones, D. I. Leech, E. B. MacKay, D. Massimino, S. Atkinson, P. J. Bacon, T. M. Brereton, L. Carvalho, T. H. Clutton-Brock, C. Duck, M. Edwards, J. M. Elliott, S. J. Hall, R. Harrington, J. W. Pearce-Higgins, T. T. Høye, L. E. Kruuk, J. M. Pemberton, T. H. Sparks, P. M. Thompson, I. White, I. J. Winfield, and S. Wanless. 2016. Phenological sensitivity to climate across taxa and trophic levels. Nature 535:241–245.

Violle, C., M. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let the concept of trait be functional! Oikos 116:882–892.

Vitasse, Y., and D. Basler. 2014. Is the use of cuttings a good proxy to explore phenological responses of temperate forests in warming and photoperiod experiments? Tree Physiology 34:174–183.

Vitasse, Y., G. Hoch, C. F. Randin, A. Lenz, C. Kollas, J. F. Scheepens, and C. Körner. 2013. Elevational adaptation and plasticity in seedling phenology of temperate deciduous tree species. Oecologia 171:663–678.

Vitasse, Y., A. Josée, A. Kremer, R. Michalet, and S. Delzon. 2009. Responses of canopy duration to temperature changes in four temperate tree species : relative contributions of spring and autumn leaf phenology. Oecologia 161:187–198.

Vitasse, Y., C. Signarbieux, and Y. H. Fu. 2018. Global warming leads to more uniform spring phenology across elevations. PNAS 115:1004–1008.

Wang, N., Z. Wu, Y. Gong, Y. Cui, and Y. H. Fu. 2025. Background climate determines the response of spring leaf-out to climate change—Results from a national-scale twig-cutting experiment. Journal of Ecology 113:2981–2991.

Wolkovich, E. M., and A. K. Ettinger. 2014. Back to the future for plant phenology research. New Phytologist 203:1021–1024.

Wright, I. J., M. Westoby, P. B. Reich, J. Oleksyn, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares, T. Chapin, J. H. C. Cornellissen, M. Diemer, J. Flexas, J. Gulias, E. Garnier, M. L. Navas, C. Roumet, P. K. Groom, B. B. Lamont, K. Hikosaka, T. Lee, W. Lee, C. Lusk, J. J. Midgley, Ü. Niinemets, H. Osada, H. Poorter, P. Pool, E. J. Veneklaas, L. Prior, V. I. Pyankov, S. C. Thomas, M. G. Tjoelker, and R. Villar. 2004. The worldwide leaf economics spectrum. Nature 428:821–827.

Wright, J. P., and A. Sutton-Grier. 2012. Does the leaf economic spectrum hold within local species pools across varying environmental conditions? Functional Ecology 26:1390–1398.

Yin, J., J. D. Fridley, M. S. Smith, and T. L. Bauerle. 2016. Xylem vessel traits predict the leaf phenology of native and non-native understorey species of temperate deciduous forests. Functional Ecology 30:206–214.

Zeng, Z. A., and E. M. Wolkovich. 2024. Weak evidence of provenance effects in spring phenology across Europe and North America. New Phytologist 242:1957–1964.

Zohner, C. M., B. M. Benito, J.-C. Svenning, and S. S. Renner. 2016. Day length unlikely to constrain climate-driven shifts in leaf-out times of northern woody plants. Nature Climate Change 6:1120–1123.



Figure 1: We collected trait data and branch cuttings from woody plants growing within eight sites, across two transects in eastern and western North America (map at left). Cuttings were used in two controlled environment (growth chamber) studies in which we applied high and low chilling, forcing, and photoperiod treatments and recorded the day of budburst. Using our paired *in situ* trait and experimental budburst data, we tested whether earlier budbursting species exhibited traits associated with more acquisitive growth strategies and smaller responses to cues and later budbursting species a more conservative growth strategy and larger responses to cues (see 'Predictions' at right).



Figure 2: Relationships between species traits (rows) and cue responses (columns) showed considerable variation across: (a-c) height, (d-f) leaf mass per area, (g-i) diameter, (j-l) wood specific density, and (m-o) leaf nitrogen content. Point colours represent different species groups, with tree species shown in red and shrub species in blue. Crosses depict the 50% uncertainty interval of the model estimates of species trait values and estimated responses to cues. Grey bands depict large relationships between a trait and cue, representing the 90% uncertainty interval, and black lines the mean response.



Figure 3: We found (a) height, (b) diameter, (c) branch wood specific density, and (d) leaf mass per area to all experience a strong interaction between latitude and transect, (e) while leaf nitrogen content showed a strong effect of latitude alone.



Figure 4: Despite previous studies finding relationships between leaf out timing and wood xylem structures, we did not find clear differences in species-level estimates of cue responses with wood structure or relative to their wood specific densities. Each cross represents the 50% uncertainty interval of (a). forcing, (b) chilling, and (c) photoperiod responses and wood specific density for each species, with colors depicting different types of wood structure. The lowest wood specific density was estimated for *Sambucus racemosa* and the highest wood specific density for *Viburnum lantanoides*.