

1 How temperature, photoperiod and evolutionary history shape
2 forest leafout

3 Deirdre Loughnan^{1,2,3} and E M Wolkovich¹

4 ¹ Department of Forest and Conservation Sciences, Faculty of Forestry, University of British Columbia,
5 2424 Main Mall Vancouver, BC, Canada, V6T 1Z4.

6
7 ² Department of Zoology, Faculty of Science, University of British Columbia, 6270 University Blvd,
8 Vancouver, BC, Canada, V6T 1Z4

9
10 ³ Hakai Institute, PO Box 25039, Campbell River, BC, Canada, V9W 0B7

11
12 Corresponding Author: Deirdre Loughnan, deirdre.loughnan@ubc.ca, 6270 University Blvd, RM 1015,
13 Vancouver, BC, V6T 1Z4

14 Coauthor: Elizabeth Wolkovich, e.wolkovich@ubc.ca

15
16 Running title: The environment and evolution shape forests

17
18 Keywords: Phenology, Budburst, Climate change, Forest communities, Community assembly, Tempo-
19 ral niche.

20

21 Abstract

22 1. How species assemble temporally has received growing attention as climate change shifts the timing
23 of major life cycle events across most systems. While theory predicts that plant species should partition
24 time to reduce resource competition—for example by leafing out at different times—few studies have
25 examined this given the complexity of environmental cues that make timings highly variable across
26 years and sites.

27

28 2. We examine species- and community-level responses in budburst across 47 woody plant species
29 across four sites in North America. By manipulating known temperature and photoperiod cues in
30 controlled environments then applying a phylogenetic Bayesian model, we estimate the effects of these
31 cues and evolutionary history in shaping leafout across species.

32

33 3. We found responses from plants sampled at different sites were highly similar, but species var-
34 ied greatly in their timings. Species-specific differences in responses to well-studied temperature and
35 photoperiod cues, however, explained only half this variation, with the remaining variation mostly
36 explained by intrinsic species differences in timing and shared evolutionary history.

37

38 Synthesis. This suggests current models of leafout phenology are missing important variation that may
39 structure species assembly within communities. Identifying these unidentified cues or traits is critical
40 to forecasting phenological shifts and understanding their effects on community dynamics.

Introduction

Climate change has led to advances of about three days per decade in the timing of species life history events—phenology (Parmesan, 2007; Thackeray *et al.*, 2016; Cohen *et al.*, 2018). Different types of events, however, are highly variable, with certain events, such as leafout, spanning a period of weeks. Understanding and explaining this variability has become increasingly important as phenology shifts in response to rising temperatures at different rates (Parmesan, 2007; Fitter & Fitter, 2002; Menzel *et al.*, 2006; Yu *et al.*, 2010; Fridley, 2012). Some degree of variability is likely due to differences in climate change itself across space—as some areas warm faster than others and generate larger phenological shifts (Hoegh-Guldberg *et al.*, 2018). Recent research suggests that species-level differences in phenology also drive some of this variation (Vitasse *et al.*, 2009; Wolkovich & Cleland, 2014; Zohner & Renner, 2014; Vitasse *et al.*, 2018). How these differences compare to other potential sources of variation, and what predicts them, is still largely unknown, but has major implications for predicting how changes in phenology will affect community dynamics and ecosystem services.

Differences in species phenologies can impact community structure and dynamics. Community ecology has long posited that species assemble in part through partitioning time, leading to unique temporal niches that allow species to coexist (Chesson *et al.*, 2004; Grime, 1977). For example, earlier germination or leafout may provide understory species early-access to soil and light resources, before larger canopy species leafout (Heberling *et al.*, 2019; Lee & Ibáñez, 2021). These different temporal niches may also trade-off different pressures, with research suggesting that in seasonal systems species that are active earlier experience greater abiotic pressures—such as spring frost—but avoid the larger competitive pressures for resources that occur later in the season when most species are fully active (Sakai & Larcher, 1987; Gotelli & Graves, 1996; Augspurger, 2009; Pau *et al.*, 2011). As climate change variably affects the timing of growth, it may alter the temporal order of species in a community and result in novel interactions between species (Cleland *et al.*, 2012; Rudolf, 2019; Tiusanen *et al.*, 2020), thus changing the competitive landscape and synchrony of interactions within communities.

While recent changes in climate are shaping differences in phenology, for many species their temporal niches may also reflect their evolutionary history. Species traits in communities today are the result of selective pressures acting over evolutionary timescales that include the historical climates and related ancestral phenotypes that were favourable under previous conditions (Wiens *et al.*, 2010). This could produce greater similarity in phenological traits of more closely related species (Davies *et al.*, 2013; Morales-Castilla *et al.*, 2024), which would impact how species respond to future climates.

Spring budburst offers an excellent system to test for species- and community-level patterns in phenology and environmental cues. Budburst of temperate woody plants is well studied and thus known to respond to environmental cues, especially temperatures in the winter and spring—referred to as chilling and forcing, respectively—and daylength or photoperiod (Chuine *et al.*, 2010; Polgar & Primack, 2011; Cooke *et al.*, 2012; Basler & Körner, 2014; Laube *et al.*, 2014). Budburst responses to these cues are generally similar across sites and populations (Zeng & Wolkovich, 2024; Aitken & Bemmels, 2016). Species-level differences, however, are generally much larger than observations of site or

population-level variation (but see Deans, 1996; Sogaard *et al.*, 2008, for examples of population-level differences). Decades of phenological research suggests these unique phenologies are likely due to different responses of species to temperature and photoperiod cues, though they may be also structured by species shared evolutionary history (Lechowicz, 1984; Davies *et al.*, 2013; Gougherty & Gougherty, 2018) and their current environments. For example, research predicts that forest understory species should vary their leafout and budburst early to both gain access to light and soil nutrients prior to canopy closure (Mahall & Bormann, 1978; Muller, 1978), as outlined above, and to reduce competition between species within the understory versus canopy.

Here we test how environmental—temperature and photoperiod—cues and evolutionary relationships shape species-level variation in phenology of forest communities across North America. Leveraging two large-scale growth chamber cutting experiments, in which we vary the temperature cues of chilling (cool temperatures in the fall and winter) and forcing (warmer temperatures, usually in the spring) with photoperiod (Fig. 1), and Bayesian phylogenetic approach to estimate budburst responses across 47 woody species. We collected samples from plants at four sites, in eastern and western North America (Fig. 1), with pairs of sites on each coast spanning 4 and 6° latitude, respectively. Our dataset includes the diverse assemblages of tree and shrub species found in eastern and western deciduous forests, allowing us to explore differences across both sites and functional groups—including understory shrubs versus canopy trees—in eastern versus western forests.

Materials and Methods

Field sampling

We combined data from two growth chamber studies using dormant branch clippings of North American deciduous woody plants. Such cutting experiments are a common and powerful approach to infer the responses of adult trees to environmental conditions (Vitasse *et al.*, 2014). In our first study, we collected samples from two eastern communities—Harvard Forest, Massachusetts, USA (42.55°N, 72.20°W) and St. Hippolyte, Quebec, Canada (45.98°N, 74.01°W), from 20-28 January, 2015 (previously reported in Flynn & Wolkovich, 2018). The second growth chamber study spanned two western communities—Smithers (54.78°N, 127.17°W) and E.C. Manning Park (49.06°N, 120.78°W), British Columbia, Canada, with sampling from 19-28 October 2019 (not previously published). Combining the datasets from these two experiments allowed us to test for spatial variation in the phenology of woody plant communities at a larger geographic scale and make stronger inferences for how different species vary in their cues.

We selected the dominant deciduous species in each forest community and maximized the number of species occurring across sites. We observed budburst for 47 species spanning eastern and western forest communities, with 28 species at our eastern transect and 22 species at our western transect, of which 3 species occurred at both regions (Fig. 1, Table S1). Of the species in our eastern transect, there were 13 shrubs and 15 trees, while our western transect included 18 shrub and 4 tree species. At all sites, we tagged 15-20 healthy, individuals for each species in the summer prior to sampling.

To reduce ontogenetic effects (Vitasse *et al.*, 2014), we only sampled mature individuals, which we assumed to be trees that were greater than 2m in height and with DBH greater than 4cm (Natural Resources Canada, 2020). We collected a single cutting from small shrub species, but depending on the size of the sampled tree, we took between 6-16 cuttings, thus collecting samples from from 6-20 tagged plants per species. All cuttings were taken from the same relative canopy position, sampling terminal branches from shrubs and the lower canopy from trees using a pole pruner from the ground. We kept samples cold and immediately placed them in erlenmeyer flasks of water upon returning from the field. We conducted the eastern study at the Arnold Arboretum, Boston, MA, USA, and the western study at the University of British Columbia, BC, Canada. We used daily weather data from local weather stations to calculate the field chilling prior to collection (see Table S2).

Growth chamber study

We used growth chambers with two levels of chilling—with either no additional chilling or 30 days at 4°C for our eastern study, and 21 or 56 days of chilling at 4°C for our western study (all dark)—after which we moved plants to one of two levels of forcing—with day and night temperatures that varied to create a cool regime of 15:5°C and a warm regime of 20:10°C—crossed with two photoperiods of either 8 or 12 hours—for a total of 8 distinct treatments in each study.

Our design was similar for both eastern and western sites (Fig. 1). We had seven to ten replicates per treatment in our eastern study and eight replicates in our western study, with the exception of *Sorbus scopulina*, for which we only had six replicates per treatment. Our eastern and western studies differed only in the timing of sample collection and thermoperiodicity—alternation in day and night temperatures—in forcing treatments. By collecting samples in our eastern study in late January, they experienced considerable field chilling. We collected the western samples in late October, with most chilling occurring under controlled conditions. In our eastern study, we also set the duration of forcing temperatures to correspond with the photoperiod treatments. While this is a widespread and common approach in phenology chamber studies, it leads to a confounding effect in the thermoperiodicity (Buonaiuto *et al.*, 2023). Thus we applied forcing temperatures for 12 hours independent of the duration of the photoperiod treatments in our western study, and controlled for this difference in our statistical approach (see below). Each experiment was conducted across 8 growth chambers. We additionally rotated cuttings within chambers approximately every two weeks to avoid any chamber effects and changed the water and re-cut each cutting to prevent callusing and keep cuttings healthy. We controlled for small differences in the number of replicates across species in our statistical approach using partial pooling (see section on statistical analysis).

We assessed phenological stages of budburst using the BBCH scale, adapted for our specific species (Finn *et al.*, 2007). We also created photographic guides to help define each stage of the BBCH scale for our specific species (Savas *et al.*, 2017; Loughnan & Wolkovich, 2024a). Every 1-3 days, we observed each sample and recorded all phenological stages up to full leaf expansion (defined as code 17 by Finn *et al.*, 2007). Here, we present the response in budburst day, defined as the day of budbreak or shoot elongation (defined as code 07 by Finn *et al.*, 2007) since forcing treatments began. Our eastern study

spanned 82 days, with over 19320 phenological observations, while the western study spanned 113 days and 47844 phenological observations.

Statistical Analysis

We tested for differences in budburst cues across species and communities using a phylogenetic mixed effects model with partial pooling (‘shrinkage’) across species. This approach accounts for both the evolutionary relatedness of our species and estimates the species-level cues and differences across sites. We obtained species phylogenetic relatedness by pruning the Smith & Brown (2018) megatree of angiosperms (Fig. 5).

In our model, we included each of our three cues as continuous variables. To account for the greater amounts of field chilling experienced in our eastern study, we converted our chilling treatments to chill portions for each individual community using local weather station data and the chillR package (v. 0.73.1, Luedeling, 2020). **Chill portions are a commonly used chill metric that is calculated using hourly climate data and a Dynamic Model in which cool temperatures lead to the accumulation of chilling, while more accurately estimates chilling accumulation under fluctuating temperatures (Luedeling, 2012, 2020).** We calculated chill portions for both the chilling experienced in the field prior to sampling and during the experiment in our 4°C chilling chambers. We also converted forcing temperatures to mean daily temperatures in each treatment to account for differences in thermoperiodicity between the two studies (Buonaiuto *et al.*, 2023). Finally, we *z*-scored each cue and site using two standard deviations to allow direct comparisons between results across cues (Gelman, 2008). In addition to the overall predictors, we included all two-way interactions between cues and between cues and sites.

Our modelling approach allowed us to combine observations of budburst (y_i) across species (sp) to estimate both the effects of species-specific differences in their responses (slope estimates, β) to cues (chilling, forcing and photoperiod) and any remaining species-specific differences in timings (via species-level intercepts, α_{sp_i}). Our full model also estimated effects of sites and all possible two-way interactions between cues, and between cues and sites, to estimate the day of budburst (\hat{y}) relative to the first day of forcing conditions.

$$\begin{aligned}\hat{y}_i = & \alpha_{sp_i} + \beta_{Manning\ park} + \beta_{Harvard\ forest} + \beta_{St.\ Hippolyte} + \\ & \beta_{chilling_{sp_i}} + \beta_{forcing_{sp_i}} + \beta_{photoperiod_{sp_i}} + \\ & \beta_{forcing \times chilling_{sp_i}} + \beta_{forcing \times photoperiod_{sp_i}} + \beta_{chilling \times photoperiod_{sp_i}} + \\ & \beta_{forcing \times Manning\ park_{sp_i}} + \beta_{forcing \times Harvard\ forest_{sp_i}} + \beta_{forcing \times St.\ Hippolyte_{sp_i}} + \\ & \beta_{chilling \times Manning\ park_{sp_i}} + \beta_{chilling \times Harvard\ forest_{sp_i}} + \beta_{chilling \times St.\ Hippolyte_{sp_i}} + \\ & \beta_{photoperiod \times Manning\ park_{sp_i}} + \beta_{photoperiod \times Harvard\ forest_{sp_i}} + \beta_{photoperiod \times St.\ Hippolyte_{sp_i}}\end{aligned}$$

190

$$y_i \sim \text{normal}(\hat{y}_i, \sigma^2)$$

191 The slopes were modelled with partial pooling at the species-level:

192

$$\beta_{forcing_{sp}} \sim \text{normal}(\mu_{forcing_{sp}}, \sigma_{forcing}^2)$$

...

$$\beta_{photoperiod \times St.Hippolyte_{sp}} \sim \text{normal}(\mu_{photoperiod \times St.Hippolyte_{sp}}, \sigma_{photoperiod \times St.Hippolyte_{sp}}^2)$$

193 We included the phylogenetic effect on species-specific differences in budburst timing (α_{sp}) as a variance
 194 covariance matrix (\mathbf{V}) in the parameterization of the normal random vector:

$$\alpha_{sp} = \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \dots \\ \alpha_n \end{bmatrix} \text{ such that } \boldsymbol{\alpha} \sim \text{multi-normal}(\mu_{\alpha}, \mathbf{V}), \quad (1)$$

(2)

195 The timing of budburst ancestrally is represented as the root value of the means of the multivariate
 196 normal distribution. The phylogenetic structuring of budburst is estimated using Pagel's lambda, a
 197 phylogenetic scaling parameter where values close to one indicate a phylogenetic signal consistent with
 198 Brownian motion and values close to zero the absence of a phylogenetic signal for a given trait (Pagel,
 199 1999; Pearse *et al.*, 2025). For more detail on this method of phylogenetic modeling, see Morales-
 200 Castilla *et al.* (2024).

201

202 We used non-informative priors for each of our model parameters (increasing priors three-fold did not
 203 demonstrably change our estimates), and validated the model code using test data prior to analysis.
 204 Our model produced \hat{R} values close to 1, well mixed chains, and n_{eff} values that exceeded 10% of the
 205 model iterations. Our model was fit using the Stan language with the rstan package (v. 2.26.23, Stan
 206 Development Team, 2018) in R (v. 4.3.1, R Development Core Team, 2017).

207

208 We present model estimates, relative to baseline or low treatment conditions. This means that we
 209 show the average daily temperature for our cooler forcing conditions (which averaged to 8.33°C for our
 210 eastern sites and 10°C for our western sites, after accounting for the differences in thermoperiodicity),
 211 with a short photoperiod (8 hours photoperiod), and the mean of lowest chill portions from each pop-
 212 ulation (which is equivalent to approximately 10 weeks of chilling at 4°C) and—when relevant—our
 213 most southern sites. We present results as means and 90% posterior uncertainty intervals from our
 214 Bayesian phylogenetic models.

215

Results

On average, species budburst occurred 30.4 days (uncertainty interval: 26.0, 35.2, all estimates are given as mean \pm 90% uncertainty intervals, henceforth ‘UI,’ and budburst dates given as relative to baseline conditions, see methods for more details) after the start of forcing and photoperiod treatments. Between the earliest and latest budbursting species, the difference in the mean day of budburst was 38.5 and 30.7 days for the eastern and western sites respectively. Budburst of individual species showed distinct differences in their timing and the relative importance of cues, but that all cues led to an advance in budburst date. Of the three cues, chilling had the largest effect, with increased chilling advancing budburst (we represent advances with negative numbers, while delays are positive; -14.8, UI: -16.7, -12.9 days per standardized chill portions) and photoperiod the smallest (-3.4, UI: -4.0, -2.7 days per standardized photoperiod, Fig. 2a, S1). But we found a large, subadditive interaction between forcing and chilling (8.4, UI: 7.0, 9.8, Fig. 2b); meaning that low chilling is offset by high forcing conditions, and vice versa (Table S3). Despite our expectations that species responses should differ across the canopy layer, with shrubs being less sensitive to cues than trees, we did not find substantial differences across these two functional groups (Fig. S2). Shrubs, like *Cornus stolonifera*, budburst early and showed small responses to chilling and forcing (Fig. 3b, d). But 36.7% of the shrubs, including *Menziesia ferruginea* and *Symphoricarpos alba*, exhibited the opposite response and budburst relatively late (Fig. 4, S2). Similarly for trees, some species did match our predictions, exhibiting stronger cues, but 23.5% of species budburst earlier than expected. While specific tree species, such as *Quercus velutina*, did have larger chilling and photoperiod responses as predicted (Fig. 3a, e), and *Fagus grandifolia* had the largest photoperiod response (Fig. 3e), overall we did not find clear differences between the cues of trees and shrubs across the four forest communities.

Overall we found species budburst was moderately phylogenetically structured (Pagel’s λ of 0.4, UI: 0.1, 0.8, estimated on the intercept). So while our results provide clear evidence for species-specific differences in individuals responses to temperature (chilling and forcing) and photoperiod, they also provide evidence for additional, unknown, latent processes that shape species-level differences in budburst timing. But the contribution of these species-specific cue differences to budburst order and timing was small compared to the overall species differences unexplained by cues (represented in the modeling framework as the intercept), with the exact magnitude for which chilling, forcing and photoperiod determine budburst date depending on the level of each cue (e.g., forcing has a much larger effect on budburst timing given a constant daytime temperature of 20°C compared to a constant daytime temperature of 15°C). Across the levels of chilling, forcing and photoperiod we studied species-specific differences in responses to cues explained only between 37.6% and 67.8% of variation for eastern species and between 47.4 and 61.7% for western species, with the intercept—which was phylogenetically structured (Fig. 5)—explaining the rest.

We also found the rank order of species’ budburst timings changed more when estimated using higher chilling (the average of which is equivalent to approximately 15 weeks chilling at 4°C), forcing (Fig. 1) and longer photoperiods than under lower chilling and forcing and shorter photoperiods (Fig S3), but remained fairly consistent. The budburst ranking for the three species that occurred in both transects

remained relatively consistent across the two communities (Fig. 3), with only *Alnus incana* in the western community experiencing a large change in rank with cues (Fig. S3). In comparing the earliest and latest budbursting species, we found relatively small differences in cues (Fig. 3). For example, an early budbursting shrub, *Lyonia ligustrina*, had chilling and forcing cues of -14.7 and -11.5 respectively, which are comparable to the cue estimates of -15.7 and -11.5 for the much later tree, *Quercus alba*. Yet the model predicts these species to have a 15.1 day difference in day of budburst, highlighting how the timing of budburst across species was also strongly related to species level differences independent of their responses to chilling, forcing and photoperiod (intercept values, see Fig. 4).

Finally, we observed small site-level effects compared to differences between cues (Fig. 2). Treatments that received a month less of chilling (low chilling), were 5°C cooler (low forcing), and had shorter photoperiods all budburst later compared to the more chilled, warmer and long photoperiod treatments across all sites and across transects (Fig. S1 a-c, shown for baseline conditions, see Table S3 for model output). Overall budburst dates did not differ between sites, though eastern sites budburst marginally earlier (day 35.8, UI: 28.3, 42.3) compared to the western sites (day 48.1, UI: 40.7, 54.5, see also Table S3 for model output). This could be due to the earlier collection date of our western species, which reduced the field chilling they received relative to our eastern samples (though see corrections for this in the statistical analysis section of our methods).

Discussion

To accurately forecast shifts in phenology and its impact on ecological communities requires understanding which environmental cues are most important at the population-, species-, functional-group and community-levels. Our results support decades of research finding that budburst timing advances in response to the well-studied major temperature cues of chilling and forcing (with the greatest responses to chilling and forcing respectively, see also Ettinger *et al.*, 2020; Flynn & Wolkovich, 2018) and longer photoperiods. But using our modelling approach allows us to estimate the extent to which these species-specific responses to these cues do—or do not—lead to differences in species timings that may structure their temporal niches within communities. Perhaps surprisingly, we found a large (one-third to one-half) of variation in species budburst timings was unexplained by different responses to chilling, forcing and photoperiod, suggesting important unexplained variation. Our results show that some of this variation is due to species evolutionary history, but not all, suggesting fundamental gaps in our model of one of the best studied and most important phenological events—woody plant budburst.

Differences in species responses to temperature and photoperiod were similar across sites and functional groups, suggesting that understanding the drivers of variation across species may be an important task to improve forecasting. Our study includes 47 of the dominant woody species from deciduous forests across North America and includes shrub and tree functional groups, with a mix of early and late budbursting. Despite the high degree of diversity and spatial scale represented in our study, we still found that these groups are not inherently different in their timings. Our results found no detectable

site-level differences across different coasts and over 6° of latitude. This contrasts with research that suggests latitudinal trends in photoperiod and temperature may impose unique selective pressures, and thus may drive site-level differences in responses to cues (Keller *et al.*, 2011). Latitudinal gradients in spring phenological shifts have been found by studies using *in situ* phenological data (Post *et al.*, 2018; Alecrim *et al.*, 2023). Such work, however, is based on long-term observations in the field, with confounding differences in study duration and start dates, variable methodologies, and geographic extent. The lack of site-level effects that we observed—using experiments to estimate responses—suggests budburst cues are not shaped by local conditions, at least at our study scale.

Community composition and interspecific variation in phenology

The variation across species in their responses to environmental cues has the potential to create large differences in species temporal niches and ecological roles. Species ranged from early to late budburst timing in our experimental conditions, spanning a similar period as observed in natural communities (Lechowicz, 1984; Richardson, A.D., O’Keefe, 2009). This suggests our experiment captured a realistic breadth in temperate forest budburst. Furthermore, all our focal species responded to each environmental cue, with large responses to chilling and comparatively small responses to photoperiod—trends consistent with previous studies (Ettinger *et al.*, 2020; Flynn & Wolkovich, 2018). Cues, however, had complex interactions that appear potentially advantageous under warming climates (Heide, 1993; Caffarra & Donnelly, 2011; Flynn & Wolkovich, 2018). The interaction between chilling and forcing may ensure that species budburst even if warmer winters cause insufficient chilling, but will require additional forcing. But despite these differences in species responses to cues, we did not find clear, generalizable trends across functional groups.

Shrubs and trees differ greatly in their physiology, filling different ecological niches. Most of our trees budburst later than our study’s shrubs, a relative order also found by previous studies (Gill *et al.*, 1998; Panchen *et al.*, 2014). But about a quarter of our trees also budburst early, with timing more similar to that of shrubs. Similarly, we found a third of shrubs budburst at similar times as the majority of our trees. These deviations from our expectations for how shrubs and trees partition their budburst timing suggests there is more nuance to these patterns. As tree species advance phenologically—resulting in earlier canopy closure and reduced light availability (Donnelly & Yu, 2019)—this subset of shrubs with small cues may have reduced fitness. Our findings suggest, however, that many shrubs will also advance phenologically and have the potential to maintain their relative temporal niche.

Community assembly in responses to cues versus evolutionary history

Our study included the three cues—chilling, forcing and photoperiod—commonly thought to determine budburst timing, but about a third of the total variation was not explained by these cues (Fig. 3), suggesting the general model of budburst may be incomplete. This, paired with the observed phylogenetic structure in budburst timing, hints at unidentified latent traits still missing from our understanding of the processes shaping spring phenology (Webb *et al.*, 2002; Davies *et al.*, 2019).

Latent traits could reflect other cue or trait relationships indicative of species growth strategies or key traits that may facilitate greater resource use, or the greater competitive abilities needed when budbursting later (Grime, 1977). For example, species that budburst later in the season may require more conservative growth strategies, possessing traits that allow them to compete for limited soil resources and light, such as low specific leaf areas or higher wood densities (Wright *et al.*, 2004; Chave *et al.*, 2009). As such, we would also expect late budbursting shrubs or subcanopy trees to possess traits associated with greater light use efficiency or shade tolerance, allowing them to grow after canopy closure and lower light availability. But our results suggest that evolutionary history and species ancestral phenotype also have some effect on species timings and cues. Species traits are shaped over ‘deep’ timescales, with previous evolution also influencing responses to the present climate. Phenological trends could thus be an indication of limited evolutionary time to adapt (Lechowicz, 1984), or shaped by selection on other important traits that are not phylogenetically structured. Incorporating both phenology and evolutionary history into a broader trait framework could thus provide insights into traits that correlate with budburst timing, and—ultimately—the current and evolutionary drivers that select for species phenotypes and possibly shape species adaptive potential to future climates.

Predicting budburst under future climates

As climate change leads to greater temperature cues, we are already observing advances in species budburst within diverse ecological communities. The greatest ecological impacts will occur in high latitude communities that are warming the most and the fastest (Hoegh-Guldberg *et al.*, 2018), making species in northern communities most likely to advance their budburst and experience the greatest changes in budburst order. Yet our ability to predict these changes is often limited by the lack of data available. The consistent trends we observed across sites supports forecasting across similar regions with limited local phenological data but similar species assemblages, allowing us to leverage existing research to mitigate the cascading effects that shifted plant phenology could have on ecological communities.

While our findings have applications to plant communities, our analytical approach is widely applicable to forecasting diverse species responses to climate change. The phylogenetic model we used here could be easily applied to other phenological events or suites of species from across the tree of life. By combining species-rich and community-wide approaches with phylogenetic relationships we can better understand the impacts of evolutionary history and current ecological processes in shaping species phenology under a changing climate.

Acknowledgements

We thank M. Yasutake, K. Slimon, S. Larter, P. Autio, and S. Collins for their help running the Western part of this experiment; D. Flynn and T. Savas for contributions to the Eastern experimental data; S. Joly for his help constructing the phylogenetic tree used in our analysis; and to J. Hopps and K. Wilson for their support during our work at E. C. Manning Park and P. Crawford and D. Belford for their support while working in Babine Provincial Park.

375 **Competing Interests Statement**

376 The authors have no competing interests.

377

378 **Author Contribution Statement**

379 Both D.L. and E.M.W. conceived the study, contributed to the analysis and code and contributed to
380 the writing and revision of the manuscript. DL led field and experimental data collection.

381

382 **Data and code availability statement**

383 All data and Stan code are freely available through the Knowledge Network for Biocomplexity (Lough-
384 nan & Wolkovich, 2024b) and at the Harvard Forest data repository.

References

- Aitken, S.N. & Bemmels, J.B. (2016) Time to get moving: assisted gene flow of forest trees. *Evolutionary Applications* **9**, 271–290.
- Alecrim, E.F., Sargent, R.D. & Forrest, J.R.K. (2023) Higher-latitude spring-flowering herbs advance their phenology more than trees with warming temperatures. *Journal of Ecology* **111**, 156–169.
- Augspurger, C.K. (2009) Spring 2007 warmth and frost : phenology , damage and refoliation in a temperate deciduous forest. *Functional Ecology* **23**, 1031–1039.
- Basler, D. & Körner, C. (2014) Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* **34**, 377–388.
- Buonaiuto, D.M., Wolkovich, E.M. & Donahue, M.J. (2023) Experimental designs for testing the interactive effects of temperature and light in ecology : The problem of periodicity. *Functional Ecology* **37**, 1747–1756.
- Caffarra, A. & Donnelly, A. (2011) The ecological significance of phenology in four different tree species: Effects of light and temperature on bud burst. *International Journal of Biometeorology* **55**, 711–721.
- Chave, J., Coomes, D., Jansen, S., Lewis, S.L., Swenson, N.G. & Zanne, A.E. (2009) Towards a worldwide wood economics spectrum. *Ecology Letters* **12**, 351–366.
- Chesson, P., Gebauer, R.L.E., Schwinning, S., Huntly, N., Wiegand, K., Ernest, M.S.K., Sher, A., Novoplansky, A. & Weltzin, J.F. (2004) Resource pulses, species interactions, and diversity maintenance in arid and semi-arid environments. *Oecologia* **141**, 236–253.
- Chuine, I., Morin, X. & Bugmann, H. (2010) Warming, photoperiods, and tree phenology. *Science* **329**, 277–278.
- Cleland, E.E., Allen, J.M., Crimmins, T.M., Dunne, J.A., Pau, S., Travers, S.E., Zavaleta, E.S. & Wolkovich, E.M. (2012) Phenological tracking enables positive species responses to climate change. *Ecology* **93**, 1765–1771.
- Cohen, J.M., Lajeunesse, M.J. & Rohr, J.R. (2018) A global synthesis of animal phenological responses to climate change. *Nature Climate Change* **8**, 224–228.
- Cooke, J.E., Eriksson, M.E. & Junttila, O. (2012) The dynamic nature of bud dormancy in trees: Environmental control and molecular mechanisms. *Plant, Cell and Environment* **35**, 1707–1728.
- Davies, T.J., Regetz, J., Wolkovich, E.M. & McGill, B.J. (2019) Phylogenetically weighted regression: A method for modelling non-stationarity on evolutionary trees. *Global Ecology and Biogeography* **28**, 275–285.
- Davies, T.J., Wolkovich, E.M., Kraft, N.J.B., Salamin, N., Allen, J.M., Ault, T.R., Betancourt, J.L., Bolmgren, K., Cleland, E.E., Cook, B.I., Crimmins, T.M., Mazer, S.J., McCabe, G.J., Pau, S., Regetz, J., Schwartz, M.D. & Travers, S.E. (2013) Phylogenetic conservatism in plant phenology. *Journal of Ecology* **101**, 1520–1530.

420 Deans, J. (1996) Frost hardiness of 16 European provenances of sessile oak growing in Scotland. *Forestry*
421 **69**, 5–12.

422 Donnelly, A. & Yu, R. (2019) Temperate deciduous shrub phenology: the overlooked forest layer.
423 *International Journal of Biometeorology* **65**, 343–355.

424 Ettinger, A.K., Chamberlain, C.J., Morales-Castilla, I., Buonaiuto, D.M., Flynn, D.F., Savas, T.,
425 Samaha, J.A. & Wolkovich, E.M. (2020) Winter temperatures predominate in spring phenological
426 responses to warming. *Nature Climate Change* **10**, 1137–1142.

427 Finn, G.A., Straszewski, A.E. & Peterson, V. (2007) A general growth stage key for describing trees
428 and woody plants. *Annals of Applied Biology* **151**, 127–131.

429 Fitter, A.H. & Fitter, R.S. (2002) Rapid changes in flowering time in British plants. *Science* **296**,
430 1689–1691.

431 Flynn, D.F.B. & Wolkovich, E.M. (2018) Temperature and photoperiod drive spring phenology across
432 all species in a temperate forest community. *New Phytologist* **219**, 1353–1362.

433 Fridley, J.D. (2012) Extended leaf phenology and the autumn niche in deciduous forest invasions.
434 *Nature* **485**, 359–362.

435 Gelman, A. (2008) Scaling regression inputs by dividing by two standard deviations. *Statistics in*
436 *Medicine* **27**, 2865–2873.

437 Gill, D.S., Amthor, J.S. & Bormann, F.H. (1998) Leaf phenology, photosynthesis, and the persistence
438 of saplings and shrubs in a mature northern hardwood forest. *Tree Physiology* **18**, 281–289.

439 Gotelli, N.J. & Graves, G.R. (1996) The temporal niche. *Null Models In Ecology*, pp. 95–112, Smith-
440 sonian Institution Press, Washington, D. C.

441 Gougherty, A.V. & Gougherty, S.W. (2018) Sequence of flower and leaf emergence in deciduous trees
442 is linked to ecological traits, phylogenetics, and climate. *New Phytologist* **220**, 121–131.

443 Grime, J.P. (1977) Evidence for the Existence of Three Primary Strategies in Plants and Its Relevance
444 to Ecological and Evolutionary Theory. *The American Naturalist* **111**, 1169–1194.

445 Heberling, J.M., Cassidy, S.T., Fridley, J.D. & Kalisz, S. (2019) Carbon gain phenologies of spring-
446 flowering perennials in a deciduous forest indicate a novel niche for a widespread invader. *New*
447 *Phytologist* **221**, 778–788.

448 Heide, O.M. (1993) Daylength and thermal time responses of budburst during dormancy release in
449 some northern deciduous trees. *Physiologia Plantarum* **88**, 531–540.

450 Hoegh-Guldberg, O., Jacob, D., Taylor, M., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante,
451 R., Ebi, K., Engelbrecht, F., Guiot, J., Hijioka, Y., Mehrotra, S., Payne, A., Seneviratne, S., Thomas,
452 A., Warren, R. & Zhou, G. (2018) Impacts of 1.5°C Global Warming on Natural and Human Systems.
453 In: Global Warming of 1.5 °C. An IPCC Special Report on the impacts of global warming of 1.5 °C

above pre-industrial levels and related global greenhouse gas emission pathways, in the context of .
Tech. rep., Cambridge University Press, Cambridge, UK and New York, NY, USA.

Keller, S.R., Soolanayakanahally, R.Y., Guy, R.D., Silim, S.N., Olson, M.S. & Tiffin, P. (2011) Climate-driven local adaptation of ecophysiology and phenology in balsam poplar, *Populus balsamifera* L. (Salicaceae). *American Journal of Botany* **98**, 99–108.

Laube, J., Sparks, T.H., Estrella, N., Höfler, J., Ankerst, D.P. & Menzel, A. (2014) Chilling outweighs photoperiod in preventing precocious spring development. *Global Change Biology* **20**, 170–182.

Lechowicz, M.J. (1984) Why Do Temperate Deciduous Trees Leaf Out at Different Times? Adaptation and Ecology of Forest Communities. *The American Naturalist* **124**, 821–842.

Lee, B.R. & Ibáñez, I. (2021) Spring phenological escape is critical for the survival of temperate tree seedlings. *Functional Ecology* **35**, 1848–1861.

Loughnan, D. & Wolkovich, E.M. (2024a) A photographic guide to spring phenology for woody plants in Western North America. *Knowledge Network for Biocomplexity (KNB)* .

Loughnan, D. & Wolkovich, E.M. (2024b) Current environments and evolutionary history shape forest temporal assembly . *Knowledge Network for Biocomplexity (KNB)* .

Luedeling, E. (2012) Climate change impacts on winter chill for temperate fruit and nut production: A review. *Scientia Horticulturae* **144**, 218–229.

Luedeling, E. (2020) *chillR: Statistical Methods for Phenology Analysis in Temperate Fruit Trees*. <https://CRAN.R-project.org/package=chillR>.

Mahall, B.E. & Bormann, F.H. (1978) A Quantitative Description of the Vegetative Phenology of Herbs in a Northern Hardwood Forest. *Botanical Gazette* **139**, 467–481.

Menzel, A., Sparks, T.H., Estrella, N., Koch, E., Aaasa, A., Ahas, R., Alm-Kübler, K., Bissolli, P., Braslavská, O., Briede, A., Chmielewski, F.M., Crepinsek, Z., Curnel, Y., Dahl, Å., Defila, C., Donnelly, A., Filella, Y., Jatczak, K., Måge, F., Mestre, A., Nordli, Ø., Peñuelas, J., Pirinen, P., Remišová, V., Scheffinger, H., Striz, M., Susnik, A., Van Vliet, A.J., Wielgolaski, F.E., Zach, S. & Züst, A. (2006) European phenological response to climate change matches the warming pattern. *Global Change Biology* **12**, 1969–1976.

Morales-Castilla, I., Davies, T.J., Legault, G., Buonaiuto, D.M., Chamberlain, C.J., Ettinger, A.K., Garner, M., Jones, F.A.M., Loughnan, D., Pearse, W.D., Sodhi, D.S. & Wolkovich, E.M. (2024) Phylogenetic estimates of species-level phenology improve ecological forecasting. *Nature Climate Change* **14**, 989–995.

Muller, R.N. (1978) The Phenology, Growth and Ecosystem Dynamics of *Erythronium americanum* in the Northern Hardwood Forest. *Ecological Monographs* **48**, 1–20.

Natural Resources Canada (2020) Forestry glossary: Sapling.

Pagel, M. (1999) Inferring the historical patterns of biological evolution. *Nature* **401**, 877–884.

489 Panchen, Z.A., Primack, R.B., Nordt, B., Ellwood, E.R., Stevens, A., Renner, S.S., Willis, C.G.,
490 Fahey, R., Whittmore, A., Du, Y. & Davis, C.C. (2014) Leaf out times of temperate woody plants
491 are related to phylogeny, deciduousness, growth habit and wood anatomy. *New Phytologist* **203**,
492 1208–1219.

493 Parmesan, C. (2007) Influences of species, latitudes and methodologies on estimates of phenological
494 response to global warming. *Global Change Biology* **13**, 1860–1872.

495 Pau, S., Wolkovich, E.M., Cook, B.I., Davies, T.J., Kraft, N.J.B., Bolmgren, K., Betancourt, J.L.
496 & Cleland, E.E. (2011) Predicting phenology by integrating ecology, evolution and climate science.
497 *Global Change Biology* **17**, 3633–3643.

498 Pearse, W.D., Davies, T.J. & Wolkovich, E.M. (2025) How to define, use, and Interpret Pagel’s lambda
499 in Ecology and Evolution. *Global Ecology and Biogeography* **34**, e70012.

500 Polgar, C.A. & Primack, R.B. (2011) Leaf-out phenology of temperate woody plants: From trees to
501 ecosystems. *New Phytologist* **191**, 926–941.

502 Post, E., Steinman, B.A. & Mann, M.E. (2018) Acceleration of phenological advance and warming
503 with latitude over the past century. *Scientific Reports* **8**, 1–8.

504 R Development Core Team (2017) R: A language and environment for statistical computing.

505 Richardson, A.D., O’Keefe, J. (2009) Phenological Differences Between Understory and Overstory.
506 *Phenology of Ecosystem Processes* (ed. A. Noormets), pp. 87–117, Springer US, New York, NY.

507 Rudolf, V.H.W. (2019) The role of seasonal timing and phenological shifts for species coexistence.
508 *Ecology Letters* **22**, 1324–1338.

509 Sakai, A. & Larcher, W. (1987) *Frost Survival of Plants: Responses and adaptation to freezing stress*.
510 Springer-Verlag, Berlin, Heidelberg.

511 Savas, T., Flynn, D.F.B. & Wolkovich, E.M. (2017) A standardized photographic guide to woody plant
512 spring phenology. *Knowledge Network for Biocomplexity (KNB)* .

513 Smith, S.A. & Brown, J.W. (2018) Constructing a broadly inclusive seed plant phylogeny. *American*
514 *journal of botany* **105**, 302–314.

515 Sogaard, G., Johnsen, O., Nilsen, J. & Junttila, O. (2008) Climatic control of bud burst in young
516 seedlings of nine provenances of Norway spruce. *Tree Physiology* **28**, 311–320.

517 Stan Development Team (2018) RStan: the R interface to Stan. R package version 2.17.3.

518 Thackeray, S.J., Henrys, P.A., Hemming, D., Bell, J.R., Botham, M.S., Burthe, S., Helaouet, P.,
519 Johns, D.G., Jones, I.D., Leech, D.I., MacKay, E.B., Massimino, D., Atkinson, S., Bacon, P.J.,
520 Brereton, T.M., Carvalho, L., Clutton-Brock, T.H., Duck, C., Edwards, M., Elliott, J.M., Hall, S.J.,
521 Harrington, R., Pearce-Higgins, J.W., Høye, T.T., Kruuk, L.E., Pemberton, J.M., Sparks, T.H.,
522 Thompson, P.M., White, I., Winfield, I.J. & Wanless, S. (2016) Phenological sensitivity to climate
523 across taxa and trophic levels. *Nature* **535**, 241–245.

- 524 Tiusanen, M., Kankaanpää, T., Schmidt, N.S. & Roslin, T. (2020) Heated rivalries: Phenological
525 variation modifies competition for pollinators among arctic plants. *Global Change Biology* **26**, 6313–
526 6325.
- 527 Vitasse, Y., Basler, D. & Way, D. (2014) Is the use of cuttings a good proxy to explore phenological
528 responses of temperate forests in warming and photoperiod experiments? *Tree Physiology* **34**, 174–
529 183.
- 530 Vitasse, Y., Delzon, S., Dufrene, E., Pontailleur, J.Y., Louvet, J.M., Kremer, A. & Michalet, R. (2009)
531 Leaf phenology sensitivity to temperature in European trees : Do within-species populations exhibit
532 similar responses? *Agricultural and Forest Meteorology* **149**, 735–744.
- 533 Vitasse, Y., Signarbieux, C. & Fu, Y.H. (2018) Global warming leads to more uniform spring phenology
534 across elevations. *PNAS* **115**, 1004–1008.
- 535 Webb, C.O., Ackerly, D.D., McPeck, M.A. & Donoghue, M.J. (2002) Phylogenies and community
536 ecology. *Annual Review of Ecology Evolution and Systematics* **33**, 475–505.
- 537 Wiens, J.J., Ackerly, D.D., Allen, A.P., Anacker, B.L., Buckley, L.B., Cornell, H.V., Damschen, E.I.,
538 Jonathan Davies, T., Grytnes, J.A., Harrison, S.P. *et al.* (2010) Niche conservatism as an emerging
539 principle in ecology and conservation biology. *Ecology Letters* **13**, 1310–1324.
- 540 Wolkovich, E.M. & Cleland, E.E. (2014) Phenological niches and the future of invaded ecosystems
541 with climate change. *AoB PLANTS* **6**, 1–16.
- 542 Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F., Cavender-Bares, J.,
543 Chapin, T., Cornelissen, J.H.C., Diemer, M., Flexas, J., Garnier, E., Groom, P.K., Gulias, J.,
544 Hikosaka, K., Lamont, B.B., Lee, T., Lee, W., Lusk, C., Midgley, J.J., Navas, M.L., Niinemets, Ü.,
545 Oleksyn, J., Osada, N., Poorter, H., Poot, P., Prior, L., Pyankov, V.I., Roumet, C., Thomas, S.C.,
546 Tjoelker, M.G., Veneklaas, E.J. & Villar, R. (2004) The worldwide leaf economics spectrum. *Nature*
547 **428**, 821–827.
- 548 Yu, H., Luedeling, E. & Xu, J. (2010) Winter and spring warming result in delayed spring phenology
549 on the Tibetan Plateau. *Proceedings of the National Academy of Sciences of the United States of*
550 *America* **107**, 22151–22156.
- 551 Zeng, Z.A. & Wolkovich, E.M. (2024) Weak evidence of provenance effects in spring phenology across
552 Europe and North America. *New Phytologist* **242**, 1957–1964.
- 553 Zohner, C.M. & Renner, S.S. (2014) Common garden comparison of the leaf-out phenology of woody
554 species from different native climates, combined with herbarium records, forecasts long-term change.
555 *Ecology Letters* **17**, 1016–1025.




Figure 1: We combined data from two controlled environment studies of temperature and photoperiod cues for budburst in temperate woody species **a**, from two forest communities in eastern North America that span a 4° latitudinal gradient and **b**, two communities in western North America that span a 6° latitudinal gradient. Combined, our dataset includes 28 tree and shrub species from eastern communities and 21 species from western communities, with three species occurring across all four populations. Both studies used a full-factorial design with two levels each of chilling treatments (consisting of a combination of field and different duration of chamber chilling), forcing and photoperiod, for a total of eight unique treatments. Observations of budburst were made during the forcing treatment, using the BBCH scale to classify stages of bud development (see Savas *et al.*, 2017; Loughnan & Wolkovich, 2024a, for guides we made for these species).




Figure 2: **a**, Posterior distributions of estimated chilling, forcing, and photoperiod responses with individual site-level effects. Symbols denote the eastern and western transects and the darker point color our more northern communities. Points represent the mean, thicker lines the 50% uncertainty interval, and the thin lines the 90% uncertainty interval. **b**, Cues interacted to produce a subadditive effect, where delays in budburst under low chilling was offset by high forcing conditions, and vice versa. To make results comparable across predictors we standardized chilling, forcing, and photoperiod via z -scores using two standard deviations, see methods for further details.




Figure 3: Estimated species **a,b** chilling, **c,d** forcing, and **e,f** photoperiod responses ordered by estimated budburst dates for both the **a,c,e**, eastern and **b,d,f** western sites based on our model estimates. Cues are plotted on differing y-axis scales to better depict species differences across cues. For each species, the points represent the mean, thinner error bars represent the 90% uncertainty interval, and thicker bars represent the 50% uncertainty interval. The three species that occur in both the eastern and western transects are depicted using darker colours. To make the results across cues directly comparable, we standardized predictors (here, chilling, forcing, and photoperiod) via z -scores using two standard deviations, see methods for further details.




Figure 4: Comparisons of estimated day of budburst based on the full model (intercept plus all cues), shown as white (open) symbols, versus the intercepts only (without effects of chilling, forcing, and photoperiod), shown as black (filled) symbols. Points are ordered according to differences in species estimated budburst dates for both the **a**, eastern **b**, and western sites under low cue conditions, depicted as circles, and high cue conditions, depicted as triangles. The abbreviated names of tree species are shown in red.




Figure 5: Species differences were accounted for by including phylogenetic effects on the species intercept in a model estimating days to budburst after the start of forcing treatments. We pruned to our species subset an existing phylogeny for flowering plants developed by Smith and Brown (2018).