
Definition

Given:


ightharpoonup a manifold  $\mathcal{M}$ ;



#### Definition

#### Given:

- $\triangleright$  a manifold  $\mathcal{M}$ ;
- ▶ a Morse function  $f: \mathcal{M} \to \mathbb{R}$  with distinct critical values;




#### Definition

#### Given:

- $\triangleright$  a manifold  $\mathcal{M}$ ;
- ▶ a Morse function  $f: \mathcal{M} \to \mathbb{R}$  with distinct critical values; the **Reeb graph** of f is the 1-dimensional simplicial complex

$$\mathcal{R}(f) = \mathcal{M}/\sim$$
.




#### Definition

#### Given:

- $\triangleright$  a manifold  $\mathcal{M}$ ;
- ▶ a Morse function  $f: \mathcal{M} \to \mathbb{R}$  with distinct critical values; the **Reeb graph** of f is the 1-dimensional simplicial complex

$$\mathcal{R}(f) = \mathcal{M}(f)$$

$$x \sim y \text{ if } f(x) = f(y) \text{ and they belong to the same connected component of } f^{-1}(f(x))$$




#### Definition

#### Given:

- ightharpoonup a manifold  $\mathcal{M}$ ;
- ▶ a Morse function  $f: \mathcal{M} \to \mathbb{R}$  with distinct critical values; the **Reeb graph** of f is the 1-dimensional simplicial complex

$$\mathcal{R}(f) = \mathcal{M}/\sim$$
.


The **segmentation map** is the quotient map



Desired algorithm

### Input:

ightharpoonup a PL manifold  ${\cal M}$ 



Desired algorithm

### Input:

a PL manifold M
 → a triangulated mesh M;



Desired algorithm

### Input:

- a PL manifold M
   → a triangulated mesh M;
- ightharpoonup a non-degenerate PL scalar field f on  ${\mathcal M}$



Desired algorithm

### Input:

- ightharpoonup a PL manifold  ${\cal M}$ 
  - $\rightsquigarrow$  a triangulated mesh  $\mathcal{M}$ ;
- ightharpoonup a non-degenerate PL scalar field f on  ${\cal M}$ 
  - $\rightsquigarrow$  a scalar value f(v) for each vertex v of  $\mathcal{M}$ .
    - pairwise different, in order to ensure non-degeneracy; this can be achieved by random perturbations



Desired algorithm

### Input:

- a PL manifold M
   → a triangulated mesh M;
- ▶ a non-degenerate PL scalar field f on  $\mathcal{M}$   $\rightsquigarrow$  a scalar value f(v) for each vertex v of  $\mathcal{M}$ .

### Output:

▶ the augmented Reeb graph  $\mathcal{R}(f)$ .

► graph + segmentation map



Desired algorithm

### Input:

- a PL manifold M
   → a triangulated mesh M;
- a non-degenerate PL scalar field f on M
   → a scalar value f(v) for each vertex v of M.



### **Output:**

▶ the augmented Reeb graph  $\mathcal{R}(f)$ .

### Time complexity:

 $\triangleright$   $O(m \cdot \log m)$ , where m is the size of the 2-skeleton of  $\mathcal{M}$ .

#vertices + #edges + #triangles

Desired algorithm

### Input:

- a PL manifold M
   → a triangulated mesh M;
- a non-degenerate PL scalar field f on M
   → a scalar value f(v) for each vertex v of M.

### Output:

▶ the augmented Reeb graph  $\mathcal{R}(f)$ .

### Time complexity:

 $ightharpoonup O(m \cdot \log m)$ , where m is the size of the 2-skeleton of  $\mathcal{M}$ .

#### Parallel.

Geometry of critical points

There are three kinds of critical points:

Geometry of critical points

There are three kinds of critical points:

▶ (local) maxima



#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
- ► (local) minima





#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
- ▶ (local) minima
- saddles







#### Geometry of critical points

There are three kinds of critical points:

► (local) maxima

---


► (local) minima



saddles



How to detect them on a PL manifold?



#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
- ► (local) minima
- saddles








### How to detect them on a PL manifold?

Given a vertex v, the star of v is the union of all simplices containing v.



#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
- ► (local) minima
- saddles








### How to detect them on a PL manifold?

Given a vertex v, the star of v is the union of all simplices containing v.



#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
- ► (local) minima
- saddles








### How to detect them on a PL manifold?

Given a vertex v, the star of v is the union of all simplices containing v.

$$Link^{+}(v) = \{x \in Link(v) : f(x) > f(v)\}$$
  

$$Link^{-}(v) = \{x \in Link(v) : f(x) < f(v)\}$$



#### Geometry of critical points

There are three kinds of critical points:

- (local) minima
- saddles








### How to detect them on a PL manifold?

Given a vertex v, the star of v is the union of all simplices containing v.

$$Link^{+}(v) = \{x \in Link(v) : f(x) > f(v)\}$$
  
$$Link^{-}(v) = \{x \in Link(v) : f(x) < f(v)\}$$



#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
  ~> Link<sup>+</sup> empty;
- ► (local) minima ~> Link<sup>-</sup> empty;
- saddles








### How to detect them on a PL manifold?

Given a vertex v, the star of v is the union of all simplices containing v.

$$Link^{+}(v) = \{x \in Link(v) : f(x) > f(v)\}$$
  
$$Link^{-}(v) = \{x \in Link(v) : f(x) < f(v)\}$$



#### Geometry of critical points

There are three kinds of critical points:

- ► (local) maxima
  ~> Link<sup>+</sup> empty;








### How to detect them on a PL manifold?


Given a vertex v, the star of v is the union of all simplices containing v.

$$\mathsf{Link}^+(v) = \{ x \in \mathsf{Link}(v) : f(x) > f(v) \}$$
  
 $\mathsf{Link}^-(v) = \{ x \in \mathsf{Link}(v) : f(x) < f(v) \}$ 



Significance of critical points


The critical points of f are closely related to the topology of the Reeb graph  $\mathcal{R}(f)$ .



Significance of critical points

The critical points of f are closely related to the topology of the Reeb graph  $\mathcal{R}(f)$ .


► Maxima and minima  $\rightsquigarrow$  nodes of valence 1 (leaves).



#### Significance of critical points

The critical points of f are closely related to the topology of the Reeb graph  $\mathcal{R}(f)$ .


- Maxima and minima → nodes of valence 1 (leaves).
- ▶ **Saddles**  $\rightsquigarrow$  nodes of valence  $\geq 2$ .



#### Significance of critical points

The critical points of f are closely related to the topology of the Reeb graph  $\mathcal{R}(f)$ .

- Maxima and minima → nodes of valence 1 (leaves).
- ▶ **Saddles**  $\rightsquigarrow$  nodes of valence  $\geq 2$ .
  - ▶ Join saddles: multiple components below.
  - ► Split saddles: multiple components above.



#### Significance of critical points

The critical points of f are closely related to the topology of the Reeb graph  $\mathcal{R}(f)$ .

- Maxima and minima → nodes of valence 1 (leaves).
- ▶ **Saddles**  $\rightsquigarrow$  nodes of valence  $\geq 2$ .
  - ▶ Join saddles: multiple components below.
  - ► Split saddles: multiple components above.

non-mutually exclusive in dimension  $\geq 3$ 

