



Sign In

SIGN IN

[Forgot Password](#)

Don't have an account yet?

SIGN UP

Sign Up

Sign Up

Already have an account? [Sign In](#)



Sign In

SIGN IN

[Forgot Password](#)

Don't have an account yet?

SIGN UP

Sign Up

SIGN UP

Already have an account? [Sign In](#)

Home

My Drug Cabinet

Settings



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Motrin IB PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



My Profile



Andy Osnard

Male, 35
Washington, DC
andy.osnard@domain.com



My Family

Add +

	Name	Age	Gender	Relationship	Pregnant?	
	Olivia	32	Female	Spouse	Yes (MM/DD)	Update
	Hugo	6	Male	Son	...	Update
	Lily	3	Female	Daughter	...	Update

Sign-in and Security

Username: andyo [Change](#)

Password: ***** [Update](#)



My Profile



Andy Osnard



Male, 35
Washington, DC
andy.osnard@domain.com

My Family

Add +



Olivia

Update

Age 32
Gender Female
Relationship Spouse
Pregnant Yes



Hugo

Update

Age 6
Gender Male
Relationship Son
Pregnant N/A



Lily

Update

Age 3
Gender Female
Relationship Daughter
Pregnant N/A

My Profile



My Family



Olivia



Hugo



Lily

Sign-in and Settings

Sign-in Username

Password: *****

Family Member

First Name

Olivia

Gender

Female



Age

32



Relationship

Spouse



Is this person pregnant?

Yes



Due Date

MM/DD/YYYY

Delete

Done

My Drug Cabinet



Filter by:

All

Search





Advil PM

Expiration: MM/DD/YYYY





Advil PM

Expiration: MM/DD/YYYY





Advil PM

Expiration: MM/DD/YYYY






Advil PM

Expiration: MM/DD/YYYY





Motrin IB PM

Expiration: MM/DD/YYYY





Advil PM

Expiration: MM/DD/YYYY





Advil PM

Expiration: MM/DD/YYYY





Advil PM

Expiration: MM/DD/YYYY





Advil PM

Expiration: MM/DD/YYYY





My Drug Cabinet

Search

Filter by:

All



Advil PM

Expiration: MM/DD/YYYY



Advil PM

- Uses
- Warnings
- Recalls

[Remove from my cabinet](#)



Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Advil PM

Uses

Warnings

Recalls

[Remove from my cabinet](#)

Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

What should I generally be aware of?

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug When using this product drowsiness will occur avoid alcoholic drinks do not drive a motor vehicle or operate machinery take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

When should I stop taking it?

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear.

When should I consult my doctor?

Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland.

When should I consult my doctor and/or pharmacist?

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug.

When should I avoid taking it?

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain.

What if I am pregnant or breastfeeding?

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.



Add Drug

Search (brand, drug Name, active ingredient, etc)

Cancel

Add Drug

Advil

- Advil Allergy Sinus Oral
- Advil Congestion Relief oral
- Advil Liqui-Gel Oral
- Advil Migraine Oral
- Advil Multi-Symptom Cold Oral
- Advil Oral
- Advil PM Oral
- Children's Advil Allergy Sinus Oral

Cancel

◀ Back to List

Advil PM Oral

Uses Side Effects Recalls

Nullam id dolor id nibh ultricies vehicula ut id elit. Donec id elit non mi porta gravida at eget metus. Vestibulum id ligula porta felis euismod semper. Sed posuere consectetur est at lobortis. Integer posuere erat a ante venenatis dapibus posuere velit aliquet.

Add to my Drug Cabinet



Advil PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY

Motrin IB PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY

Advil PM

Expiration: MM/DD/YYYY



Advil PM



Motrin IB PM