



Sign In

SIGN IN

[Forgot Password](#)

Don't have an account yet?

[SIGN UP](#)

Sign Up

Sign Up

Already have an account? [Sign In](#)



Sign In

SIGN IN

[Forgot Password](#)

Don't have an account yet?

[SIGN UP](#)

Sign Up

SIGN UP

Already have an account? [Sign In](#)

Welcome to Rex!

Thank you for signing up. To take full advantage of the features that ReX has to offer, please complete your profile and add your family members. This information will better help us in providing you warnings and recalls about drugs in your virtual drug cabinet.

Go to My Profile

My Profile



Andy



Male, 35
Washington, DC
andy@domain.com

[Change Password](#)

My Family

[Add +](#)

| Name | | Age | Gender | Pregnant? | |
|------|--------|-----|--------|-------------|--|
| | Olivia | 32 | Female | Yes (MM/DD) | |
| | Hugo | 6 | Male | ... | |
| | Lily | 3 | Female | ... | |

My Profile



Update Profile

First Name

Andy

Gender

Male

Age

32

State

DC

City

Washington

Email

andy@domain.com







CANCEL

UPDATE

My Profile



My Family

| Name | | |
|---|--------|---|
|  | Olivia |  |
|  | Hugo |  |
|  | Lily |  |

Add +

Change Password

Current Password

New Password

Confirm New Password

CANCEL

UPDATE

My Profile



Andy



Male, 35
Washington, DC
andy@domain.com

[Change Password](#)

My Family

[Add +](#)



Olivia



| | |
|----------|--------|
| Age | 32 |
| Gender | Female |
| Pregnant | Yes |



Hugo



| | |
|----------|------|
| Age | 6 |
| Gender | Male |
| Pregnant | N/A |



Lily



| | |
|----------|--------|
| Age | 3 |
| Gender | Female |
| Pregnant | N/A |

My Profile



Family Member

First Name

Olivia

This is too short

Gender



Age

32



Is this person pregnant?



This is a conditional field (hidden unless pregnant question is yes)

Due Date

Delete

Done

My Family



Olivia



Hugo



Lily

Add +

Update

Update

Update

Sign-in and Settings

Sign-in Username

Update

Password: *****

Update

My Drug Cabinet



Filters

Search





Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY

Takes me
directly to
the Recall
tab for that
drug



Advil PM

Expiration: MM/DD/YYYY



Motrin IB PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY

My Drug Cabinet

Filters

| BRAND NAME | ACTIVE INGREDIENT | DRUG TYPE | PREGNANCY DRUG CLASS |
|------------|-------------------|------------------|---------------------------|
| Advil | Acetaminophen | Over-the-counter | Class A |
| Bayer | Aspirin | Prescription | Class B |
| Motrin | Ibuprofen | | Class C |
| | | | Class D |
| | | | Class X |
| | | | Clear All |

CANCEL

APPLY

My Drug Cabinet

Search

Filters



Floating

Advil PM

Expiration: MM/DD/YYYY



My Drug Cabinet

Filters

BRAND NAME

Advil

Bayer

Motrin

ACTIVE INGREDIENT

Acetaminophen

Aspirin

Ibuprofen

DRUG TYPE

Over-the-counter

Prescription

PREGNANCY DRUG CLASS

Class A

Class B

Class C

Class D

Class E

Advil PM

Expiration: MM/DD/YYYY

- Uses
- Warnings
- Recalls
- Pregnancy & Breast Feeding



Update Expiration Date

Remove from My Cabinet



Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Advil PM

- Uses
- Warnings
- Recalls
- Pregnancy

Expiration: MM/DD/YYYY



Update Expiration Date

Remove from My Cabinet



Pregnancy/Breastfeeding

This drug is...curabitur
non metus auctor fring
luctus, nisi erat porttito

Are you sure you want to remove this drug from you cabinet?

CANCEL

REMOVE

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Advil PM

- Uses
- Warnings
- Recalls
- Pregnancy

Expiration: MM/DD/YYYY



Update Expiration Date

Remove from My Cabinet



Pregnancy/Breastfeeding

This drug is...curabitur
non metus auctor fring
luctus, nisi erat porttito

a at eget metus. Donec ullamcorper nulla
t eros. Duis mollis, est non commodo

Expiration Date

MM/DD/YYYY

Date format is wrong

CANCEL

UPDATE

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Advil PM

Expiration: MM/DD/YYYY

Uses

Warnings

Recalls

Pi >



Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Donec id elit non mi porta gravida at eget metus. Integer posuere erat a ante venenatis dapibus posuere velit aliquet.

Advil PM

Expiration: MM/DD/YYYY

- Uses
- Warnings
- Recalls
- Pregnancy & Breast Feeding



Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

What should I generally be aware of?

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug When using this product drowsiness will occur avoid alcoholic drinks do not drive a motor vehicle or operate machinery take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

When should I stop taking it?

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear.

When should I consult my doctor?

Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland.

When should I consult my doctor and/or pharmacist?

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug.

When should I avoid taking it?

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain.

What if I am pregnant or breastfeeding?

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Codeine

Expiration: MM/DD/YYYY

- Uses
- Warnings
- Recalls
- Pregnancy & Breast Feeding



Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

What should I generally be aware of?

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug When using this product drowsiness will occur avoid alcoholic drinks do not drive a motor vehicle or operate machinery take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

When should I stop taking it?

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear.

When should I consult my doctor?

Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland.

When should I consult my doctor and/or pharmacist?

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug.

When should I avoid taking it?

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain.

What if I am pregnant or breastfeeding?

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.



Add Drug

Search (brand, drug Name, active ingredient, etc)

Cancel

Add Drug

Advil

- Advil Allergy Sinus Oral
- Advil Congestion Relief oral
- Advil Liqui-Gel Oral
- Advil Migraine Oral
- Advil Multi-Symptom Cold Oral
- Advil Oral
- Advil PM Oral
- Children's Advil Allergy Sinus Oral

Cancel

◀ Back to List

Advil PM Oral

Uses Warnings Recalls

Nullam id dolor id nibh ultricies vehicula ut id elit. Donec id elit non mi porta gravida at eget metus. Vestibulum id ligula porta felis euismod semper. Sed posuere consectetur est at lobortis. Integer posuere erat a ante venenatis dapibus posuere velit aliquet.

Add to my Drug Cabinet