



Sign In

SIGN IN

[Forgot Password](#)

Don't have an account yet?

SIGN UP

Sign Up

Sign Up

Already have an account? [Sign In](#)



Sign In

SIGN IN

[Forgot Password](#)

Don't have an account yet?

[SIGN UP](#)

Sign Up

SIGN UP

Already have an account? [Sign In](#)

Getting Started: Welcome!


Welcome to Rex!

Etiam porta sem malesuada magna mollis euismod. Etiam porta sem malesuada magna mollis euismod. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Donec ullamcorper nulla non metus auctor fringilla. Nullam quis risus eget urna mollis ornare vel eu leo.

Vivamus sagittis lacus vel augue laoreet rutrum faucibus dolor auctor. Sed posuere consectetur est at lobortis. Praesent commodo cursus magna, vel scelerisque nisl consectetur et. Donec ullamcorper nulla non metus auctor fringilla. Vivamus sagittis lacus vel augue laoreet rutrum faucibus dolor auctor. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus.



NEXT



Home

My Drug Cabinet

Settings



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Motrin IB PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



My Profile



Andy Osnard

Male, 35
Washington, DC
andy.osnard@domain.com



My Family

Add +

Name		Age	Gender	Pregnant?	
	Olivia	32	Female	Yes (MM/DD)	Update
	Hugo	6	Male	...	Update
	Lily	3	Female	...	Update

Sign-in and Security

Username: andyo [Change](#)

Password: ***** [Update](#)

My Profile



Andy Osnard



Male, 35
Washington, DC
andy.osnard@domain.com

My Family

[Add +](#)



Olivia

[Update](#)

Age	32
Gender	Female
Pregnant	Yes



Hugo

[Update](#)

Age	6
Gender	Male
Pregnant	N/A



Lily

[Update](#)

Age	3
Gender	Female
Pregnant	N/A

My Profile



My Family



Olivia



Hugo



Lily

Sign-in and Settings

Sign-in Username

Password: *****



Family Member

First Name

Olivia

This is too short

Gender



Age

32



Is this person pregnant?



Due Date

Delete

Done

This is a conditional field (hidden unless pregnant question is yes)

My Drug Cabinet



Filters

Search





Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY

Takes me directly to the Recall tab for that drug



Advil PM

Expiration: MM/DD/YYYY



Motrin IB PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY



Advil PM

Expiration: MM/DD/YYYY

My Drug Cabinet

Filters

BRAND NAME	ACTIVE INGREDIENT	DRUG TYPE	PREGNANCY DRUG CLASS
Advil	Acetaminophen	Over-the-counter	Class A
Bayer	Aspirin	Prescription	Class B
Motrin	Ibuprofen		Class C
			Class D
			Class X
			Clear All

CANCEL

APPLY

My Drug Cabinet

Search

Filters



Floating

Advil PM

Expiration: MM/DD/YYYY



My Drug Cabinet

Filters

BRAND NAME

Advil

Bayer

Motrin

ACTIVE INGREDIENT

Acetaminophen

Aspirin

Ibuprofen

DRUG TYPE

Over-the-counter

Prescription

PREGNANCY DRUG CLASS

Class A

Class B

Class C

Class D

Class E

Advil PM

- Uses
- Warnings
- Recalls
- Pregnancy

[Remove from my cabinet](#)



Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Advil PM

- Uses
- Warnings
- Recalls
- Pregnancy
- 

Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

Indications and Usage

Uses for relief of occasional sleeplessness when associated with minor aches and pains helps you fall asleep and stay asleep.

Dosage and Administration

Do not take more than directed. Adults and children 12 years and over: take 2 capsules at bedtime. Do not take more than 2 capsules in 24 hours

Purpose

Nighttime sleep-aid Pain reliever.

Donec id elit non mi porta gravida at eget metus. Integer posuere erat a ante venenatis dapibus posuere velit aliquet.

Advil PM

Uses

Warnings

Recalls

Pregnancy

[Remove from my cabinet](#)

Pregnancy/Breastfeeding Warning

This drug is....curabitur blandit tempus porttitor. Donec id elit non mi porta gravida at eget metus. Donec ullamcorper nulla non metus auctor fringilla. Morbi leo risus, porta ac consectetur ac, vestibulum at eros. Duis mollis, est non commodo luctus, nisi erat porttitor ligula, eget lacinia odio sem nec elit.

What should I generally be aware of?

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug When using this product drowsiness will occur avoid alcoholic drinks do not drive a motor vehicle or operate machinery take with food or milk if stomach upset occurs the risk of heart attack or stroke may increase if you use more than directed or for longer than directed Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

When should I stop taking it?

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better pain gets worse or lasts more than 10 days sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. redness or swelling is present in the painful area any new symptoms appear.

When should I consult my doctor?

Ask a doctor before use if stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma you are taking a diuretic you have a breathing problem such as emphysema or chronic bronchitis you have glaucoma you have trouble urinating due to an enlarged prostate gland.

When should I consult my doctor and/or pharmacist?

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness taking any other antihistamines taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug.

When should I avoid taking it?

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep in children under 12 years of age right before or after heart surgery with any other product containing diphenhydramine, even one used on skin if you have sleeplessness without pain.

What if I am pregnant or breastfeeding?

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.



Add Drug

Search (brand, drug Name, active ingredient, etc)

Cancel

Add Drug

Advil

- Advil Allergy Sinus Oral
- Advil Congestion Relief oral
- Advil Liqui-Gel Oral
- Advil Migraine Oral
- Advil Multi-Symptom Cold Oral
- Advil Oral
- Advil PM Oral
- Children's Advil Allergy Sinus Oral

Cancel

Back to List

Advil PM Oral

UsesWarningsRecalls

Nullam id dolor id nibh ultricies vehicula ut id elit. Donec id elit non mi porta gravida at eget metus. Vestibulum id ligula porta felis euismod semper. Sed posuere consectetur est at lobortis. Integer posuere erat a ante venenatis dapibus posuere velit aliquet.

Add to my Drug Cabinet