

ACM/IDS 104 APPLIED LINEAR ALGEBRA

PROBLEM SET 2

Please submit your solution as a [single PDF file](#), that contains both the written-up and published code parts, via [Gradescope](#) by **9pm Tuesday, October 17**. An example of the submission process is shown here: https://www.gradescope.com/get_started#student-submission

- For theoretical problems, please use a pen, not a pencil: it is hard to read scanned submission written by a pencil.
- For coding problems, please convert your MATLAB livescripts (.mlx) to PDF by selecting **Live Editor** → **Save** → **Export to PDF** and merge them with the rest of your solution.
- After uploading your submission to Gradescope, please label all pages.

Problem 1. (10 POINTS) SUBSPACES

Which of the following subsets $W \subset V$ are *subspaces* of the vector spaces V ? Explain why.

- (a) (2 points) $V = \mathbb{M}_{n \times n}$ and W is a subset of all singular matrices, $W = \{A \mid \det A = 0\}$.
 (b) (2 points) The *trace* of a square matrix $A \in V = \mathbb{M}_{n \times n}$ is the sum of its diagonal entries:

$$\text{tr} A = \sum_{i=1}^n a_{ii}. \quad (1)$$

Let W be a subset of all trace zero matrices, $W = \{A \mid \text{tr} A = 0\}$.

- (c) (2 points) Let $V = C([0, 1])$ be the vector space of all continuous functions on the interval $[0, 1]$, and $W = \{f(x) \in V \mid f(0)f(1) = 1\}$.
 (d) (2 points) The vector space V is again $C([0, 1])$. The subset $W \subset V$ consists of all functions satisfying the following condition:

$$f\left(\frac{1}{2}\right) = \int_0^1 f(t) dt. \quad (2)$$

- (e) (2 points) A *planar vector field* is a function which assigns a vector

$$v(x, y) = \begin{bmatrix} v_1(x, y) \\ v_2(x, y) \end{bmatrix} \quad (3)$$

to each point $(x, y) \in \mathbb{R}^2$. Let V be the vector space of all planar vector fields. A planar vector field is called *incompressible* if it has zero divergence:

$$\nabla \cdot v = \frac{\partial v_1}{\partial x} + \frac{\partial v_2}{\partial y} = 0. \quad (4)$$

Let $W \subset V$ be the subset of all incompressible vector fields.

Problem 2. (10 POINTS) POLYNOMIALS

Let $\mathcal{P}^{(2)}$ be the vector space of quadratic polynomials. Let

$$p_1(x) = x^2 - 3, \quad p_2(x) = 2 - x, \quad p_3(x) = (x + 1)^2. \quad (5)$$

- (a) (4 points) Determine whether p_1, p_2 and p_3 are linearly independent or not.
 (b) (2 points) Do they span $\mathcal{P}^{(2)}$?
 (c) (4 points) Do they form a basis of $\mathcal{P}^{(2)}$? If yes, find the coordinates of $q(x) = 1$ in that basis.

Problem 3. (10 POINTS) FIBONACCI SEQUENCES

When learning abstract concepts, such as vector spaces, it is handy to have some weird examples at hand that illustrate the concept. These examples will a) develop you abstract thinking and b) motivate

you to look for vector spaces in other examples that you will encounter in your study or research¹. In lectures, we discussed one such example: $(Q, \oplus, *)$. Here is another one.

The Fibonacci sequence is a famous² sequence in mathematics and engineering:

$$\mathbf{f}^* = (1, 1, 2, 3, 5, 8, 13, \dots). \quad (6)$$

The first two numbers of the sequence are $x_1 = x_2 = 1$ and each subsequent number is the sum of the previous two:

$$x_n = x_{n-1} + x_{n-2}, \quad n = 3, \dots \quad (7)$$

The sequence \mathbf{f}^* in (6) is the original Fibonacci sequence. Let us define a generalized Fibonacci sequence \mathbf{f} as follows:

$$\begin{aligned} \mathbf{f} &= (x_1, x_2, x_3, \dots) \\ x_1 \text{ and } x_2 &\text{ are arbitrary real numbers (not necessarily 1)} \\ x_n &= x_{n-1} + x_{n-2}, \quad n = 3, \dots \end{aligned} \quad (8)$$

That is, in a generalized Fibonacci sequence we can start from any first two numbers. Let \mathcal{F} denote the set of all generalized Fibonacci sequences. Let's define two natural operations on \mathcal{F} :

$$\begin{aligned} \text{Addition: } \mathbf{f}_1 &= (x_1, x_2, \dots), \quad \mathbf{f}_2 = (y_1, y_2, \dots), \quad \mathbf{f}_1 + \mathbf{f}_2 = (x_1 + y_1, x_2 + y_2, \dots) \\ \text{Scalar multiplication: } \mathbf{f} &= (x_1, x_2, \dots), \quad \alpha \in \mathbb{R}, \quad \alpha \mathbf{f} = (\alpha x_1, \alpha x_2, \dots). \end{aligned} \quad (9)$$

- (a) (4 points) Show that the set of generalized Fibonacci sequences \mathcal{F} is a vector space.
- (b) (4 points) Find its dimension and a basis.
- (c) (2 points) Find the coordinates of the original sequence \mathbf{f}^* in that basis.

Problem 4. (10 POINTS) KERNEL, IMAGE, COKERNEL, COIMAGE

Find bases for the kernel, image, cokernel, and coimage of the following matrix:

$$A = \begin{bmatrix} 1 & 2 & \dots & n \\ n+1 & n+2 & \dots & 2n \\ \vdots & \vdots & & \vdots \\ n^2 - n + 1 & n^2 - n + 2 & \dots & n^2 \end{bmatrix} \quad (10)$$

Hint: The proof of the Fundamental Theorem of Linear Algebra (see notes, pages 20-21) may be helpful, but not necessary.

Problem 5. (10 POINTS) FUNDAMENTAL MATRIX SUBSPACES.

Complete Problem 5 in PS2.mlx.

¹And as soon as you identify that a set is actually a vector space, you can employ the powerful machinery of linear algebra to attack the underlying problem.

²Read https://en.wikipedia.org/wiki/Fibonacci_number for information and watch <https://youtu.be/02wU-HT7FiM> for inspiration.