

Is Gravity Entropic?

Numerical Verification of the Entropic Gravity Hypothesis
on Causal Diamond Lattices

Ryan Osacra

Murrieta, California

*Framework: the Osacra–Gravaser unified theory;
numerical evidence: the CEFÉ (Causal Entropic Field Engine) lattice*

August 2026

Abstract

We state the minimal axioms of entropic gravity — holographic area scaling of information, equipartition on screens, and a universal entropy gradient — and subject every testable consequence to numerical measurement on a causal-diamond lattice simulation of a free real scalar field (the CEFÉ engine, an exact implementation of the Srednicki covariance-matrix method). We find: (i) the entanglement entropy obeys an area law with exponent $\alpha = 2.09 \pm 0.04$ and coefficient $\kappa = 0.314$, matching Srednicki’s 1993 result $\kappa = 0.30$; (ii) driven by the *measured* entropy alone, the Verlinde equipartition chain produces an inverse-square force with exponent -2.07 , with an $O(10)$ renormalization of Newton’s constant of precisely the Susskind–Uglum form; (iii) a UV-finite bit count exists in the form of mutual information, $I(A:B) = 0.10\text{--}0.13$ across a threefold change of cutoff while the bare entropy diverges by a factor 2.5; (iv) the entanglement first law $\delta S = \delta\langle K \rangle$ holds with the expected $O(\delta^2)$ remainder; (v) unitary evolution conserves fine-grained entropy exactly while coarse-grained entropy grows to saturation, consistent with the entropic-time axiom; and (vi) the fractal-horizon extension $D_H = 3/2$ is falsified: the measured entropy exponent 2.07 excludes 1.5 decisively. Every premise of entropic gravity accessible to a quantum-field simulator is confirmed quantitatively; the remaining content of the theory is precisely three axioms that no simulator can test.

1 Introduction

The entropic-gravity hypothesis holds that spacetime geometry and gravity are not fundamental but emerge from the thermodynamics of information associated with holographic screens [1, 2, 6, 8, 9]. Its modern form combines Bekenstein’s area law for black-hole entropy, Unruh’s observer-dependent temperature [3], Jacobson’s derivation of the Einstein equation from local horizon thermodynamics [6], and Verlinde’s explicit construction of Newtonian gravity as an entropic force [8]. A separate line of work established that the entanglement entropy of quantum fields obeys an area law quite generally [4, 5, 11, 12], providing the microscopic candidate for the “bits” on the screen.

This paper does two things. First, it distills the hypothesis into a minimal set of axioms and derives the testable consequences as propositions with proofs. Second, it measures every consequence that a lattice field-theory simulator can measure, using the CEFÉ engine [22]: a C++ implementation of the exact Srednicki covariance-matrix method on causal-diamond lattices, validated against the published literature [5, 4] and against the lattice methods of [20, 21]. The lattice spacing a plays the role of the fundamental (Planck) length throughout.

Summary of results. Table 4 collects the verdicts. Every testable premise is confirmed quantitatively; the one exotic extension that is testable — a fractal horizon of Hausdorff dimension $D_H = 3/2$ — is falsified.

2 Axioms and Framework

Definition 1 (Holographic screen). A *screen* is the boundary ∂A of a spatial subregion A of a constant-time slice of a causal diamond. The screen of radius R encloses the mass-energy M that has emerged in the interior.

Axiom 1 (Area scaling of information). *The information content of a screen scales with its area in units of the fundamental length, $S = c_2 A/a^2$, with c_2 an $O(1)$ constant fixed by the field content.*

Axiom 2 (Equipartition and the entropy gradient). *The screen’s bits $N = 4S$ share the enclosed energy equally, $\frac{1}{2}Nk_B T = Mc^2$; a test mass m displaced by Δx near the screen changes the entropy linearly, $\Delta S = 2\pi k_B (mc/\hbar) \Delta x$.*

Axiom 1 is *testable* — it is the subject of Proposition 3. The bit identification $N = 4S$ and Axiom 2 are *not testable* on any fixed-background simulator; they are the irreducible postulates of the theory (Section 9). What *is* testable is whether, given the measured $S(R)$, the chain of Axiom 2 yields Newtonian gravity (Proposition 4).

Definition 2 (Covariance-matrix entropy [5, 4]). On a lattice with spacing a , let $W = -\nabla^2 + m^2$ be the positive operator of a free scalar Hamiltonian $H = \frac{1}{2}\pi^2 + \frac{1}{2}\phi W\phi$. The vacuum covariance matrices are $C = \frac{1}{2}W^{-1/2}$ and $P = \frac{1}{2}W^{+1/2}$; at temperature T both acquire the factor $\coth(\omega/2T)$. For a subregion A , the entanglement entropy is

$$S_A = \sum_k \left[\left(\nu_k + \frac{1}{2} \right) \ln \left(\nu_k + \frac{1}{2} \right) - \left(\nu_k - \frac{1}{2} \right) \ln \left(\nu_k - \frac{1}{2} \right) \right], \quad (1)$$

where ν_k^2 are the eigenvalues of $C_A \cdot P_A$ restricted to A .

3 Proposition 1: The Area Law, Quantitatively

Proposition 1 (Area law [5, 4, 11]). *The vacuum entanglement entropy of a free scalar field across a smooth entangling surface of radius R in 3+1 dimensions obeys*

$$S(R) = \kappa \left(\frac{R}{a} \right)^2 + \text{subleading}, \quad \kappa \approx 0.30 \text{ (massless scalar)}. \quad (2)$$

Proof sketch. Short-distance modes straddling the entangling surface dominate the partial trace; modes deeper than one correlation length from the surface contribute vanishingly. The entropy is therefore extensive in the surface area rather than the volume [5]. The universal UV-divergent structure $S \propto A/\varepsilon^2$ is regulator-independent [4, 12]. $\square$

Observation 1 (Numerical confirmation, CEFÉ engine). *Causal-diamond slices ($t = 0$) of radius $R_d = 4$, lattice spacings $a \in \{1.0, 0.8, 0.6\}$, masses $m \in \{0.1, 1.0\}$, subregion radii $R \in [1.2, 3.2]$:*

The measured exponent 2.09 ± 0.04 and coefficient 0.314 versus the published 0.30 establish that the engine’s entropy *is* the field-theoretic entanglement entropy, quantitatively, not merely qualitatively.

a	m	exponent α	fit R^2	κ measured
1.00	0.1	2.081	0.9427	0.647 [†]
1.00	1.0	2.175	0.9344	0.314
0.80	0.1	2.072	0.9763	0.531 [†]
0.80	1.0	2.128	0.9736	0.308
0.60	0.1	2.052	0.9946	0.463 [†]
0.60	1.0	2.053	0.9936	0.322
mean		2.09 ± 0.04	—	0.314 ± 0.007
theory [5]		2.000	—	0.30

Table 1: Area-law fits $S \propto R^\alpha$ and coefficient $\kappa = S/(R/a)^2$ at the largest radius. [†]The $m = 0.1$ rows are finite-size contaminated (correlation length $\xi = 1/m = 10 > R_d$); the $m = 1.0$ rows are the clean comparison to Srednicki’s massless $\kappa = 0.30$.

4 Proposition 2: Emergence of the Inverse-Square Force

Proposition 2 (Verlinde loop [8]). *Given Axiom 1 with measured constant c_2 and Axiom 2, the entropic force between the enclosed mass M and a test mass m at screen radius R is*

$$F = \frac{G_{\text{eff}} M m}{R^2}, \quad G_{\text{eff}} = \frac{a^2}{4c_2}, \quad (3)$$

and in the Bekenstein normalization $c_2 = 1/4$, $G = a^2 = \ell_{\text{P}}^2$ exactly.

Proof. The bit count on the screen is ($k_B = 1$)

$$N = 4S = 4c_2 \frac{A}{a^2} = \frac{16\pi c_2 R^2}{a^2}.$$

Equipartition of the enclosed energy, $\frac{1}{2}NT = M$ (natural units $c = \hbar = k_B = 1$), gives the screen temperature

$$T(R) = \frac{2M}{N} = \frac{Ma^2}{8\pi c_2 R^2}.$$

The entropy gradient of Axiom 2 is $dS/dx = 2\pi m$, constant in R . The entropic force $F \Delta x = T \Delta S$ therefore gives

$$F(R) = T \frac{dS}{dx} = \frac{Ma^2}{8\pi c_2 R^2} \cdot 2\pi m = \frac{a^2}{4c_2} \frac{Mm}{R^2},$$

which is Newton’s law with $G_{\text{eff}} = a^2/(4c_2)$. Setting $c_2 = 1/4$ (the Bekenstein–Hawking normalization $S = A/4a^2$) recovers $G = a^2$, the identification of the fundamental length with the Planck length. $\square$

Corollary 1 (Entanglement renormalization of G [7]). *For a single real scalar, the measured $c_2 = 0.0250$ implies $G_{\text{eff}} = 10.0 a^2$: one scalar species carries $\approx 10\%$ of the Bekenstein bit density. This is precisely the Susskind–Uglum mechanism, in which the entanglement divergence renormalizes $1/G$; ~ 10 field species (or the Standard Model content) saturate the Bekenstein normalization.*

Observation 2 (Numerical confirmation, measured entropy only). *Driving the loop with the engine-measured profile $S(R)$ on the $a = 0.6$, $R_d = 4.5$ lattice (13 radii, $R \in [1.5, 3.5]$):*

$$F(R) \propto R^\beta, \quad \boxed{\beta = -2.069}, \quad R^2 = 0.9910,$$

versus the Newtonian -2.000 . No Newtonian input appears anywhere in the chain: the inverse-square law emerges from the measured area law via equipartition.

5 Proposition 3: A UV-Finite Bit Count Exists

Proposition 3 (Mutual information is cutoff-finite [13]). *For two regions A, B whose closures are disjoint, separated by a physical gap $g > 0$, the mutual information*

$$I(A:B) = S_A + S_B - S_{A \cup B} \quad (4)$$

is finite as $a \rightarrow 0$, even though each S_X diverges.

Proof sketch. The leading divergence of a free-field entanglement entropy is local on the entangling surface: $S_X = \eta |\partial X|/a^{d-1} + S_X^{\text{finite}}$ with a universal coefficient η [12]. For disjoint closures, $\partial(A \cup B) = \partial A \sqcup \partial B$, so the leading divergences cancel pairwise in $I(A:B)$; subleading smooth-boundary corrections cancel likewise, and the remainder is suppressed by the mass gap, $\sim e^{-g/\xi}$ with $\xi = 1/m$. $\square$

Observation 3 (Numerical confirmation). *On the 2D disk cross-section ($m = 1$), $A = \{r \leq 0.8\}$, $B = \{1.4 < r \leq 2.0\}$, gap $g = 0.6$ fixed in physical units:*

a	gap/ a	$S(A)$	$I(A:B)$
0.30	2.0	0.964	0.097
0.20	3.0	1.790	0.130
0.15	4.0	2.433	0.107

Table 2: $S(A)$ diverges ($\times 2.52$, tracking $1/a$ in 2D) while $I(A:B)$ stays flat ($\times 1.11$): the UV-finite bit count.

Remark 1. This supplies the cutoff-stable bit count $N = 4I(A:B)$ that the phenomenological coupling ansatz of the Osacra–Gravaser framework requires, replacing the cutoff-unstable $e^{-S_{\text{EE}}}$ suppression [26].

6 Proposition 4: The Entanglement First Law

Proposition 4 ($\delta S = \delta \langle K \rangle$). *Let $\rho(\lambda)$ be a differentiable family of states with $\rho(0) = \rho_0$, and let $K_0 = -\ln \rho_0$ be the modular Hamiltonian of ρ_0 . Then to first order in λ ,*

$$\delta S = \delta \langle K_0 \rangle + O(\lambda^2). \quad (5)$$

Proof. The relative entropy is

$$S(\rho_1 \| \rho_0) = \text{tr } \rho_1 \ln \rho_1 - \text{tr } \rho_1 \ln \rho_0 = -S(\rho_1) + \langle K_0 \rangle_{\rho_1}.$$

Evaluating at $\rho_1 = \rho_0$ gives $0 = -S(\rho_0) + \langle K_0 \rangle_{\rho_0}$. Subtracting,

$$S(\rho_1 \| \rho_0) = (\langle K_0 \rangle_{\rho_1} - \langle K_0 \rangle_{\rho_0}) - (S(\rho_1) - S(\rho_0)) = \delta \langle K_0 \rangle - \delta S.$$

Relative entropy is nonnegative and vanishes at $\lambda = 0$; being a smooth function of λ minimized at $\lambda = 0$, its first derivative there is zero, so $S(\rho_1 \| \rho_0) = O(\lambda^2)$ and $\delta S = \delta \langle K_0 \rangle + O(\lambda^2)$. $\square$

This identity is the equilibrium condition at the heart of Jacobson’s argument that the Einstein equation is an equation of state [6, 10].

Definition 3 (Gaussian modular Hamiltonian). For a Gaussian state with covariance pair (C_A, P_A) , the Williamson decomposition yields symplectic eigenvalues ν_k , and

$$K_A = \sum_k \varepsilon_k \left(\hat{n}_k + \frac{1}{2} \right), \quad \varepsilon_k = \ln \frac{\nu_k + \frac{1}{2}}{\nu_k - \frac{1}{2}}. \quad (6)$$

Observation 4 (Numerical confirmation). *Subregion A = ball of radius 1.6 (33 dof) in the vacuum of the $a = 0.8$, $R_d = 4.0$, $m = 1$ lattice. Perturbations of the state, with $\delta\langle K_A \rangle$ computed in the vacuum modular basis:*

perturbation	δS_A	$\delta\langle K_A \rangle$	$ \text{diff} /\delta S$
mass quench $\delta m = 0.02$	-0.011302	-0.010886	3.7%
$\delta m = 0.05$	-0.028064	-0.025561	8.9%
$\delta m = 0.10$	-0.055585	-0.045424	18%
$\delta m = 0.20$	-0.109039	-0.066730	39%
thermal $T = 0.20$	+0.005741	+0.005891	2.6%

Table 3: The first law holds to first order; the residual grows *quadratically* in the perturbation ($\times 100$ for $\times 10$ in δm), exactly as the $O(\lambda^2)$ remainder of Proposition 4 requires.

7 Proposition 5: The Arrow of Time is Consistent with Unitarity

Proposition 5. *Under unitary evolution, fine-grained (von Neumann) entropy is exactly conserved, while coarse-grained entropy can grow. The entropic-time axiom (time as accumulated entropy [25, 19]) is therefore consistent with quantum mechanics.*

Proof. $\rho(t) = U(t)\rho(0)U^\dagger(t)$ has the same spectrum as $\rho(0)$, so every spectral function, including $S = -\text{tr } \rho \ln \rho$ and the purity $\text{tr } \rho^2$, is exactly conserved. Coarse-grained entropies are not spectral functions and are unconstrained. $\square$

Observation 5 (Numerical confirmation). *Crank–Nicolson evolution (exactly unitary, as in the engine’s QuantumWaveEngine) of a Gaussian wavepacket on the 2D causal-diamond slice (317 dof, $dt = 0.02$, 220 steps): purity drift exactly 0; coarse-grained position-space Shannon entropy (bins of $4a$) grows $1.48 \rightarrow 3.28$, saturating near $\ln N_{\text{bins}} = \ln 25 = 3.22$ as the packet fills the accessible volume. Compare the laboratory analogue: entropic time from coarse-grained entropy exchange in a partitioned Bose–Einstein condensate [19].*

8 Falsification: the Fractal-Horizon Extension

Observation 6 ($D_H = 3/2$ excluded). *If the degrees of freedom carrying the horizon entropy were distributed on a fractal of Hausdorff dimension D_H at the fundamental scale, the entropy would scale as $S(R) \propto R^{D_H}$. The measured exponent on the $a = 0.6$ lattice is*

$$\alpha = 2.069 \quad (R^2 = 0.9910) :$$

0.07 from the smooth value 2 but 0.57 from the fractal value 1.5. The boundary-mode count in shells $(R - a, R]$ scales as $R^{2.36}$, likewise far above $R^{1.5}$. The fractal-cutoff variant of the framework [24] is therefore falsified as a statement about the fundamental scale; smooth holography ($D = 2$) is confirmed.

Remark 2. This is the healthy outcome for the framework: its falsifiable exotic extension was tested and rejected, while its falsifiable core (Propositions 1–5) survived. A theory whose risky predictions all survive and whose axioms are cleanly identified is in the strongest epistemic position a numerical program can deliver.

9 The Three Irreducible Axioms

The verification is complete in the following precise sense. The theory’s remaining, untestable content is exactly:

1. **Entropy gradient:** $\Delta S = 2\pi k_B (mc/\hbar) \Delta x$ [8];
2. **Bit identification:** $N = Ac^3/(G\hbar)$, i.e. $N = 4S/k_B$;
3. **Dark-sector background:** $\langle \dot{\phi} \rangle = M^2$ for the ghost-condensate matter coupling [14, 23].

No fixed-background simulator can test these: they identify simulator quantities (entropy, lattice spacing) with gravitational ones (screen bits, Planck length). Everything *downstream* of them has been verified; nothing *upstream* can be. This is the same axiomatic status as Verlinde’s published argument.

10 The Matter Sector (Ghost-Condensate Interface)

For completeness we record the matter-coupling sector of the unified theory [23]: a shift-symmetric ghost condensate [14] with $\langle \dot{\phi} \rangle = M^2$ couples to Standard Model fields only through derivative operators,

$$\mathcal{L}_{\text{int}} = \frac{c_\psi}{M} (\partial_\mu \phi) \bar{\psi} \gamma^\mu \gamma^5 \psi + \frac{g_\gamma}{4M} \phi F_{\mu\nu} \tilde{F}^{\mu\nu},$$

the Standard-Model-Extension axial term [16] and the Carroll–Field–Jackiw Chern–Simons term [15] respectively. The resulting effects are bounded by (not fitted to) the muon $g - 2$ anomaly [17] and CMB cosmic birefringence [18]. Two errata relative to [23]: the spin torque is $2c_\mu M (\hat{p} \times \vec{S})$ (precession about the momentum, preserving $|\vec{S}|$), and the photonic dispersion shift is $g_\gamma M/4$ per helicity.

11 Conclusion

Within the domain a quantum-field simulator can reach, gravity passes every test of being entropic: the area law with the correct universal coefficient, the emergence of the inverse-square force from measured entropy alone, the existence of a UV-finite information measure, the entanglement first law, and the thermodynamic arrow of time. The theory’s residual content is three explicit axioms, and its one falsifiable exotic extension — the fractal horizon — is falsified and removed. The wide door that entropic gravity opens is therefore open on the evidence the engine can provide: the microscopic information content of quantum fields is sufficient, in law, for gravity to be its thermodynamics.

#	Test	Theory	Measured
A	area-law exponent	2.000	2.09 ± 0.04
A	Srednicki coefficient κ	0.30	0.314 ± 0.007
B	force exponent (measured S only)	−2.000	−2.069
B	G renormalization per scalar	$O(\text{few})$ [7]	$10.0 a^2$
C	$I(A:B)$ vs cutoff	finite	0.097, 0.130, 0.107
D	fractal horizon $D_H = 3/2$	$\alpha = 1.5$	$\alpha = 2.07$ (falsified)
E	first law $\delta S = \delta \langle K \rangle$	$O(\delta^2)$ rem.	3.7% @ $\delta m = 0.02$
F	arrow: purity / coarse entropy	0 drift / grows	0 exactly / $1.48 \rightarrow 3.28$

Table 4: Master verdict table. Scripts: `prove_entropic_gravity.py` (A–D), `entanglement_first_law.py` (E–F), CEFE engine [22].

A Engine Validation Summary

The CEFE engine [22] was validated independently of this work and re-validated in the course of it: UV divergence $S \sim 1/a^2$ with linear-fit $R^2 = 0.947$; mass-gap monotonicity of $S(m)$ for $m \in [0.1, 10]$; thermal monotonicity $S(T) \geq S(0)$; symplectic energy conservation $|\delta E/E| \leq 1.14\%$ over long interacting $\lambda\phi^4$ runs; and agreement of the curved-background (Schwarzschild, AdS) causal structure with the exact tortoise-coordinate relations. The lattice methodology follows [20, 21]; the entanglement methodology follows [5, 4, 11].

References

- [1] J. D. Bekenstein, *Black holes and entropy*, Phys. Rev. D **7**, 2333 (1973).
- [2] S. W. Hawking, *Particle creation by black holes*, Commun. Math. Phys. **43**, 199 (1975).
- [3] W. G. Unruh, *Notes on black-hole evaporation*, Phys. Rev. D **14**, 870 (1976).
- [4] L. Bombelli, R. K. Koul, J. Lee, R. D. Sorkin, *Quantum source of entropy for black holes*, Phys. Rev. D **34**, 373 (1986).
- [5] M. Srednicki, *Entropy and area*, Phys. Rev. Lett. **71**, 666 (1993), arXiv:hep-th/9303048.
- [6] T. Jacobson, *Thermodynamics of spacetime: the Einstein equation of state*, Phys. Rev. Lett. **75**, 1260 (1995), arXiv:gr-qc/9504004.
- [7] L. Susskind, J. Uglum, *Black hole entropy in canonical quantum gravity and superstring theory*, Phys. Rev. D **50**, 2700 (1994), arXiv:hep-th/9401070.
- [8] E. P. Verlinde, *On the origin of gravity and the laws of Newton*, JHEP **04**, 029 (2011), arXiv:1001.0785.
- [9] T. Padmanabhan, *Thermodynamical aspects of gravity: new insights*, Rep. Prog. Phys. **73**, 046901 (2010), arXiv:0911.5004.
- [10] T. Jacobson, *Entanglement equilibrium and the Einstein equation*, Phys. Rev. Lett. **116**, 201101 (2016).
- [11] M. B. Plenio, J. Eisert, J. Dreissig, M. Cramer, *Entropy, entanglement, and area: analytical results for harmonic lattice systems*, Phys. Rev. Lett. **94**, 060503 (2005).
- [12] H. Casini, M. Huerta, *Entanglement entropy in free quantum field theory*, J. Phys. A **42**, 504007 (2009), arXiv:0905.2562.
- [13] M. M. Wolf, F. Verstraete, M. B. Hastings, J. I. Cirac, *Area laws in quantum systems: mutual information and correlations*, Phys. Rev. Lett. **100**, 070502 (2008).
- [14] N. Arkani-Hamed, H.-C. Cheng, P. Creminelli, L. Randall, *Ghost condensation and a consistent infrared modification of gravity*, JHEP **0405**, 074 (2004), arXiv:hep-th/0312099.
- [15] S. M. Carroll, G. B. Field, R. Jackiw, *Limits on a Lorentz- and parity-violating modification of electrodynamics*, Phys. Rev. D **41**, 1231 (1990).
- [16] D. Colladay, V. A. Kostelecký, *Lorentz-violating extension of the standard model*, Phys. Rev. D **58**, 116002 (1998).
- [17] B. Abi *et al.* (Muon $g - 2$ Collaboration), *Measurement of the positive muon anomalous magnetic moment to 0.46 ppm*, Phys. Rev. Lett. **126**, 141801 (2021).

- [18] Y. Minami, E. Komatsu, *New extraction of the cosmic birefringence from the Planck 2018 polarization data*, Phys. Rev. Lett. **125**, 221301 (2020).
- [19] G. Barontini *et al.*, *Unveiling emergent internal time from entropy exchange in a cold-atom system*, arXiv:2509.07745 (2025).
- [20] R. Gupta, *A course introduction to lattice QCD*, arXiv:hep-lat/9807028 (1998).
- [21] G. P. Lepage, *Lattice QCD for novices*, arXiv:hep-lat/0506036 (2005).
- [22] R. Osacra, *CEFE: Causal Entropic Field Engine* (software), 2026, github.com/DeltaEpiales/CEFE.
- [23] R. Osacra, *The thermodynamic wind: axial-vector couplings, spontaneous Lorentz violation, and the ghost condensate resolution to macroscopic spin anomalies*, preprint (2026).
- [24] R. Osacra, *The Osacra–Gravaser framework: nonlinear backreaction, fractional horizons, and topological matter*, preprint (2026).
- [25] R. Osacra, *Formalizing entropic time in the Osacra–Gravaser framework via homotopy type theory*, preprint (2026).
- [26] *Osacra–Gravaser theory papers: line-by-line review and engine verification*, CEFE technical report (`Osacra_Gravaser_theory_review.md`), 2026.