

# **Studies with Improved Renormalization Group Techniques**

by

**Gregory James Petropoulos**

B.S., University of Connecticut, 2010

M.S., University of Colorado, 2013

A thesis submitted to the  
Faculty of the Graduate School of the  
University of Colorado in partial fulfillment  
of the requirements for the degree of  
Doctor of Philosophy  
Department of Physics

2015

This thesis entitled:  
Studies with Improved Renormalization Group Techniques  
written by Gregory James Petropoulos  
has been approved for the Department of Physics

---

Anna Hasenfratz

---

Prof. Thomas DeGrand

---

Prof. Ethan Neil

---

Prof. Senarath de Alwis

---

Prof. Thomas A. Manteuffel

Date \_\_\_\_\_

The final copy of this thesis has been examined by the signatories, and we find that both the content and the form meet acceptable presentation standards of scholarly work in the above mentioned discipline.

Petropoulos, Gregory James (Ph.D., Physics)

Studies with Improved Renormalization Group Techniques

Thesis directed by Prof. Anna Hasenfratz

## Dedication

## **Acknowledgements**

This thesis was supported by an award from the Department of Energy (DOE) Office of Science Graduate Fellowship Program (DOE SCGF). The DOE SCGF Program was made possible in part by the American Recovery and Reinvestment Act of 2009. The DOE SCGF program is administered by the Oak Ridge Institute for Science and Education for the DOE. ORISE is managed by Oak Ridge Associated Universities (ORAU) under DOE contract number DE-AC05-06OR23100. All opinions expressed in this presentation are the author's and do not necessarily reflect the policies and views of DOE, ORAU, or ORISE.

# Contents

## Chapter

<b>1</b>	Wilson Flow MCRG	1
1.1	Introduction . . . . .	1
1.2	Wilson Flow Optimized MCRG . . . . .	2
1.3	12 Flavor Results . . . . .	3

<b>Bibliography</b>	9
---------------------	---

## Appendix

## Tables

### Table

## Figures

### Figure

- 1.1 In Wilson Flow MCRG we use the Wilson Flow (blue) to approach the renormalized trajectory. An optimization step similar to that used in MCRG allows us to locate the flow time that gets us closest to the renormalized trajectory (orange). We then block our lattice using a fixed block transformation (green). . . . . 6
- 1.2 Optimization of the Wilson flow time  $t_f$  with fixed  $\alpha = 0.5$ , for  $\beta_F = 4.5$ . The uncertainties on the data points are dominated by averaging over the different observables. 6
- 1.3 The bare step-scaling function  $s_b$  predicted by three-lattice matching with  $6^4$ ,  $12^4$  and  $24^4$  lattices blocked down to  $3^4$ , comparing three different renormalization schemes. The error bars come from the standard deviation of predictions using the different observables. . . . . 7
- 1.4 As in Fig. 1.3, the bare step-scaling function  $s_b$  for three different renormalization schemes from three-lattice matching, now using  $8^4$ ,  $16^4$  and  $32^4$  lattices blocked down to  $4^4$ . . . . . 7
- 1.5 The bare step-scaling function  $s_b$  for scheme 1, comparing three-lattice matching using different volumes:  $6^4$ ,  $12^4$  and  $24^4$  lattices blocked down to  $3^4$  (black  $\times$ s) as well as  $8^4$ ,  $16^4$  and  $32^4$  lattices blocked down to  $4^4$  (blue bursts) and  $2^4$  (red crosses). 8



## Chapter 1

### Wilson Flow MCRG

#### 1.1 Introduction

For the past several years many lattice groups have been involved in studying strongly-coupled near-conformal gauge–fermion systems. Some of these models may be candidates for new physics beyond the standard model, while others are simply interesting non-perturbative quantum field theories. Because the dynamics of these lattice systems are unfamiliar, it is important to study them with several complementary techniques. Not only does this allow consistency checks, it can also provide information about the most efficient and reliable methods to investigate near-conformal lattice theories.

Monte Carlo Renormalization Group (MCRG) two-lattice matching is one of several analysis tools that we are using to investigate SU(3) gauge theories with many massless fermion flavors. This technique predicts the step-scaling function  $s_t$  in the bare parameter space. In a previous work [99] we proposed an improved MCRG method that exploits the Wilson flow to obtain a bare step-scaling function that corresponds to a unique discrete  $\beta$  function. We briefly review our Wilson-flow-optimized MCRG (WMCRG) procedure in Sections 1.2. It is important to note that we are investigating a potential infrared fixed point (IRFP) where the coupling is irrelevant: its running slows and eventually stops. This is challenging to distinguish from a near-conformal system where the gauge coupling runs slowly but does not flow to an IRFP. The observation of a backward flow that survives extrapolation to the infinite-volume limit could provide a clean signal. In Section 1.3 we report WMCRG results for SU(3) gauge theory with  $N_f = 12$  flavors of massless

fermions in the fundamental representation.

This 12-flavor model has been studied by many groups, including Refs. [10, 39, 46, 7, 66, 33, 28, 80, 86, 6, 48, 49, 76, 27, 5, 69, 68, 26]. Using new ensembles of 12-flavor gauge configurations generated with exactly massless fermions, our improved WMCRG technique predicts a conformal IRFP where the step-scaling function vanishes. As with every method, it is essential to study the systematic effects. For WMCRG the most important systematic effects are due to the finite volume and limited number of blocking steps. While we are not able to carry out a rigorous infinite-volume extrapolation, the observed zero of the bare step-scaling function is present for all investigated lattice volumes and renormalization schemes, and agrees with the earlier MCRG results of Ref. [66]. The results of our complementary  $N_f = 12$  investigations of finite-temperature phase transitions [107, 69], the Dirac eigenmode number [27, 26], and finite-size scaling [68] are also consistent with the existence of an infrared fixed point and IR conformality.

## 1.2 Wilson Flow Optimized MCRG

As an alternative to optimizing the RG blocking transformation, and thus changing the renormalization scheme at each coupling  $\beta_F$ , here we propose to use the Wilson flow to move the lattice system as close as possible to the renormalized trajectory of a fixed renormalization scheme.

The Wilson flow is a continuous smearing transformation [92] that can be related to the  $\overline{\text{MS}}$  running coupling in perturbation theory [87]. Refs. [51, 50] recently used the Wilson flow to compute a renormalized step-scaling function in a way similar to Schrödinger functional methods. While this approach appears very promising, it is based on perturbative relations that are only fully reliable at weak coupling. Here we do not use this perturbative connection, instead applying the Wilson flow as a continuous smearing that removes UV fluctuations. The Wilson flow moves the system along a surface of constant lattice scale in the infinite-dimensional action-space; it is not a renormalization group transformation and does not change the IR properties of the system.

Our goal is to use a one-parameter Wilson flow transformation to move the lattice system as close as possible to the renormalized trajectory of our fixed RG blocking transformation. This

is shown in figure 1.1. We proceed by carrying out two-lattice matching after applying the Wilson flow for a flow time  $t_f$  on all lattice volumes. (The Wilson flow is run only on the unblocked lattices, not in between RG blocking steps.) As above, since we can block our lattices only a few times, we must optimize  $t_f$  by requiring that consecutive RG blocking steps yield the same  $\Delta\beta_F$ , as shown in 1.2. As for traditional MCRG, increasing the number of blocking steps reduces the dependence on the optimization parameter; in the limit  $n_b \rightarrow \infty$ , our results would be independent of  $t_f$ .

With Wilson-flowed MCRG we can efficiently determine bare step-scaling functions that correspond to unique RG  $\beta$  functions. The uniqueness of the  $\beta$  function is a result of using a fixed block transformation. In this work our block transformation consists of nHYP smearing the unblocked lattice and then multiplying adjacent links in the same direction.

The ability to study a unique  $\beta$  function opens up interesting directions for future studies. By comparing different  $\beta$  functions around the perturbative gaussian FP, we can study scaling violations in the lattice system. In IR-conformal systems, we can investigate the scheme-dependence of the  $\beta$  function near the IRFP, an issue explored in perturbation theory by Ref. [104].

### 1.3 12 Flavor Results

Our WMCRG results for the 12-flavor system are obtained on gauge configurations generated with exactly massless fermions. Our lattice action uses nHYP-smearred staggered fermions as described in Ref. [28], and to run with  $m = 0$  we employ anti-periodic boundary conditions in all four directions. All of our analyses are carried out at couplings weak enough to avoid the unusual strong-coupling “ $\mathcal{S}^4$ ” phase discussed by Refs. [28, 69].

We perform three-lattice matching with volumes  $6^4$ – $12^4$ – $24^4$  and  $8^4$ – $16^4$ – $32^4$ . Three-lattice matching is based on two sequential two-lattice matching steps, to minimize finite-volume effects [66]. Both two-lattice matching steps are carried out on the same final volume  $V_f$ . We denote the number of blocking steps on the largest volume by  $n_b$ , and tune the length of the initial Wilson flow by requiring that the last two blocking steps predict the same step-scaling function. Using the  $8^4$ – $16^4$ – $32^4$  data we determine the bare step-scaling function for  $n_b = 3$  and  $V_f = 4^4$  as

well as  $n_b = 4$  and  $V_f = 2^4$ , while the  $6^4$ – $12^4$ – $24^4$  data set is blocked to a final volume  $V_f = 3^4$  ( $n_b = 3$ ). This allows us to explore the effects of both the final volume and the number of blocking steps. We investigate three renormalization schemes by changing the HYP smearing parameters in our blocking transformation [99]: scheme 1 uses smearing parameters (0.6, 0.2, 0.2), scheme 2 uses (0.6, 0.3, 0.2) and scheme 3 uses (0.65, 0.3, 0.2).

Figs. 1.3, 1.4 and 1.5 present representative results for 12 flavors. All of the bare step-scaling functions clearly show  $s_b = 0$ , signalling an infrared fixed point, for every  $n_b$ ,  $V_f$  and renormalization scheme. Appropriately for an IR-conformal system, the location of the fixed point is scheme dependent. We observe that the fixed point moves to stronger coupling as the HYP smearing parameters in the RG blocking transformation increase.

When we block our  $8^4$ ,  $16^4$  and  $32^4$  lattices down to a final volume  $V_f = 2^4$  (corresponding to  $n_b = 4$ ), the observables become very noisy, making matching more difficult. The problem grows worse as the HYP smearing parameters increase, and our current statistics do not allow reliable three-lattice matching for  $V_f = 2^4$  in schemes 2 and 3. To resolve this issue, we are accumulating more statistics in existing  $32^4$  runs, and generating additional  $32^4$  ensembles at more values of the gauge coupling  $\beta_F$ . These additional data will also improve our results for scheme 1, which we show in Fig. 1.5. Different volumes and  $n_b$  do not produce identical results in scheme 1, suggesting that the corresponding systematic effects are still non-negligible. We can estimate finite-volume effects by comparing  $n_b = 3$  with  $V_f = 3^4$  and  $V_f = 4^4$ . Systematic effects due to  $n_b$  can be estimated from  $n_b = 4$  and  $V_f = 2^4$ , but this is difficult due to the noise in the  $2^4$  data. Even treating the spread in the results shown in Fig. 1.5 as a systematic uncertainty, we still obtain a clear zero in the bare step-scaling function, indicating an IR fixed point.

In this chapter we have shown how the Wilson-flow-optimized MCRG two-lattice matching procedure proposed in Ref. [99] improves upon traditional lattice renormalization group techniques. By optimizing the flow time for a fixed RG blocking transformation, WMCRG predicts a bare step-scaling function  $s_b$  that corresponds to a unique discrete  $\beta$  function. Applying WMCRG to new 12-flavor ensembles generated with exactly massless fermions, we observe an infrared fixed point

in  $s_b$ . The fixed point is present for all investigated lattice volumes, number of blocking steps and renormalization schemes, even after accounting for systematic effects indicated by Fig. 1.5. This result reinforces the IR-conformal interpretation of our complementary  $N_f = 12$  studies of phase transitions [107, 69], the Dirac eigenmode number [27, 26], and finite-size scaling [68].




Figure 1.1: In Wilson Flow MCRG we use the Wilson Flow (blue) to approach the renormalized trajectory. An optimization step similar to that used in MCRG allows us to locate the flow time that gets us closest to the renormalized trajectory (orange). We then block our lattice using a fixed block transformation (green).




Figure 1.2: Optimization of the Wilson flow time  $t_f$  with fixed  $\alpha = 0.5$ , for  $\beta_F = 4.5$ . The uncertainties on the data points are dominated by averaging over the different observables.




Figure 1.3: The bare step-scaling function  $s_b$  predicted by three-lattice matching with  $6^4$ ,  $12^4$  and  $24^4$  lattices blocked down to  $3^4$ , comparing three different renormalization schemes. The error bars come from the standard deviation of predictions using the different observables.




Figure 1.4: As in Fig. 1.3, the bare step-scaling function  $s_b$  for three different renormalization schemes from three-lattice matching, now using  $8^4$ ,  $16^4$  and  $32^4$  lattices blocked down to  $4^4$ .




Figure 1.5: The bare step-scaling function  $s_b$  for scheme 1, comparing three-lattice matching using different volumes:  $6^4$ ,  $12^4$  and  $24^4$  lattices blocked down to  $3^4$  (black  $\times$ s) as well as  $8^4$ ,  $16^4$  and  $32^4$  lattices blocked down to  $4^4$  (blue bursts) and  $2^4$  (red crosses).



## Bibliography

- [1]
- [2] David Adams. Fourth root prescription for dynamical staggered fermions. Phys. Rev. D, 72:114512, Dec 2005.
- [3] Yasumichi Aoki, Tatsumi Aoyama, Masafumi Kurachi, Toshihide Maskawa, Kohtaroh Miura, Kei-ichi Nagai, Hiroshi Ohki, Enrico Rinaldi, Akihiro Shibata, Koichi Yamawaki, and Takeshi Yamazaki. Light composite scalar in eight-flavor QCD on the lattice. 2014.
- [4] Yasumichi Aoki, Tatsumi Aoyama, Masafumi Kurachi, Toshihide Maskawa, Kei-ichi Nagai, Hiroshi Ohki, Enrico Rinaldi, Akihiro Shibata, Koichi Yamawaki, and Takeshi Yamazaki. Light composite scalar in twelve-flavor QCD on the lattice. Phys. Rev. Lett., 111:162001, 2013.
- [5] Yasumichi Aoki, Tatsumi Aoyama, Masafumi Kurachi, Toshihide Maskawa, Kei-ichi Nagai, Hiroshi Ohki, Enrico Rinaldi, Akihiro Shibata, Koichi Yamawaki, and Takeshi Yamazaki. The scalar spectrum of many-flavour QCD. 2013.
- [6] Yasumichi Aoki, Tatsumi Aoyama, Masafumi Kurachi, Toshihide Maskawa, Kei-ichi Nagai, Hiroshi Ohki, Akihiro Shibata, Koichi Yamawaki, and Takeshi Yamazaki. Lattice study of conformality in twelve-flavor QCD. Phys. Rev., D86:054506, 2012.
- [7] T. Appelquist, G. T. Fleming, M. F. Lin, E. T. Neil, and D. Schaich. Lattice Simulations and Infrared Conformality. Phys. Rev., D84:054501, 2011.
- [8] Thomas Appelquist, Richard Brower, Simon Catterall, George Fleming, Joel Giedt, Anna Hasenfratz, Julius Kuti, Ethan Neil, and David Schaich. Lattice Gauge Theories at the Energy Frontier. 2013.
- [9] Thomas Appelquist, George T. Fleming, and Ethan T. Neil. Lattice study of the conformal window in QCD-like theories. Phys. Rev. Lett., 100:171607, 2008.
- [10] Thomas Appelquist, George T. Fleming, and Ethan T. Neil. Lattice Study of Conformal Behavior in SU(3) Yang-Mills Theories. Phys. Rev., D79:076010, 2009.
- [11] Thomas Appelquist and Ethan T. Neil. Lattice gauge theory beyond the standard model. pages 699–729, 2009.
- [12] Janos Balog, Ferenc Niedermayer, and Peter Weisz. Logarithmic corrections to  $O(a^{*2})$  lattice artifacts. Phys. Lett., B676:188–192, 2009.

- [13] Janos Balog, Ferenc Niedermayer, and Peter Weisz. The Puzzle of apparent linear lattice artifacts in the 2d non-linear sigma-model and Symanzik's solution. Nucl. Phys., B824:563–615, 2010.
- [14] Tom Banks and A. Zaks. On the Phase Structure of Vector-Like Gauge Theories with Massless Fermions. Nucl. Phys., B196:189, 1982.
- [15] Claude Bernard, Maarten Golterman, Yigal Shamir, and Stephen R. Sharpe. Comment on: chiral anomalies and rooted staggered fermions [phys. lett. b 649 (2007) 230]. Physics Letters B, 649(23):235 – 240, 2007.
- [16] Gyan Bhanot.  $Su(3)$  lattice gauge theory in 4 dimensions with a modified wilson action. Physics Letters B, 108(45):337 – 340, 1982.
- [17] Gyan Bhanot and Michael Creutz. Variant actions and phase structure in lattice gauge theory. Phys. Rev. D, 24:3212–3217, Dec 1981.
- [18] J. Binney, N.J. Dowrick, A.J. Fisher, and M.E.J. Newman. The Theory of Critical Phenomena: An Introduction to the Renormalization Group. Oxford University Press, Oxford, 1992.
- [19] T. Blum, C. DeTar, Urs M. Heller, Leo Krkkinen, K. Rummukainen, and D. Toussaint. Thermal phase transition in mixed action  $\{SU\}$  (3) lattice gauge theory and wilson fermion thermodynamics. Nuclear Physics B, 442(12):301 – 316, 1995.
- [20] A. Bode. Two loop expansion of the schrödinger functional coupling  $sf$  in  $\{SU\}$  (3) lattice gauge theory. Nuclear Physics B - Proceedings Supplements, 63(13):796 – 798, 1998. Proceedings of the  $\{XVth\}$  International Symposium on Lattice Field Theory.
- [21] Szabolcs Borsanyi, Stephan Durr, Zoltan Fodor, Christian Hoelbling, Sandor D. Katz, S. Krieg, T. Kurth, L. Lellouch, T. Lippert, C. McNeile, and K. K. Szabo. High-precision scale setting in lattice QCD. JHEP, 1209:010, 2012.
- [22] William E. Caswell. Asymptotic Behavior of Nonabelian Gauge Theories to Two Loop Order. Phys. Rev. Lett., 33:244, 1974.
- [23] Simon Catterall and Francesco Sannino. Minimal walking on the lattice. Phys.Rev., D76:034504, 2007.
- [24] Anqi Cheng, Anna Hasenfratz, Yuzhi Liu, Gregory Petropoulos, and David Schaich. Finite size scaling of conformal theories in the presence of a near-marginal operator. 2013.
- [25] Anqi Cheng, Anna Hasenfratz, Yuzhi Liu, Gregory Petropoulos, and David Schaich. Step scaling studies using the gradient flow running coupling. 2014, in preparation.
- [26] Anqi Cheng, Anna Hasenfratz, Gregory Petropoulos, and David Schaich. Determining the mass anomalous dimension through the eigenmodes of Dirac operator. PoS, LATTICE 2013:088, 2013.
- [27] Anqi Cheng, Anna Hasenfratz, Gregory Petropoulos, and David Schaich. Scale-dependent mass anomalous dimension from Dirac eigenmodes. JHEP, 1307:061, 2013.

- [28] Anqi Cheng, Anna Hasenfratz, and David Schaich. Novel phase in  $SU(3)$  lattice gauge theory with 12 light fermions. Phys. Rev., D85:094509, 2012.
- [29] Christian B. Lang Christof Gattringer. Quantum Chromodynamics on the Lattice. Springer, 2010.
- [30] Michael Creutz. Chiral anomalies and rooted staggered fermions. Physics Letters B, 649(23):230 – 234, 2007.
- [31] Michael Creutz. Reply to: comment on: chiral anomalies and rooted staggered fermions [phys. lett. b 649 (2007) 230] [phys. lett. b 649 (2007) 235]. Physics Letters B, 649(23):241 – 242, 2007.
- [32] Thomas DeGrand. Lattice studies of QCD-like theories with many fermionic degrees of freedom. 2010.
- [33] Thomas DeGrand. Finite-size scaling tests for spectra in  $SU(3)$  lattice gauge theory coupled to 12 fundamental flavor fermions. Phys. Rev., D84:116901, 2011.
- [34] Thomas DeGrand and Anna Hasenfratz. Remarks on lattice gauge theories with infrared-attractive fixed points. Phys.Rev., D80:034506, 2009.
- [35] Thomas DeGrand, Yigal Shamir, and Benjamin Svetitsky. Gauge theories with fermions in the two-index symmetric representation. PoS, LATTICE2011:060, 2011.
- [36] Thomas DeGrand, Yigal Shamir, and Benjamin Svetitsky. Infrared fixed point in  $SU(2)$  gauge theory with adjoint fermions. Phys.Rev., D83:074507, 2011.
- [37] Luigi Del Debbio, Biagio Lucini, Agostino Patella, Claudio Pica, and Antonio Rago. Mesonic spectroscopy of Minimal Walking Technicolor. Phys.Rev., D82:014509, 2010.
- [38] T. DeGrand & C DeTar. Lattice Methods for Quantum Chromodynamics. World Scientific, 2006.
- [39] A. Deuzeman, M. P. Lombardo, and E. Pallante. Evidence for a conformal phase in  $SU(N)$  gauge theories. Phys. Rev., D82:074503, 2010.
- [40] Albert Deuzeman, Maria Paola Lombardo, Tiago Nunes da Silva, and Elisabetta Pallante. The bulk transition of QCD with twelve flavors and the role of improvement. 2012.
- [41] Savas Dimopoulos and Leonard Susskind. Mass Without Scalars. Nucl.Phys., B155:237–252, 1979.
- [42] Michael Dine. Tasi lectures on the strong cp problem.
- [43] Estia Eichten and Kenneth D. Lane. Dynamical Breaking of Weak Interaction Symmetries. Phys.Lett., B90:125–130, 1980.
- [44] F. Englert and R. Brout. Broken Symmetry and the Mass of Gauge Vector Mesons. Phys.Rev.Lett., 13:321–323, 1964.
- [45] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Nogradi, and Chris Schroeder. Nearly conformal gauge theories on the lattice. Int.J.Mod.Phys., A25:5162–5174, 2010.

- [46] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, and Chris Schroeder. Twelve massless flavors and three colors below the conformal window. Phys. Lett., B703:348–358, 2011.
- [47] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, Chris Schroeder, and Chik Him Wong. Can the nearly conformal sextet gauge model hide the Higgs impostor? Phys. Lett., B718:657–666, 2012.
- [48] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, Chris Schroeder, and Chik Him Wong. Confining force and running coupling with twelve fundamental and two sextet fermions. PoS, Lattice 2012:025, 2012.
- [49] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, Chris Schroeder, and Chik Him Wong. Conformal finite size scaling of twelve fermion flavors. PoS, Lattice 2012:279, 2012.
- [50] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, and Chik Him Wong. The gradient flow running coupling scheme. PoS, Lattice 2012:050, 2012.
- [51] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, and Chik Him Wong. The Yang-Mills gradient flow in finite volume. JHEP, 1211:007, 2012.
- [52] Zoltan Fodor, Kieran Holland, Julius Kuti, Daniel Negradi, and Chik Him Wong. Can a light Higgs impostor hide in composite gauge models? PoS, LATTICE 2013:062, 2014.
- [53] Patrick Fritzsche and Alberto Ramos. The gradient flow coupling in the Schrödinger Functional. JHEP, 1310:008, 2013.
- [54] J. Gasser and H. Leutwyler. Chiral perturbation theory to one loop. Ann. Phys., 158:142, 1984.
- [55] Howard Georgi and David B. Kaplan. Composite Higgs and Custodial SU(2). Phys.Lett., B145:216, 1984.
- [56] Joel Giedt. Confining force and running coupling with twelve fundamental and two sextet fermions. PoS, Lattice 2012:006, 2012.
- [57] Joel Giedt. Lattice gauge theory and physics beyond the standard model. PoS, Lattice 2012:006, 2012.
- [58] Sheldon L. Glashow. Partial-symmetries of weak interactions. Nuclear Physics, 22(4):579 – 588, 1961.
- [59] Maarten Golterman. Applications of chiral perturbation theory to lattice QCD. pages 423–515, 2009.
- [60] David Gross and Frank Wilczek. Ultraviolet behavior of non-abelian gauge theories. Phys. Rev. Lett., 30:1343–1346, Jun 1973.
- [61] G.S. Guralnik, C.R. Hagen, and T.W.B. Kibble. Global Conservation Laws and Massless Particles. Phys.Rev.Lett., 13:585–587, 1964.
- [62] A. Hasenfratz, R. Hoffmann, and F. Knechtli. The Static potential with hypercubic blocking. Nucl.Phys.Proc.Suppl., 106:418–420, 2002.

- [63] Anna Hasenfratz. Investigating the critical properties of beyond-qcd theories using monte carlo renormalization group matching. Phys. Rev. D, 80:034505, Aug 2009.
- [64] Anna Hasenfratz. Conformal or Walking? Monte Carlo renormalization group studies of SU(3) gauge models with fundamental fermions. Phys. Rev., D82:014506, 2010.
- [65] Anna Hasenfratz. MCRG study of 12 fundamental flavors with mixed fundamental-adjoint gauge action. PoS, Lattice 2011:065, 2011.
- [66] Anna Hasenfratz. Infrared fixed point of the 12-fermion SU(3) gauge model based on 2-lattice MCRG matching. Phys. Rev. Lett., 108:061601, 2012.
- [67] Anna Hasenfratz, Anqi Cheng, Gregory Petropoulos, and David Schaich. Mass anomalous dimension from Dirac eigenmode scaling in conformal and confining systems. PoS, Lattice 2012:034, 2012.
- [68] Anna Hasenfratz, Anqi Cheng, Gregory Petropoulos, and David Schaich. Finite size scaling and the effect of the gauge coupling in 12 flavor systems. PoS, LATTICE 2013:075, 2013.
- [69] Anna Hasenfratz, Anqi Cheng, Gregory Petropoulos, and David Schaich. Reaching the chiral limit in many flavor systems. 2013.
- [70] Anna Hasenfratz, Roland Hoffmann, and Stefan Schaefer. Hypercubic smeared links for dynamical fermions. JHEP, 0705:029, 2007.
- [71] Anna Hasenfratz and Francesco Knechtli. Flavor symmetry and the static potential with hypercubic blocking. Phys. Rev., D64:034504, 2001.
- [72] Anna Hasenfratz, David Schaich, and Aarti Veernala. Nonperturbative beta function of eight-flavor SU(3) gauge theory. 2014.
- [73] Peter W. Higgs. Broken symmetries and the masses of gauge bosons. Phys. Rev. Lett., 13(16):508–509, October 1964.
- [74] P.W. Higgs. Broken symmetries, massless particles and gauge fields. Physics Letters, 12(2):132 – 133, 1964.
- [75] Christopher T. Hill and Elizabeth H. Simmons. Strong dynamics and electroweak symmetry breaking. Physics Reports, 381(46):235 – 402, 2003.
- [76] Etsuko Itou. Properties of the twisted Polyakov loop coupling and the infrared fixed point in the SU(3) gauge theories. PTEP, 2013:083B01, 2013.
- [77] Y. Iwasaki, K. Kanaya, S. Kaya, S. Sakai, and T. Yoshie. Phase structure of lattice QCD for general number of flavors. Phys.Rev., D69:014507, 2004.
- [78] Xiao-Yong Jin and Robert D. Mawhinney. Lattice QCD with Eight Degenerate Quark Flavors. PoS, LATTICE2008:059, 2008.
- [79] Xiao-Yong Jin and Robert D. Mawhinney. Lattice QCD with 8 and 12 degenerate quark flavors. PoS, LAT2009:049, 2009.

- [80] Xiao-Yong Jin and Robert D. Mawhinney. Lattice QCD with 12 Degenerate Quark Flavors. PoS, Lattice 2011:066, 2012.
- [81] David B. Kaplan, Howard Georgi, and Savas Dimopoulos. Composite Higgs Scalars. Phys.Lett., B136:187, 1984.
- [82] D.B. Kaplan. Chiral symmetry and lattice fermions.
- [83] John Kogut and Leonard Susskind. Hamiltonian formulation of wilson’s lattice gauge theories. Phys. Rev. D, 11:395–408, Jan 1975.
- [84] Andreas S. Kronfeld. Lattice gauge theory with staggered fermions: How, where, and why (not). PoS, LAT2007:016, 2007.
- [85] Kenneth Lane. Two lectures on technicolor.
- [86] C.-J. David Lin, Kenji Ogawa, Hiroshi Ohki, and Eigo Shintani. Lattice study of infrared behaviour in  $SU(3)$  gauge theory with twelve massless flavours. JHEP, 1208:096, 2012.
- [87] Martin Luscher. Properties and uses of the Wilson flow in lattice QCD. JHEP, 1008:071, 2010.
- [88] Martin Luscher. Trivializing maps, the Wilson flow and the HMC algorithm. Commun. Math. Phys., 293:899–919, 2010.
- [89] M. Lscher and P. Weisz. Computation of the action for on-shell improved lattice gauge theories at weak coupling. Physics Letters B, 158(3):250 – 254, 1985.
- [90] Adam Martin. Technicolor signals at the lhc.
- [91] Shinya Matsuzaki and Koichi Yamawaki. Holographic techni-dilaton at 125 GeV. Phys. Rev., D86:115004, 2012.
- [92] R. Narayanan and H. Neuberger. Infinite N phase transitions in continuum Wilson loop operators. JHEP, 0603:064, 2006.
- [93] Ethan T. Neil. Exploring Models for New Physics on the Lattice. PoS, Lattice 2011:009, 2011.
- [94] H.B. Nielsen and M. Ninomiya. Absence of neutrinos on a lattice: (i). proof by homotopy theory. Nuclear Physics B, 185(1):20 – 40, 1981.
- [95] H.B. Nielsen and M. Ninomiya. Absence of neutrinos on a lattice: (ii). intuitive topological proof. Nuclear Physics B, 193(1):173 – 194, 1981.
- [96] H.B. Nielsen and M. Ninomiya. A no-go theorem for regularizing chiral fermions. Physics Letters B, 105(23):219 – 223, 1981.
- [97] Paula Perez-Rubio and Stefan Sint. Non-perturbative running of the coupling from four flavour lattice QCD with staggered quarks. PoS, Lattice 2010:236, 2010.
- [98] Michael E. Peskin and Dan V. Schroeder. An Introduction To Quantum Field Theory (Frontiers in Physics). Westview Press, 1995.

- [99] Gregory Petropoulos, Anqi Cheng, Anna Hasenfratz, and David Schaich. PoS, Lattice 2012:051, 2012.
- [100] Gregory Petropoulos, Anqi Cheng, Anna Hasenfratz, and David Schaich. Improved Lattice Renormalization Group Techniques. PoS, LATTICE 2013:079, 2013.
- [101] H. David Politzer. Reliable Perturbative Results for Strong Interactions? Phys.Rev.Lett., 30:1346–1349, 1973.
- [102] C. Quigg. Spontaneous symmetry breaking as a basis of particle mass. Rept. Prog. Physics, pages 1019–1054, 2007.
- [103] C. Quigg. Unanswered questions in the electroweak theory. Annual Review of Nuclear and Particle Science, pages 505–555, 2009.
- [104] Thomas A. Ryttov and Robert Shrock. An Analysis of Scheme Transformations in the Vicinity of an Infrared Fixed Point. Phys.Rev., D86:085005, 2012.
- [105] Abdus Salam and John Clive Ward. Electromagnetic and weak interactions. Phys. Lett., 13:168–171, 1964.
- [106] Francesco Sannino. Conformal Dynamics for TeV Physics and Cosmology. Acta Phys.Polon., B40:3533–3743, 2009.
- [107] David Schaich, Anqi Cheng, Anna Hasenfratz, and Gregory Petropoulos. Bulk and finite-temperature transitions in SU(3) gauge theories with many light fermions. PoS, Lattice 2012:028, 2012.
- [108] Robert Shrock. Some recent results on models of dynamical electroweak symmetry breaking. pages 227–241, 2007.
- [109] Stefan Sint. On the schrödinger functional in {QCD}. Nuclear Physics B, 421(1):135 – 156, 1994.
- [110] Jan Smit. Introduction to Quantum Fields on a Lattice. Cambridge University Press, 2002.
- [111] Rainer Sommer. Scale setting in lattice QCD. PoS, LATTICE 2013:015, 2014.
- [112] Leonard Susskind. Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory. Phys.Rev., D20:2619–2625, 1979.
- [113] R.H. Swendsen. Phys. Rev. Lett., 42:859, 1979.
- [114] K. Symanzik. Continuum limit and improved action in lattice theories : (ii).  $o(n)$  non-linear sigma model in perturbation theory. Nuclear Physics B, 226(1):205 – 227, 1983.
- [115] Fatih Tekin, Rainer Sommer, and Ulli Wolff. The Running coupling of QCD with four flavors. Nucl. Phys., B840:114–128, 2010.
- [116] Steven Weinberg. A Model of Leptons. Phys.Rev.Lett., 19:1264–1266, 1967.
- [117] Steven Weinberg. Implications of Dynamical Symmetry Breaking. Phys.Rev., D13:974–996, 1976.

- [118] Steven Weinberg. Implications of Dynamical Symmetry Breaking: An Addendum. Phys.Rev., D19:1277–1280, 1979.
- [119] P. Weisz. Continuum limit improved lattice action for pure yang-mills theory (i). Nuclear Physics B, 212(1):1 – 17, 1983.
- [120] Kenneth Wilson. Confinement of quarks. Phys. Rev. D, 10:2445–2459, Oct 1974.