

How Different Landscape Elements Limit the Breeding Habitat of Meadow Bird Species

Authors: Vliet, Roland E. van der, Dijk, Jerry van, and Wassen, Martin J.

Source: *Ardea*, 98(2) : 203-209

Published By: Netherlands Ornithologists' Union

URL: <https://doi.org/10.5253/078.098.0210>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

How different landscape elements limit the breeding habitat of meadow bird species

Roland E. van der Vliet^{1,2,*}, Jerry van Dijk¹ & Martin J. Wassen¹

Van der Vliet R.E., van Dijk J. & Wassen M.J. 2010. How different landscape elements limit the breeding habitat of meadow bird species. *Ardea* 98: 203–209.

Meadow bird species in the Netherlands have shown a long-term decline in numbers. This has been generally attributed to agricultural intensification, but in addition an increase of disturbing elements may have played a role in the decline. In this review we compiled data from literature to explore to what extent breeding by meadow birds (Eurasian Oystercatcher *Haematopus ostralegus*, Northern Lapwing *Vanellus vanellus*, Black-tailed Godwit *Limosa limosa*) is limited by disturbing landscape elements. We considered elements that can be considered 'view-obstructing' (like trees or houses), and those that can be considered 'flat' (canal, road or railway). Our review shows that breeding birds keep a distance to both types of landscape elements, with distances ranging over 1 km for highways and village edges. Roads with high traffic intensity caused the largest disturbance. Eurasian Oystercatcher was most tolerant to the presence of landscape elements, and Black-tailed Godwit least tolerant.

Key words: Black-tailed Godwit, Eurasian Oystercatcher, grassland ecosystem, habitat loss, Northern Lapwing, meadow bird, Netherlands, openness of landscape, view obstruction

¹Environmental Sciences, Copernicus Institute, Utrecht University, P.O. Box 80115, 3508 TC Utrecht, The Netherlands; ²present address: Tauw, P.O. Box 3015, 3502 GA Utrecht, The Netherlands;

*corresponding author (roland.vandervliet@tauw.nl)



In recent decades, populations of almost all meadow bird species in The Netherlands have declined, resulting in most of them being placed on the Dutch red list of threatened bird species, including Black-tailed Godwit *Limosa limosa* and Common Redshank *Tringa totanus* (van Beusekom *et al.* 2005). This decline has been attributed to several factors, including collecting eggs, hunting and alterations of habitat (Klomp 1951, Wymenga & Engelmoer 2001). However, it is now generally accepted that the decline has been mostly caused by a continuing intensification of agriculture in the breeding areas impairing reproductive success, for instance by an increased intensity of farmers' activities, increasing livestock densities, earlier and more frequent mowing, and lowering of groundwater levels (Beintema & Müskens 1987, Kruk 1993, Vickery *et al.* 2001, Newton 2004).

Another reason for the decline that has gained much less attention is that opportunities for safe breed-

ing by meadow birds are hampered by an increase of elements that obstruct visibility in the otherwise open habitat (Dyrce *et al.* 1981, van der Vliet *et al.* 2008). Klomp (1954) showed that Northern Lapwings were most effective in deterring potential predators in a treeless landscape, and calculated that Lapwings needed a field of at least 5 ha in size if lined by trees with a height of 6–10 m.

In this paper, we review effects of landscape elements on the breeding distribution of meadow birds, focusing on the species for which sufficient data have been published, Eurasian Oystercatcher *Haematopus ostralegus*, Northern Lapwing *Vanellus vanellus*, and Black-tailed Godwit *Limosa limosa*. Meadow birds keep a distance to elements in the landscape, thus resulting locally in lowered meadow bird densities. The distance where the population density is depressed due to the presence of a disturbing source is called the disturbance distance (cf. Veen 1973, van der Zande *et al.* 1980, Gill

et al. 1996, Reijnen *et al.* 1996). The maximum disturbance distance is defined as the largest distance where a negative effect is still measurable. We use this maximum disturbance distance as an indicator of the effect that a landscape element may have on suitable habitat and on the distribution of meadow birds.

We aim to distinguish between two sources of disturbance, (1) elements that hamper visibility (view-obstructing elements), and (2) flat elements that cause disturbance by the presence of potential predators without impairing visibility (canals, roads and railways). By using data on traffic intensity we separate effects of the flat elements proper and the presence of 'predators'. Thus, based on a literature review, this paper gives an indication to what extent different sources of disturbance limit the distribution of breeding meadow birds. Forman & Alexander (1998) reviewed the disturbance caused by roads on wildlife. We use their framework to discuss our findings.

METHODS

To quantify the effects of elements in the landscape on meadow bird populations, we searched literature sources for quoted disturbance distances, using both widely available and grey literature. We assumed that distances to field borders were negligible compared to the distance to landscape elements. We aimed to find figures for the disturbance distance for the three species separately but unfortunately not all sources made this distinction. Many studies were carried out in western Netherlands, where most Dutch meadow birds breed (Bijlsma *et al.* 2001). In most study sites, habitat was homogeneous, although some were heterogeneous with respect to meadow vegetation type. We only included studies that had considered this heterogeneity in their conclusions. Also, most studies distinguished between roads with or without a supporting view-obstructing element. If they did, we also made this distinction in our categories of elements. Studies calculated distances by either interpreting numbers of breeding meadow birds along a transect perpendicular to the disturbing source, or by interpreting field data afterwards.

Effects by landscape elements on disturbance distances were statistically tested for each species separately and for the species pooled. View obstruction and traffic intensity were the independent variables. Based on the original publications, traffic intensity was scored as 'low' (values 0 and 1), 'intermediate' (value 2), and 'high' (value 3). Note that both view-obstructing and

flat elements can have a high intensity of traffic. Unfortunately, inclusion of species as an independent variable led to strong deviations from normality, even after log-transformation of the data.

The effect of view obstruction and traffic intensity on the average disturbance distance was tested by ANOVAs. A Bonferroni post-hoc test was used to identify significant differences between levels of traffic intensity. Data were log-transformed as data assumptions were not met, but for Eurasian Oystercatcher log-transformation did not lead to improvement. Non-parametric tests were therefore used for this species to test main effects. Because these tests yielded similar results as the ANOVA, we only present the results of the latter. To test whether the presence of disturbing elements caused lower densities within 500 m from the element, we used a series of χ^2 -tests. As a preliminary log-linear analysis showed no significant interactions between variables, separate χ^2 -tests were made. This could only be performed for all species pooled due to the small amount of data for each species separately. To test for species-specific avoidance distances, we included species as a third independent variable.

RESULTS

Visual inspection of Table 1 shows a large variation in the distance that meadow birds keep from view-obstructing and flat elements. The largest disturbance distance (>1 km) was reported for the high-intensively used highways. However, flat elements with low traffic intensity like cycle paths also resulted in disturbance distances of 100 m or more. Among view-obstructing elements, 'edge of a village or city' caused the largest disturbance distance. Forest edges, hedges or reed beds, – elements without traffic – showed disturbance distances of up to 350 m (Table 1). Traffic intensity was the only parameter that influenced breeding distance significantly ($\chi^2 = 8.85$, $P = 0.012$). At the lowest traffic intensity, disturbance distance was less than 500 m in 71% of the cases, whereas at intermediate and high traffic intensities this value was as low as 30% and 18%, respectively.

Eurasian Oystercatcher showed the smallest disturbance distance in nine between-species comparisons, Northern Lapwing in three comparisons, and Black-tailed Godwit in none of the comparisons (Table 1).

ANOVA-tests confirmed that Northern Lapwing and Black-tailed Godwit were significantly affected by traffic intensity, but Eurasian Oystercatcher was not (Table 2). In Northern Lapwing, post-hoc tests showed

Table 1. Maximum disturbance distance (m) from landscape elements for three species of meadow bird in The Netherlands during the breeding season. (A) References do not distinguish between species; (B) References distinguish between species. Figures between parentheses indicate approximate car traffic density, in numbers per day. Species abbreviations: Oys = Eurasian Oystercatcher, Lap = Northern Lapwing, God = Black-tailed Godwit. View obstructing: 0 = no, 1 = yes. Traffic intensity: 0 and 1 = low, 2 = intermediate, 3 = high. - = no data available.

Landscape element A	View obstructing	Traffic intensity	Distance for three species combined			Reference
Forest edge	1	0	100			Altenburg & Wymenga 1987
Tree line, hedge	1	0	50			Altenburg & Wymenga 1987
Reed bed/reedy edge	1	0	50			Brandsma 1991
'group of trees'	1	0	250–350			van 't Veer & Scharringa 2008
High dam without tree line	1	0	50			Altenburg & Wymenga 1987
'building'	1	1	175			van 't Veer & Scharringa 2008
Edge village/city	1	1	100			Altenburg & Wymenga 1987
	1	1	> 500			Wind 1977
	1	1	1200			Veen 1973
Canal without tree line	0	0	50			Altenburg & Wymenga 1987
Cycle path without tree line	0	1	100			Altenburg & Wymenga 1987
Minor/municipal road	0	1	800–900			Veen 1973
Minor road with tree line	1	1	100			Altenburg & Wymenga 1987
Highway (5000–10,000)	0	2	150			Altenburg & Wymenga 1987
	0	2	1200			Veen 1973
Railway	0	2	650			Veen 1973

Landscape element B	View obstructing	Traffic intensity	Distance for species separately			Reference
			Oys	Lap	God	
Plantation	1	0	-	300	250	van der Zande <i>et al.</i> 1980
Single farm house	1	1	150–250	250–350	350–450	Wind 1978
			630	470–630	470–875	van der Zande <i>et al.</i> 1980
Minor/municipal road	0	1	0	480	720	van der Zande <i>et al.</i> 1980
			200–300	350–450	250–350	Wind 1978
Provincial road (5000)	0	2	0	750	750	van der Zande 1975
Provincial road (10,000)	0	2	550–650	600–700	>1000	Wind 1978
Provincial road (5000; with ribbon-development)	1	2	550–650	600–700	>1000	Wind 1978
Provincial road (5000; with scattered farms)	1	2	100–200	450–550	750–850	Wind 1978
Highway (5000–10,000)	0	2	0	625	625	van der Zande <i>et al.</i> 1980
			> 625	625	625	Verstrael <i>et al.</i> 1983
			1700	120	230	Reijnen <i>et al.</i> 1996
Highway (50,000)	0	3	0	2000	1000	van der Zande 1975, van der Zande <i>et al.</i> 1980
			3530	560	930	Reijnen <i>et al.</i> 1996
			-	1800–2100	1800–2100	Verstrael <i>et al.</i> 1983
Railway	0	2	-	250	375	Grontmij 1981

significant differences in disturbance distance between the three categories, with shorter distances at low intensities and larger distances at high intensities. In Black-tailed Godwit, distances were longer at high traffic intensities in comparison to intermediate and low traffic intensity. Disturbance distances were not affected by view-obstructing elements (Table 2). When all species were pooled, no significant effect of any of the landscape element characteristics were found (Table 2).

DISCUSSION

Possible ecological explanations

Our review supports the notion that Northern Lapwing and Black-tailed Godwit keep distances from elements with elevated traffic intensity. Forman & Alexander (1998) list several factors that can cause disturbance by roads for breeding birds: pollutants, noise, direct human presence, and loss of visibility. Below we discuss the relevance of these four factors for the disturbance distances that we found. (1) Pollutants like lead have an effect over distances that are much smaller than the disturbance distances than we compiled from literature (cf. Reijnen *et al.* 1996 and Table 1). Furthermore, an effect of pollutants does not explain distances as observed for natural elements like forests. This factor does not seem to have much weight in explaining the distribution of meadow birds. (2) Reijnen *et al.* (1996) postulated that noise, rather than visual stimuli, explained disturbance distances by roads. Indeed, we found the largest disturbance distance for highways, which can be considered the noisiest landscape element (Table 1). Despite this, noise may not be the only explanation for meadow bird species as they use visual cues as well as sounds in e.g. their display behaviour. (3) Similarly, disturbance by man is an important cause of disturbance given that the largest disturbance distances were associated with highest traffic intensity (Table 1). Also,

objects with the lowest traffic intensity, like minor roads and cycle paths, have smaller disturbance distances than for instance highways (Table 1). However, as human disturbance is often a combination of noise and traffic it is difficult to distinguish between the most important underlying reason. (4) Because meadow birds give priority to detect predators (van der Vliet *et al.* 2008), lack of visibility to scan for predators, must also partially explain disturbance distances for meadow birds – in line with the fourth explanation given by Forman & Alexander (1998). Indeed, elements not used by man equally had large disturbance distances.

In landscapes with view-obstructing elements, meadow birds prefer breeding further away from perches used by avian predators, resulting in decreased predation rates (Berg *et al.* 1992, Johansson 2001, contra Ottvall *et al.* 2005, Wallander *et al.* 2006). For predators, a landscape with elements is attractive due to availability of suitable nesting sites and, for avian predators, perching opportunities (Preston 1957, Galbraith 1989, Andr  n 1992, Brandsma 1992). Furthermore, predators may use landscape elements as a hiding place when attacked by meadow birds (Elliot 1985). Finally, an open landscape may make avian predators more susceptible to stronger wind blows, which reduces the success rate of their attacks on potential prey (Quinn & Cresswell 2004). It is therefore not surprising that predation rates for meadow birds have been found to be especially high in a landscape with scattered trees, tree lines or other elements (Wymenga & Engelmoer 2001). More research is needed to elucidate the ecological understanding of disturbance distances caused by landscape elements to meadow birds, e.g. the differences between element types, and also the influence of elements for each species separately.

Data variability and uncertainty

There are several weaknesses in the data sets we used (Table 1; also van der Zande *et al.* 1980, Altenburg &

Table 2. Effects of view obstruction and traffic intensity of landscape elements on breeding distances of Eurasian Oystercatcher, Northern Lapwing, Black-tailed Godwit and all species pooled (tested by ANOVAs). Significant effects are highlighted in bold.

	Eurasian Oystercatcher* (n = 13)		Northern Lapwing** (n = 16)		Black-tailed Godwit (n = 16)		All species pooled** (n = 45)	
	F	P	F	P	F	P	F	P
Obstruction	0.003	0.956	0.058	0.813	0.329	0.577	1.172	0.285
Traffic intensity	0.579	0.580	5.067	0.025	5.212	0.023	0.432	0.652
Obstruction × Traffic intensity	0.127	0.730	0.326	0.579	0.845	0.376	0.018	0.894

*Data assumptions violated; ** log-transformed.

Wymenga 1987). Most importantly, in most studies meadow birds were studied as one group for which the response by an element was considered the same for all species (e.g. Veen 1973, Altenburg & Wymenga 1987, Brandsma 1991). However, in reality the impact of the element is likely to be species specific (e.g. van der Zande *et al.* 1980, Reijnen *et al.* 1996). Another point of concern is a very high variability in disturbance distances. This effect is most obvious for the disturbance caused by roads. For instance for Eurasian Oystercatcher, estimates for disturbance distance from highways with 5000 to 10,000 cars a day vary from 0 to 1700 meters (Table 1). Furthermore, it should be noted that several studies did not take into account effects by direct human disturbance. Finally, in some studies, data have been collected in relatively small areas, where other factors, not quantified or perhaps overlooked, may have played a role.

Implications for conservation of the landscape

Our review shows that still little is known how elements in the landscape influence the breeding distribution of meadow birds. Highways resulted in the highest reported disturbance distances (Table 1). For the western part of The Netherlands, 16% less pairs of Black-tailed Godwit were present because of the disturbance distance along highways (Reijnen *et al.* 1997). We presume that with the ongoing alteration of the landscape even fewer pairs are breeding as other elements add to unsuitable habitat. Larsen & Madsen (2000) similarly demonstrated the influence of view-obstructing elements in open grassland habitats for wintering Pink-footed Goose *Anser brachyrhynchus*. They calculated an effective loss of 68% of the total field area due to the presence of view-obstructing elements.

In this respect, the extreme site fidelity shown by meadow birds to the previous year's breeding site, especially when a breeding attempt has been successful (Klomp 1954, Thompson & Hale 1989, Groen 1993, Johansson 2001), may be a disadvantageous trait in a changing landscape. It may lead to a higher extinction risk of local meadow bird populations (Kruk *et al.* 1998). In contrast, in an unchanged landscape, extreme site fidelity may be beneficiary from an evolutionary point-of-view, as the birds are familiar with local environmental factors and, possibly, conspecifics (Johansson 2001). Because of site fidelity, not only continuity of the management of agricultural fields (Beintema & Rijk 1988, Beintema *et al.* 1995) but also the conservation of an open landscape appears to be essential for the conservation of meadow birds (van der Vliet *et al.* 2008). We consider that the adverse effect of

development on openness, leading to a reduced predator visibility, may become a more important factor for meadow birds in the near future.

On the positive side, meadow birds are known to have adapted to changes in their breeding habitat. Mulder (1972) and Beintema *et al.* (1985) showed that meadow birds were able to respond to the intensification of agricultural management by breeding earlier in response to an earlier start of agricultural activities. Likewise, many pairs of Eurasian Oystercatcher and Northern Lapwing breed in arable fields nowadays (Klomp 1954, Imboden 1970, Kooiker 1984, Galbraith 1989, Berg *et al.* 1992), although these are still shunned by grassland specialists like Black-tailed Godwit (Mulder 1972). Incidentally, Eurasian Oystercatcher and Northern Lapwing are generally also the least affected by the presence of landscape elements, whereas Black-tailed Godwit seems to be most affected. It seems that meadow birds may also be able to cope with the loss of predator visibility. De Molenaar *et al.* (2000), for instance, found that birds that arrived later at their breeding area chose their breeding sites close to road illumination, because the earlier birds already bred at sites the farthest from this object. In this way, the later birds, having the choice between no breeding or breeding at a less favourable site, showed to have adapted to a certain extent to the presence of the disturbing object.

Many policy measures have been taken by the Dutch government to preserve meadow bird species, and their habitat, because they are internationally important. Such measures have often been directed to the improvement of nesting and recruitment success, and include for instance creation of reserves, facilitation of nest protection schemes by volunteers, payment to farmers for successfully protecting nests while carrying out agricultural activities, and payment for delay of agricultural activities during the breeding season. However, up to now such measures did not stop the decline of meadow bird species (e.g. Kleijn *et al.* 2001, Berendse *et al.* 2004, Schekkerman 2008). One of the reasons might be that the landscape context has so far not been taken into account, making an effort to improve the suitability of breeding sites less successful. For meadow bird conservation, priority should be given to the protection of open and quiet landscapes (without much human disturbance). If such landscapes are to be disclosed, sources of disturbance of the landscape, stemming from a loss of openness or from the presence of human beings, should perhaps be concentrated. It would make one edge of the meadow bird habitat disturbed but would leave the remainder open and

undisturbed. We argue that, before taking measures directed to the benefit of breeding meadow birds, the already present human disturbance in and the openness of the landscape should be considered.

ACKNOWLEDGEMENTS

We thank Margien Bootsma, Tom van der Have, Arie van Noordwijk, Max Rietkerk, Peter de Ruiter, Eddy Wymenga and two anonymous referees for their useful comments on earlier drafts of this text.

REFERENCES

- Altenburg W. & Wymenga E. 1987. Natuurwetenschappelijk onderzoek voor de evaluatie van het beheersplan Midden-Opsterland II. De vegetatie en weidevogels in relatie tot het graslandgebruik. Directie Beheer Landbouwgronden, Utrecht, The Netherlands.
- Andrén H. 1992. Corvid density and nest predation in relation to forest fragmentation: a landscape perspective. *Ecology* 73: 794–804.
- Beintema A.J. & Müskens G.J.D.M. 1987. Nesting success of birds breeding in Dutch agricultural grasslands. *J. Appl. Ecol.* 24: 743–758.
- Beintema A.J. & Rijk P.J. 1988. Kosten en baten van reservaat-beheer en beheersovereenkomsten in een aantal weidevogelgebieden. LEI, 's-Gravenhage/Rijksinstituut voor Natuurbeheer, Arnhem, the Netherlands.
- Beintema A.J., Beintema-Hietbrink R.J. & Müskens G.J.D.M. 1985. A shift in timing of breeding in meadow birds. *Ardea* 73: 83–89.
- Beintema A.J., Moedt O. & Ellinger D. 1995. Ecologische atlas van de Nederlandse weidevogels. Schuyt & Co, Haarlem, the Netherlands.
- Berendse F., Chamberlain D., Kleijn D. & Schekkerman H. 2004. Declining biodiversity in agricultural landscapes and the effectiveness of agri-environment schemes. *Ambio* 33: 499–502.
- Berg Å., Lindberg T. & Källebrink K.G. 1992. Hatching success of Lapwings on farmland: differences between habitats and colonies of different sizes. *J. Anim. Ecol.* 61: 469–476.
- Bijlsma R.G., Hustings F. & Camphuysen C.J. 2001. Common and scarce birds of the Netherlands / Algemene en schaarse vogels van Nederland. Avifauna van Nederland 2. GMB Uitgeverij, Haarlem/KNNV Uitgeverij, Utrecht, the Netherlands.
- Brandsma O.H. 1991. Weidevogelbeheer in het relatienotage-bied Giethoorn-Wanneperveen. *Levende Natuur* 92: 5–12.
- Brandsma O.H. 1992. Onderzoek weidevogelbeheer en bodem-fauna in het reservaatgebied Giethoorn-Wanneperveen V (1991). Directie Beheer Landbouwgronden, Utrecht/ Provincie Overijssel, Zwolle, the Netherlands.
- de Molenaar J.G., Jonkers D.A. & Sanders M.E. 2000. Road illumination and nature 3. Local influence of road lights on a Black-tailed Godwit (*Limosa l. limosa*) population. *Alterra, Wageningen, the Netherlands*.
- Dyrce A., Witkowski J. & Okulewicz J. 1981. Nesting of 'timid' waders in the vicinity of 'bold' ones as an antipredator adaptation. *Ibis* 123: 542–545.
- Elliot R.D. 1985. The exclusion of avian predators from aggregations of nesting Lapwings (*Vanellus vanellus*). *Anim. Behav.* 33: 308–314.
- Forman R.T.T. & Alexander L.E. 1998. Roads and their major ecological effects. *Ann. Rev. Ecol. Syst.* 29: 207–231.
- Galbraith H. 1989. Arrival and habitat use by Lapwings *Vanellus vanellus* in the early breeding season. *Ibis* 131: 377–388.
- Gill J.A., Sutherland W.J. & Watkinson A.R. 1996. A method to quantify the effects of human disturbance on animal populations. *J. Appl. Ecol.* 33: 786–792.
- Groen N.M. 1993. Breeding site tenacity and natal philopatry in the Black-tailed Godwit *Limosa l. limosa*. *Ardea* 81: 107–113.
- Grontmij 1981. Ecologische onderzoeken in het 'oude land' naar aanleiding van de aanleg van de Flevolijn. Grontmij, Zeist, the Netherlands.
- Imboden Chr. 1970. Zur Ökologie einer Randzonen-Population des Kiebitz *Vanellus vanellus* in der Schweiz. *Ornithologische Beobachter* 67: 41–58.
- Johansson T. 2001. Habitat selection, nest predation and conservation biology in a Black-tailed Godwit (*Limosa limosa*) population. PhD thesis, Uppsala University, Sweden.
- Kleijn D., Berendse F., Smit R. & Gilissen N. 2001. Agri-environment schemes do not effectively protect biodiversity in Dutch agricultural landscapes. *Nature* 413: 723–725.
- Klomp H. 1951. Over de achteruitgang van de Kievit, *Vanellus vanellus* (L.), in Nederland en gegevens over het legmechanisme en het eiproductie-vermogen. *Ardea* 39: 143–182.
- Klomp H. 1954. De terreinkeus van de Kievit, *Vanellus vanellus* (L.). *Ardea* 42: 1–139.
- Kooiker G. 1984. Brutökologische Untersuchungen an einer Population des Kiebitzes (*Vanellus vanellus*). *Vogelwelt* 105: 121–137.
- Kruk M. 1993. Meadow bird conservation on modern commercial dairy farms in the western peat district of the Netherlands: possibilities and limitations. PhD-thesis, Rijksuniversiteit Leiden, the Netherlands.
- Kruk M., Noordervliet M.A.W. & ter Keurs W.J. 1998. Natal philopatry in the Black-tailed Godwit *Limosa limosa* L. and its possible implications for conservation. *Ring. Migrat.* 19: 13–16.
- Larsen J.K. & Madsen J. 2000. Effects of wind turbines and other physical elements on field utilization by pink-footed geese (*Anser brachyrhynchus*): a landscape perspective. *Landscape Ecol.* 15: 755–764.
- Mulder Th. 1972. De Grutto (*Limosa limosa* (L.)) in Nederland. Aantallen, verspreiding, terreinkeuze, trek en overwintering. Wetenschappelijke mededelingen 90. KNNV, Hoogwoud, the Netherlands.
- Newton I. 2004. The recent declines of farmland bird populations in Britain: an appraisal of causal factors and conservation actions. *Ibis* 146: 579–600.
- Ottvall R., Larsson K. & Smith H.G. 2005. Nesting success in Redshank *Tringa totanus* breeding on coastal meadows and the importance of habitat features used as perches by avian predators. *Bird Study* 52: 289–296.
- Preston F.W. 1957. The look-out perch as a factor in predation by Crows. *Wilson Bull.* 69: 368–370.

- Quinn J.L. & Cresswell W. 2004. Predator hunting behaviour and prey vulnerability. *J. Anim. Ecol.* 73: 143–154.
- Reijnen R., Foppen R. & Meeuwsen H. 1996. The effects of traffic on the density of breeding birds in Dutch agricultural grasslands. *Biol. Conserv.* 75: 255–260.
- Reijnen R., Foppen R. & Veenbaas G. 1997. Disturbance by traffic of breeding birds: evaluation of the effect and considerations in planning and managing road corridors. *Biodiv. Conserv.* 6: 567–581.
- Schekkerman H. 2008. Precocial problems: shorebird chick performance in relation to weather, farming, and predation. PhD-thesis, University of Groningen, the Netherlands.
- Thompson P.S. & Hale W.G. 1989. Breeding site fidelity and natal philopatry in the Redshank *Tringa totanus*. *Ibis* 131: 214–224.
- van Beusekom R., Huigen P., Hustings F., de Pater K. & Thissen J. (eds) 2005. Rode Lijst van de Nederlandse Broedvogels. Tirion, Baarn.
- van der Vliet R.E., Schuller E. & Wassen M.J. 2008. Avian predators in a meadow landscape: consequences of their occurrence for breeding open-area birds. *J. Avian Biol.* 39: 523–529.
- van der Zande A.N. 1975. De beïnvloeding van weidevogels door wegen. Master Thesis, Rijksuniversiteit Leiden, Leiden, the Netherlands.
- van der Zande A.N., ter Keurs W.J. & van der Weijden W.J. 1980. The impact of roads on the densities of four bird species in an open field habitat – evidence of a long-distance effect. *Biol. Conserv.* 18: 299–321.
- van 't Veer R. & Scharringa C.J.G. 2008. Weidevogelonderzoek Laag Holland 2006. Landschap Noord-Holland, Castricum, the Netherlands.
- Veen J. 1973. The disturbance of meadow bird populations. *Stedebouw & Volkshuisvesting* 54: 16–26.
- Verstrael T., ter Keurs W., van der Zande A. & van der Weijden W. 1983. De verstoring van weidevogelpopulaties door wegen. *Vogeljaar* 31: 138–151.
- Vickery J.A., Tallowin J.R., Feber R.E., Asteraki E.J., Atkinson P.W., Fuller R.J., & Brown V.K. 2001. The management of lowland neutral grasslands in Britain: effects of agricultural practices on birds and their food resources. *J. Appl. Ecol.* 38: 647–664.
- Wallander J., Isaksson D. & Lenberg T. 2006. Wader nest distribution and predation in relation to man-made structures on coastal pastures. *Biol. Conserv.* 132: 343–350.
- Wind H.B. 1977. De invloed van wegen en boerderijen op de verspreiding en het voorkomen van weidevogels. Master Thesis, Landbouw Hogeschool, Wageningen, the Netherlands.
- Wind H. 1978. De invloed van wegen en boerderijen op de verspreiding van weidevogels. *Vanellus* 31: 72–76.
- Wymenga E. & Engelman M. 2001. Takomst foar de Skries. Altenburg & Wymenga, Veenwouden, the Netherlands.

SAMENVATTING

Weidevogels hebben een voorkeur voor een open landschap met zo min mogelijk landschapselementen die hun blikveld in de weg staan. Op deze manier kunnen predatoren op tijd worden opgemerkt en weggejaagd. Ook platte landschapselementen, zoals wegen, zouden een verstoring van het blikveld kunnen opleveren door de bewegingen van de gebruikers van die elementen op de achtergrond. Het resultaat van deze verstoringen zou kunnen zijn dat weidevogels een afstand tot dergelijke landschapselementen aanhouden bij de selectie van een geschikte plek om te broeden, met als gevolg een verminderde dichtheid in vergelijking tot een situatie waarbij deze elementen er niet zouden zijn geweest. Wij hebben in de literatuur gezocht of hiervoor aanwijzingen zijn te vinden voor Scholekster *Haematopus ostralegus*, Kievit *Vanellus vanellus* en Grutto *Limosa limosa*. De resultaten van dit literatuuronderzoek laten zien dat deze soorten inderdaad een afstand aanhouden tot beide typen landschapselementen. Voor snelwegen en stads- en dorpsranden kan die afstand meer dan een kilometer bedragen. De grootste verstoringafstand werd gevonden voor wegen met een hoge gebruikersdruk. Maar ook natuurlijke elementen, zoals bosranden en rietvelden, hebben een negatieve invloed op vestigingsmogelijkheden voor weidevogels. De soorten verschillen wel in hun tolerantie ten opzichte van de vestigingsmogelijkheid in zulke gebieden. De Scholekster is het meest tolerant, de Grutto het minst. Wij beargumenteren waarom de inrichting van het landschap aandacht verdient tijdens de planning van beleid.

Corresponding editor: Julia Stahl

Received: 8 July 2008; provisionally accepted 5 May 2009; accepted 2 September 2010