

Calibration of computer models

A Closer Look at the Discrepancy Function

Pierre BARBILLON

Fall 2023, école ETICS

université
PARIS-SACLAY

AgroParisTech
Talents d'une planète soutenable

INRAE

Outline

- 1 Validation
- 2 Robust Calibration
- 3 Model Selection
 - Bayes Factor
 - Mixture model
- 4 Posterior Inclusion Probabilities of input variables in the discrepancy

Outline

1 Validation

2 Robust Calibration

3 Model Selection

- Bayes Factor
- Mixture model

4 Posterior Inclusion Probabilities of input variables in the discrepancy

Validation in [Bayarri et al., 2007] by examining the Discrepancy

Estimate tolerance bounds by computing: For a fixed level γ , the tolerance bounds $\tau = \tau(\mathbf{x})$ are then computed such that $\gamma \cdot 100\%$ of the samples satisfy:

- for pure simulator predictions $\left| \hat{f}(\mathbf{x}_{new}, \hat{\theta}) - \zeta^{(i)}(\mathbf{x}_{new}) \right| < \tau$
- for bias-corrected predictions $\left| \hat{\zeta}(\mathbf{x}_{new}) - \zeta^{(i)}(\mathbf{x}_{new}) \right| < \tau$

where

- $\hat{f}(\mathbf{x}_{new}, \hat{\theta}) = m_D(\mathbf{x}_{new}, \hat{\theta})$ where $\hat{\theta}$ may refer to the posterior mean,
- $\zeta^{(i)}(\mathbf{x}_{new}) = F^{(i)}(\mathbf{x}_{new}, \theta^{(i)}) + \delta^{(i)}(\mathbf{x}_{new})$,
- $F^{(i)}(\mathbf{x}_{new}, \theta^{(i)})$ and $\delta^{(i)}(\mathbf{x}_{new})$ ($1 \leq i \leq N$) are obtained by an MCMC algorithm sampling from the joint posterior predictive distribution,
- $\hat{\zeta}(\mathbf{x}_{new}) = \frac{1}{N} \sum_{i=1}^N \left(F^{(i)}(\mathbf{x}_{new}, \theta^{(i)}) + \delta^{(i)}(\mathbf{x}_{new}) \right)$.

Note that the discrepancy can be estimated from the posterior model and discrepancy sampling on a set of new locations: \mathbf{X}_{new} :

$$\hat{\delta}_{\hat{\theta}} = \hat{\zeta}_{new} - \hat{f}(\mathbf{x}_{new}, \hat{\theta}).$$

Outline

1 Validation

2 Robust Calibration

3 Model Selection

- Bayes Factor
- Mixture model

4 Posterior Inclusion Probabilities of input variables in the discrepancy

L_2 Calibration

Defined in [Tuo and Wu, 2016]:

$$\theta_{L_2} = \underset{\Theta}{\operatorname{argmin}} \|\delta_{\theta}(\cdot)\|_{L_2(\mathcal{X})} = \underset{\Theta}{\operatorname{argmin}} \left(\int_{\mathcal{X}} (\zeta(\mathbf{x}) - f(\mathbf{x}, \theta))^2 d\mathbf{x} \right)^{1/2}.$$

[Tuo et al., 2015] proposes to first obtain an estimate $\hat{\zeta}$ of the reality ζ via a Gaussian stochastic process and then plug it into the minimization problem to get $\hat{\theta}_{L_2}$. Consistent estimation $\hat{\theta}_{L_2} \rightarrow \theta_{L_2}$ provided that $\hat{\zeta}$ is good approximation. An alternative least square :

$$\hat{\theta}_{LS} = \underset{\Theta}{\operatorname{argmin}} \left(\sum_{i=1}^{n_e} (y_i^e - f(\mathbf{x}^e, \theta))^2 \right)$$

[Tuo et al., 2015] proves the convergence $\hat{\theta}_{LS} \rightarrow \theta_{L_2}$
 [Wong et al., 2017] uses LS calibration as a plug-in estimator for estimating the discrepancy function via a nonparametric regression

Scaled Gaussian Process

[Gu and Wang, 2018]

$$\begin{aligned}
 y_i^e &= f(\mathbf{x}_i^e, \boldsymbol{\theta}) + \mu^\delta(\mathbf{x}_i^e) + \delta_z(\mathbf{x}_i^e) + \epsilon_i \\
 \mu^\delta(\mathbf{x}) &= \sum_{i=1}^q h(\mathbf{x})\beta_i \\
 \delta_z(\cdot) &\sim GP(0, \sigma_\delta^2 c_\delta(\cdot, \cdot)) \text{ s.t. } \int_{\mathcal{X}} \delta_z(\mathbf{x})^2 d\mathbf{x} = Z \\
 Z &\sim p_{\delta_z}(\cdot), \quad p_{\delta_z}(Z) \propto f_Z(Z = z|\lambda) \cdot p_\delta(z|\boldsymbol{\theta}, \Psi)
 \end{aligned}$$

where $p_\delta(z|\boldsymbol{\theta}, \Psi)$ is the implicit prior on Z for a GP on the discrepancy.
 Then if f_Z constant \Rightarrow Model is equivalent to KOH model.

Comparison GP with SGP




Figure 2. Fifty samples from the GaSP and discretized S-GaSP are graphed in the left and right panels, respectively, where x_i is equally spaced in $[0, 1]$. For both processes, we let $\mu^\delta = 0$, $\sigma_\delta^2 = 1$ and $\gamma^\delta = 1/2$. In the discretized S-GaSP, $\mathbf{x}_i^C = \mathbf{x}_i$ for $i = 1, \dots, N_C$, $N_C = n$ and $\lambda = n/2$ are assumed..

from [Gu and Wang, 2018].

Outline

1 Validation

2 Robust Calibration

3 **Model Selection**

- Bayes Factor
- Mixture model

4 Posterior Inclusion Probabilities of input variables in the discrepancy

Outline

1 Validation

2 Robust Calibration

3 **Model Selection**

- **Bayes Factor**

- Mixture model

4 Posterior Inclusion Probabilities of input variables in the discrepancy

Model Comparison

[Damblin et al., 2016]

- $\mathcal{H}_0 : \zeta(\cdot) = f(\cdot, \theta^*)$ for a “true” θ^* :

$$y_i^e = f(\mathbf{x}_i^e, \theta^*) + \epsilon_i^0,$$

where $\epsilon_i^0 \stackrel{iid}{\sim} \mathcal{N}(0, \lambda_0^2)$.

- \mathcal{H}_1 : **Code discrepancy** term $\delta(\mathbf{x})$ s.t. $\zeta(\mathbf{x}) = f(\mathbf{x}, \theta^*) + \delta(\mathbf{x})$:

$$y_i^e = f(\mathbf{x}_i^e, \theta^*) + \delta(\mathbf{x}_i^e) + \epsilon_i^1 \quad \text{where } \delta(\cdot) \sim \mathcal{GP}(0, \sigma_\delta^2 \Sigma_\psi(\cdot, \cdot))$$

$$\text{and } \epsilon_i^1 \stackrel{iid}{\sim} \mathcal{N}(0, \lambda_1^2)$$

Bayes Factor

$$B_{0,1}(\mathbf{y}^e) := \frac{p(\mathbf{y}^e | \mathcal{H}_0)}{p(\mathbf{y}^e | \mathcal{H}_1)} \quad \text{where} \quad p(\mathbf{y}^e | \mathcal{H}_j) = \int_{\mathbf{p}_j} p(\mathbf{y}^e | \mathbf{p}_j, \mathcal{H}_j) \pi(\mathbf{p}_j) d\mathbf{p}_j.$$

Intrinsic Bayes Factor

[Berger and Pericchi, 1996]

Main issue: Evidence $p(\mathbf{y}^e | \mathcal{H}_j)$ sensitive to priors $\pi(\mathbf{p}_j)$.

- Need to use compatible priors [Celeux et al., 2006] or objective priors [Casella and Moreno, 2006],
- but marginal likelihood ill-defined (up to arbitrary constant) for improper priors (as objective priors often are).

Idea: using a part of data to obtain a proper prior:

- Partial Bayes Factor:

$$B_{0,1}(\mathbf{y}^e(-m) | \mathbf{y}^e(m)) = \frac{\int l(\mathbf{p}_0; \mathbf{y}^e(-m) | \mathbf{y}^e(m)) \pi(\mathbf{p}_0 | \mathbf{y}^e(m)) d\mathbf{p}_0}{\int l(\mathbf{p}_1; \mathbf{y}^e(-m) | \mathbf{y}^e(m)) \pi(\mathbf{p}_1 | \mathbf{y}^e(m)) d\mathbf{p}_1} = \frac{B_{0,1}(\mathbf{y}^e)}{B_{0,1}(\mathbf{y}^e(m))}.$$

- $B_{0,1}(\mathbf{y}^e(-m) | \mathbf{y}^e(m))$ well-defined for $|m| \geq n_0$ large enough:
- Intrinsic Bayes factor obtained by averaging over all $\mathbf{y}^e(m)$ s :

$$B_{0,1}^A(\mathbf{y}^e) = \frac{B_{0,1}(\mathbf{z})}{C(n, n_0)} \sum_{|m|=n_0} B_{0,1}(\mathbf{y}^e(m))^{-1}.$$

IBF computation under linearization of the code

Linear assumption: $f(\mathbf{x}, \theta) = g(\mathbf{x})^\top \theta$, with $g(\mathbf{x}) \in \mathbb{R}^d$.

Prior choices and consequences:

- Model \mathcal{H}_0 boils down to:

$$\mathcal{H}_0 : \mathbf{y}^e \sim \mathcal{N}(G\theta_0; \lambda_0^2 \mathbf{I}_{n_e}); \quad \mathbf{p}_0 = (\theta_0, \lambda_0^2)$$

where $G = [g(\mathbf{x}_1^e), \dots, g(\mathbf{x}_{n_e}^e)]^\top$ the $n_e \times p$ design matrix.

→ Under Jeffreys prior: $\pi(\mathbf{p}_0) \propto \lambda_0^{-2}$, $p(\mathbf{y}^e | \mathcal{H}_0)$ explicit.

- Model \mathcal{H}_1 boils down to:

$$\mathcal{H}_1 : \mathbf{y}^e \sim \mathcal{N}(G\theta_1; \sigma_\delta^2 V_{k,\psi}); \quad \mathbf{p}_1 = (\theta_1, \sigma_\delta^2, \psi, k)$$

$$V_{k,\psi}(i, j) = k\delta_{i,j} + e^{-\|\mathbf{x}_i - \mathbf{x}_j\|^2 / \psi^2} \quad k = \lambda_1^2 \sigma^{-2}.$$

- Prior choice: $\pi(\mathbf{p}_1) \propto \pi(\psi|k)\pi(k)\sigma^{-2}$ with proper priors for $\pi(\psi|k)\pi(k)$,
- Integration of $p(\mathbf{y}^e | \mathbf{p}_1, \mathcal{H}_1)$: explicit over $(\theta_1, \sigma_\delta^2)$, by Gaussian quadrature over (ψ, k) .

Computation of the IBF

Proposition

If $\pi(\mathbf{p}_1) = \pi(\theta_1, \sigma_\delta^2, \psi, k) = \pi(\psi|k)\pi(k)/\sigma_\delta^2$, $\pi(\psi, k)$ is proper and $m = d + 1$ then

$$B_{0,1}^A(\mathbf{y}^e) = \frac{B_{0,1}(\mathbf{z})}{C(n, n_0)} \sum_{|m|=n_0} B_{0,1}(\mathbf{y}^e(m))^{-1} = B_{0,1}(\mathbf{y}^e)$$

In the following,

$$\begin{aligned}\pi(\psi|k) &= \mathcal{U}([0, 1]), \\ \pi(k) &= Be(1, 3).\end{aligned}$$


Synthetic data

Data simulated according to model \mathcal{H}_1 , with $\delta \sim GP(0, \sigma_\delta^2 \Sigma_\psi)$:

$$\mathbf{x}_i^e = \left(\frac{i}{n_e} \right)_{1 \leq i \leq n}, \quad n_e = 30, \quad \sigma_\delta^2 = 0.1, \quad k = 0.1.$$



From left to right

- constant trend $g(\mathbf{x}) = 1$; $\theta_1 = 1$,
- linear trend $g(\mathbf{x}) = (1, x)$; $\theta_1 = (1, 1)$,
- quadratic trend $g(\mathbf{x}) = (1, x, x^2)$; $\theta_1 = (1, 1, 1)$.
- Bayes factor $B_{0,1}^A$ expected to decrease with ψ .



Boxplots of $B_{0,1}^A(\mathbf{y}^e)$ values over 100 simulations with constant, linear and quadratic trends (left to right)

Confounding Effect



$\psi = 0.2$ left and $\psi = 0.7$ right

- ψ, k, σ_δ^2 estimated by maximum likelihood.
- For $\psi = 0.7$, discrepancy indistinguishable from quadratic trend!

Case description

- Industrial computer code predicting the **productivity** of an electric power plant, based on measurements (temperature, pressure, discharge, . . .) throughout the plant
- $n_e = 24$ available field measures (results of periodic testing) to validate code


Main code features:

- $p = 20$ input variables ($\mathbf{x} \in \mathbb{R}^{20}$)
- $d = 2$ parameters: heat transfer coefficient of the condenser, yield of the main turbine 2 .
- Two outputs of interest (electric power, condenser pressure), seen here as two separate codes
- Code **linearized** in neighbourhood of reference value θ^* :

$$f(\mathbf{x}_i, \theta) \approx f(\mathbf{x}_i, \theta^*) + h(\mathbf{x}_i)^\top (\theta - \theta^*),$$


where $h(\mathbf{x}_i) = \nabla_\theta f(\mathbf{x}_i, \theta^*)$ evaluated numerically through finite difference

Calibrated code predictions vs measures



Pressure

$$B_{0,1}^A = 2 \times 10^{-18}$$



Power

$$B_{0,1}^A = 3 \times 10^{-3}$$

- Bias reduced by calibration, but not suppressed
- strong evidence for code discrepancy

Outline

1 Validation

2 Robust Calibration

3 **Model Selection**

- Bayes Factor
- **Mixture model**

4 Posterior Inclusion Probabilities of input variables in the discrepancy

Model selection as a mixture problem

[Kamary et al., 2019] inspired by [Kamary et al., 2014].

Model selection problem:

$$\mathfrak{M}_0 : y_i^e = f(\mathbf{x}_i^e, \boldsymbol{\theta}_0) + \epsilon_i^0$$

$$\mathfrak{M}_1 : y_i^e = f(\mathbf{x}_i^e, \boldsymbol{\theta}_1) + \delta(\mathbf{x}_i^e) + \epsilon_i^1.$$

where $\epsilon_i^0 \stackrel{iid}{\sim} \mathcal{N}(0, \lambda_0^2)$ and $\epsilon_i^1 \stackrel{iid}{\sim} \mathcal{N}(0, \lambda_1^2)$

converted into a mixture model:

$$\mathfrak{M}_\alpha : y_i \sim \alpha \left(\ell_{\mathfrak{M}_0}(\boldsymbol{\theta}_0, \lambda_0^2; y_i^e, \mathbf{x}_i^e) \right) + (1 - \alpha) \left(\ell_{\mathfrak{M}_1}(\boldsymbol{\theta}_1, \lambda_1^2, \delta; y_i^e, \mathbf{x}_i^e) \right).$$

- Model \mathfrak{M}_α is defined under the hypothesis that the likelihood of the model \mathfrak{M}_1 is conditioned on δ .
- δ is considered as a parameter of \mathfrak{M}_1 .
- Conditionnally on δ , the y_i 's are considered independent.
- Posterior distribution on α will provide a decision rule for \mathfrak{M}_0 against \mathfrak{M}_1 .

Hypotheses and prior distribution

- Linear code: $f(\mathbf{x}, \boldsymbol{\theta}) = g(\mathbf{x})^\top \boldsymbol{\theta}$.

- GP prior for discrepancy function:

$$\delta(\cdot) \sim \mathcal{GP}(0, \sigma_\delta^2 \Sigma_\psi(\cdot, \cdot)).$$

- Some parameters are common, $\boldsymbol{\theta}$ and λ^2 so a common prior distribution is chosen for both.

$$\mathfrak{M}_\alpha : y_i^e \sim \alpha \left(\ell_{\mathfrak{M}_0}(\boldsymbol{\theta}, \lambda^2; y_i^e, \mathbf{x}_i^e) \right) + (1 - \alpha) \left(\ell_{\mathfrak{M}_1}(\boldsymbol{\theta}, \lambda^2, \delta; y_i^e, \mathbf{x}_i^e) \right).$$

Posterior distribution

Theorem

Let $g : \mathbb{R}^p \rightarrow \mathbb{R}^d$ be a finite-valued function and vector $\mathbf{x}_1^e, \dots, \mathbf{x}_n^e$ such that the rank of $\{g(\mathbf{x}_1^e), \dots, g(\mathbf{x}_n^e)\}$ is d . The posterior distribution associated with the prior $\pi(\boldsymbol{\theta}, \lambda^2) = 1/\lambda^2$ and with the likelihood is proper when

- for any $0 < k < 1$, the hyperparameter σ_δ^2 of the discrepancy prior distribution is reparameterized as $\sigma_\delta^2 = \lambda^2/k$ and so $\Sigma_\psi = (\lambda^2/k)\text{Corr}_{\psi_\delta}$ when $\text{Corr}_{\psi_\delta}$ is the correlation function of δ .
- the mixture weight α has a proper beta prior $\mathcal{B}(a_0, a_0)$;
- ψ_δ has a proper Beta prior $\mathcal{B}(b_1, b_2)$.
- proper distribution is used on k .

Metropolis within Gibbs

Algorithm 1: Metropolis-within-Gibbs algorithm

for $t=1, \dots, T$ do

- a) $\delta^{(t)}$ is sampled from $\pi(\delta | \mathbf{y}^e, \mathbf{X}^e, \boldsymbol{\theta}^{(t-1)}, \lambda^{(t-1)}, k^{(t-1)}, \psi_{\delta}^{(t-1)}, \alpha^{(t-1)})$ as follows.

- a.1) For $i = 1, \dots, n; j = 0, 1$, generate auxiliary variable $\nu_i^{(t)}$ from

$$\mathbb{P}(\nu_i = j | \mathbf{y}_i^e, \mathbf{x}_i^e, \delta^{(t-1)}, \boldsymbol{\theta}^{(t-1)}, \lambda^{(t-1)}, k^{(t-1)}, \psi_{\delta}^{(t-1)}) .$$

- a.2) Generate $\delta^{(t)}$ according to the conditional posterior distribution

$$\delta^{(t)} | \mathbf{y}^e, \mathbf{X}^e, \nu^{(t)} = 1, \boldsymbol{\theta}^{(t-1)}, \lambda^{(t-1)}, k^{(t-1)}, \psi_{\delta}^{(t-1)}, \alpha^{(t-1)} \sim \mathcal{N}_n(\hat{\mu}_{\delta}, \hat{\Sigma}_{\delta}) .$$

- b) Generate $\boldsymbol{\theta}^{(t)} | \mathbf{y}^e, \mathbf{X}^e, \nu^{(t)}, \delta^{(t)}, \lambda^{(t-1)}, k^{(t-1)}, \alpha^{(t-1)} \sim \mathcal{N}_d(\hat{\mu}_{\boldsymbol{\theta}}, \hat{\Sigma}_{\boldsymbol{\theta}})$.

- c) Generate $\lambda^{(t)} | \mathbf{y}^e, \mathbf{X}^e, \nu^{(t)}, \delta^{(t-1)}, \boldsymbol{\theta}^{(t)}, k^{(t-1)}, \alpha^{(t-1)} \sim \mathcal{IG}(\hat{a}_{\lambda}, \hat{b}_{\lambda})$.

- d) Generate $\alpha^{(t)} | \mathbf{y}^e, \mathbf{X}^e, \nu^{(t)}, \delta^{(t)}, \boldsymbol{\theta}^{(t)}, \lambda^{(t)}, k^{(t-1)} \sim \text{Beta}(n - m + a_0, m + a_0)$.

- e) Generate $k^{(t)}$ from a random walk Metropolis-Hastings algorithm conditionally to $(\mathbf{y}^e, \mathbf{X}^e, \nu^{(t)}, \delta^{(t)}, \boldsymbol{\theta}^{(t)}, \lambda^{(t)}, \alpha^{(t)}, \psi_{\delta}^{(t-1)})$.

- f) Generate $\psi_{\delta}^{(t)}$ from a random walk Metropolis-Hastings algorithm conditionally to $(\mathbf{y}^e, \mathbf{X}^e, \nu^{(t)}, \delta^{(t)}, \boldsymbol{\theta}^{(t)}, \lambda^{(t)}, \alpha^{(t)}, k^{(t)})$.

Synthetic example \mathfrak{M}_0

Code is a quadratic function.

50 datasets of size $n = 30$ from $\mathfrak{M}_0 : y_i^e = g(\mathbf{x}_i^e)^\top \boldsymbol{\theta}^* + \epsilon_i$.

Priors as in the theorem, $\alpha \sim \text{Beta}(1, 1)$, $\delta \sim \mathcal{GP}(0_n, \Sigma_\psi)$, $\psi_\delta \sim \text{Beta}(1, 1)$ and $k \sim \text{Beta}(1, 1)$.

Number of MCMC iterations is 10^4 with a burn-in of 10^3 iterations




Figure: Posterior mean estimates of $\boldsymbol{\theta}, \lambda^2$, Posterior densities of α , Posterior prediction of the code.

Synthetic example \mathfrak{M}_1

Code is a quadratic function.

50 samples of size 50 simulated from \mathfrak{M}_1 when ψ_δ^* varies between 0.01 and 0.9, $\delta^*(x) \sim \mathcal{GP}(0_n, \Sigma_\psi)$, $\lambda^{2*} = 0.1$ and $k^* = 0.1$.

Priors as in the theorem, $\alpha \sim \text{Beta}(1, 1)$, $\delta \sim \mathcal{GP}(0_n, \Sigma_\psi)$, $\psi_\delta \sim \text{Beta}(1, 1)$ and $k \sim \text{Beta}(1, 1)$.

Number of MCMC iterations is 10^4 with a burn-in of 10^3 iterations.




Figure: Posterior mean estimates for α and λ^2 .


Hydraulic application: Garonne river

- TELEMAC 2D models the flow of the Garonne between Tonneins and la Réole:



$$h_i = f(q_i, \mathbf{K}_s),$$

with:

- \mathbf{h}_i water heights,
- $\mathbf{K}_s = (K_{s1}, \dots, K_{s5})$ Strickler coefficients (5 friction coefficients)
- q_i river flow at Tonneins
- Linearization of the model around a reference value for the Strickler coefficient (limited to the most influential ones).
- Only 7 data points available.



Results



Observation nb.	1	2	3	4	5	6	7
Bias probability	0.513	0.473	0.452	0.448	0.451	0.472	0.514

Table: Probability of a code bias for each observation in Marmande

Outline

- 1 Validation
- 2 Robust Calibration
- 3 Model Selection
 - Bayes Factor
 - Mixture model
- 4 Posterior Inclusion Probabilities of input variables in the discrepancy

Variable selection in the discrepancy function

[Joseph and Yan, 2015]

- perform a sensitivity analysis on the discrepancy function,
- two-step procedure: i) find an optimal $\hat{\theta}$, ii) run an SA on the discrepancy (consequence of the fixed $\hat{\theta}$)

[Barbillon et al., 2021]

- run an MCMC algorithm to obtain posterior distribution,
- post-process the posterior samples to compute probabilities of inclusion for each of the input variable in the discrepancy.



Barbillon, P., Forte, A., and Paulo, R. (2021).
Screening the discrepancy function of a computer model.
[arXiv preprint arXiv:2109.02726](#).



Bayarri, M. J., Berger, J. O., Paulo, R., Sacks, J., Cafeo, J. A., Cavendish, J., Lin, C.-H., and Tu, J. (2007).
A framework for validation of computer models.
[Technometrics](#), 49(2):138–154.



Berger, J. O. and Pericchi, L. R. (1996).
The intrinsic bayes factor for model selection and prediction.
[Journal of the American Statistical Association](#), 91(433):109–122.



Casella, G. and Moreno, E. (2006).
Objective Bayesian variable selection.
[Journal of the American Statistical Association](#), 101(473):157–167.



Celeux, G., Marin, J.-M., and Robert, C. (2006).
Sélection bayésienne de variables en régression linéaire.
[Journal de la société française de statistique](#), 147(1):59–79.



Damblin, G., Keller, M., Barbillon, P., Pasanisi, A., and Parent, É. (2016).
Bayesian model selection for the validation of computer codes.
[Quality and Reliability Engineering International](#), 32(6):2043–2054.



Gu, M. and Wang, L. (2018).

Scaled gaussian stochastic process for computer model calibration and prediction.

[SIAM/ASA Journal on Uncertainty Quantification](#), 6(4):1555–1583.



Joseph, V. R. and Yan, H. (2015).

Engineering-driven statistical adjustment and calibration.

[Technometrics](#), 57(2):257–267.



Kamary, K., Keller, M., Barbillon, P., Goeury, C., and Éric Parent (2019).

Computer code validation via mixture model estimation.



Kamary, K., Mengersen, K., Robert, C. P., and Rousseau, J. (2014).

Testing hypotheses via a mixture estimation model.

[arXiv preprint arXiv:1412.2044](#).



Tuo, R., Wu, C. J., et al. (2015).

Efficient calibration for imperfect computer models.

[The Annals of Statistics](#), 43(6):2331–2352.



Tuo, R. and Wu, J. (2016).

A theoretical framework for calibration in computer models: parametrization, estimation and convergence properties.

[SIAM/ASA Journal on Uncertainty Quantification](#), 4(1):767–795.



Wong, R. K., Storlie, C. B., and Lee, T. (2017).

A frequentist approach to computer model calibration.

Journal of the Royal Statistical Society: Series B (Statistical Methodology),
79(2):635–648.