

Equivariant dendroidal sets and simplicial operads

Peter Bonventre, Luís A. Pereira

October 24, 2017

Abstract

bla bla, generalizing [CM13a](#)
[3].

Bla, also obtain an equivariant notion of Reedy category

Contents

1 Equivariant dendroidal sets

1.1 Preliminaries

Definition 1.1. A map $f : S_0 \rightarrow T_0$ in Ω is called a *face map* if it is injective on underlying sets. A face map is *inner* if it is of the form $T_0 \setminus E \hookrightarrow T_0$, where E is a subset of the set of inner edges of T . Fixing such a subset E , let $\Phi_{\text{Inn}}^E(T_0)$ denote the poset (under inclusion) of all inner face maps $S_0 \hookrightarrow T_0$ such that $E \subseteq T_0 \setminus S_0$.

Definition 1.2. Given $S_0 \in \Omega$, $T \in \Omega_G$, and a map of forests $f : S_0 \rightarrow T$, let T_0 denote the component of T containing the image of S_0 .

We say f is a (non-equivariant) (*inner*) *face map* if $f : S_0 \rightarrow T_0$ is a non-equivariant (inner) face map.

Given a subsequence E of inner edges, let $\Phi_{\text{Inn}}^E(T)$ denote the subposet of faces such that miss all of E .

Definition 1.3. Fix $T \in \Omega_G$, and a component T_0 of T , with $H := \text{Stab}_G(T_0)$.

Let $\Phi(T)$ denote the poset (under inclusion) of face maps whose image is strictly contained in T_0 . Given an inner edge $e \in T$, let $\Phi^{Ge}(T) = \Phi(T) \setminus (T_0 \setminus He)$. Define the *Ge-horn* of T to be the subdendroidal set

$$\Lambda^{Ge}[T] := \text{colim}_{\Phi^{Ge}(T)} \Omega[G \cdot S_0]. \quad (1.4)$$

Definition 1.5. A face map $f : S_0 \rightarrow T$ is called *orbital* if $f(S_0) \subseteq T$ is K -closed, where $K = \text{Stab}_G(f(r_s))$, for r_s the root of S_0 .

Fixing a component T_0 of T , let $\Phi_o(T)$ denote the poset (under inclusion) of orbital face maps whose image is strictly contained in T_0 . Define the *orbital boundary* of T to be the subdendroidal set

$$\partial_o \Omega[T] := \text{colim}_{\Phi_o(T)} \Omega[G \cdot S_0]. \quad (1.6)$$

Given an inner edge $e \in T$, let $\Phi_{\text{Orb}}^{Ge}(T) := \Phi_o(T) \setminus (T_0 \setminus He)$ where $H = \text{Stab}_G(T_0)$ (that is; those faces S_0 such that either $T \setminus (G \cdot f(S_0) \cup Ge) \neq \emptyset$, or outer faces removing a stump). Define the *Ge-orbital horn* to be the subdendroidal set

$$\Lambda_o^{Ge}[T] := \text{colim}_{\Phi_{\text{Orb}}^{Ge}(T)} \Omega[G \cdot S_0]. \quad (1.7)$$

Definition 1.8. Given T , T_0 , and H as above, let $U_0 \subseteq T_0$ be a (non-equivariant) subtree, and let $K = \text{Stab}_H(U_0)$. Suppose we have another subdendroidal set $X \subseteq \Omega[T]$ which contains all outer faces of U , and an edge $e \in U_0$. We say that $He \subseteq U_0$ is a *characteristic edge orbit* of $X \subseteq \Omega[T] \supseteq \Omega[G \cdot U_0]$ if we have

$$e \in R_0 \in \Phi_{\text{Inn}}(U) \cap X \text{ if and only if } R_0/\bar{K}e \in \Phi_{\text{Inn}}(U) \cap X, \quad (1.9)$$

where $\bar{K} = \text{Stab}_K(R_0)$.

Proposition 1.10. *Given T , T_0 , H , U_0 , and X as above, we have that He is a characteristic edge orbit of $X \subseteq \Omega[T] \supseteq \Omega[G \cdot U]$ implies that $X \rightarrow \mathcal{X} \cup \Omega[G \cdot U]$ is inner G -anodyne.*

Proof. If $\Omega[U]$ is already contained in X , we are done. Assuming otherwise, it suffices to show that for all $C \subseteq C'$ K -closed concave subsets of $\Phi_{\text{Inn}}^{He}(U_0) \setminus X$, the map

$$X \cup G \cdot_K \left(\bigcup_{E \in C} \Omega[U_0 \setminus E] \right) \rightarrow X \cup G \cdot_K \left(\bigcup_{E' \in C'} \Omega[U_0 \setminus E'] \right) \quad (1.11)$$

is inner G -anodyne. Indeed, once $C = \Phi_{\text{Inn}}^{He}(U_0) \setminus X$, we have the pushout

$$\begin{array}{ccc} G \cdot_K (\Lambda^{He}[U_0]) & \longrightarrow & X \cup G \cdot_K \left(\bigcup_{E \in \Phi_{\text{Inn}}^{He}(U_0)} \Omega[U_0 \setminus E] \right) \\ \downarrow & & \downarrow \\ G \cdot_K \Omega[U_0] \simeq \Omega[G \cdot U_0] & \longrightarrow & X \cup \Omega[G \cdot U_0]. \end{array} \quad (1.12)$$

Moreover, it suffices to consider $C' = C \cup H \cdot D$ for $D \subseteq U_0$, where without loss of generality $e \in D$ and $U_0 \setminus D$ is not in the domain. Let $\bar{K} = \text{Stab}_H(D)$.

We first claim that $\Lambda^{\bar{K}e}[U_0 \setminus D]$ is in the domain. If S_0 is an outer face of $U_0 \setminus D$, then S_0 factors through an outer face of U_0 , and so S_0 is in X . Further, if $S_0 = U_0 \setminus (D \cup E)$ with $E \cap Ke = \emptyset$, then concavity implies that S_0 is in the domain, as required.

Second, we claim that no face $S_0 = U_0 \setminus D \cup \bar{e}$, with $\bar{e} \subseteq Ke$, is in the domain. Suppose $U_0 \setminus D \cup \bar{e}$ is contained in some $U_0 \setminus E$ already attached. Then, since $E \cap Ke = \emptyset$, we have $U_0 \setminus D \subseteq U_0 \setminus E$, so $U_0 \setminus D$ is also in the domain, a contradiction. Further, if $U_0 \setminus D \cup \bar{e}$ is in X , then so is $U_0 \setminus D \cup Ke$, and hence so is $U_0 \setminus D$ (by definition of characteristic edge orbit), also a contradiction.

Now, all faces $U_0 \setminus D \cup \bar{e}$ with $\emptyset \subsetneq \bar{e} \subseteq Ke$ have stabilizer \bar{K} (else $D \cap Ke \neq \emptyset$). Thus the desired map is the pushout of

$$G \cdot_{\bar{K}} (\Lambda^{\bar{K}e}[U_0 \setminus D] \rightarrow \Omega[U_0 \setminus D]), \quad (1.13)$$

and hence is anodyne, as required. \square

come back

Lemma 1.14. *Suppose U_0 is a minimal outer face of T not in $\Lambda_0^{Ge}[T]$, and suppose $e \in U_0$. Then He is a characteristic edge orbit for $\Lambda_0^{Ge}[T] \subseteq \Omega[T] \supseteq [G \cdot U_0]$.*

Proof. Consider an inner face $U_0 \setminus D$ of U_0 , with $\bar{K} = \text{Stab}_H(U_0 \setminus D) \leq K$, and suppose $U_0 \setminus D \cup \bar{K}e \in \Lambda_0^{Ge}[T]$. Then

$$K_r = \text{Stab}_H(r_U) \leq \text{Stab}_H(U_0 \setminus D \cup \bar{K}e) \leq \text{Stab}_H(U_0) \leq K_r, \quad (1.15)$$

so the outer face U_0 is K_r -closed and not in $\Lambda_0^{Ge}[T]$, and hence we conclude that U_0 must be T_0 . However, if $\Lambda_0^{Ge}[T] \neq \Lambda^{Ge}[T]$, T_0 is not a minimal outer face not in $\Lambda_0^{Ge}[T]$. Thus, in these cases, He is a characteristic edge orbit vacuously. If in fact $\Lambda_0^{Ge}[T] = \Lambda^{Ge}[T]$, then $T \simeq G \cdot T_0$, so $H = \bar{K} = K_r = \{e\}$. Now, since $U_0 \setminus D \cup e \in \Lambda_0^{Ge}[T]$, $D \setminus e \neq \emptyset$, so $U_0 \setminus D \in \Lambda_0^{Ge}[T]$. \square

Proposition 1.16. $\Lambda_o^{Ge}[T] \rightarrow \Omega[T]$ is inner G -anodyne, and hence the (hyper)saturated class of the orbital horn inclusions is contained in the (hyper)saturated class of the horn inclusions.

Proof. Let $\text{Out}_o(T)$ be the poset of outer faces U_0 of T which are not in $\Lambda_o^{Ge}[T]$. It suffices to show that for any G -closed convex subsets $B \subseteq B'$ of $\text{Out}_o(T)$, the map

$$\Lambda_o^{Ge}[T] \cup \bigcup_{R_0 \in B} \Omega[G \cdot R_0] \rightarrow \Lambda_o^{Ge}[T] \cup \bigcup_{R'_0 \in B'} \Omega[G \cdot R'_0] \quad (1.17)$$

is inner G -anodyne. Again, suffices to show when $B' = B \cup \{U_0\}$. Let $K = \text{Stab}_G(U_0)$, and without loss of generality assume $e \in U_0$. The case $B = \emptyset$ is the previous lemma. For general B , we know the domain contains all outer faces of U_0 by convexity. Now, let $U_0 \setminus D$ be an inner face, with stabilizer \bar{K} . Then $U_0 \setminus D \cup \bar{K}e$ is in the domain if either $U_0 \setminus D \cup \bar{K}e$ is contained in some $\Omega[G \cdot R_0]$, or $U_0 \setminus D \cup \bar{K}e \in \Lambda_o^{Ge}[T]$. But then U_0 is contained in R_0 since both are outer faces, or again we apply the previous lemma. Both lead to contradictions, and thus we may conclude that He is a characteristic edge orbit for the domain relative to $\Omega[T] \supseteq \Omega[G \cdot U_0]$. Thus, the result holds by Proposition 1.10. \square

1.2 Actual Stuff

Notation 1.18. Given subgroups $H_i \leq G$, $0 \leq i \leq k$ such that $H_0 \geq H_i$, $1 \leq i \leq k$ we write $C_{u_i H_0/H_i}$ for the G -corolla encoding the H_0 -set $H_0/H_1 \sqcup \dots \sqcup H_0/H_k$.

Following the discussion preceding [4, Prop. 3.6.8], we will call a class of maps of \mathbf{dSet}^G *hypersaturated* if it is closed under pushouts, transfinite composition, retracts, and satisfies the following cancellation property: if in

$$A \xrightarrow{f} B \xrightarrow{g} C$$

both f and gf are in the class, then so is g .

The following is an equivariant generalization of [CM13a, Props. 2.4 and 2.5].

Proposition 1.19. *The following sets of maps generate the same hypersaturated class:*

- the G -inner horn inclusions $\Lambda^{Ge}[T] \rightarrow \Omega[T]$ for $T \in \Omega_G$ and Ge an inner edge orbit;
- the G -inner orbital horn inclusions $\Lambda_o^{Ge}[T] \rightarrow \Omega[T]$ for $T \in \Omega_G$ and Ge an inner edge orbit;
- the G -segal core inclusions $Sc[T] \rightarrow \Omega[T]$ for $T \in \Omega_G$.

HERE

Remark 1.20. Setting $G = e$ and slicing over the stick tree η in the previous result one recovers the more well known claim that the hypersaturation (in fact, saturation) of the simplicial inner horns $\{\Lambda^i[n] \rightarrow \Delta[n]: 0 < i < n\}$ coincides with the hypersaturation of the simplicial Segal core inclusions $\{Sc[n] \rightarrow \Delta[n]\}$.

Remark 1.21. We will also make use of a variant of the previous remark for the hypersaturation of *all* simplicial horns. Namely, we claim that the hypersaturation of all simplicial horns $\{\Lambda^i[n] \rightarrow \Delta[n]: 0 \leq i \leq n\}$ coincides with the hypersaturation of all vertex inclusion maps $\{\Delta[0] \rightarrow \Delta[n]\}$. Indeed, call the latter hypersaturation S . An easy argument shows that the Segal core inclusions $\{Sc[n] \rightarrow \Delta[n]\}$ are in S and thus so are all inner horn inclusions. On the other hand, the skeletal filtration of the left horns $\Lambda^0[n]$ is built exclusively out of left horn inclusions, and thus since $\Delta[0] = \Lambda^0[1] \rightarrow \Delta[1]$ is in S so are all left horn inclusions $\Lambda^0[n] \rightarrow \Delta[n]$. The case of right horn inclusions $\Lambda^n[n] \rightarrow \Delta[n]$ is dual.

HERE

The following is the equivariant generalization of [CM13a, Thm. 3.5].

Proposition 1.22. *Let $X \rightarrow Y$ be a map between G - ∞ -operads. The following are equivalent:*

(a) *for all G -corollas C_A and $H \leq G$ the maps*

$$k(\Omega[C_A], X) \rightarrow k(\Omega[C_A], Y), \quad k(\Omega[G/H \cdot \eta], X) \rightarrow k(\Omega[G/H \cdot \eta], Y)$$

are weak equivalences in \mathbf{sSet} ;

(b) *for all G -trees T the maps*

$$k(\Omega[T], X) \rightarrow k(\Omega[T], Y)$$

are weak equivalences in \mathbf{sSet} ;

(c) *for all normal G -dendroidal sets A , the maps*

$$k(A, X) \rightarrow k(A, Y)$$

are weak equivalences in \mathbf{sSet} ;

(d) *$f: X \rightarrow Y$ is a weak equivalence in \mathbf{dSet}^G .*

HERE

Definition 1.23. Let X be a G - ∞ -operad. A G -profile on X is a map

$$\partial\Omega[C] \rightarrow X$$

for some G -corolla $C \in \Sigma_G$.

More explicitly, a G -profile consists of:

- subgroups $H_i \leq G$, $0 \leq i \leq k$ such that $H_0 \geq H_i$ for $1 \leq i \leq k$;
- objects $x_i \in X(\eta)^{H_i}$ for $0 \leq i \leq k$.

To simplify notation, we will prefer to denote a G -profile as $(x_1, \dots, x_k; x_0)$, and refer to it as a C -profile.

Definition 1.24. Given a G - ∞ -operad and a C -profile $(x_1, \dots, x_k; x_0)$ we define the space of maps $X(x_1, \dots, x_k; x_0)$ to be given by the pullback

$$\begin{array}{ccc} X(x_1, \dots, x_k; x_0) & \longrightarrow & \text{Hom}(\Omega[C], X) \\ \downarrow & & \downarrow \\ G \cdot \eta & \xrightarrow{(x_1, \dots, x_k; x_0)} & \prod_{0 \leq i \leq k} X(\eta)^{H_i} \end{array}$$

Noting that there are equivalences of categories (the first of which is an isomorphism)

$$(\mathbf{dSet}_G)/G \cdot \eta \simeq \mathbf{sSet}^{B_G} \simeq \mathbf{sSet},$$

one sees that $X(x_1, \dots, x_k; x_0)$ can indeed be regarded as a simplicial set (in fact, this is a Kan complex).

Definition 1.25. Let $f: X \rightarrow Y$ be a map of G - ∞ -operads.

The map f is called *fully faithful* if, for each C -profile $(x_1, \dots, x_k; x_0)$ one has that

$$X(x_1, \dots, x_k; x_0) \rightarrow Y(f(x_1), \dots, f(x_k); f(x_0))$$

is weak equivalence in \mathbf{sSet} .

The map f is called *essentially surjective* if for each subgroup $H \leq G$ the map of categories $\tau(\iota^*(X^H)) \rightarrow \tau(\iota^*(Y^H))$ are essentially surjective.

The following is the equivariant generalization of [3, Thm. 3.11 and Remark 3.12]. CM13a

Theorem 1.26. *A map $f: X \rightarrow Y$ of G - ∞ -operads is fully faithful iff for all G -corollas $C \in \Sigma_G$ the commutative squares of Kan complexes*

$$\begin{array}{ccc} k(\Omega[C], X) & \longrightarrow & k(\Omega[C], Y) \\ p \downarrow & & \downarrow q \\ k(\partial\Omega[C], X) & \xrightarrow{f_*} & k(\partial\Omega[C], Y) \end{array} \quad (1.27) \quad \boxed{\text{COMSQ EQ}}$$

are homotopy pullback squares.

Hence, f is a weak equivalence in \mathbf{dSet}^G iff f is both fully faithful and essentially surjective.

Proof. Noting that the 0-simplices of $k(\partial\Omega[C], X)$ are precisely the C -profiles (x_1, \dots, x_k, x_0) , fully faithfulness can be reinterpreted as saying that all fiber maps $p^{-1}(x_1, \dots, x_k, x_0) \rightarrow q^{-1}(f(x_1), \dots, f(x_k), f(x_0))$ are weak equivalences. But since p, q are Kan fibrations, this is equivalent to the condition that (1.27) is a homotopy pullback (see [3, Lemma 3.9]), and the first half follows. COMSQ EQ

For the second half, note first that the bottom map in (1.27) can be rewritten as

$$\prod_{0 \leq i \leq k} k(G/H_i \cdot \eta, X) \rightarrow \prod_{0 \leq i \leq k} k(G/H_i \cdot \eta, Y).$$

Assume first that f is a weak equivalence. Proposition 1.22 then implies that the horizontal maps in (1.27) are weak equivalences, so that the square is a pull back square, and thus f is fully faithful. That f is essentially surjective follows from the identity $k(G/H \cdot \eta, X) = k(\iota^*(Z^H))$, so that $\tau(\iota^*(X^H)) \rightarrow \tau(\iota^*(Y^H))$ is essentially surjective at the level of maximal groupoids, and this suffices for essential surjectivity. TFAE PROP

Assume now that f is fully faithful and essentially surjective. Since (1.27), Proposition 1.22 now implies that one needs only check that the maps of Kan complexes

$$k(G/H \cdot \eta, X) \rightarrow k(G/H \cdot \eta, Y) \quad \text{or} \quad k(\iota^*(X^H)) \rightarrow k(\iota^*(Y^H)) \quad (1.28) \quad \boxed{\text{KANMAP EQ}}$$

are weak equivalences. As before, essential surjectivity is equivalent to the fact that the maps (1.28) induce surjections on connected components. Hence, it now suffices to show that for each 0-simplex $x \in X^H$ the top map of loop spaces in

$$\begin{array}{ccc} \Omega(k(\iota^*X^H), x) & \longrightarrow & \Omega(k(\iota^*Y^H), f(x)) \\ \downarrow & & \downarrow \\ X(x; x) & \longrightarrow & Y(f(x); f(x)) \end{array} \quad (1.29) \quad \boxed{\text{OMEGASQ EQ}}$$

is a weak equivalence. Noting that the bottom map in (1.29) is a weak equivalence since F is fully faithful and that the vertical maps are the inclusion of the connected components corresponding to automorphisms of x in $\tau(\iota^*X^H)$. It thus suffices to check that the top map in (1.29) is an isomorphism on π_0 , and this follows since the map of categories $\tau(\iota^*(X^H)) \rightarrow \tau(\iota^*(Y^H))$ is fully faithful. OMEGASQ EQ

HERE

2 Equivariant simplicial dendroidal sets

The results in [3, §4] concerning the simplicial Reedy model structure all generalized mutatis mutandis. CM13a

Proposition 2.1. *Suppose \mathcal{C} admits two model structures (C, W_1, F_1) and (C, W_2, F_2) with the same class of cofibrations, and assume further that both model structures are cofibrantly generated and admit left Bousfield localizations with respect to any set of maps.*

Then there is a smallest common left Bousfield localization (C, W, F) and for any (C, W, F) -local c, d objects one has that $c \rightarrow d$ is in W iff it is in W_1 iff it is in W_2 .

Proof. The model structure (C, W, F) can be obtained by either localizing (C, W_1, F_1) with regards to the generating trivial cofibrations of (C, W_2, F_2) or vice-versa. That the two processes yield the same model structure follows from the universal property of left Bousfield localizations [5, Prop. 3.4.18]. The claim concerning local c, d follows from the local Whitehead theorem [5, Thm. 3.3.8], stating that the local equivalences between local objects match the initial weak equivalences. \square

HERE

The prototypical example of Proposition 2.1 is given by the category \mathbf{ssSet} of bisimplicial sets together with the two possible Reedy structures (over the Kan model structure on \mathbf{sSet}).

Explicitly, writing the levels of $X \in \mathbf{ssSet}$ as $X_{n,m}$ one can either form a Reedy model structure with respect to the *horizontal index* n or with respect to the *vertical index* m . In either case, the generating cofibrations are then given by the maps

$$(\partial\Delta[n] \rightarrow \Delta[n]) \sqcup (\partial\Delta[m] \rightarrow \Delta[m]), \quad n, m \geq 0.$$

Further, in the horizontal Reedy model structure the generating trivial cofibrations are the maps

$$(\partial\Delta[n] \rightarrow \Delta[n]) \sqcup (\Lambda^j[m] \rightarrow \Delta[m]), \quad n \geq 0, m \geq j \geq 0.$$

while for the vertical Reedy model structure the generating trivial cofibrations are the maps

$$(\Lambda^i[n] \rightarrow \Delta[n]) \sqcup (\partial\Delta[m] \rightarrow \Delta[m]), \quad n \geq i \geq 0, m \geq 0.$$

We caution the reader about a possible hiccup

HERE kk

Proposition 2.2. *Suppose that $X, Y \in \mathbf{ssSet}$ are horizontal Reedy fibrant. Then:*

- (i) *for each fixed m all vertex maps $X_{\bullet, m} \rightarrow X_{\bullet, 0}$ are trivial Kan fibrations;*
- (ii) *any vertical Reedy fibrant replacement \tilde{X} of X is in fact fibrant in the joint Reedy model structure;*
- (iii) *a map $X \rightarrow Y$ is a horizontal weak equivalence iff it is a joint weak equivalence;*
- (iv) *the canonical map $X_{n, 0} \rightarrow X_{n, n}$ is a Kan equivalence.*

Proof. (i) follows since the trivial cofibrations for the horizontal Reedy structure include all the maps of the form $(\partial\Delta[n] \rightarrow \Delta[n]) \sqcup (\Delta[0] \rightarrow \Delta[m])$.

(ii) follows since by (i) \tilde{X} is then local (with the vertical Reedy model structure as the initial model structure) with respect to all maps of the form $\Delta[0] \times (\Delta[0] \rightarrow \Delta[m])$, and thus by Remark 1.21 it is fibrant in the joint Reedy model structure (add a remark about this).

(iii) follows from (ii) since the localizing maps $X \rightarrow \tilde{X}, Y \rightarrow \tilde{Y}$ are horizontal equivalences.

For (iv), note first that the diagonal functor $\Delta: \mathbf{ssSet} \rightarrow \mathbf{sSet}$ is left Quillen for either the horizontal or vertical Reedy structures (and thus also for the joint Reedy structure). But noting that all objects are cofibrant, and regarding $X_{n, 0}$ as a bisimplicial set that is vertically constant, the claim follows by noting that by (i) the map $X_{n, 0} \rightarrow X$ is a horizontal weak equivalence in \mathbf{ssSet} . \square

HERE HERE

Proposition 2.3. *There is a model structure on \mathbf{sdSet} such that*

- *cofibrations are the Reedy cofibrations*

- *weak equivalences are generated by the levelwise dendroidal equivalences and the levelwise topological equivalences.*

Proof. Since both Δ^{op} and Ω^{op} are generalized Reedy, one can attempt to build such a model structure in two ways.

The first way, regarding the category as $\mathbf{sSet}^{\Omega^{op}}$, starts with generating cofibrations

$$(\partial\Omega[T] \rightarrow \Omega[T]) \sqcup (\partial\Delta^n \rightarrow \Delta^n)$$

and generating trivial cofibrations

$$(\partial\Omega[T] \rightarrow \Omega[T]) \sqcup (\Lambda_i^n \rightarrow \Delta^n)$$

and then localizes at the maps

$$\Lambda^e\Omega[T] \rightarrow \Omega[T].$$

The second way, regarding the category as $\mathbf{dSet}^{\Delta^{op}}$, starts with the same generating cofibrations and generating trivial cofibrations

$$J_{\mathbf{dSet}} \sqcup (\partial\Delta^n \rightarrow \Delta^n),$$

where $J_{\mathbf{dSet}}$ is a set of generating trivial cofibrations for \mathbf{dSet} , and then localizes at

$$(\partial\Omega[T] \rightarrow \Omega[T]) \sqcup (\Lambda_i^n \rightarrow \Delta^n).$$

It suffices to check that the two procedures produce the same model structure. Since the cofibrations coincide, and clearly the latter procedure has at least as many weak equivalences as the former, it suffices to check that local objects in the former are in fact also local in the latter. For this the non trivial claim (using the characterization of fibrations between fibrant objects in \mathbf{dSet}) is that the former model structure also localized the maps $(\ast \rightarrow J) \sqcup (\partial\Delta^n \rightarrow \Delta^n)$. But the original result easily implies that we have in fact also localized the maps $(\ast \rightarrow J) \times \Delta^n$ and a 2-out-of-3 argument then finishes the proof. \square

HERE HERE

A Equivariant Reedy model structures

In [BM08] Berger and Moerdijk extend the notion of Reedy category so as to allow for categories \mathbb{R} with non-trivial automorphism groups $\text{Aut}(r)$ for $r \in \mathbb{R}$. For such \mathbb{R} and suitable model category \mathcal{C} they then show that there is a *Reedy model structure* on $\mathcal{C}^{\mathbb{R}}$ that is defined by modifying the usual characterizations of Reedy cofibrations, weak equivalences and fibrations (see [1, Thm. 1.6] or Theorem A.8 below) to be determined by the $\text{Aut}(r)$ -projective model structures on $\mathcal{C}^{\text{Aut}(r)}$ for each $r \in \mathbb{R}$. REEDYADM THM

The purpose of this appendix is to show that, under suitable conditions, this can also be done by replacing the $\text{Aut}(r)$ -projective model structures on $\mathcal{C}^{\text{Aut}(r)}$ with the more general $\mathcal{C}_{\mathcal{F}_r}^{\text{Aut}(r)}$ model structures for $\{\mathcal{F}_r\}_{r \in \mathbb{R}}$ a nice collection of families of subgroups of each $\text{Aut}(r)$.

To do so, we first need some essential notation. For each map $r \rightarrow r'$ in a category \mathbb{R} we will write $\text{Aut}(r \rightarrow r')$ for its automorphism group in the arrow category and write

$$\text{Aut}(r) \xleftarrow{\pi_r} \text{Aut}(r \rightarrow r') \xrightarrow{\pi_{r'}} \text{Aut}(r') \quad (\text{A.1}) \quad \boxed{\text{PIDEFR EQ}}$$

for the obvious projections. We now introduce our equivariant generalization of the “generalized Reedy categories” of [1, Def. 1.1]. BM08

GENRED DEF

Definition A.2. A *generalized Reedy category structure* on a small category \mathbb{R} consists of wide subcategories \mathbb{R}^+ , \mathbb{R}^- and a degree function $|\cdot|: \text{ob}(\mathbb{R}) \rightarrow \mathbb{N}$ such that:

- (i) non-invertible maps in \mathbb{R}^+ (resp. \mathbb{R}^-) raise (lower) degree; isomorphisms preserve degree;
- (ii) $\mathbb{R}^+ \cap \mathbb{R}^- = \text{Iso}(\mathbb{R})$;
- (iii) every map f in \mathbb{R} factors as $f = f^+ \circ f^-$ with $f^+ \in \mathbb{R}^+$, $f^- \in \mathbb{R}^-$, and this factorization is unique up to isomorphism.

Let $\{\mathcal{F}_r\}_{r \in \mathbb{R}}$ be a collection of families of subgroups of the groups $\text{Aut}(r)$. The collection $\{\mathcal{F}_r\}$ is called *Reedy-admissible* if:

- (iv) for all maps $r \twoheadrightarrow r'$ in \mathbb{R}^- one has $\pi_{r'}(\pi_r^{-1}(H)) \in \mathcal{F}_{r'}$ for all $H \in \mathcal{F}_r$.

We note that condition (iv) above should be thought as of a constraint on the pair $(\mathbb{R}, \{\mathcal{F}_r\})$. The original setup of [1] then deals with the case where $\{\mathcal{F}_r\} = \{\{e\}\}$ is the collection of trivial families. Indeed, our setup recovers the setup in [1], as follows. BM08

Example A.3. When $\{\mathcal{F}_r\} = \{\{e\}\}$, Reedy-admissibility coincides with axiom (iv) in [1, Def. 1.1], stating that if $\theta \circ f^- = f^-$ for some $f^- \in \mathbb{R}^-$ and $\theta \in \text{Iso}(\mathbb{R})$ then θ is an identity.

Example A.4. For any generalized Reedy category \mathbb{R} , the collection $\{\mathcal{F}_{\text{all}}\}$ of the families of all subgroups of $\text{Aut}(r)$ is Reedy-admissible.

Example A.5. Let G be a group and set $\mathbb{R} = G \times (0 \rightarrow 1)$ with $\mathbb{R} = \mathbb{R}^+$. Then any pair $\{\mathcal{F}_0, \mathcal{F}_1\}$ of families of subgroups of G is Reddy-admissible.

Similarly, set $\mathbb{S} = G \times (0 \leftarrow 1)$ with $\mathbb{S} = \mathbb{S}^-$. Then a pair $\{\mathcal{F}_0, \mathcal{F}_1\}$ of families of subgroups of G is Reddy-admissible iff $\mathcal{F}_0 \supset \mathcal{F}_1$.

Example A.6. Letting \mathbb{S} denote any generalized Reedy category in the sense of [1, Def. 1.1] and G a group, we set $\mathbb{R} = G \times \mathbb{S}$ with $\mathbb{R}^+ = G \times \mathbb{S}^+$ and $\mathbb{R}^- = G \times \mathbb{S}^-$. Further, for each $s \in \mathbb{S}$ we write \mathcal{F}_s^Γ for the family of G -graph subgroups of $G \times \text{Aut}_{\mathbb{S}}(s)$, i.e., those subgroups $K \leq G \times \text{Aut}_{\mathbb{S}}(s)$ such that $K \cap \text{Aut}_{\mathbb{S}}(s) = \{e\}$. BM08

Reedy admissibility of $\{\mathcal{F}_s^\Gamma\}$ follows since for every degeneracy map $s \twoheadrightarrow s'$ in \mathbb{S}^- one has that the homomorphism $\pi_s: \text{Aut}_{\mathbb{S}}(s \twoheadrightarrow s') \rightarrow \text{Aut}_{\mathbb{S}}(s)$ is injective (we note that this is equivalent to axiom (iv) in [1, Def. 1.1] for \mathbb{S}). BM08

Our primary example of interest will come by setting $\mathbb{S} = \Omega^{op}$ in the previous example. In fact, in this case we will also be interested in certain subfamilies $\{\mathcal{F}_T\}_{T \in \Omega} \subset \{\mathcal{F}_T^\Gamma\}_{T \in \Omega}$.

Example A.7. Let $\mathbb{R} = G \times \Omega^{op}$ and let $\{\mathcal{F}_T\}_{T \in \Omega}$ be the family of graph subgroups determined by a weak indexing system \mathcal{F} . Then $\{\mathcal{F}_T\}$ is Reedy-admissible. To see this, recall first that each $K \in \mathcal{F}_T$ encodes an H -action on $T \in \Omega$ for some $H \leq G$ so that $G \cdot_H T$ is a \mathcal{F} -tree. Given a face map $f: T' \hookrightarrow T$, the subgroup $\pi_{T'}^{-1}(K)$ is then determined by the largest subgroup $\bar{H} \leq H$ such that T' inherits the \bar{H} -action from T along f (so that f becomes a \bar{H} -map), so that $\pi_{T'}(\pi_{T'}^{-1}(K))$ encodes the \bar{H} -action on T' . Thus, we see that Reedy-admissibility is simply the sieve condition for the induced map of G -trees $G \cdot_{\bar{H}} T' \rightarrow G \cdot_H T$.

We now state the main result. We will assume throughout that \mathcal{C} is a model category such that for any group G and family of subgroups \mathcal{F} , the category \mathcal{C}^G admits the \mathcal{F} -model structure (for example, this is the case whenever \mathcal{C} is a cofibrantly generated cellular model category in the sense of [7]).

REEDYADM THM

Theorem A.8. Let \mathbb{R} be generalized Reedy and $\{\mathcal{F}_r\}_{r \in \mathbb{R}}$ a Reedy-admissible collection of families. Then there is a $\{\mathcal{F}_r\}$ -**Reedy model structure** on $\mathcal{C}^{\mathbb{R}}$ such that a map $A \rightarrow B$ is

- a (trivial) cofibration if $A_r \coprod_{L_r A} L_r B \rightarrow B_r$ is a (trivial) \mathcal{F}_r -cofibration in $\mathcal{C}^{\text{Aut}(r)}$, $\forall r \in \mathbb{R}$;
- a weak equivalence if $A_r \rightarrow B_r$ is a \mathcal{F}_r -weak equivalence in $\mathcal{C}^{\text{Aut}(r)}$, $\forall r \in \mathbb{R}$;
- a (trivial) fibration if $A_r \rightarrow B_r \times_{M_r B} M_r A$ is a (trivial) \mathcal{F}_r -fibration in $\mathcal{C}^{\text{Aut}(r)}$, $\forall r \in \mathbb{R}$.

The proof of this result is given at the end of the section after establishing some routine generalizations of the key lemmas in [1] (indeed, the true novelty in this appendix is the Reedy-admissibility condition in part (iv) of Definition A.2).

We first recall the following, cf. [2, Props. 6.5 and 6.6] (we note that [2, Prop. 6.6] can be proven in terms of fibrations, and thus does not depend on special assumptions on \mathcal{C}).

Proposition A.9. Let $\phi: G \rightarrow \bar{G}$ be a homomorphism and $\mathcal{F}, \bar{\mathcal{F}}$ families of subgroups of G, \bar{G} . Then the leftmost (resp. rightmost) adjunction below is a Quillen adjunction

$$\bar{G} \cdot_G (-): \mathcal{C}_{\mathcal{F}}^G \rightleftarrows \mathcal{C}_{\bar{\mathcal{F}}}^{\bar{G}} : \text{res}_{\bar{G}}^{\bar{G}} \quad \text{res}_{\bar{G}}^{\bar{G}}: \mathcal{C}_{\bar{\mathcal{F}}}^{\bar{G}} \rightleftarrows \mathcal{C}_{\mathcal{F}}^G : \text{Hom}_G(\bar{G}, -)$$

provided that for $H \in \mathcal{F}$ it is $\phi(H) \in \bar{\mathcal{F}}$ (resp. for $\bar{H} \in \bar{\mathcal{F}}$ it is $\phi^{-1}(\bar{H}) \in \mathcal{F}$).

RESGEN COR

Corollary A.10. For any homomorphism $\phi: G \rightarrow \bar{G}$, the functor $\text{res}_{\bar{G}}^{\bar{G}}: \mathcal{C}_{\bar{\mathcal{F}}}^{\bar{G}} \rightarrow \mathcal{C}_{\mathcal{F}}^G$ preserves all four classes of genuine cofibrations, trivial cofibrations, fibrations and trivial fibrations.

The following formalizes an argument implicit in the proof of [1, Lemma 5.2].

Definition A.11. Consider a commutative diagram

$$\begin{array}{ccc} A & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \\ B & \longrightarrow & Y \end{array}$$

(A.12) BLA EQ

in $\mathcal{C}^{\mathbb{R}}$. A collection of maps $f_s: B_s \rightarrow X_s$ for $|s| \leq n$ that induce a lift of the restriction of (A.12) to $\mathcal{C}^{\mathbb{R}_{\leq n}}$ will be called a n -partial lift.

BLALIFT LEM

Lemma A.13. Let \mathcal{C} be any bicomplete category, and consider a commutative diagram as in (A.12). Then any $(n-1)$ -partial lift uniquely induces commutative diagrams

$$\begin{array}{ccc} A_r \sqcup_{L_r A} L_r B & \longrightarrow & X_r \\ \downarrow & \nearrow & \downarrow \\ B_r & \longrightarrow & Y_r \times_{M_r Y} M_r X \end{array}$$

(A.14) BLALIFT EQ

in $\mathcal{C}^{\text{Aut}(r)}$ for each r such that $|r| = n$. Furthermore, extensions of the $(n-1)$ -partial lift to a n -partial lift are in bijection with choices of $\text{Aut}(r)$ -equivariant lifts of the diagrams (A.14) for r ranging over representatives of the isomorphism classes of r with $|r| = n$.

In the next result, by $\{\mathcal{F}_r\}$ -cofibration/trivial cofibration/fibration/trivial fibration we mean a map as described in Theorem A.8, regardless of whether such a model structure exists.

BLALIFT COR

Corollary A.15. *Let \mathbb{R} be generalized Reedy and $\{\mathcal{F}_r\}$ an arbitrary family of subgroups of $\text{Aut}(r)$, $r \in \mathbb{R}$. Then a map in $\mathcal{C}^{\mathbb{R}}$ is a $\{\mathcal{F}_r\}$ -cofibration (resp. trivial cofibration) iff it has the left lifting property with respect to all $\{\mathcal{F}_r\}$ -trivial fibrations (resp. fibrations), and vice-versa for the right lifting property.*

GINJ LEM

Lemma A.16. *Let \mathbb{S} be a generalized Reedy with $\mathbb{S} = \mathbb{S}^+$, K a group, and $\pi: \mathbb{S} \rightarrow K$ a functor. Then if a map $A \rightarrow B$ in $\mathcal{C}^{\mathbb{S}}$ is such that for all $s \in \mathbb{S}$ the maps $A_s \sqcup_{L_s A} L_s B \rightarrow B_s$ are (resp. trivial) $\text{Aut}(s)$ -cofibrations one has that $\text{Lan}_{\pi: \mathbb{S} \rightarrow K}(A \rightarrow B)$ is a (trivial) K -cofibration.*

Proof. By adjunction, one needs only show that for any K -fibration $X \rightarrow Y$ in \mathcal{C}^K , the map $\pi^*(X \rightarrow Y)$ has the right lifting property against all maps $A \rightarrow B$ in $\mathcal{C}^{\mathbb{S}}$ as in the statement. By Corollary A.15, it thus suffices to check that the maps

$$(\pi^* X)_s \rightarrow (\pi^* Y)_s \times_{M_s \pi^* Y} M_s \pi^* X$$

are $\text{Aut}(s)$ -fibrations. But since $M_s Z = *$ (recall $\mathbb{S} = \mathbb{S}^+$) this map is just $X \rightarrow Y$ with the $\text{Aut}(s)$ -action induced by $\pi: \text{Aut}(s) \rightarrow K$, hence Corollary A.10 finishes the proof. \square

GINJMIN LEM

Lemma A.17. *Let \mathbb{S} be a generalized Reedy with $\mathbb{S} = \mathbb{S}^-$, K a group, and $\pi: \mathbb{S} \rightarrow K$ a functor. Then if a map $X \rightarrow Y$ in $\mathcal{C}^{\mathbb{S}}$ is such that for all $s \in \mathbb{S}$ the maps $X_s \rightarrow Y_s \times_{M_s Y} M_s X$ are (resp. trivial) $\text{Aut}(s)$ -fibrations one has that $\text{Ran}_{\pi: \mathbb{S} \rightarrow K}(A \rightarrow B)$ is a (trivial) K -fibration.*

Proof. This follows dually to the previous proof. \square

Remark A.18. Lemmas A.16 and A.17 generalize key parts of the proofs of [BM08, Lemmas 5.3 and 5.5]. The duality of their proofs reflects the duality in Corollary A.10.

Remark A.19. Lemma A.16 will be applied when $K \leq \text{Aut}_{\mathbb{R}}(r)$, and $\mathbb{S} = K \rtimes \mathbb{R}^+(r)$ for \mathbb{R} a given generalized Reedy category and $r \in \mathbb{R}$. Similarly, Lemma A.17 will be applied when $\mathbb{S} = K \rtimes \mathbb{R}^-(r)$. It is straightforward to check that in the \mathbb{R}^+ (resp. \mathbb{R}^-) case maps in \mathbb{S} can be identified with squares as on the left (right)

$$\begin{array}{ccc} r' & \xrightarrow{+} & r \\ + \downarrow & & \downarrow \simeq \\ r'' & \xrightarrow{+} & r \end{array} \quad \begin{array}{ccc} r & \xrightarrow{-} & r' \\ \simeq \downarrow & & \downarrow - \\ r & \xrightarrow{-} & r'' \end{array} \quad (\text{A.20})$$

such that the maps labelled $+$ are in \mathbb{R}^+ , maps labelled $-$ are in \mathbb{R}^- , the horizontal maps are non-invertible, and the maps labeled \simeq are automorphisms in K .

In particular, there is thus a *domain* (resp. *target*) functor $d: \mathbb{S} \rightarrow \mathbb{R}$ ($t: \mathbb{S} \rightarrow \mathbb{R}$), and our interest is in maps $d^* A \rightarrow d^* B$ ($t^* A \rightarrow t^* B$) in $\mathcal{C}^{\mathbb{S}}$ induced from maps $A \rightarrow B$ in $\mathcal{C}^{\mathbb{R}}$ so that

$$\text{Lan}_{\pi} d^*(A \rightarrow B) = (L_r A \rightarrow L_r B) \quad \text{Ran}_{\pi} t^*(A \rightarrow B) = (M_r A \rightarrow M_r B)$$

We are now in a position to prove the following, which are the essence of Theorem A.8.

Lemma A.21. *Let \mathbb{R} be generalized Reedy and $\{\mathcal{F}_r\}_{r \in \mathbb{R}}$ a Reedy-admissible family.*

Suppose $A \rightarrow B$ be a $\{\mathcal{F}_r\}$ -Reedy cofibration. Then the maps $A_r \rightarrow B_r$ are all $\{\mathcal{F}_r\}$ -weak equivalences iff so are the maps $A_r \sqcup_{L_r A} L_r B \rightarrow B_r$.

Proof. It suffices to check by induction on n that the analogous claim with the restriction $|r| \leq n$ also holds. The $n = 0$ case is obvious. Otherwise, letting r range over representatives of the isomorphism classes of r with $|r| = n$, it suffices to check that for each $H \in \mathcal{F}_r$ the map $A_r \rightarrow B_r$ is a H -genuine weak equivalence iff so is $A_r \sqcup_{L_r A} L_r B \rightarrow B_r$.

One now applies Lemma A.16 with $K = H$ and $\mathbb{S} = H \rtimes \mathbb{R}^+(r)$ to the map $d^* A \rightarrow d^* B$. Note that \mathcal{F} -trivial cofibrations are always genuine trivial cofibrations, for any family, so

REEDYTRCOF LEM

that the trivial cofibrancy requirements are immediate from Corollary A.10. It thus follows that the maps labelled \sim RESGEN COR

$$\begin{array}{ccc} L_r A & \xrightarrow{\sim} & L_r B \\ \downarrow & & \downarrow \\ A_r & \xrightarrow{\sim} & L_T B \sqcup_{L_T A} A_T \longrightarrow B_r \end{array}$$

are H -genuine trivial cofibrations, finishing the proof. □

Lemma A.22. *Let \mathbb{R} be generalized Reedy and $\{\mathcal{F}_r\}_{r \in \mathbb{R}}$ a Reedy-admissible family.*

Let $X \rightarrow Y$ be a $\{\mathcal{F}_r\}$ -Reedy fibration. Then the maps $X_r \rightarrow Y_r$ are all $\{\mathcal{F}_r\}$ -weak equivalences iff so are the maps $X_r \rightarrow Y_r \times_{M_r Y} M_r X$.

Proof. One repeats the same induction argument on $|r|$. In the induction step, it suffices to verify that, for each r with $|r| = n$ and $H \in \mathcal{F}_r$, the map $X_r \rightarrow Y_r$ is a H -genuine weak equivalence iff so is $X_r \rightarrow Y_r \times_{M_r Y} M_r X$. GINJMIN LEM

One now applies Lemma A.17 with $K = H$ and $\mathbb{S} = H \rtimes \mathbb{R}^-(r)$ to the map $t^* A \rightarrow t^* B$. Note that for each $(r \twoheadrightarrow r') \in \mathbb{S}$ one has $\text{Aut}_{\mathbb{S}}(r \twoheadrightarrow r') = \pi_r^{-1}(H)$ (where π_r is as in (A.1)), so that the trivial fibrancy requirement in Lemma A.17 follows from $\{\mathcal{F}_r\}$ being Reedy-admissible. It follows that the maps labelled \sim PIDEFF EQ

$$\begin{array}{ccc} X_r & \longrightarrow & Y_r \times_{M_r Y} M_r X \xrightarrow{\sim} Y_r \\ & & \downarrow \\ & & M_r X \xrightarrow{\sim} M_r Y \end{array}$$

are H -genuine trivial fibrations, finishing the proof. □

Remark A.23. The proofs of Lemmas A.21 and A.22 are similar, but not dual, since Lemma A.22 uses Reedy-admissibility while Lemma A.21 does not. This reflects the difference in the proofs of [1, Lemmas 5.3 and 5.5] as discussed in [1, Remark 5.6], albeit with a caveat. REEDYTRCOF LEM

Setting $K = \{e\}$ in Lemma A.16 yields that $\lim_{\mathbb{S}}(A \rightarrow B)$ is a cofibration provided that $A \rightarrow B$ is a genuine Reedy cofibration, i.e. a Reedy cofibration for $\{\mathcal{F}_{\text{all}}\}$ the families of all subgroups. On the other hand, the proof of [1, Lemma 5.3] argues that $\lim_{\mathbb{S}}(A \rightarrow B)$ is a cofibration provided that $A \rightarrow B$ is a projective Reedy cofibration, i.e. a Reedy cofibration for $\{\{e\}\}$ the trivial families (note that all projective cofibrations are genuine cofibrations, so that our claim is more general). Since the cofibration half of the projective analogue of Corollary A.10 only holds if ϕ is a monomorphism, the argument in the proof of [1, Lemma 5.3] also includes an injectivity check that is not needed for our proof of Lemma A.21. REEDYTRCOF LEM

proof of Theorem A.8. Lemmas A.21 and A.22 say that the characterizations of trivial cofibrations (resp. trivial fibrations) in the statement of Theorem A.8 are correct, i.e. that they describe the maps that are both cofibrations (resp. fibrations) and weak equivalences. REEDYADM THM

We refer to the model category axioms in [6, Def. 1.1.3]. Both 2-out-of-3 and the retract axioms are immediate (recall that retracts commute with limits/colimits). The lifting axiom follows from Corollary A.15 while the task of building factorizations $X \rightarrow A \rightarrow Y$ of a given map $X \rightarrow Y$ follows by a similar standard argument by iteratively factorizing the maps BLALIFT COR

$$X_r \sqcup_{L_r X} L_r A \rightarrow Y_r \times_{M_r Y} M_r A$$

in $\mathcal{C}^{\text{Aut}(r)}$, thus building both A and the factorization inductively (see, e.g., the proof of [1, Thm. 1.6]). □

References

- [1] C. Berger and I. Moerdijk. On an extension of the notion of Reedy category. *Math. Z.*, 269(3-4):977–1004, 2011. BM08

- | | |
|--|--|
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">BP17</div> | [2] P. Bonventre and L. A. Pereira. Genuine equivariant operads. arXiv preprint: 1707.02226, 2017. |
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">CM13a</div> | [3] D.-C. Cisinski and I. Moerdijk. Dendroidal Segal spaces and ∞ -operads. <i>J. Topol.</i> , 6(3):675–704, 2013. |
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">HHM16</div> | [4] G. Heuts, V. Hinich, and I. Moerdijk. On the equivalence between Lurie’s model and the dendroidal model for infinity-operads. <i>Adv. Math.</i> , 302:869–1043, 2016. |
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">Hi03</div> | [5] P. S. Hirschhorn. <i>Model categories and their localizations</i> , volume 99 of <i>Mathematical Surveys and Monographs</i> . American Mathematical Society, Providence, RI, 2003. |
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">Hov98</div> | [6] M. Hovey. Monoidal model categories. arXiv preprint: 9803002, 1998. |
| <div style="border: 1px solid black; padding: 2px; display: inline-block;">Ste16</div> | [7] M. Stephan. On equivariant homotopy theory for model categories. <i>Homology Homotopy Appl.</i> , 18(2):183–208, 2016. |