

Genuine equivariant operads

Peter Bonventre, Luís A. Pereira

June 21, 2017

Abstract

We build new algebraic structures, which we call genuine equivariant operads, which can be thought of as a hybrid between equivariant operads and coefficient systems. We then prove an Elmendorf-Piacenza type theorem stating that equivariant operads, with their graph model structure, are equivalent to genuine equivariant operads with their projective model structure.

As an application, we build explicit models for the N_∞ -operads of Blumberg and Hill.

Contents

1	Introduction	2
1.1	Main results	6
1.2	Outline	8
1.3	Future Work	9
2	Preliminaries	10
2.1	Grothendieck fibrations	10
2.2	Wreath product over finite sets	11
2.3	Monads and adjunctions	14
3	Planar and tall maps	15
3.1	Planar structures	16
3.2	Outer faces, tall maps, and substitution	21
3.3	Equivariant leaf-root and vertex functors	26
3.4	Planar strings	30
4	Genuine equivariant operads	34
4.1	A monad on spans	34
4.2	The genuine equivariant operad monad	38
4.3	Comparison with (regular) equivariant operads	41
4.4	Indexing systems and partial genuine operads	44
5	Free extensions and the existence of model structures	46
5.1	Labeled planar strings	47
5.2	The category of extension trees	52
5.3	Filtrations of free extensions	56
5.4	Proof of Theorems I and II	59

6	Cofibrancy and Quillen equivalences	61
6.1	Families of subgroups	61
6.2	Pushout powers	63
6.3	G -graph families and G -trees	67
6.4	Cofibrancy and the proof of Theorem MAINQUILLENEQUIV THM	71
6.5	N_∞ -operads	74
A	Transferring Kan extensions	74

1 Introduction

A surprising feature of topological algebra is that the category of (connected) topological commutative monoids is quite small, consisting only of products of Eilenberg-MacLane spaces (e.g. [8, 4K.6]). Instead, the more interesting structures are those monoids which are commutative and associative only up to homotopy, and moreover up to “all higher homotopies”. To capture these more subtle algebraic notions, Boardman-Vogt [BV73] and May [May72] developed the theory of *operads*. Informally, an operad \mathcal{O} consists of sets/spaces $\mathcal{O}(n)$ of “ n -ary operations” carrying a Σ_n -action recording “reordering the inputs of the operations”, and a suitable notion of “composition of operations”. The purpose of the theory is then the study of “objects X with operations indexed by \mathcal{O} ”, referred to as *algebras*, with the notions of monoid, commutative monoid, Lie algebra, algebra with a module, and more, all being recovered as algebras over some fixed operad in an appropriate category. Of special importance are the E_∞ -operads, introduced by May in [May72], which are “homotopical replacements” for the commutative operad and encode the aforementioned “commutative monoids up to homotopy”. In particular, while an E_∞ -algebra structure on X does not specify unique maps $X^n \rightarrow X$, it nonetheless specifies such maps “uniquely up to homotopy”.

E_∞ -operads are characterized by the homotopy type of their levels $\mathcal{O}(n)$: \mathcal{O} is E_∞ iff each $\mathcal{O}(n)$ is Σ_n -free and contractible. That is, for each subgroup $\Gamma \leq \Sigma_n$ one has

$$\mathcal{O}(n)^\Gamma \sim \begin{cases} * & \Gamma = \{*\}, \\ \emptyset & \Gamma \neq \{*\}. \end{cases}$$

Notably, when studying the homotopy theory of operads in topological spaces the preferred notion of weak equivalence is usually that of “naive equivalence”, with a map of operads $\mathcal{O} \rightarrow \mathcal{O}'$ deemed a weak equivalence if each of the maps $\mathcal{O}(n) \rightarrow \mathcal{O}'(n)$ is a weak equivalence of spaces upon forgetting the Σ_n -actions. In this context, E_∞ -operads are then equivalent to the commutative operad **Comm** and, moreover, any cofibrant replacement of **Comm** is E_∞ . However, naive equivalences differ from the equivalences in “genuine equivariant homotopy theory”, where a map of G -spaces $X \rightarrow Y$ is deemed a G -equivalence only if the induced fix point maps $X^H \rightarrow Y^H$ are weak equivalences for all $H \leq G$. This contrast hints at a number of novel subtleties that appear in the study of equivariant operads, which we now discuss.

Firstly, noting that for a G -operad \mathcal{O} (i.e. an operad \mathcal{O} together with a G -action commuting with all the structure) the n -th level $\mathcal{O}(n)$ has a $G \times \Sigma_n$ -action, one might guess that a map of G -operads $\mathcal{O} \rightarrow \mathcal{O}'$ should be called a weak equivalence if each of the maps $\mathcal{O}(n) \rightarrow \mathcal{O}'(n)$ is a G -equivalence after forgetting the Σ_n -actions, i.e. if the maps

$$\mathcal{O}(n)^H \xrightarrow{\sim} \mathcal{O}'(n)^H, \quad H \leq G \leq G \times \Sigma_n, \tag{1.1} \quad \boxed{\text{NAIVEOPEQ EQ}}$$

are weak equivalences of spaces. However, the notion of equivalence suggested in (1.1) turns out to not be “genuine enough”. To see why, we first consider a homotopical replacement for **Comm** using this theory: if one simply equips an E_∞ -operad \mathcal{O} with a trivial G -action, the resulting G -operad has fixed points for each subgroup $\Gamma \leq G \times \Sigma_n$ determined by

$$\mathcal{O}(n)^\Gamma \sim \begin{cases} * & \text{if } \Gamma \leq G, \\ \emptyset & \text{otherwise.} \end{cases} \tag{1.2} \quad \boxed{\text{NAIVEGEINFTY EQ}}$$

However, as first noted by Costenoble-Waner in [CW91], in their study of equivariant infinite loop spaces, the G -trivial E_∞ -operads of (1.2) do not provide the correct replacement of \mathbf{Comm} in the G -equivariant context. Rather, that replacement is provided instead by the G - E_∞ -operads, characterized by the fixed point conditions

$$\mathcal{O}(n)^\Gamma \sim \begin{cases} * & \text{if } \Gamma \cap \Sigma_n = \{*\}, \\ \emptyset & \text{otherwise.} \end{cases} \quad (1.3) \quad \text{GENGEINFTY EQ}$$

In contrasting (1.2) and (1.3), we note that the subgroups $\Gamma \leq G \times \Sigma_n$ such that $\Gamma \cap \Sigma_n = \{*\}$ are readily shown to be precisely the graphs of partial homomorphisms $G \geq H \rightarrow \Sigma_n$, and that $\Gamma \leq G$ iff Γ is the graph of such a trivial homomorphism. However, the notion of weak equivalence described in (1.1) fails to distinguish (1.2) and (1.3), and indeed it is possible to build maps $\mathcal{O} \rightarrow \mathcal{O}'$ where \mathcal{O} is a G -trivial E_∞ -operad (as in (1.2)) and \mathcal{O}' is a G - E_∞ -operad (as in (1.3)). Therefore, in order to distinguish such operads, one needs to replace the notion of weak equivalence in (1.1) with the notion of *graph equivalence*, so that $\mathcal{O} \rightarrow \mathcal{O}'$ is considered a weak equivalence only if

$$\mathcal{O}(n)^\Gamma \xrightarrow{\sim} \mathcal{O}'(n)^\Gamma, \quad \Gamma \leq G \times \Sigma_n, \Gamma \cap \Sigma_n = \{*\}. \quad (1.4) \quad \text{GENEOPEQ EQ}$$

are weak equivalences.

As mentioned above, the original evidence [CW91] that (1.3), rather than (1.2), provides the best up to homotopy replacement for \mathbf{Comm} in the equivariant context comes from the study of equivariant infinite loop spaces. For our purposes, however, we instead focus on the perspective of Blumberg-Hill in [BH15], which concerns the Hill-Hopkins-Ravenel norm maps featured in the solution of the Kervaire invariant problem [HHR].

Given a G -spectrum R and finite G -set X with n elements, the corresponding *norm* is a G -spectrum $N^X R$ whose underlying spectrum is $R^{\wedge X} \simeq R^{\wedge n}$ but equipped with a mixed G -action that combines the actions on R and X in the natural way. Moreover, for any \mathbf{Comm} -algebra R , i.e. strictly commutative G -ring spectrum, ring multiplication further induces so called *norm maps*

$$N^X R \rightarrow R. \quad (1.5) \quad \text{NORMMAPS EQ}$$

Furthermore, by reducing structure on R the maps (1.5) are also defined when X is only a H -set for some subgroup $H \leq G$, and the maps (1.5) then satisfy a number of natural equivariance and associativity conditions. Crucially, we note that the more interesting of these associativity conditions involve H -sets for varying H (for an example packaged in operadic language, see (1.10) below).

The key observation at the source of the work in [3] is then that, operadically, norm maps are encoded by the graph fixed points as in (1.4). More explicitly, noting that a H -set X with n elements is encoded by a partial homomorphism $G \geq H \rightarrow \Sigma_n$, one obtains an associated graph subgroup $\Gamma_X \leq G \times \Sigma_n$, $\Gamma_X \cap \Sigma_n = \{*\}$, well defined up to conjugation. It then follows that for R an \mathcal{O} -algebra, maps of the form (1.5) are parametrized by the fixed point space $\mathcal{O}(n)^{\Gamma_X}$. The flaw with the G -trivial E_∞ -operad described in (1.2) is then that it lacks all norm maps other than those for H -trivial X , thus lacking some of the data encoded by \mathbf{Comm} . Further, from this perspective one may regard the more naive notion of weak equivalence in (1.1), according to which (1.2) and (1.3) are equivalent, as studying “operads without norm maps” (in the sense that equivalences ignore norm maps), while the equivalences (1.4) study “operads with norm maps”.

Our first main result, Theorem 1, establishes the existence of a model structure on operads with weak equivalences the graph equivalences of (1.4). Though our analysis goes significantly further, again guided by Blumberg-Hill’s work in [BH15].

The main novelty of [3] is the definition, for each finite group G , of a finite lattice of new types of equivariant operads, which they dub N_∞ -operads. The minimum type of N_∞ -operads is that of the G -trivial E_∞ -operads in (1.2) while the maximal type is that of the G - E_∞ -operads in (1.3). The remaining types, which interpolate between G -trivial E_∞ and

$G\text{-}E_\infty$, can hence be thought of as encoding varying degrees of “up to homotopy equivariant commutativity”. More concretely, each type of N_∞ -operad is determined by a collection $\mathcal{F} = \{\mathcal{F}_n\}_{n \geq 0}$ where each \mathcal{F}_n is itself a collection of graph subgroups of $G \times \Sigma_n$, with an operad \mathcal{O} being called a $N\mathcal{F}$ -operad if it satisfies the fixed point condition

$$\mathcal{O}(n)^\Gamma \sim \begin{cases} * & \text{if } \Gamma \in \mathcal{F}_n, \\ \emptyset & \text{otherwise.} \end{cases} \quad (1.6)$$

NFINFTY EQ

Such collections \mathcal{F} are, however, far from arbitrary, with much of the work in [BH15, §3] spent cataloging a number of closure conditions that such \mathcal{F} must satisfy. The simplest of these conditions state that each \mathcal{F}_n is a *family*, i.e. closed under subgroups and conjugation. These first conditions, which are common in equivariant homotopy theory, are a simple consequence of each $\mathcal{O}(n)$ being a space. However, the remaining conditions, all of which involve \mathcal{F}_n for various n simultaneously and are a consequence of operadic multiplication, are both novel and subtle. In loose terms, these conditions, which are more easily described in terms of the H -sets X associated to graph subgroups, concern closure of those under disjoint union, cartesian product, subobjects, and an entirely novel key condition called *self-induction*. The precise conditions are collected in [3, Def. 3.22], which also introduces the term *indexing system* for a \mathcal{F} satisfying all such conditions. The main result of [3, §4] is then that whenever a $N\mathcal{F}$ -operad \mathcal{O} as in (1.6) exists, the associated collection \mathcal{F} must be an indexing system. However, the converse statement, that given any indexing system \mathcal{F} such an \mathcal{O} can be produced, was left as a conjecture.

One of the key motivating goals of the present work was to verify this conjecture of Blumberg-Hill, which we obtain in Corollary IV, and, moreover, to produce models of $N\mathcal{F}$ -operads that are as explicit as possible.

To motivate our approach, we first recall the solution of a closely related but simpler problem: that of building universal spaces for families of subgroups. Given a family \mathcal{F} of subgroups of G (i.e. a collection closed under conjugation and subgroups), a *universal space* X for \mathcal{F} , also called a $E\mathcal{F}$ -space, is a space with fixed points X^H characterized just as in (1.6). In particular, whenever \mathcal{O} is a $N\mathcal{F}$ -operad, each $\mathcal{O}(n)$ is necessarily a $E\mathcal{F}_n$ -space. The existence of $E\mathcal{F}$ -spaces for any choice of the family \mathcal{F} is best understood in light of Piacenza’s classical result from [16] (with the key insight dating back to Elmendorf in [7]) stating that there is a Quillen equivalence (where \mathbf{O}_G is the *orbit* category, formed by the G -sets G/H)

$$\begin{array}{ccc} \mathbf{Top}^{\mathbf{O}_G^{op}} & \xrightleftharpoons[\iota_*]{\iota^*} & \mathbf{Top}^G \\ (G/H \mapsto Y(G/H)) & \longmapsto & Y(G) \\ (G/H \mapsto X^H) & \longleftarrow & X \end{array} \quad (1.7)$$

COFADJINT EQ

where the weak equivalences (and fibrations) on \mathbf{Top}^G are detected on all fixed points and the weak equivalences (and fibrations) on the category $\mathbf{Top}^{\mathbf{O}_G^{op}}$ of *coefficient systems* are detected at each presheaf level. Noting that the fixed point characterization of $E\mathcal{F}$ -spaces define an obvious object $\delta_{\mathcal{F}} \in \mathbf{Top}^{\mathbf{O}_G^{op}}$ by $\delta_{\mathcal{F}}(G/H) = *$ if $H \in \mathcal{F}$ and $\delta_{\mathcal{F}}(G/H) = \emptyset$ otherwise, $E\mathcal{F}$ -spaces can then be built as $\iota^*(C\delta_{\mathcal{F}}) = C\delta_{\mathcal{F}}(G)$, where C denotes cofibrant replacement in $\mathbf{Top}^{\mathbf{O}_G^{op}}$. Moreover, we note that, as in [7], these cofibrant replacements can be built via explicit simplicial realizations.

The overarching goal of this paper is then that of proving the analogue of Elmendorf-Piacenza’s Theorem (1.7) in the context of operads with norm maps (i.e. with equivalences as in (1.4)), which we state as our main result, Theorem III. However, in trying to formulate such a result one immediately runs into a fundamental issue: it is unclear which category should take the role of the coefficient systems $\mathbf{Top}^{\mathbf{O}_G^{op}}$ in that context. This last remark likely requires justification. Indeed, it may at first seem tempting to simply employ one of the known formal generalizations of Elmendorf-Piacenza’s result (see, e.g. [20, Thm.

3.17]) which simply replace \mathbf{Top} on either side of (1.7) with a more general model category \mathcal{V} . However, if one applies such a result when $\mathcal{V} = \mathbf{Op}$ to establish a Quillen equivalence $\mathbf{Op}^{\mathcal{O}^{op}} \rightleftarrows \mathbf{Op}^G$, the fact that the levels of each $\mathcal{P} \in \mathbf{Op}^{\mathcal{O}^{op}}$ correspond only to those fixed-point spaces appearing in (1.1) would require working in the context of operads *without* norm maps, and thereby forgo the ability to distinguish the many types of $N\mathcal{F}$ -operads.


In order to work in the context of operads with norm maps we will need to replace $\mathbf{Top}^{\mathcal{O}^{op}}$ with a category \mathbf{Op}_G of new algebraic objects we dub *genuine equivariant operads* (as opposed to (regular) equivariant operads \mathbf{Op}^G). Each genuine equivariant operad $\mathcal{P} \in \mathbf{Op}_G$ will consist of a list of spaces indexed in the same way as in (1.4) along with obvious restriction maps and, more importantly, suitable *composition maps*. Precisely identifying the required composition maps is one of the main challenges of this theory, and again we turn to [3] for motivation.

When analyzing the proofs of the results in [3, §4] concerning the closure properties for indexing systems \mathcal{F} a common motif emerges: when performing an operadic composition

$$\begin{aligned} \mathcal{O}(n) \times \mathcal{O}(m_1) \times \cdots \times \mathcal{O}(m_n) &\longrightarrow \mathcal{O}(m_1 + \cdots + m_n) \\ (f, g_1, \cdots, g_n) &\longmapsto f(g_1, \cdots, g_n) \end{aligned} \quad (1.8)$$

careful choices of fixed point conditions on the operations f, g_1, \cdots, g_n yield a fixed point condition on the composite operation $f(g_1, \cdots, g_n)$. The desired multiplication maps for a genuine equivariant operad $\mathcal{P} \in \mathbf{Op}_G$ will then abstract such interactions between multiplication and fixed points for a equivariant operad $\mathcal{O} \in \mathbf{Op}^G$. However, these interactions can be challenging to write down explicitly and, indeed, the arguments in [3, §4] do not quite provide the sort of unified conceptual approach to these needed for our purposes. The cornerstone of the current work was then the joint discovery by the authors of such a conceptual framework: equivariant trees.

Non-equivariantly, it has long been known that the combinatorics of operadic composition is best visualized by means of tree diagrams. For instance, the tree




encodes the operadic composition

$$\mathcal{O}(3) \times \mathcal{O}(2) \times \mathcal{O}(3) \times \mathcal{O}(0) \rightarrow \mathcal{O}(5)$$

where the inputs $\mathcal{O}(3), \mathcal{O}(2), \mathcal{O}(3), \mathcal{O}(0)$ correspond to the nodes (i.e. circles) in the tree, with arity given by number of incoming edges (i.e. edges immediately above) and the output $\mathcal{O}(5)$ has arity given by counting leaves (i.e. edges at the top, not capped by a node). Similarly, the role of equivariant trees is, in the context of equivariant operads, to encode such operadic compositions together with fixed point compatibilities. A detailed introduction to equivariant trees can be found in [15, §4], where the second author develops the theory of equivariant dendroidal sets (which is a parallel approach to equivariant operads), though here we include a single representative example. Let $G = \{\pm 1, \pm i, \pm j, \pm k\}$ denote the group of quaternionic units and $G \geq H \geq K \geq L$ denote the subgroups $H = \langle j \rangle$, $K = \langle -1 \rangle$, $L = \{1\}$. There is then a G -tree T with *expanded representation* given by the two trees on the left

below and *orbital representation* given by the (single) tree on the right.



D6SMALLER EQ

We note that G acts on the expanded representation of T as indicated by the edge labels (so that the edges a, b, c, d have stabilizers L, K, K, H respectively), and the orbital representation is obtained by collapsing the edge orbits of the expanded representation. As explained in [Fe17, Example 4.9], T then encodes the fact that for any equivariant operad $\mathcal{O} \in \mathbf{Op}^G$ the composition $\mathcal{O}(2) \times \mathcal{O}(3)^{\times 2} \rightarrow \mathcal{O}(6)$ restricts to a fixed point composition

$$\mathcal{O}(H/K)^H \times \mathcal{O}(K/L \sqcup K/K)^K \rightarrow \mathcal{O}(H/L \sqcup H/K)^H \quad (1.10)$$

INTFIXPTCOMP EQ

where $\mathcal{O}(X)$ for an H -set (resp. K -set) X denotes $\mathcal{O}(|X|)$ together with a suitably mixed H -action (K -action). We note that the inputs $\mathcal{O}(H/K)^H, \mathcal{O}(K/L \sqcup K/K)^K$ in (1.10) correspond to the nodes of the orbital representation in (1.9), though in contrast to the non-equivariant case arity is now determined by both incoming and outgoing edge *orbits*, while the output $\mathcal{O}(H/L \sqcup H/K)^H$ is similarly determined by both the leaf and root edge *orbits*. The existence of maps of the form (1.10) is essentially tantamount to the subtlest closure property for indexing systems \mathcal{F} , self-induction (cf. [3, Def. 3.20]), and similar descriptions exist for all other closure properties, as detailed by the second author in [15, §9].

We can now at last give a full informal description of the category \mathbf{Op}_G featured in our main result, Theorem III. A genuine equivariant operad $\mathcal{P} \in \mathbf{Op}_G$ has levels $\mathcal{P}(X)$ for each H -set X , $H \leq G$, that mimic the role of the fixed points $\mathcal{O}(X)^H \simeq \mathcal{O}(|X|)^{\Gamma_X}$ for $\mathcal{O} \in \mathbf{Op}^G$. More explicitly, there are restriction maps $\mathcal{P}(X) \rightarrow \mathcal{P}(X|_K)$ for $K \leq L$, isomorphisms $\mathcal{P}(X) \simeq \mathcal{P}(gX)$ where gX denotes the conjugate gHg^{-1} -set, and composition maps given by

$$\mathcal{P}(H/K) \times \mathcal{P}(K/L \sqcup K/K) \rightarrow \mathcal{P}(H/L \sqcup H/K)$$

in the case of the abstraction of (1.10), and more generally by

$$\begin{aligned} & \mathcal{P}(H/K_1 \sqcup \dots \sqcup H/K_n) \times \mathcal{P}(K_1/L_{11} \sqcup \dots \sqcup K_1/L_{1m_1}) \times \dots \times \mathcal{P}(K_n/L_{n1} \sqcup \dots \sqcup K_n/L_{nm_n}) \\ & \quad \downarrow \\ & \mathcal{P}(H/L_{11} \sqcup \dots \sqcup H/L_{1m_1} \sqcup \dots \sqcup H/L_{n1} \sqcup \dots \sqcup H/L_{nm_n}). \end{aligned} \quad (1.11)$$

GENGENMULT EQ

Lastly, these composition maps must satisfy associativity, unitality, compatibility with restriction maps, and equivariance conditions, as encoded by the theory of G -trees. Rather than making such compatibilities explicit, it will be more convenient and effective for our purposes to simply define genuine equivariant operads intrinsically in terms of G -trees.

1.1 Main results

We now discuss our main results.

Recall that $\mathbf{Op}^G(\mathcal{V}) = (\mathbf{Op}(\mathcal{V}))^G$ denotes G -objects in $\mathbf{Op}(\mathcal{V})$.

Theorem I. *Let (\mathcal{V}, \otimes) denote either (\mathbf{sSet}, \times) or $(\mathbf{sSet}_*, \wedge)$.*

Then there exists a model structure on $\mathbf{Op}^G(\mathcal{V})$ such that $\mathcal{O} \rightarrow \mathcal{O}'$ is a weak equivalence (resp. fibration) if all the maps

$$\mathcal{O}(n)^\Gamma \rightarrow \mathcal{O}'(n)^\Gamma \quad (1.12)$$

GENEOEQMT EQ

for $\Gamma \leq G \times \Sigma_n, \Gamma \cap \Sigma_n = \{*\}$, are weak equivalences (fibrations) in \mathcal{V} .

More generally, for $\mathcal{F} = \{\mathcal{F}_n\}_{n \geq 0}$ with \mathcal{F}_n an arbitrary collection of subgroups of $G \times \Sigma_n$ there exists a model structure on $\text{Op}_G^{\mathcal{F}}(\mathcal{V})$, which we denote $\text{Op}_{\mathcal{F}}^G(\mathcal{V})$, with weak equivalences (resp. fibrations) determined by (I.12) for $\Gamma \in \mathcal{F}_n$.

Lastly, analogous semi-model structures $\text{Op}_G^G(\mathcal{V})$, $\text{Op}_{\mathcal{F}}^G(\mathcal{V})$ exist provided that (\mathcal{V}, \otimes) : (i) is a strongly cofibrantly generated model category; (ii) is a closed monoidal model category; (iii) has cellular fixed points; (iv) has cofibrant symmetric pushout powers.

Theorem I is proven in §7.7, where the additional conditions are also discussed.

Our next result concerns the model structure on the category $\text{Op}_G(\mathcal{V})$ of genuine equivariant operads introduced in this paper, the construction of which we now briefly outline. Firstly, the levels of each $\mathcal{P} \in \text{Op}_G(\mathcal{V})$, i.e. the H -sets in (II.11), are encoded by a category Σ_G of G -corollas, introduced in §3.3, which generalizes the usual category Σ of finite sets and isomorphisms. We then define G -symmetric sequences by $\text{Sym}_G(\mathcal{V}) = \mathcal{V}^{\Sigma_G}$ and whenever \mathcal{V} is a closed symmetric monoidal category with diagonals (cf. Remark 2.18), we define in §4.2 a free genuine equivariant operad monad \mathbb{F}_G on $\text{Sym}_G(\mathcal{V})$ whose algebras form the desired category $\text{Op}_G(\mathcal{V})$.

Moreover, inspired by the analogues $\text{Top}_{\mathcal{F}}^{\text{op}} \rightleftarrows \text{Top}_{\mathcal{F}}^G$ of the Elmendorf-Piacenza equivalence where $\text{Top}_{\mathcal{F}}^{\text{op}}$ are partial coefficient systems determined by a family \mathcal{F} , we show in §4.4 that (a slight generalization of) Blumberg-Hill's indexing systems \mathcal{F} give rise to sieves $\Sigma_{\mathcal{F}} \hookrightarrow \Sigma_G$ and partial symmetric sequences $\text{Sym}_{\mathcal{F}}(\mathcal{F}) = \mathcal{V}^{\Sigma_{\mathcal{F}}^{\text{op}}}$ which are suitably compatible with the monad \mathbb{F}_G , thus giving rise to categories $\text{Op}_{\mathcal{F}}(\mathcal{V})$ of partial genuine equivariant operads.

Theorem II. *Let (\mathcal{V}, \otimes) denote either (sSet, \times) or (sSet_*, \wedge) . Then the projective model structure on $\text{Op}_G(\mathcal{V})$ exists. Explicitly, a map $\mathcal{P} \rightarrow \mathcal{P}'$ is a weak equivalence (resp. fibration) if all maps*

$$\mathcal{P}(C) \rightarrow \mathcal{P}'(C) \quad (1.13)$$

are weak equivalences (fibrations) in \mathcal{V} for each $C \in \Sigma_G$.

More generally, for \mathcal{F} a weak indexing system, a projective model structure on $\text{Op}_{\mathcal{F}}(\mathcal{V})$ exists. Explicitly, weak equivalences (resp. fibrations) are determined by (I.13) for $C \in \Sigma_{\mathcal{F}}$.

Lastly, analogous semi-model structures $\text{Op}_G^G(\mathcal{V})$, $\text{Op}_{\mathcal{F}}^G(\mathcal{V})$ exist provided that (\mathcal{V}, \otimes) : (i) is a strongly cofibrantly generated model category; (ii) is a closed monoidal model category; (iii) has cellular fixed points; (iv) has cofibrant symmetric pushout powers; (v) (\mathcal{V}, \otimes) has diagonals.

Theorem II is proven in §7.7. We note that the condition that (\mathcal{V}, \otimes) has diagonals (cf. Remark 2.18), which is not needed in Theorem I, is required to build $\text{Op}_G(\mathcal{V})$, $\text{Op}_{\mathcal{F}}(\mathcal{V})$.

The following is our main result.

Theorem III. *Let (\mathcal{V}, \otimes) denote either (sSet, \times) or (sSet_*, \wedge) .*

Then the adjunctions, where in the more general rightmost case \mathcal{F} is a weak indexing system,

$$\text{Op}_G(\mathcal{V}) \xrightleftharpoons[\iota_*]{\iota^*} \text{Op}_G^G(\mathcal{V}), \quad \text{Op}_{\mathcal{F}}(\mathcal{V}) \xrightleftharpoons[\iota_*]{\iota^*} \text{Op}_{\mathcal{F}}^G(\mathcal{V}). \quad (1.14)$$

are Quillen equivalences.

Moreover, analogous Quillen equivalences of semi-model structures $\text{Op}_G^G(\mathcal{V})$, $\text{Op}_{\mathcal{F}}^G(\mathcal{V})$ exist provided that (\mathcal{V}, \otimes) : (i) is a strongly cofibrantly generated model category; (ii) is a closed monoidal model category; (iii) has cellular fixed points; (iv) has cofibrant symmetric pushout powers; (v) (\mathcal{V}, \otimes) has diagonals; (vi) (\mathcal{V}, \otimes) has cartesian fixed points.

Theorem III is proven in §6.4.

Lastly, our techniques also confirm the main conjecture of [3], as we discuss in §6.5. Moreover, we note that our models for \mathcal{NF} -operads are given by explicit bar constructions.

Corollary IV. For $\mathcal{V} = \mathbf{sSet}$ or \mathbf{Top} and $\mathcal{F} = \{\mathcal{F}_n\}_{n \geq 0}$ any weak indexing system, $N\mathcal{F}$ -operads exist. That is, there exist explicit operads \mathcal{O} such that

$$\mathcal{O}(n)^\Gamma \sim \begin{cases} * & \text{if } \Gamma \in \mathcal{F}_n \\ \emptyset & \text{otherwise.} \end{cases} \quad (1.15)$$

In particular, the map $\mathrm{Ho}(N_\infty\text{-Op}) \rightarrow \mathcal{I}$ in [BH15, Cor. 5.6] is an equivalence of categories.

1.2 Outline

come back here

There are multiple distinct steps along the path to proving our main result. Specifically, we need better understanding and control of the interplay between the subtle equivariant structures and the combinatorics which underly operads. To that end, we rebuild operads in a way which allows us to exploit the equivariant generalization Ω_G of the dendroidal category Ω discussed in [Pe17].

We begin by observing that the free operad $\mathbb{F}X$ generated by a symmetric sequence $X \in \mathbf{Sym}(\mathcal{V})$, discussed in [Spitz201, BM03, Rub17] etc., can be repackaged as a left Kan extension out of Ω .

$$\begin{array}{ccc} \Omega^{op} & \xrightarrow{N_X} & \mathcal{V} \\ \downarrow \mathrm{lr} & \searrow \mathbb{F}X & \\ \Sigma^{op} & & \end{array} \qquad \begin{array}{ccc} \Omega_G^{op} & \xrightarrow{N_Y} & \mathcal{V} \\ \downarrow \mathrm{lr} & \searrow \mathbb{F}_G Y & \\ \Sigma_G^{op} & & \end{array}$$

Here, lr is the “leaf-root” or “valence” functor, and N_X sends T to $\prod_{v \in V(T)} X(v)$.

This can be equivariantly generalized, replacing Ω and Σ with Ω_G and Σ_G , as seen on the right-hand-side. This yields our “genuine G -operad” monad \mathbb{F}_G , acting on the new category of G -sequences $Y \in \mathbf{Sym}_G(\mathcal{V}) = \mathcal{V}^{\Sigma_G^{op}}$. Now, having defined our notion of “coefficient systems”, we can begin further analysis. As is often the case, we desire a clear understanding of free \mathbb{F}_G -extensions (5.1).

First, we produce multiple categorical manipulations of this functor. We modify both Ω_G and \mathbb{F}_G to construct a description of coproducts of \mathbb{F}_G -algebras as a similar looking left Kan extension. Then, we “add in relations” found in our free \mathbb{F}_G -extension by inserting new maps into this modified Ω_G , via a realization of a simplicial category of strings of such maps. Finally, using this description, we can build a filtration of such free extensions.

Secondly, we consider the homotopical characteristics of \mathbb{F}_G , in particular a detailed study of the interactions of cofibrancy with composition. We also compare \mathbb{F} and \mathbb{F}_G , categorically and homotopically, in order to display the Quillen equivalence.

We now give a brief overview of the remaining sections.

Section 2: Preliminaries. Here, we identify the major constructions we will be using throughout the paper, as well as establish notion and background material.

Section 3: Planar and tall maps. Our machinery requires our trees to record an additional structure, namely a planarization. In this section, we define this notion for G -trees, and explore the notion of “substitution”, a second operad-like structure on the class of trees, which many of our constructions employ.

Section 4: The genuine equivariant operad monad. Defining \mathbb{F}_G as an endofunctor is fairly straightforward, as is defining composition and unit maps. However, showing that this data forms a monad requires a more subtle analysis. We define a more general monad N on the category of weak spans $\mathbf{WSpan}(\Sigma_G, \mathcal{V})$, and simplify the discussion through the flexibility of this more general category. It is also here that the concept of “planar strings” of maps are introduced.

Section 5: Free extensions and the existence of model structures. In this section, we develop the technology to produce free T -extensions for a particular class of monads T , utilizing the realizations of simplicial categories of planar strings. After providing

a filtration of such constructions, we prove the existence of various model structures on our categories of algebras.

Section 6: Cofibrancy and Quillen equivalences. This section contains the bulk of the homotopical analysis. It is here that we show that weak indexing systems \mathcal{F} provide sufficient structure so that their associated \mathcal{F} -model structures are well-behaved, in particular proving the desired Quillen equivalences. We end this section, and this paper, by providing two proofs of the N_∞ realization conjecture.

Much of the machinery in Sections 4 and 5 are built in large categorical generality, designed such that the intuitive descriptions of, for example, the free operad monad and operations on forests, are verified with strict categorical rigor; as such, they may be of broader interest.

1.3 Future Work

come back here

While this paper focuses solely on *single coloured* operads, we expect the theory to extend immediately to the coloured setting. One key difference is that coloured genuine G -operads will have a *coefficient system* of colours: evaluation at a G -corolla will include the additional data of a colour for each G -orbit of edges, and the isotropy of the colours must match that of its associated edge. This feature is inspired by those seen in other models for G -operads, on which we now elaborate.

As stated earlier, the main goal of the authors' current project is to detail the full story of the homotopy theory of equivariant operads. This paper and [Pe17] form the basis of a generalization of the work of Cisinski-Moerdijk-Weiss, describing the homotopy theory of non-equivariant operads, into the G -equivariant context. Importantly, the notion of “multicoloured genuine G -operads” will provide the natural (and most highly structured) target for the strictification, or “homotopy operad” construction, on the G - ∞ -operads of [Pe17].

Moreover, sequels will extend the definitions of “dendroidal complete Segal spaces” and “Segal pre-operads”, as well as the comparison Diagram (\star) of Quillen equivalences from [CM13b], reproduced here below, to the G -equivariant context.

$$\begin{array}{ccc}
 \text{PreOp} & \xleftarrow{\quad} & \text{sOp} \\
 \downarrow & & \downarrow hcN_d \\
 \text{sdSet} & \xleftarrow{\quad} & \text{dSet}
 \end{array}
 \quad (\star) \quad \boxed{\text{CM_EQ}}$$

We note that, as in [Pe17], there are both multiple possible *categorical* generalizations (i.e. dSet^G versus dSet_G , or Op^G versus Op_G) and multiple possible *model categorical* generalizations (i.e. $\text{dSet}_{\mathcal{F}}^G$ for varying weak indexing systems \mathcal{F}) for each corner of (\star) . Analogous to the main result of this paper, we expect all compatible notions at each corner to be Quillen equivalent (i.e. $\text{dSet}_{\mathcal{F}}^G \simeq_Q \text{dSet}_G^{\mathcal{F}} \simeq_Q \text{dSet}_{\mathcal{F}}^{\mathcal{F}}$). In future papers, we will study this precise question, as well as comparisons between the different models.

We end by interpreting these different models of G -operads, focusing on the right half of (\star) . We recall that, intuitively, ∞ -operads can be thought of as operads where composition is “weakly defined”, and similarly G -coefficient systems as spaces with a “relaxed” fixed-point condition. In this fashion, genuine G -operads can be thought of as G -operads where composition is still rigidly defined, but with relaxed fixed-point conditions. Comparatively, the G - ∞ -operads of [Pe17] have rigid fixed-point conditions but weak composition. The remaining missing link is then the suitable notion of G - ∞ -operad (that is, a suitable model structure) in the true pre-sheaf category dSet_G , in which both composition and fixed-point conditions

are weak. This would yield an expanded equivariant right half of \mathbb{K}_* , given by the following.

$$\begin{array}{ccc}
 \mathrm{Op}_{\mathcal{F}}^G(\mathcal{V}) & \xrightarrow{i_*} & \mathrm{Op}_G^{\mathcal{F}}(\mathcal{V}) \\
 \downarrow \scriptstyle hcN^G & \nearrow \scriptstyle Ho_G & \downarrow \scriptstyle hcN_G \\
 \mathrm{dSet}_{\mathcal{F}}^G & \xrightarrow{i_*} & \mathrm{dSet}_G^{\mathcal{F}}
 \end{array}
 \begin{array}{c}
 \text{Ho}^G \text{ (dashed)} \\
 \text{Ho}_G \text{ (dashed)}
 \end{array}$$

We expect that all solid arrows are Quillen equivalences. However, analogous to the work of Cisinski-Moerdijk, direct vertical comparisons between G -operads and G -dendroidal sets will be challenging, inspiring further need to explore other notions of G - ∞ -operads in the previous paragraph.

We also record our finding of an unexpected complication, that G - ∞ -operads in dSet^G themselves satisfy a rigid fixed point condition, but their *genuine* homotopy operad does *not*, in particular that

$$\mathrm{Ho}_G(X) \neq \mathrm{Ho}_G i_*(X) \neq i_* \mathrm{Ho}^G(X)$$

for G - ∞ -operads X .

A full exploration of this and the other above topics will be discussed in sequels.

Mention colored operad versions, equivalence with dendroidal sets

2 Preliminaries

This section lists some elementary concepts and results that will be used throughout the paper, but may not be entirely standard.

2.1 Grothendieck fibrations

Recall that a functor $\pi: \mathcal{E} \rightarrow \mathcal{B}$ is called a *Grothendieck fibration* if for every arrow $f: b' \rightarrow b$ in \mathcal{B} and $e \in \mathcal{E}$ such that $\pi(e) = b$, there exists a cartesian arrow $f^*e \rightarrow e$ lifting f , meaning that for any choice of solid arrows

$$\begin{array}{ccc}
 e'' & \xrightarrow{\quad} & e \\
 \searrow \scriptstyle \exists! \text{ dashed} & & \nearrow \\
 & f^*e &
 \end{array}
 \qquad
 \begin{array}{ccc}
 b'' & \xrightarrow{\quad} & b \\
 \searrow & & \nearrow \scriptstyle f \\
 & b' &
 \end{array}$$

such that the rightmost diagram commutes and $e'' \rightarrow e$ lifts $b'' \rightarrow b$ there exists a unique dashed arrow $e'' \rightarrow f^*e$ lifting $b'' \rightarrow b'$ and making the leftmost diagram commute.

In most contexts the cartesian arrows $f^*e \rightarrow e$ are assumed to be defined only up to unique isomorphism, but in all examples considered in this paper we will in fact be able so identify preferred choices of cartesian arrows to which we will refer to as *pullbacks*. Moreover, pullbacks will be compatible with composition and units in the obvious way, i.e. $g^*f^*e = (fg)^*e$ and $id_b^*e = e$. As a terminological note, one sometimes encounters the term *split fibration* to refer to a Grothendieck fibration together with such a choice of pullbacks, though we will rarely have need to make the distinction outside of the present discussion.

A map of Grothendieck fibrations (resp. split fibrations) is then a commutative diagram

$$\begin{array}{ccc}
 \mathcal{E} & \xrightarrow{\delta} & \bar{\mathcal{E}} \\
 \pi \searrow & & \swarrow \bar{\pi} \\
 & \mathcal{B} &
 \end{array}
 \tag{2.1}$$

GROTHFIBMAP EQ

such that δ preserves cartesian arrows (pullbacks).

There is a well known equivalence between Grothendieck fibrations over \mathcal{B} and contravariant pseudo-functors $\mathcal{B}^{op} \rightarrow \mathbf{Cat}$ with split fibrations corresponding to (regular) contravariant functors. We now recall how this works in the (covariant) split case.

Definition 2.2. Given a diagram category \mathcal{B} and functor \mathcal{E}_\bullet .

$$\begin{aligned} \mathcal{B} &\xrightarrow{\mathcal{E}_\bullet} \mathbf{Cat} \\ b &\longmapsto \mathcal{E}_b \end{aligned} \quad (2.3)$$

the (covariant) Grothendieck construction $\mathcal{B} \ltimes \mathcal{E}_\bullet$ has objects pairs (b, e) with $b \in \mathcal{B}$, $e \in \mathcal{E}_b$ and arrows $(b, e) \rightarrow (b', e')$ given by pairs

$$(f: b \rightarrow b', g: g_*(e) \rightarrow e'),$$

where $f_*: \mathcal{E}_b \rightarrow \mathcal{E}_{b'}$ is a shorthand for the functor $\mathcal{E}_\bullet(f)$.

Note that the chosen pushforward of (b, e) along $b \rightarrow b'$ is then (b', f_*e) .

One useful property of Grothendieck constructions is that right Kan extensions can be computed using fibers, i.e., given a functor $F: \mathcal{E} \rightarrow \mathcal{V}$ into a complete category \mathcal{V} one has

$$\mathbf{Ran}_\pi F(b) \simeq \lim F|_{b \downarrow \mathcal{E}} \simeq \lim F|_{\mathcal{E}_b} \quad (2.4)$$

FIBERKAN EQ

where the first identification is the usual pointwise formula for Kan extensions (cf. [McL, X.3.1]) and the second identification follows by noting that due to the existence of cartesian arrows the fibers \mathcal{E}_b are initial (in the sense of [McL, IX.3]) in the undercategories $b \downarrow \mathcal{E}$. In fact, a little more is true: a choice of cartesian arrows yields a right adjoint to the inclusion $\mathcal{E}_b \hookrightarrow b \downarrow \mathcal{E}$, so that \mathcal{E}_b is a coreflexive subcategory of $b \downarrow \mathcal{E}$, a well known sufficient condition for initiality. In practice, we will also need a generalization of the Kan extension formula (2.4) for maps of Grothendieck fibrations as in (2.1). Keeping the notation therein, given an $\bar{e} \in \bar{\mathcal{E}}$ we will write $\bar{e} \downarrow_\pi \mathcal{E} \hookrightarrow \bar{e} \downarrow \mathcal{E}$ for the full subcategory of those pairs $(e, f: \bar{e} \rightarrow \delta(e))$ such that $\bar{\pi}(f) = \bar{\pi}(\bar{e})$.

Proposition 2.5. *Given a map of Grothendieck fibrations each subcategory $\bar{e} \downarrow_\pi \mathcal{E}$ is an initial subcategory of $\bar{e} \downarrow \mathcal{E}$ so that for each functor $\mathcal{E} \rightarrow \mathcal{V}$ with \mathcal{V} complete one has*

$$\mathbf{Ran}_\delta F(\bar{e}) \simeq \lim F|_{\bar{e} \downarrow \mathcal{E}} \simeq \lim F|_{\bar{e} \downarrow_\pi \mathcal{E}}. \quad (2.6)$$

FIBERKANMAP EQ

Proof. One readily checks that the assignment $(e, f: \bar{e} \rightarrow \delta(e)) \mapsto ((\pi(f)^* e, \bar{e} \rightarrow \delta\pi(f)^*(e)))$ (where $\delta\pi(f)^* = \bar{\pi}^*(f)\delta$) is right adjoint to the inclusion $\bar{e} \downarrow_\pi \mathcal{E} \hookrightarrow \bar{e} \downarrow \mathcal{E}$, so that the claim follows by coreflexivity (note that if not in the split case pullbacks may be chosen arbitrarily). \square

We also record the following, the proof of which is straightforward.

Proposition 2.7. *Suppose that $\mathcal{E} \rightarrow \mathcal{B}$ is a (split) Grothendieck fibration. Then so is the map of functor categories $\mathcal{E}^c \rightarrow \mathcal{B}^c$ for any category \mathcal{C} as well as the map $\bar{\mathcal{E}} \rightarrow \bar{\mathcal{B}}$ in any pullback of categories*

$$\begin{array}{ccc} \bar{\mathcal{E}} & \longrightarrow & \mathcal{E} \\ \downarrow & & \downarrow \\ \bar{\mathcal{B}} & \longrightarrow & \mathcal{B}. \end{array}$$

2.2 Wreath product over finite sets

Throughout we will let \mathbf{F} denote the usual skeleton of the category of (ordered) finite sets and all set maps. Explicitly, its objects are the finite sets $\{1, 2, \dots, n\}$ for $n \geq 0$.

Definition 2.8. For a category \mathcal{C} , we write $\mathbf{F} \wr \mathcal{C} = (\mathbf{F}^{op} \ltimes \mathcal{C}^{\times \bullet})^{op}$ for the opposite of the Grothendieck construction (cf. Definition 2.2) of the functor

$$\begin{aligned} \mathbf{F}^{op} &\longrightarrow \mathbf{Cat} \\ I &\longmapsto \mathcal{C}^{\times I} \end{aligned}$$

Explicitly, the objects of $\mathbf{F} \wr \mathcal{C}$ are tuples $(c_i)_{i \in I}$ and a map $(c_i)_{i \in I} \rightarrow (d_j)_{j \in J}$ consists of a pair

$$(\phi: I \rightarrow J, (f_i: c_i \rightarrow d_{\phi(i)})_{i \in I}),$$

henceforth abbreviated as $(\phi, (f_i))$.

Remark 2.9. Let $(c_i)_{i \in I} \in \mathbf{F} \wr \mathcal{C}$ and write λ for the partition $I = \lambda_1 \sqcup \dots \sqcup \lambda_k$ such that $1 \leq i_1, i_2 \leq n$ are in the same class iff $c_{i_1}, c_{i_2} \in \mathcal{C}$ are isomorphic. Writing $\Sigma_\lambda = \Sigma_{\lambda_1} \times \dots \times \Sigma_{\lambda_k}$ and picking representatives $i_j \in \lambda_j$, the automorphism group of $(c_i)_{i \in I}$ is given by

$$\mathrm{Aut}((c_i)_{i \in I}) \simeq \Sigma_\lambda \wr \prod_i \mathrm{Aut}(c_i) \simeq \Sigma_{|\lambda_1|} \wr \mathrm{Aut}(c_{i_1}) \times \dots \times \Sigma_{|\lambda_k|} \wr \mathrm{Aut}(c_{i_k}). \quad (2.10)$$

Notation 2.11. Using the coproduct functor $\mathbf{F}^{\wr 2} = \mathbf{F}^{\wr \{0,1\}} = \mathbf{F} \wr \mathbf{F} \xrightarrow{\sqcup} \mathbf{F}$ (where $\sqcup_{i \in I} J_i$ is ordered lexicographically) and the singleton $\{1\} \in \mathbf{F}$ one can regard the collection of categories $\mathbf{F}^{\wr n+1} \wr \mathcal{C} = \mathbf{F}^{\wr \{0, \dots, n\}} \wr \mathcal{C}$ for $n \geq -1$ as a coaugmented cosimplicial object in \mathbf{Cat} . As such, we will denote by

$$\delta^i: \mathbf{F}^{\wr n} \wr \mathcal{C} \rightarrow \mathbf{F}^{\wr n+1} \wr \mathcal{C}, \quad 0 \leq i \leq n$$

the cofaces obtained by inserting simpletons $\{1\} \in \mathbf{F}$ and by

$$\sigma^i: \mathbf{F}^{\wr n+2} \wr \mathcal{C} \rightarrow \mathbf{F}^{\wr n+1} \wr \mathcal{C}, \quad 0 \leq i \leq n$$

the codegeneracies obtained by applying the coproduct $\mathbf{F}^{\wr 2} \xrightarrow{\sqcup} \mathbf{F}$ to adjacent \mathbf{F} coordinates.

Further, note that there are identifications $\mathbf{F} \wr \delta^i = \delta^{i+1}$, $\mathbf{F} \wr \sigma^i = \sigma^{i+1}$.

Remark 2.12. If \mathcal{V} has all finite coproducts then injections and fold maps assemble into a functor as on the left below. Dually, if \mathcal{V} has all finite products then projections and diagonals assemble into a functor as on the right.

$$\begin{array}{ccc} \mathbf{F} \wr \mathcal{V} & \xrightarrow{\sqcup} & \mathcal{V} \\ (v_i)_{i \in I} & \longmapsto & \sqcup_{i \in I} v_i \end{array} \quad \begin{array}{ccc} (\mathbf{F} \wr \mathcal{V}^{op})^{op} & \xrightarrow{\prod} & \mathcal{V} \\ (v_i)_{i \in I} & \longmapsto & \prod_{i \in I} v_i \end{array} \quad (2.13) \quad \boxed{\text{WREATHPROD EQ}}$$

Moreover, these functors satisfy a number of additional coherence conditions. Firstly, there is a natural isomorphism α as on the left below

$$\begin{array}{ccc} \mathbf{F}^{\wr 2} \wr \mathcal{V} & \xrightarrow{\mathbf{F} \wr \sqcup} & \mathbf{F} \wr \mathcal{V} \xrightarrow{\sqcup} \mathcal{V} \\ \sigma^0 \downarrow & \nearrow \alpha & \parallel \\ \mathbf{F} \wr \mathcal{V} & \xrightarrow{\sqcup} & \mathcal{V} \end{array} \quad \begin{array}{ccc} \mathcal{V} & & \\ \delta^0 \downarrow & \searrow & \\ \mathbf{F} \wr \mathcal{V} & \xrightarrow{\prod} & \mathcal{V} \end{array} \quad (2.14) \quad \boxed{\text{COHER EQ}}$$

that encodes both changes in parenthesizing of coproducts and removal of initial objects (note that the empty tuple $()_{i \in \emptyset} \in \mathbf{F} \wr \mathcal{V}$ maps under \sqcup to an initial object of \mathcal{V}). Additionally, we are free to assume that the triangle on the right of (2.14) strictly commutes, i.e. that “unary coproducts” of simpletons (v) are given simply by v itself. α is then associative in the sense that the composite natural isomorphisms between the two functors $\mathbf{F}^{\wr 3} \wr \mathcal{V} \rightarrow \mathcal{V}$ in

the diagrams below coincide.

$$\begin{array}{ccc}
 F^{\iota 3} \wr \mathcal{V} & \xrightarrow{F^{\iota 2} \wr \Pi} & F^{\iota 2} \wr \mathcal{V} & \xrightarrow{F \wr \Pi} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} \\
 \sigma^0 \downarrow & & \sigma^0 \downarrow & & \swarrow \alpha & & \parallel \\
 F^{\iota 2} \wr \mathcal{V} & \xrightarrow{F \wr \Pi} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & & \\
 \sigma^1 \downarrow & & \swarrow \alpha & & \parallel & & \\
 F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & & & &
 \end{array}
 \quad
 \begin{array}{ccc}
 F^{\iota 3} \wr \mathcal{V} & \xrightarrow{F^{\iota 2} \wr \Pi} & F^{\iota 2} \wr \mathcal{V} & \xrightarrow{F \wr \Pi} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} \\
 \sigma^0 \downarrow & & \swarrow F \wr \alpha & & \parallel & & \parallel \\
 F^{\iota 2} \wr \mathcal{V} & \xrightarrow{F \wr \Pi} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & & \\
 \sigma^0 \downarrow & & \swarrow \alpha & & \parallel & & \\
 F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & & & &
 \end{array}
 \quad (2.15)$$

COHER2 EQ

Similarly, α is also unital in the sense that both of the following diagrams strictly commute or, more precisely, if the composite natural transformation in either diagram is the identity for the functor $\Pi: F \wr \mathcal{V} \rightarrow \mathcal{V}$.

$$\begin{array}{ccc}
 F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & \xrightarrow{\text{id}} & \mathcal{V} \\
 \delta^0 \downarrow & & \delta^0 \downarrow & & \parallel \\
 F^{\iota 2} \wr \mathcal{V} & \xrightarrow{F \wr \Pi} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} \\
 \sigma^0 \downarrow & & \swarrow \alpha & & \parallel \\
 F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & &
 \end{array}
 \quad
 \begin{array}{ccc}
 F \wr \mathcal{V} & \xrightarrow{\text{id}} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} \\
 \delta^1 \downarrow & & \parallel & & \parallel \\
 F^{\iota 2} \wr \mathcal{V} & \xrightarrow{F \wr \Pi} & F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} \\
 \sigma^0 \downarrow & & \swarrow \alpha & & \parallel \\
 F \wr \mathcal{V} & \xrightarrow{\Pi} & \mathcal{V} & &
 \end{array}
 \quad (2.16)$$

COHER3 EQ

SIGMA_WR_REM

Remark 2.17. More generally, if \mathcal{V} is an arbitrary symmetric monoidal category, one instead has a functor $\Sigma \wr \mathcal{V} \xrightarrow{\otimes} \mathcal{V}$ (where as usual $\Sigma \hookrightarrow F$ denotes finite sets and isomorphisms) satisfying the obvious analogues of (2.14), (2.15), (2.16), as is readily shown using the standard coherence results for symmetric monoidal categories (moreover, we note that α itself encodes all associativity, unital and symmetry isomorphisms, with the right side of (2.14) and (2.16) being mere common sense desiderata for “unary products”).

It is likely no surprise that the converse is also true, i.e. that a functor $\Sigma \wr \mathcal{V} \xrightarrow{\otimes} \mathcal{V}$ satisfying the analogues of (2.14), (2.15), (2.16) endows \mathcal{V} with a symmetric monoidal structure. We will however have no direct need to use this fact, and as such include only a few pointers concerning the associativity pentagon axiom (the hardest condition to check) that the interested reader may find useful. Firstly, it becomes convenient to write expressions such as $(A \otimes B) \otimes C$ instead as $(A \otimes B) \otimes (C)$, so as to encode notationally the fact that this is the image of $((A, B), (C)) \in F^{\iota 2} \wr \mathcal{V}$ under the top map in (2.14). The associativity isomorphisms are hence given by the composites $(A \otimes B) \otimes (C) \xrightarrow{\sim} A \otimes B \otimes C \xleftarrow{\sim} (A) \otimes (B \otimes C)$ obtained by combining $\alpha_{((A, B), (C))}$ and $\alpha_{((A), (B, C))}$. The pentagon axiom is then checked by combining *six* instances of each of the squares in (2.15) (i.e. twelve squares total), most of which are obvious except for the fact that the $(A \otimes B) \otimes (C \otimes D)$ vertex of the pentagon contributes two pairs of squares rather than just one, with each pair corresponding to the two alternate expressions $((A \otimes B)) \otimes ((C) \otimes (D))$ and $((A) \otimes (B)) \otimes ((C \otimes D))$.

FINSURJ_REM

Remark 2.18. In lieu of the two previous remarks, and writing $F_s \hookrightarrow F$ for the subcategory of surjections, we define a *symmetric monoidal category with fold maps* as a category \mathcal{V} together with a functor $F_s \wr \mathcal{V} \xrightarrow{\otimes} \mathcal{V}$ satisfying the analogues of (2.14), (2.15), (2.16). Further, the dual of such \mathcal{V} is called a *symmetric monoidal category with diagonals*.

Maybe mention that if both folds and inclusions then cocartesian

Remark 2.19. Replacing F_s in the previous remark with the subcategory $F_i \hookrightarrow F$ of injections yields the notion of a *symmetric monoidal category with injection maps* or, dually, *symmetric monoidal category with diagonals*.

maybe mention that “cat with diagonals” is the same as semi-cartesian

GENUINE_OP_MONAD_SECTION

We end this section by collecting some straightforward lemmas that will be used in §4.

FWRGROTH LEM

Lemma 2.20. *If $\mathcal{E} \rightarrow \mathcal{B}$ a (split) Grothendieck fibration then so is $F \wr \mathcal{E} \rightarrow F \wr \mathcal{B}$.*

Moreover, if $\mathcal{E} \rightarrow \bar{\mathcal{E}}$ is a map of (split) Grothendieck fibrations over \mathcal{B} then $F \wr \mathcal{E} \rightarrow F \wr \bar{\mathcal{E}}$ is a map of (split) Grothendieck fibrations over $F \wr \mathcal{B}$.

Proof. Given a map $(\phi, (f_i)): (b'_i)_{i \in I} \rightarrow (b_j)_{j \in J}$ in $F \wr \mathcal{B}$ and object $(e_j)_{j \in J}$ one readily checks that its pullback can be defined by $(f_{\phi(i)}^* e_{\phi(i)})_{i \in I}$. \square

WREATPRODLIM LEM

Lemma 2.21. *Suppose that \mathcal{V} is a bicomplete category such that coproducts commute with limits in each variable. If the leftmost diagram*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{G} & \mathcal{V} \\ k \downarrow & \nearrow \eta & \nearrow H \\ \mathcal{D} & & \end{array} \quad \begin{array}{ccccc} F \wr \mathcal{C} & \xrightarrow{F \wr G} & F \wr \mathcal{V} & \xrightarrow{u} & \mathcal{V} \\ F \wr k \downarrow & \nearrow F \wr \eta & \nearrow F \wr H & \nearrow u \circ F \wr H & \\ F \wr \mathcal{D} & & & & \end{array} \quad (2.22) \quad \text{WRRAN EQ}$$

is a right Kan extension diagram then so is the composite of the rightmost diagram.

Dually, if in \mathcal{E} products commute with colimits in each variable, and the leftmost diagram

$$\begin{array}{ccc} \mathcal{C}^{op} & \xrightarrow{G} & \mathcal{V} \\ k^{op} \downarrow & \nearrow \epsilon & \nearrow H \\ \mathcal{D}^{op} & & \end{array} \quad \begin{array}{ccccc} (F \wr \mathcal{C})^{op} & \xrightarrow{(F \wr G)^{op}} & (F \wr \mathcal{V})^{op} & \xrightarrow{\Pi} & \mathcal{V} \\ (F \wr k)^{op} \downarrow & \nearrow (F \wr \epsilon)^{op} & \nearrow (F \wr H)^{op} & \nearrow \Pi \circ (F \wr H)^{op} & \\ (F \wr \mathcal{D})^{op} & & & & \end{array} \quad (2.23) \quad \text{WRLAN EQ}$$

is a left Kan extension diagram then so is the composite of the rightmost diagram.

Proof. Unpacking definitions using the pointwise formula for Kan extensions ([McL11, X.3.1]), the claim concerning (2.22) amounts to showing that for each $(d_i) \in F \wr \mathcal{D}$ one has natural isomorphisms

$$\lim_{((d_i) \rightarrow (kc_j)) \in ((d_i) \downarrow F \wr \mathcal{C})} \left(\coprod_j G(c_j) \right) \simeq \coprod_i \lim_{(d_i \rightarrow kc_i) \in d_i \downarrow \mathcal{C}} (G(c_i)). \quad (2.24) \quad \text{POINTKAN EQ}$$

Proposition 2.5 now applies to the map $F \wr \mathcal{C} \rightarrow F \wr \mathcal{D}$ of Grothendieck fibrations over F and one readily checks that $(d_i) \downarrow_\pi F \wr \mathcal{C} \simeq \prod_i (d_i \downarrow \mathcal{C})$ so that

$$\lim_{((d_i) \rightarrow (kc_j)) \in ((d_i) \downarrow F \wr \mathcal{C})} \left(\coprod_j G(c_j) \right) \simeq \lim_{((d_i) \rightarrow (kc_i)) \in \prod_i (d_i \downarrow \mathcal{D})} \left(\coprod_i G(c_i) \right)$$

and the isomorphisms (2.24) now follow from the assumption that coproducts commute with limits in each variable. \square

2.3 Monads and adjunctions

In §4 we will make use of the following straightforward results concerning the transfer of monads along adjunctions (note that L (resp. R) denotes the left (right) adjoint).

Proposition 2.25. *Let $L: \mathcal{C} \rightleftarrows \mathcal{D}: R$ be an adjunction and T a monad on \mathcal{D} . Then*

- (i) *RTL is a monad and R induces a functor $R: \text{Alg}_T(\mathcal{D}) \rightarrow \text{Alg}_{RTL}(\mathcal{C})$;*
- (ii) *if $LRTL \xrightarrow{\epsilon} TL$ is an isomorphism one further has an induced adjunction*

$$L: \text{Alg}_{RTL}(\mathcal{C}) \rightleftarrows \text{Alg}_T(\mathcal{D}): R.$$

Proposition 2.26. *Let $L: \mathcal{C} \rightleftarrows \mathcal{D}: R$ be an adjunction, T a monad on \mathcal{C} , and suppose further that*

$$LR \xrightarrow{\epsilon} id_{\mathcal{D}}, \quad LT \xrightarrow{\eta} LTRL$$

are natural isomorphisms (so that in particular \mathcal{D} is a reflexive subcategory of \mathcal{C}).

Then

MONADADJ1 PROP

MONADADJ PROP

(i) LTR is a monad, with multiplication and unit given by

$$LTR LTR \xrightarrow{\eta^{-1}} LTTR \rightarrow LTR, \quad id_{\mathcal{D}} \xrightarrow{\epsilon^{-1}} LR \rightarrow LTR;$$

(ii) $d \in \mathcal{D}$ is a LTR -algebra iff Rd is a T -algebra;

(iii) there is an induced adjunction

$$L: \mathbf{Alg}_T(\mathcal{C}) \rightleftarrows \mathbf{Alg}_{LTR}(\mathcal{D}): R.$$

Any monad T on \mathcal{C} induces obvious monads $T^{\times l}$ on $\mathcal{C}^{\times l}$. More generally, and letting I denote the identity monad, a partition $\{1, \dots, l\} = \lambda_a \sqcup \lambda_i$, which we denote by λ , determines a monad $T^{\times \lambda} = T^{\times \lambda_a} \times I^{\times \lambda_i}$ on \mathcal{C} . Here “a” stands for “active” and “i” for “inert”.

Such monads satisfy a number of compatibility conditions. Firstly, if $\lambda'_a \subset \lambda_a$ there is a monad map $T^{\times \lambda'} \Rightarrow T^{\times \lambda}$, and we write $\lambda' \leq \lambda$. Moreover, writing $\alpha^*: \mathcal{C}^{\times m} \rightarrow \mathcal{C}^{\times l}$ for the forgetful functor induced by a map $\alpha: \{1, \dots, l\} \rightarrow \{1, \dots, m\}$, one has an equality $T^{\times \alpha^* \lambda} \alpha^* = \alpha^* T^{\times \lambda}$, where $\alpha^* \lambda$ is the pullback partition. The following is straightforward.

Proposition 2.27. *Suppose \mathcal{C} has finite coproducts and write $\alpha_!: \mathcal{C}^{\times l} \rightarrow \mathcal{C}^{\times m}$ for the left adjoint of α^* . Then the map*

$$T^{\times \alpha^* \lambda} \Rightarrow \alpha^* T^{\times \lambda} \alpha_! \quad (2.28)$$

adjoint to the identity $T^{\times \alpha^ \lambda} \alpha^* = \alpha^* T^{\times \lambda}$ is a map of monads on $\mathcal{C}^{\times l}$.*

Hence, since $T^{\times \lambda} \alpha_!$ is a right $\alpha^ T^{\times \lambda} \alpha_!$ -module, it is also a right $T^{\times \lambda'}$ whenever $\lambda' \leq \alpha^* \lambda$. Finally, the natural map*

$$\alpha_! T^{\times \alpha^* \lambda} \Rightarrow T^{\times \lambda} \alpha_! \quad (2.29)$$

is a map of right $T^{\times \alpha^ \lambda}$ -modules, and thus also a map of right $T^{\times \lambda'}$ -modules whenever $\lambda' \leq \alpha^* \lambda$.*

Remark 2.30. We unpack the content of (2.29) when $\alpha: \{1, \dots, l\} \rightarrow *$ is the unique map to the singleton $*$, in which case we write $\alpha_! = \coprod$. We thus have commutative diagrams

$$\begin{array}{ccc} \coprod_{j \in \lambda_a} TTA_j \sqcup \coprod_{j \in \lambda_i} A_j & \longrightarrow & T(\coprod_{j \in \lambda_a} TA_j \sqcup \coprod_{j \in \lambda_i} A_j) \\ \downarrow & & \downarrow \\ \coprod_{j \in \lambda_a} TA_j \sqcup \coprod_{j \in \lambda_i} A_j & \longrightarrow & T(\coprod_{j \in \lambda_a} A_j \sqcup \coprod_{j \in \lambda_i} A_j) \end{array} \quad (2.31)$$

for each collection $(A_j)_{j \in \ell}$ in \mathcal{C} , where the vertical maps come from the right $T^{\times \lambda}$ -module structure. Writing $\check{\sqcup}$ for the coproduct of T -algebras and recalling the canonical identifications $\check{\sqcup}_{k \in K} (TA_k) \simeq T(\check{\sqcup}_{k \in K} A_k)$, (2.31) shows that the right $T^{\times \lambda}$ -module structure on $T \circ \check{\sqcup}$ codifies the multiplication maps

$$\check{\sqcup}_{j \in \lambda_a} TTA_j \check{\sqcup} \check{\sqcup}_{j \in \lambda_i} TA_j \rightarrow \check{\sqcup}_{j \in \lambda_a} TA_j \check{\sqcup} \check{\sqcup}_{j \in \lambda_i} TA_j.$$

3 Planar and tall maps

Throughout we will assume that the reader is familiar with the category Ω of trees. A good introduction to Ω is given by [13, §3], where arrows are described both via the “colored operad generated by a tree” and by identifying explicit generating arrows, called faces and degeneracies. Alternatively, Ω can also be described using the algebraic model of *broad posets* introduced by Weiss in [21] and further worked out by the second author in [15, §5]. This latter will be our “official” model, though a detailed understanding of broad posets is needed only to follow our formal discussion of planar structures in §5.1, and the reader willing to accept the results therein should be able to read the remainder of the paper.

Given a finite group G , there is also a category Ω_G of G -trees, jointly discovered by the authors and first discussed by the second author in [Pe17, §4.3, §5.3], which we now recall. Firstly, we let Φ denote the category of forests, i.e. “formal coproducts of trees”. A broad poset description of Φ is found in [15, §5.2], but here we prefer the alternative definition $\Phi = \mathbf{F} \wr \Omega$. The category of G -forests is then Φ^G , i.e. the category of G -objects in Φ . Identifying the G -orbit category as the subcategory $\mathbf{O}_G \hookrightarrow \mathbf{F}^G$ of those sets with transitive actions, Ω_G can then be described as given by the pullback of categories

$$\begin{array}{ccc} \Omega_G & \longrightarrow & \Phi^G \\ \downarrow & & \downarrow \\ \mathbf{O}_G & \longrightarrow & \mathbf{F}^G, \end{array} \quad (3.1) \quad \text{OGDEF EQ}$$

which is a repackaging of [Pe17, Prop. 5.46]. Explicitly, a G -tree T is then a tuple $T = (T_x)_{x \in X}$ with $X \in \mathbf{O}_G$ together with isomorphisms $T_x \rightarrow T_{gx}$ that are suitably associative and unital.

3.1 Planar structures

The specific model for the orbit category \mathbf{O}_G used in (3.1) has extra structure not found in the usual model (i.e. that of the G -sets G/H for $H \leq G$), namely the fact that each $X \in \mathbf{O}_G$ comes with a canonical total order (the underlying set of X being one of the sets $\{1, \dots, n\}$).


We will find it convenient to use a model of Ω with similar extra structure, given by planar structures on trees. Intuitively, a planar structure on a tree is the data of a planar representation of the tree, and definitions of *planar trees* along those lines are found throughout the literature. However, to allow for precise proofs of some key results concerning the interaction of planar structures with the maps in Ω (namely Propositions 3.28 and 3.47) we will instead use a combinatorial definition of planar structure in the context of broad posets.

In what follows a tree will be a *dendroidally ordered broad poset* as in [We12, [Pe17, Def. 5.9].

Definition 3.2. Let $T \in \Omega$ be a tree. A *planar structure* of T is an extension of the descendency partial order \leq_d to a total order \leq_p such that:

- *Planar*: if $e \leq_p f$ and $e \not\leq_d f$ then $g \leq_d f$ implies $e \leq_p g$.

Example 3.3. An example of a planar structure on a tree T follows, with \leq_p encoded by the number labels.



Intuitively, given a planar depiction of a tree T , $e \leq_d f$ holds when the downward path from e passes through f and $e \leq_p f$ holds if either $e \leq_d f$ or if the downward path from e is to the left of the downward path from f (as measured at the node where the paths intersect).

It is visually clear that a planar depiction of a tree amounts to choosing a total order for each of the sets of *input edges* of each node (i.e. those edges immediately above that node).

While we will not need to make this last statement precise, we will nonetheless find it convenient to show that Definition 3.2 is equivalent to such choices of total orders for each of the sets of input edges. To do so, we first introduce some notation.

INPUTPATH NOT

Notation 3.5. Let $T \in \Omega$ be a tree and $e \in T$ an edge. We will denote

$$I(e) = \{f \in T : e \leq_d f\}$$

and refer to this poset as the *input path* of e .

We will repeatedly use the following, which is a consequence of [Pe17, Cor. 5.26].

INCOMPNOTOP

Lemma 3.6. If $e \leq_d f$, $e \leq_d f'$, then f, f' are \leq_d -comparable.

INPUTPATHS PROP

Proposition 3.7. Let $T \in \Omega$ be a tree. Then

- (a) for any $e \in T$ the finite poset $I(e)$ is totally ordered;
- (b) the poset (T, \leq_d) has all joins, denoted \vee . In fact, $\bigvee_i e_i = \min(\bigcap_i I(e_i))$.

Proof. (a) is immediate from Lemma 3.6. To prove (b) we note that $\min(\bigcap_i I(e_i))$ exists by (a), and that this is clearly the join $\bigvee_i e_i$. \square

Notation 3.8. Let $T \in \Omega$ be a tree and suppose that $e <_d b$. We will denote by $b_e^\dagger \in T$ the predecessor of b in $I(e)$.

Proposition 3.9. Suppose e, f are \leq_d -incomparable edges of T and write $b = e \vee f$. Then

- (a) $e <_d b$, $f <_d b$ and $b_e^\dagger \neq b_f^\dagger$;
- (b) $b_e^\dagger, b_f^\dagger \in b^\dagger$. In fact $\{b_e^\dagger\} = I(e) \cap b^\dagger$, $\{b_f^\dagger\} = I(f) \cap b^\dagger$;
- (c) if $e' \leq_d e$, $f' \leq_d f$ then $b = e' \vee f'$ and $b_{e'}^\dagger = b_e^\dagger$, $b_{f'}^\dagger = b_f^\dagger$.

Proof. (a) is immediate: the condition $e = b$ (resp. $f = b$) would imply $f \leq_d e$ (resp. $e \leq_d f$) while the condition $b_e^\dagger = b_f^\dagger$ would provide a predecessor of b in $I(e) \cap I(f)$.

For (b), note that any relation $a <_d b$ factors as $a \leq_d b_a^* <_d b$ for some unique $b_a^* \in b^\dagger$, where uniqueness follows from Lemma 3.6. Choosing $a = e$ implies $I(e) \cap b^\dagger = \{b_e^*\}$ and letting a range over edges such that $e \leq_d a <_d b$ shows that b_e^* is in fact the predecessor of b .

To prove (c) one reduces to the case $e' = e$, in which case it suffices to check $I(e) \cap I(f') = I(e) \cap I(f)$. But if it were otherwise there would exist an edge a satisfying $f' \leq_d a <_d f$ and $e \leq_d a$, and this would imply $e \leq_d f$, contradicting our hypothesis. \square

Proposition 3.10. Let $c = e_1 \vee e_2 \vee e_3$. Then $c = e_i \vee e_j$ iff $c_{e_i}^\dagger \neq c_{e_j}^\dagger$.

Therefore, all ternary joins in (T, \leq_d) are binary, i.e.

$$c = e_1 \vee e_2 \vee e_3 = e_i \vee e_j \tag{3.11}$$


TERNJOIN EQ

for some $1 \leq i < j \leq 3$, and (3.11) fails for at most one choice of $1 \leq i < j \leq 3$.

Proof. If $c_{e_i}^\dagger \neq c_{e_j}^\dagger$, then $c = \min(I(e_i) \cap I(e_j)) = e_i \vee e_j$, whereas the converse follows from Proposition 3.9(a).

The “therefore” part follows by noting that $c_{e_1}^\dagger, c_{e_2}^\dagger, c_{e_3}^\dagger$ can not all coincide, or else c would not be the minimum of $I(e_1) \cap I(e_2) \cap I(e_3)$. \square

Example 3.12. In the following example $b = e \vee f$, $c = e \vee f \vee g$, $c_e^\dagger = c_f^\dagger = b$.



Notation 3.13. Given a set S of size n we write $\text{Ord}(S) \simeq \text{Iso}(S, \{1, \dots, n\})$. We will freely abuse notation by regarding its objects as pairs (S, \leq) where \leq is a total order in S .

Proposition 3.14. *Let $T \in \Omega$ be a tree. There is a bijection*

$$\begin{aligned} \{\text{planar structures } (T, \leq_p)\} &\xrightarrow{\cong} \prod_{(a^\dagger \leq a) \in V(T)} \text{Ord}(a^\dagger) \\ \leq_p &\longmapsto (\leq_p \upharpoonright_{a^\dagger}) \end{aligned} \quad (3.15) \quad \text{PLANAR EQ}$$

Proof. We will keep the notation of Proposition 3.9 throughout: e, f are \leq_d -incomparable edges and we write $b = e \vee f$.

We first show that (3.15) is injective, i.e. that the restrictions $\leq_p \upharpoonright_{a^\dagger}$ determine if $e <_p f$ holds or not. If $b_e^\dagger <_p b_f^\dagger$, the relations $e \leq_d b_e^\dagger <_p b_f^\dagger \geq_d f$ and Definition 3.2 imply it must be $e <_p f$. Dually, if $b_f^\dagger <_p b_e^\dagger$ then $f <_p e$. Thus $b_e^\dagger <_p b_f^\dagger \Leftrightarrow e <_p f$ and hence (3.15) is indeed injective.

To check that (3.15) is surjective, it suffices (recall that e, f are assumed \leq_d -incomparable) to check that defining $e \leq_p f$ to hold iff $b_e^\dagger < b_f^\dagger$ holds in b^\dagger yields a planar structure.

Antisymmetry and the total order conditions are immediate, and it thus remains to check the transitivity and planar conditions. Transitivity of \leq_p in the case $e' <_p e <_p f$ and the planar condition, which is the case $e <_p f \geq_d f'$, follow from Proposition 3.9(c). Transitivity of \leq_p in the case $e <_p f \leq_d f'$ follows since either $e \leq_d f'$ or else e, f' are \leq_d -incomparable, in which case one can apply 3.9(c) with the roles of f, f' reversed.

It remains to check transitivity in the hardest case, that of $e <_p f <_p g$ with e, f, g pairwise incomparable. We write $c = e \vee f \vee g$. By the “therefore” part of Proposition 3.10, either (i) $e \vee f <_d c$, in which case Proposition 3.10 implies $c = e \vee g$, $c_e^\dagger = c_f^\dagger$ and transitivity follows; (ii) $f \vee g <_d c$, which follows just as (i); (iii) $e \vee f = f \vee g = c$, in which case $c_e^\dagger < c_f^\dagger < c_g^\dagger$ in c^\dagger so that $c_e^\dagger \neq c_g^\dagger$ and by Proposition 3.10 it is also $c = e \vee g$ and transitivity follows. \square

CLOSURE REM


Remark 3.16. Proposition 3.14 states in particular that \leq_p is the closure of the relations in \leq_d and on the vertices a^\dagger under the planar condition in Definition 3.2.

The discussion of the substitution procedure in §3.2 will be significantly simplified by working with a model for the category Ω possessing exactly one representative of each possible planar structure on each tree or, more precisely, if the only isomorphisms preserving the planar structure are the identities. On the other hand, exclusively using such a model for Ω throughout would, among other issues, make the discussion of faces in §3.2 rather awkward. We now outline our conventions to address such issues.

Let Ω^p , the category of *planarized trees*, denote the category with objects pairs $T_{\leq_p} = (T, \leq_p)$ of trees together with a planar structure and morphisms the *underlying* maps of trees (so that the planar structures are ignored). There is a full subcategory $\Omega^s \hookrightarrow \Omega^p$, whose objects we call *standard models*, of those T_{\leq_p} whose underlying set is one of the sets $\underline{n} = \{1, 2, \dots, n\}$ and for which \leq_p coincides with the canonical order.

STANDMODEL EX

Example 3.17. Some examples of standard models, i.e. objects of Ω^s , follow (further, (3.4) can also be interpreted as such an example).




Here T_1 and T_2 are isomorphic to each other but not isomorphic to any other standard model in Ω^s while both C and U are the unique objects in their isomorphism classes.

Given $T_{\leq_p} \in \Omega^p$ there is an obvious standard model $T_{\leq_p}^s \in \Omega^s$ given by replacing each edge by its order following \leq_p . Indeed, this defines a retraction $(-)^s: \Omega^p \rightarrow \Omega^s$ and a natural transformation $\sigma: id \Rightarrow (-)^s$ given by isomorphisms preserving the planar structure (in fact, the pair $((-)^s, \sigma)$ is unique characterized by this property).

FORESTPLAN REM

Remark 3.19. Definition 3.2 readily extends to the broad poset definition of forests $F \in \Phi$ in [15, Def. 5.27], with the analogue of Proposition 3.14 then stating that a planar structure is equivalent to total orderings of the nodes of F together with a total ordering of its set of roots. There are thus two competing notions of standard forests: the [15, Def. 5.27] model Φ^s whose objects are planar forest structures on one of the standard sets $\{1, \dots, n\}$ and (following the discussion at the start of §5) the model $\mathbf{F} \wr \Omega^s$, whose objects are tuples, indexed by a standard set, of planar tree structures on standard sets. An illustration follows.



TWOPLAFCORCONV EQ

It is however clear that there is a *canonical* isomorphism $\Phi^s \simeq \mathbf{F} \wr \Omega^s$ (with the two side of (3.20) representing the same planar forest). Moreover, while the similarly defined categories Φ^p and $\mathbf{F} \wr \Omega^p$ are only equivalent (rather than isomorphic), their retractions onto $\Phi^s \simeq \mathbf{F} \wr \Omega^s$ are compatible, and we will thus henceforth not distinguish between Φ^s and $\mathbf{F} \wr \Omega^s$.

PLANARCONV CON

Convention 3.21. From now on we write simply Ω , Ω_G to denote the categories Ω^s , Ω_G^s of standard models (where planar structures are defined in the underlying forest as in Remark 3.19). Therefore, whenever a construction produces an object/diagram in Ω^p , Ω_G^p (of trees, G -trees) we always implicitly reinterpret it by using the standardization functor $(-)^s$.


Similarly, any finite set or orbital finite G -set together with a total order is implicitly reinterpreted as an object in \mathbf{F} , \mathbf{O}_G^p .

Example 3.22. To illustrate our convention, consider the trees in Example 3.17.

There are subtrees $F_1 \subset F_2 \subset U$ where F_1 is the subtree with edge set $\{1, 2, 6, 7\}$ and F_2 is the subtree with edge set $\{1, 2, 3, 6, 7\}$, both with inherited tree and planar structures. Applying $(-)^s$ to the inclusion diagram on the left below then yields a diagram as on the right.




Similarly, let $\leq_{(12)}$ and $\leq_{(45)}$ denote alternate planar structures for U exchanging the orders of the pairs 1, 2 and 4, 5, so that one has objects $U_{\leq_{(12)}}$, $U_{\leq_{(45)}}$ in Ω^p . Applying $(-)^s$ to the diagram of underlying identities on the left yields the permutation diagram on the right.



Example 3.23. An additional reason to leave the use of $(-)^s$ implicit as detailed in Convention 3.21 is that when depicting G -trees it is preferable to choose edge labels that describe the action rather than the planarization (which is already implicit anyway).

For example, when $G = \mathbb{Z}/4$, $\tilde{G} = \mathbb{Z}/3$, in both diagrams below the orbital representation on the left represents the isomorphism class consisting only of the two trees $T_1, T_2 \in \Omega_G$ on

the right.



In general, isomorphism classes are of course far bigger. The interested reader may show that there are $3 \cdot 3! \cdot 2 \cdot 3! \cdot 3!$ trees in the isomorphism class of the tree depicted in (I.9).

The attentive reader may have noted that it follows from Proposition 2.7 that both vertical maps in (B.1) are split Grothendieck fibrations. We now introduce some terminology.

Definition 3.24. The map $r: \Omega_G \rightarrow \mathcal{O}_G$ in (B.1) is called the *root functor*.

Further, fiber maps (i.e. maps inducing identities, i.e. ordered bijections, on $r(-)$) are called *rooted maps* and pullbacks with respect to r are called *root pullbacks*.

To motivate the terminology, note first that unpacking definitions shows that $r(T)$ is the ordered set of tree components of $T \in \Omega_G$, which coincides with the ordered set of roots. The exact choice of name is further meant to accentuate the connection with another key functor which we call the *leaf-root functor*, described in §3.3.

Further, unpacking the construction in (B.1), one sees that the pullback of the G -tree $T = (T_x)_{x \in X}$ with structure maps $T_x \rightarrow T_{gx}$ along the map $\varphi: Y \rightarrow X$ is simply the G -tree $(T_{\varphi(y)})_{y \in Y}$ with structure maps $T_{\varphi(y)} \rightarrow T_{g\varphi(y)} = T_{\varphi(gy)}$.

Example 3.25. Let $G = \{\pm 1, \pm i, \pm j, \pm k\}$, $H = \langle j \rangle$ and $K = \langle -1 \rangle$. Figure I illustrates the pullbacks of two G -trees T and S along the twist map $\tau: G/H \rightarrow G/H$ and the unique map $\pi: G/H \rightarrow G/G$ (or, more precisely, noting that in our model the underlying set of G/H is actually $\{1, 2\}$, τ is the permutation (12)). We note that the stabilizers of a, b, c are $\{1\}, K, H$ for T and K, H, G for S . The top depictions of τ^*T , $\pi^*(S)$ then use the edge orbit generators suggested by T, S while the bottom depictions choose generators that are minimal with regard to the planar structure, so that in τ^*T it is $d = ic$, $e = ib$, $f = ia$ and in π^*S it is $e = ib'$, $d = ia'$.

Definition 3.26. A map $S \xrightarrow{\varphi} T$ in Ω preserving the planar structure \leq_p is called a *planar map*.

More generally, a map $F \rightarrow G$ in one of the categories Φ, Φ^G, Ω^G of forests, G -forests, G -trees is called a *planar map* if it is an independent map (cf. [Pe17, Def. 5.28]) compatible with the planar structures \leq_p .

Remark 3.27. The need for the independence condition is justified by [Pe17, Lemma 5.33] and its converse, since non independent maps do not reflect \leq_a -comparability.

However, we note that in the case of Ω_G independence admits simpler characterizations: φ is independent iff φ is injective on each edge orbit iff φ is injective on the root orbit.

Proposition 3.28. Let $F \xrightarrow{\varphi} G$ be an independent map in Φ (or Ω, Ω_G, Φ_G). Then there is a unique factorization

$$F \xrightarrow{\sim} \bar{F} \rightarrow G$$

such that $F \xrightarrow{\sim} \bar{F}$ is an isomorphism and $\bar{F} \rightarrow G$ is planar.

Proof. We need to show that there is a unique planar structure $\leq_{\bar{F}}$ on the underlying forest of F making the underlying map a planar map. Simplicity of the broad poset G ensures




Figure 1: Root pullbacks

FIGURE

that for any vertex $e^\dagger \leq e$ of F the edges in $\varphi(e^\dagger)$ are all distinct while independence of φ likewise ensures that the edges in $\varphi(r_F)$ are distinct. By (the forest version of) Proposition 3.14 the only possible planar structure \leq_p^F is the one which orders each set e^\dagger and the root tuple r_F according to their images. The claim that φ is then planar follows from Remark 3.16 together with the fact ([15, Lemma 5.33]) that φ reflects \leq_d -comparability. \square

Remark 3.29. Proposition 3.28 says that planar structures can be pulled back along independent maps. However, they can not always be pushed forward. As an example, in the notation of (3.18), consider the map $C \rightarrow T_1$ defined by $1 \mapsto 1, 2 \mapsto 4, 3 \mapsto 2, 4 \mapsto 5$.

3.2 Outer faces, tall maps, and substitution

One of the key ideas needed for our description of operads is the notion of substitution of tree nodes, a process that we will prefer to repackage in terms of maps of trees.

In preparation for that discussion, we first recall some basic definitions and results concerning outer subtrees and tree grafting, as in [15, §5].

Definition 3.30. Let $T \in \Omega$ be a tree and $e_1 \cdots e_n = \underline{e} \leq e$ a broad relation in T .

We define the *planar outer face* $T_{\underline{e} \leq e}$ to be the subtree with underlying set those edges $f \in T$ such that

$$f \leq_d e, \quad \forall i f \not\leq_d e_i, \quad (3.31)$$

generating broad relations the relations $f^\dagger \leq f$ for those $f \in T$ satisfying (3.31) but $\forall i f \neq e_i$, and planar structure pulled back from T (in the sense of Remark 3.29).

Remark 3.32. If one forgoes the requirement that $T_{\underline{e} \leq e}$ be equipped with the pullback planar structure, the inclusion $T_{\underline{e} \leq e} \hookrightarrow T$ is usually called simply an *outer face*.

We now recap some basic results.

Proposition 3.33. *Let $T \in \Omega$ be a tree.*

- (a) $T_{\underline{e} \leq e}$ is a tree with root e and edge tuple \underline{e} ;
(b) there is a bijection

$$\{\text{planar outer faces of } T\} \leftrightarrow \{\text{broad relations of } T\};$$

- (c) if $R \rightarrow S$ and $S \rightarrow T$ are outer face maps then so is $R \rightarrow T$;
(d) any pair of broad relations $\underline{g} \leq v$, $\underline{f}v \leq e$ induces a grafting pushout diagram

$$\begin{array}{ccc} \eta & \xrightarrow{v} & T_{\underline{g} \leq v} \\ v \downarrow & & \downarrow \\ T_{\underline{f}v \leq e} & \longrightarrow & T_{\underline{f}\underline{g} \leq e}. \end{array} \quad (3.34) \quad \boxed{\text{GRAPTPUSH EQ}}$$

Further, $T_{\underline{f}\underline{g} \leq e}$ is the unique choice of pushout that makes the maps in (3.34) planar. $\boxed{\text{GRAPTPUSH EQ}}$

Proof. We first show (a). That $T_{\underline{e} \leq e}$ is indeed a tree is the content of [Pe17, Prop. 5.20]: more precisely, $T_{\underline{e} \leq e} = (T^{\leq e})_{< \underline{e}}$ in the notation therein. That the root of $T_{\underline{e} \leq e}$ is e is clear and that the root tuple is \underline{e} follows from [Pe17, Remark 5.23].

(b) follows from (a), which shows that $\underline{e} \leq e$ can be recovered from $T_{\underline{e} \leq e}$. $\boxed{\text{Pe17}}$

(c) follows from the definition of outer face together with [Pe17, Lemma 5.33], which states that the \leq_d relations on S, T coincide.

Since by (b) and (c) both $T_{\underline{g} \leq v}$ and $T_{\underline{f}v \leq e}$ are outer faces of $T_{\underline{f}\underline{g} \leq e}$, the first part of (d) is a restatement of [Pe17, Prop. 5.15], while the additional planarity claim follows by Proposition 5.14 together with the vertex identification $V(T_{\underline{f}\underline{g} \leq e}) = V(T_{\underline{f}v \leq e}) \sqcup V(T_{\underline{g} \leq v})$. \square

Definition 3.35. A map $S \xrightarrow{\varphi} T$ in Ω is called a *tall map* if

$$\varphi(l_S) = l_T, \quad \varphi(r_S) = r_T,$$

where $l_{(-)}$ denotes the (unordered) leaf tuple and $r_{(-)}$ the root.

The following is a restatement of [Pe17, Cor. 5.24]

Proposition 3.36. Any map $S \xrightarrow{\varphi} T$ in Ω has a factorization, unique up to unique isomorphism,

$$S \xrightarrow{\varphi^t} U \xrightarrow{\varphi^u} T$$

as a tall map followed by an outer face (in fact, $U = T_{\varphi(l_S) \leq r_S}$).

We recall that a face $F \rightarrow T$ is called inner if is obtained by iteratively removing inner edges, i.e. edges other than the root or the leaves. In particular, it follows that a face is inner iff it is tall. The usual face-degeneracy decomposition thus combines with Proposition 3.36 to give the following. $\boxed{\text{TALLOUTERDEC PROP}}$

Corollary 3.37. Any map $S \xrightarrow{\varphi} T$ in Ω has a factorization, unique up to unique isomorphisms,

$$S \xrightarrow{\varphi^-} U \xrightarrow{\varphi^i} V \xrightarrow{\varphi^u} T \quad (3.38) \quad \boxed{\text{TRIPLEFACT EQ}}$$

as a degeneracy followed by an inner face followed by an outer face.

Proof. The factorization (3.38) can be built by first performing the degeneracy-face decomposition and then performing the tall-outer decomposition on the face map. \square

We will find it convenient throughout to regard the groupoid Σ of finite sets as the subcategory $\Sigma \hookrightarrow \Omega$ consisting of *corollas* (i.e. trees with a single vertex) and isomorphisms.


Notation 3.39. Given a tree $T \in \Omega$ there is a unique corolla $\text{lr}(T) \in \Sigma$ and planar tall map $\text{lr}(T) \rightarrow T$, which we call the *leaf-root* of T (this name is motivated by the equivariant analogue, discussed in §3.3). Explicitly, the number of leaves of $\text{lr}(T)$ matches that of T , together with the inherited order. $\boxed{\text{LRVERT SEC}}$

We now turn to discussing the substitution operation. We start with an example, focused on the closely related notion of iterated graftings of trees (as described in (3.34)).

Example 3.40. The trees U_1, U_2, \dots, U_6 on the left below can be grafted to obtain the tree U in the middle. More precisely (among other possible grafting orders), one has

$$U = (((((U_6 \sqcup_a U_2)) \sqcup_a U_1) \sqcup_b U_3) \sqcup_d U_5) \sqcup_c U_4 \quad (3.41)$$

UFORMULA EQ




(3.42)

SUBSDATUMTREES EQ

We now consider the tree T , which is built by converting each U_i into the corolla $\text{lr}(U_i)$, and then performing the same grafting operations as in (3.41). T can then be regarded as encoding the combinatorics of the iterated grafting in (3.41), with alternative ways to reorder operations in (3.41) in bijection with ways to assemble T out of its nodes.

One can now therefore think of the iterated grafting (3.41) as being instead encoded by the tree T together with the (unique) planar tall maps φ_i below.



(3.43)

SUBSDATUMTREES2 EQ

From this perspective, U can then be thought of as obtained from T by *substituting* each of its nodes with the corresponding U_i . Moreover, the φ_i assemble to a planar tall map $\varphi: T \rightarrow U$ (such that $a_i \mapsto a, b \mapsto b, \dots, e \mapsto e$), which likewise encodes the same information.

One of the fundamental ideas that shape our perspective on operads is then that data for substitution of nodes as in (3.43) can equivalently be repackaged using planar tall maps.

Definition 3.44. Let $T \in \Omega$ be a tree.

UBSTITUTIONDATUM

A T -substitution datum is a tuple $(U_{e^\dagger \leq e})_{(e^\dagger \leq e) \in V(T)}$ together with tall maps $T_{e^\dagger \leq e} \rightarrow U_{e^\dagger \leq e}$. Further, a map of T -substitution data $(U_{e^\dagger \leq e}) \rightarrow (V_{e^\dagger \leq e})$ is a tuple of tall maps $(U_{e^\dagger \leq e} \rightarrow V_{e^\dagger \leq e})$ compatible with the substitution maps.

Lastly, a substitution datum is called a *planar T -substitution datum* if the chosen maps are planar (so that $\text{lr}(U_{e^\dagger \leq e}) = T_{e^\dagger \leq e}$) and a morphism of planar data is called a planar morphism if it consists of a tuple of planar maps.

We denote the category of (resp. planar) T -substitution data by $\text{Sub}(T)$ (resp. $\text{Sub}_p(T)$).

Definition 3.45. Let $T \in \Omega$ be a tree. The *Segal core poset* $\text{Sc}(T)$ is the poset with objects the single edge subtrees η_e and vertex subtrees $T_{e^\dagger \leq e}$, ordered by inclusion.

Remark 3.46. Note that the only maps in $\text{Sc}(T)$ are inclusions of the form $\eta_a \subset T_{e^\dagger \leq e}$. In particular, there are no pairs of composable non-identity relations in $\text{Sc}(T)$.

Given a T -substitution datum $\{U_{\{e^\dagger \leq e\}}\}$ we abuse notation by writing

$$U_{(-)} : \text{Sc}(T) \rightarrow \Omega$$

for the functor $\eta_a \mapsto \eta$, $T_{e^\dagger \leq e} \mapsto U_{e^\dagger \leq e}$ and sending the inclusions $\eta_a \subset T_{e^\dagger \leq e}$ to the composites

$$\eta \xrightarrow{a} T_{e^\dagger \leq e} \rightarrow U_{e^\dagger \leq e}.$$

Proposition 3.47. Let $T \in \Omega$ be a tree. There is an isomorphism of categories

$$\begin{aligned} \text{Sub}_p(T) &\xrightleftharpoons{\quad} T \downarrow \Omega^{\text{pt}} \\ (U_{e^\dagger \leq e}) &\longmapsto (T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}) \\ (U_{\varphi(e^\dagger) \leq \varphi(e)}) &\longleftarrow (T \xrightarrow{\varphi} U) \end{aligned} \tag{3.48}$$

SUBDATAUNDERPLAN EQ

where $T \downarrow \Omega^{\text{pt}}$ denotes the category of planar tall maps under T and $\text{colim}_{\text{Sc}(T)} U_{(-)}$ is chosen in the unique way that makes the inclusions of the $U_{e^\dagger \leq e}$ planar.

Proof. We first show in parallel that: (i) $\text{colim}_{\text{Sc}(T)} U_{(-)}$, which we denote U_T , exists; (ii) for the datum $(T_{e^\dagger \leq e})$, it is $T = \text{colim}_{\text{Sc}(T)} T_{(-)}$; (iii) $V(U_T) = \coprod_{V(T)} V(U_{e^\dagger \leq e})$; (iv) the induced map $T \rightarrow U_T$ is planar tall.

The argument is by induction on the number of vertices of T , with the base cases of T with 0 or 1 vertices being immediate, since then T is the terminal object of $\text{Sc}(T)$. Otherwise, one can choose a non trivial grafting decomposition so as to write $T = R \cup_e S$, resulting in identifications $\text{Sc}(R) \subset \text{Sc}(T)$, $\text{Sc}(S) \subset \text{Sc}(T)$ so that $\text{Sc}(R) \cup \text{Sc}(S) = \text{Sc}(T)$ and $\text{Sc}(R) \cap \text{Sc}(S) = \{\eta_e\}$. The existence of $U_T = \text{colim}_{\text{Sc}(T)} U_{(-)}$ is thus equivalent to the existence of the pushout below (where the rightmost diagram merely simplifies notation).

$$\begin{array}{ccc} \eta & \xrightarrow{e} & \text{colim}_{\text{Sc}(R)} U_{(-)} \\ \downarrow e & & \downarrow \\ \text{colim}_{\text{Sc}(S)} U_{(-)} & \dashrightarrow & \text{colim}_{\text{Sc}(T)} U_{(-)} \end{array} \quad \begin{array}{ccc} \eta & \xrightarrow{e} & U_R \\ \downarrow e & & \downarrow \\ U_S & \dashrightarrow & U_T \end{array} \tag{3.49}$$

ASSEMBLYGRAFT EQ

By induction, U_R and U_S exist for any $U_{(-)}$, equal R and S in the case $U_{(-)} = T_{(-)}$, $V(U_R) = \coprod_{V(R)} V(U_{e^\dagger \leq e})$ and likewise for S (so that there are unique choices of U_R, U_S making the inclusions of $U_{e^\dagger \leq e}$ planar), and the maps $R \rightarrow \text{colim}_{\text{Sc}(R)} U_{(-)}$, $S \rightarrow \text{colim}_{\text{Sc}(S)} U_{(-)}$ are planar tall. But it now follows that (3.49) is a grafting pushout diagram (cf. (3.34)), so that the pushout indeed exists. The conditions $T = \text{colim}_{\text{Sc}(T)} T_{(-)}$, $V(U_T) = \coprod_{V(T)} V(U_{e^\dagger \leq e})$, and that $T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}$ is planar tall follow.

The fact that the two functors in (3.48) are inverse to each other is clear from the same inductive argument. \square

Corollary 3.50. *Let $T \in \Omega$ be a tree. The formulas in (3.48) give an isomorphism of categories*

$$\text{Sub}(T) \xrightarrow{\sim} T \downarrow \Omega^t \quad (3.51)$$

SUBDATAUNDERNONPL EQ

where $T \downarrow \Omega^t$ denotes the category of tall maps under T .

Proof. This is a consequence of Proposition 3.28 together with the previous result. Indeed, Proposition 3.14 can be restated as saying that isomorphisms $T \rightarrow T'$ are in bijection with substitution data consisting of isomorphisms, and thus bijectiveness of $\text{Sub}(T) \rightarrow T \downarrow \Omega^t$ reduces to that in the previous result. \square

Remark 3.52. As noted in the proof of Proposition 3.47, writing $U = \text{colim}_{\text{sc}(T)} U_{(-)}$, one has

$$V(U) = \coprod_{(e^\dagger \leq e) \in V(T)} V(U_{e^\dagger \leq e}). \quad (3.53)$$

VERTEXDECOMP EQ

Alternatively, (3.53) can be regarded as a map $\varphi^*: V(U) \rightarrow V(T)$ induced by the planar tall map $\varphi: T \rightarrow U$. Explicitly, $\varphi^*(U_{u^\dagger \leq u})$ is the unique $T_{t^\dagger \leq t}$ such that there is an inclusion of outer faces $U_{u^\dagger \leq u} \hookrightarrow U_{t^\dagger \leq t}$, so that φ^* indeed depends contravariantly on the tall map φ .

Remark 3.54. Suppose that $e \in T$ has input path $I_T(e) = (e = e_n < e_{n-1} < \dots < e_0)$. It is intuitively clear that for a tall map $\varphi: T \rightarrow U$ the input path of $\varphi(e)$ is built by gluing input paths in the $U_{t^\dagger \leq t}$. More precisely (and omitting φ for readability), one has

$$I_U(e_n) \simeq I_{n-1}(e_n) \sqcup_{e_{n-1}} I_{n-2}(e_{n-1}) \sqcup_{e_{n-2}} \dots \sqcup_{e_1} I_1(e_0).$$

where $I_k(-)$ denotes the input path in $U_{t^\dagger \leq t}$. More formally, this follows from the characterization of predecessors in Proposition 3.9(b).

We end this section with a couple of lemmas that will allow us to reverse the substitution procedure of Proposition 3.47 and will be needed in §5.2.

Proposition 3.55. *Let $U \in \Omega$ be a tree. Then:*

- (i) *given non stick outer subtrees U_i such that $V(U) = \coprod_i V(U_i)$ there is a unique tree T and planar tall map $T \rightarrow U$ such that the sets $\{U_i\}$, $\{U_{e^\dagger \leq e}\}$ coincide;*
- (ii) *given multiplicities $m_e \geq 1$ for each edge $e \in U$, there is a unique planar degeneracy $\rho: T \rightarrow U$ such that $\rho^{-1}(e)$ has m_e elements;*
- (iii) *planar tall maps $T \rightarrow U$ are in bijection with collections $\{U_i\}$ of outer subtrees such that $V(U) = \coprod_i V(U_i)$ and U_j is not an inner edge of any U_i whenever $U_j \simeq \eta$ is a stick.*

Proof. We first show (i) by induction on the number of subtrees U_i . The base case $\{U_i\} = \{U\}$ is immediate, setting $T = \text{lr}(U)$. Otherwise, U must not be a corolla and letting e be an edge that is both an inner edge of U and a root of some U_i , and one can form a grafting pushout diagram

$$\begin{array}{ccc} \eta & \xrightarrow{e} & U^{\leq e} \\ e \downarrow & & \downarrow \\ U_{\not\leq e} & \longrightarrow & U \end{array} \quad (3.56)$$

DECOMPPROOF EQ

where $U^{\leq e}$ (resp. $U_{\not\leq e}$) are the outer faces consisting of the edges $u \leq_d e$ (resp. $u \not\leq_d e$). Since there is a unique U_i containing the vertex $e^\dagger \leq e$, it follows from the definition of outer face that there is a nontrivial partition $\{U_i\} = \{U_i | U_i \hookrightarrow V\} \sqcup \{U_i | U_i \hookrightarrow W\}$. Existence of $T \rightarrow U$ now follows from the induction hypothesis. For uniqueness, the condition that no U_i is a stick guarantees that T possesses a single inner edge mapping to e , and thus admits a compatible decomposition as in (3.56), so that uniqueness too follows from the induction hypothesis.

For (ii), we argue existence by nested induction on the number of vertices $|V(U)|$ and the sum of the multiplicities m_e . The base case $|V(U)| = 0$, i.e., $U = \eta$ is immediate. Otherwise, writing $m_e = m'_e + 1$, one can form a decomposition (3.56) where either $|V(V)|, |V(W)| < |V(U)|$ or one of V, W is η , so that $T \rightarrow U$ can be built via the induction hypothesis. For uniqueness, note first that by [Fe17, Lemma 5.33] each pre-image $\rho^{-1}(e)$ is linearly ordered and by the “further” claim in [Fe17, Cor. 5.39] the remaining broad relations are precisely the pre-image of the non-identity relations in U , showing that underlying broad poset of the tree T is unique up to isomorphism. Strict uniqueness is then Proposition 3.28.

(iii) follows by combining (i) and (ii). Indeed, any planar tall map $T \rightarrow U$ has a unique decomposition $T \twoheadrightarrow \bar{T} \hookrightarrow U$ as a planar degeneracy followed by a planar inner face, and each of these maps is classified by the data in (b) and (a). \square

Lemma 3.57. *Suppose $T_1, T_2 \hookrightarrow T$ are two outer faces with at least one common edge e . Then there exists a unique outer face $T_1 \cup T_2$ such that $V(T_1 \cup T_2) = V(T_1) \cup V(T_2)$.*

Proof. If either T is a corolla or one of T_1, T_2 consists only of the root or a leaf stick subtrees the result is obvious. Otherwise, one can necessarily choose e to be an inner edge of T , in which case all of three of T_1, T_2, T admit compatible decompositions as in (3.56) and the result follows by induction on $|V(T)|$. \square

3.3 Equivariant leaf-root and vertex functors

This section introduces two functors that are the very center of our definition of the category \mathcal{Op}_G of genuine equivariant operads: the leaf-root and vertex functors.

We start by recalling a key class of maps of G -trees.

Definition 3.58. Let $S = (S_y)_{y \in Y}$ and $T = (T_x)_{x \in X}$ be G -trees. A map of G -trees

$$\varphi = (\phi, (\varphi_y)): S \rightarrow T$$

is called a *quotient* if each of the constituent tree maps

$$\varphi_y: S_y \rightarrow T_{\phi(y)}$$

is an isomorphism of trees.

The category of G -trees and quotients is denoted Ω_G^0 (this notation is justified in §3.4).

Remark 3.59. Quotients can alternatively be described as the cartesian arrows for the Grothendieck fibration $\Omega_G \xrightarrow{t} \mathcal{O}_G$. We note that this differs from the notion of root pullbacks, which are the *chosen* cartesian arrows, and include only those quotients such that each $\varphi_y: S_y \rightarrow T_{\phi(y)}$ is a planar isomorphism, i.e., an identity.

Definition 3.60. The *G -symmetric category*, whose objects we call *G -corollas*, is the full subcategory $\Sigma_G \hookrightarrow \Omega_G^0$ of those G -trees $C = (C_x)_{x \in X}$ such that some (and thus all) C_x is a corolla $C_x \in \Sigma \hookrightarrow \Omega$ (cf. Notation 3.39).


Definition 3.61. The *leaf-root functor* is the functor $\Omega_G^0 \xrightarrow{\text{lr}} \Sigma_G$ defined by

$$\text{lr}((T_x)_{x \in X}) = (\text{lr}(T_x))_{x \in X}.$$

Remark 3.62. The leaf-root functor extends to a functor $\text{lr}: \Omega_G^t \rightarrow \Sigma_G$, where Ω_G^t is the category of tall maps, defined exactly as in Definition 3.58, but not to a functor defined on all arrows in Ω_G . However, we will mostly be concerned with the restriction $\Omega_G^0 \xrightarrow{\text{lr}} \Sigma_G$.

Remark 3.63. Generalizing the remark in Notation 3.39, $\text{lr}(T)$ can alternatively be characterized as being the *unique* G -corolla which admits an also unique tall planar map $\text{lr}(T) \rightarrow T$.

Moreover, $\text{lr}(T)$ can usually be regarded as the “smallest inner face” of T , obtained by removing all the inner edges, although this characterization fails when $T = (\eta_x)_{x \in X}$ is a stick G -tree. Some examples with $G = \mathbb{Z}/4$ follow.




Remark 3.64. Since planarizations can not be pushed forward along tree maps (cf. Remark 3.29) the leaf-root functor $\text{lr}: \Omega_G^0 \rightarrow \Sigma_G$ is not a Grothendieck fibration, but instead only a map of Grothendieck fibrations over \mathbf{O}_G (for the obvious root functor $r: \Sigma_G \rightarrow \mathbf{O}_G$).

Definition 3.65. Given $T = (T_x)_{x \in X} \in \Omega_G$ we define its set of *vertices* to be $V(T) = \coprod_{x \in X} V(T_x)$ and its set of *G-vertices* to be the orbit set $V(T)/G$.

Furthermore, we will regard $V(T)$ as an object of \mathbf{F} by using the induced planar order (with $e^\dagger \leq e$ ordered according to e) and likewise $V_G(T)$ will be regarded as an object of \mathbf{F} by using the lexicographic order: i.e. vertex equivalence classes $[e^\dagger \leq e]$ are ordered according to the planar order \leq_p of the smallest representative $ge, g \in G$.

Remark 3.66. Following Remark 3.52, a tall map $\varphi: T \rightarrow U$ of G -trees induces a G -equivariant map $\varphi^*: V(U) \rightarrow V(T)$ and thus also a map of orbits $\varphi^*: V_G(U) \rightarrow V_G(T)$. We note, however, that φ^* is not in general compatible with the order on $V_G(-)$ even if φ is planar, as is indeed the case even in the non-equivariant setting.

A minimal example follows.



In $V(T)$ the vertices are ordered as $a < c < d$ while in $V(U)$ they are ordered as $a < e < c < d$ but the map $\varphi^*: V(U) \rightarrow V(T)$ is given by $a \mapsto a, c \mapsto c, d \mapsto d, e \mapsto d$.

Notation 3.67. Given $T = (T_x)_{x \in X} \in \Omega_G$ and $(e^\dagger \leq e) \in V(T)$ we write $T_{e^\dagger \leq e}$ as a shorthand for $T_{x, e^\dagger \leq e}$, where $e \in T_x$.

Further, each element $V_G(T)$ corresponds to an unique edge orbit Ge for e not a leaf. We will prefer to write G -vertices as v_{Ge} , and write

$$T_{v_{Ge}} = (T_{f^\dagger \leq f})_{f \in Ge} \quad (3.68)$$

where Ge inherits the planar order.


We note that $T_{v_{Ge}}$ is always a G -corolla, leading to the following definition.

Definition 3.69. The G -vertex functor is the functor

$$\begin{aligned} \Omega_G^0 &\xrightarrow{V_G} \mathbf{F}_s \wr \Sigma_G \\ T &\longmapsto (T_{v_{Ge}})_{v_{Ge} \in V_G(T)}, \end{aligned} \tag{3.70} \quad \boxed{\text{VFUNCTOR EQ}}$$

where \mathbf{F}_s is the category of finite sets and surjections of Remark [FINSURJ REM 2.18](#).

Remark 3.71. In the non-equivariant case the vertex functor can be defined to land instead in $\Sigma \wr \Sigma$. The need to introduce the $\mathbf{F} \wr (-)$ construction comes from the fact that in general quotient maps do not preserve the number of G -vertices. For a simple example, let $G = \{\pm 1, \pm i, \pm j, \pm k\}$ and consider the pullback map $\varphi: \pi^*S \rightarrow S$ of Example [3.25](#) determined by the assignments $a \mapsto a$, $b \mapsto b$, $c \mapsto c$, $d \mapsto ia$, $e \mapsto ib$, and presented below in orbital notation.



We note that $T = \pi^*S$ has three G -vertices v_{Gc} , v_{Gb} , v_{Ge} while S has only two G -vertices v_{Gc} and v_{Gb} . $V_G(\varphi)$ then maps the two G -corollas $T_{v_{Gb}}$ and $T_{v_{Ge}}$ isomorphically onto $S_{v_{Gb}}$ and the G -corolla $T_{v_{Gc}}$ by a non-isomorphism quotient onto $S_{v_{Gc}}$.

The following elementary statement will play an important auxiliary role.

Lemma 3.72. The G -vertex functor

$$\Omega_G^0 \xrightarrow{V_G} \mathbf{F}_s \wr \Sigma_G$$

sends pullbacks over \mathbf{O}_G (i.e. root pullbacks) to pullbacks over $\mathbf{F}_s \wr \mathbf{O}_G$ (cf. Lemma [FWRGROTH LEM 2.20](#)).


Proof. Note first that an arrow $(\phi, (\varphi_i)): (C_i)_{i \in I} \rightarrow (C'_j)_{j \in J}$ is a pullback for the split fibration $\mathbf{F}_s \wr \Sigma_G \rightarrow \mathbf{F}_s \wr \mathbf{O}_G$ iff each of the constituent arrows $\varphi_i: C_i \rightarrow C'_{\phi(i)}$ are pullbacks for the split fibration $\Sigma_G \rightarrow \mathbf{O}_G$.

The pullback $\psi^*T \xrightarrow{\bar{\psi}} T$ of $T = (T_x)_{x \in X} \in \Omega_{G,0}$ over $\psi: Y \rightarrow X$ has the form $(T_{\psi(y)})_{y \in Y} \rightarrow (T_x)_{x \in X}$ and it now suffices to check that each of the vertex maps $(\psi^*T)_{v_{Ge}} \rightarrow T_{v_{G\bar{\psi}(e)}}$ is itself a pullback. By [\(3.68\)](#), this is the statement that for $f \in Ge$ the induced map

$$(\psi^*T)_{f \leq f} \rightarrow T_{\bar{\psi}(f) \leq \bar{\psi}(f)} \tag{3.73} \quad \boxed{\text{VGPULL EQ}}$$

is an identity (i.e. planar isomorphism), and letting y be such that $f \in T_{\psi(y)}$ one sees that [\(3.73\)](#) is the identity $T_{\psi(y), f \leq f} = T_{x, \bar{\psi}(f) \leq \bar{\psi}(f)}$, where $x = \psi(y)$, finishing the proof. \square

Example 3.74. The following depicts one of the maps [\(3.73\)](#) for the pullback $\tau^*T \rightarrow T$ appearing in Example [3.25](#).



Note that $(\tau^*T)_{v_{Ge}} = \rho^*T_{v_{Gb}}$ for ρ the map $\{e, je, ie, ke\} \rightarrow \{b, jb, ib, kb\}$ defined by $e \mapsto b$ so that, accounting for orders, ρ is the block permutation $\rho = (13)(24)$.

We are now in a position to generalize Definition [3.44](#).

TUTIONDATUMG DEF

Definition 3.75. Let $T \in \Omega_G$ be a G -tree.

A (resp. planar) T -substitution datum is a tuple $(U_{f^\dagger \leq f})_{V(T)}$ of G -trees together with

- (i) associative and unital G -action maps $U_{f^\dagger \leq f} \rightarrow U_{gf^\dagger \leq g f}$;
- (ii) (resp. planar) tall maps $T_{f^\dagger \leq f} \rightarrow U_{f^\dagger \leq f}$ compatible with the G -action maps.

Further, a map of (resp. planar) T -substitution data $(U_{f^\dagger \leq f}) \rightarrow (V_{f^\dagger \leq f})$ is a compatible tuple of (resp. planar) tall maps $(U_{f^\dagger \leq f} \rightarrow V_{f^\dagger \leq f})$.

We denote the category of (resp. planar) T -substitution data by $\mathbf{Sub}(T)$ (resp. $\mathbf{Sub}_p(T)$).

Recall that a map of G -trees is called **rootfull def** *rooted* if it induces an ordered isomorphism on the root orbit (cf. Definition 3.24).

SUBSGREF DEF

Remark 3.76. Writing $U_{v_{Ge}}^r = (U_{f^\dagger \leq f})_{f \in Ge}$ a T -substitution datum can equivalently be encoded by the tuple $(U_{v_{Ge}}^r)_{V_G(T)}$ together with *rooted* tall maps $T_{v_{Ge}} \rightarrow U_{v_{Ge}}^r$. The need to include r (which stands for “rooted”) in the notation is explained by Remark 3.81. **WHYR REM**

Further, the T -substitution datum is planar iff the so are the maps $T_{v_{Ge}} \rightarrow U_{v_{Ge}}^r$.

UBSDATUMCONV REM

Remark 3.77. Writing $T = (T_x)_{x \in X}$ as usual one obtains (non-equivariant) T_x -substitution data $U_{x,(-)}$ for each T_x . We again write $U_{x,(-)}: \mathbf{Sc}(T_x) \rightarrow \Omega$ and note that these are compatible with the G -action in the sense that the obvious diagram

$$\begin{array}{ccc} \mathbf{Sc}(T_x) & \xrightarrow{U_{x,(-)}} & \Omega \\ & \searrow g & \nearrow U_{gx,(-)} \\ & \mathbf{Sc}(T_{gx}) & \end{array} \quad (3.78) \quad \text{EQUIVSCMAP EQ}$$

commutes. Writing $\mathbf{Sc}(T) = \coprod_x \mathbf{Sc}(T_x)$, (3.78) is then equivalent to a functor $G \ltimes \mathbf{Sc}(T) \rightarrow \Omega$, where $G \ltimes \mathbf{Sc}(T)$ is the Grothendieck construction for the G -action (which, explicitly, adds arrows $\eta_a \rightarrow \eta_{ga}$, $T_{e^\dagger \leq e} \rightarrow T_{ge^\dagger \leq ge}$ to $\mathbf{Sc}(T)$ that satisfy obvious compatibilities). **EQUIVSCMAP EQ**

In the following we write $\text{colim}_{\mathbf{Sc}(T)} U_{(-)}$ to mean $(\text{colim}_{\mathbf{Sc}(T_x)} U_{x,(-)})_{x \in X}$ or, in other words, we take the colimit in $\Phi = \mathbf{F}\Omega$ rather than in Ω (as is needed since Ω lacks coproducts).

TAUNDERPLANG COR

Corollary 3.79. Let $T \in \Omega_G$ be a G -tree. There are isomorphisms of categories

$$\begin{array}{ccc} \mathbf{Sub}_p(T) & \xrightarrow{\sim} & T \downarrow \Omega_G^{\text{pt}} \\ (U_{f^\dagger \leq f})_{V(T)} & \longmapsto & (T \rightarrow \text{colim}_{\mathbf{Sc}(T)} U_{(-)}) \end{array} \quad \begin{array}{ccc} \mathbf{Sub}(T) & \xrightarrow{\sim} & T \downarrow \Omega_G^{\text{rt}} \\ (U_{f^\dagger \leq f})_{V(T)} & \longmapsto & (T \rightarrow \text{colim}_{\mathbf{Sc}(T)} U_{(-)}) \end{array} \quad (3.80) \quad \text{SUBDATAUNDERPLANG EQ}$$

where $T \downarrow \Omega_G^{\text{pt}}$ (resp. $T \downarrow \Omega_G^{\text{rt}}$) is the category of planar tall (resp. rooted tall) maps under T .

Proof. This is a direct consequence of the non-equivariant analogues Proposition 3.47 and Corollary 3.50 applied to each individual T_x together with the equivariance analysis in Remark 3.77. **SUBDATAUNDERPLAN COR** **SUBDATUMCONV REM** **SUBDATAUNDERPLAN PROP** \square

WHYR REM

Remark 3.81. Writing $U = \text{colim}_{\mathbf{Sc}(T)} U_{(-)}$, it follows from the non-equivariant results Proposition 3.47 and Corollary 3.50 that each inclusion map $U_{f^\dagger \leq f} \rightarrow U$ is planar, so that there is no conflict with Notation 3.67. **SUBDATAUNDERPLAN PROP** **SUBDATAUNDERPLAN COR** **GVERI NOT**

However, some care is needed concerning the $U_{v_{Ge}}^r$ appearing in the reformulation of substitution data given in Remark 3.76. Letting $\varphi: T \rightarrow U$ be the induced map, one sees that while $U_{v_{Ge}}^r$ and $U_{v_{G\varphi(e)}}$ have the same constituent trees (with the latter defined by Notation 3.67), the roots of $U_{v_{Ge}}^r$ are ordered by Ge while those of $U_{v_{G\varphi(e)}}$ are ordered by $G\varphi(e)$. More succinctly, it is then $U_{v_{Ge}}^r = \varphi_{Ge}^* U_{v_{G\varphi(e)}}$ for $\varphi_{Ge}: Ge \rightarrow G\varphi(e)$ the induced map. **SUBSGREF DEF** **GVERI NOT**

Lastly, we note that such distinctions are unnecessary for planar data, since then the φ_{Ge} are ordered isomorphisms (i.e. identities), so that $U_{v_{Ge}}^r = U_{v_{G\varphi(e)}}$.

PULLCOMP REM

Remark 3.82. The isomorphisms in Corollary 3.79 are compatible with root pullbacks of trees. More concretely, as in the proof of Lemma 3.72 each pullback $\psi: \psi^*T \rightarrow T$ determines pullback maps $\bar{\psi}_{Ge}: (\psi^*T)_{v_{Ge}} \rightarrow T_{v_{G\bar{\psi}(e)}}$, which we now note are pullbacks over the maps $\bar{\psi}_{Ge}: Ge \rightarrow G\bar{\psi}(e)$ in \mathcal{O}_G . The definition of pullback then allows us to uniquely fill any diagram (where we reformulate substitution data as in Remark 3.76)

$$\begin{array}{ccc} (\psi^*T)_{v_{Ge}} & \dashrightarrow & \bar{\psi}_{Ge}^* U_{v_{G\bar{\psi}(e)}}^r \\ \downarrow & & \downarrow \\ T_{v_{G\bar{\psi}(e)}} & \longrightarrow & U_{v_{G\bar{\psi}(e)}}^r \end{array}$$

defining the left vertical functors (with the right functors defined analogously) in the commutative diagrams below.

$$\begin{array}{ccc} \text{Sub}_p(\psi^*T) \xrightarrow{\quad} \psi^*T \downarrow \Omega_G^{\text{pt}} & \text{Sub}(\psi^*T) \xrightarrow{\quad} \psi^*T \downarrow \Omega_G^{\text{rt}} & \\ (\bar{\psi}_{Ge}^*) \uparrow & \uparrow \psi^* & (\bar{\psi}_{Ge}^*) \uparrow \\ \text{Sub}_p(T) \xrightarrow{\quad} T \downarrow \Omega_G^{\text{pt}} & \text{Sub}(T) \xrightarrow{\quad} T \downarrow \Omega_G^{\text{rt}} & \end{array} \quad (3.83)$$

SUBDATAUNDERPLANG2 EQ

3.4 Planar strings

We now use the leaf-root and vertex functors to repackage our substitution results in a format that will be more convenient for our discussion of operads in §4.

Definition 3.84. The category Ω_G^n of *planar n -strings* is the category whose objects are strings

$$T_0 \xrightarrow{\varphi_1} T_1 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_n} T_n \quad (3.85)$$

STRINGOBJ EQ

where $T_i \in \Omega_G$ and the φ_i are tall planar maps, while arrows are commutative diagrams

$$\begin{array}{ccccccc} T_0 & \xrightarrow{\varphi_1} & T_1 & \xrightarrow{\varphi_2} & \dots & \xrightarrow{\varphi_n} & T_n \\ \pi_0 \downarrow & & \pi_1 \downarrow & & & & \pi_n \downarrow \\ T'_0 & \xrightarrow{\varphi'_1} & T'_1 & \xrightarrow{\varphi'_2} & \dots & \xrightarrow{\varphi'_n} & T'_n \end{array} \quad (3.86)$$

PTNARROW EQ

where each π_i is a quotient map.

Notation 3.87. Since compositions of planar tall arrows are planar tall and identity arrows are planar tall it follows that Ω_G^\bullet forms a simplicial object in \mathbf{Cat} , with faces given by composing and degeneracies by inserting identities.

Further setting $\Omega_G^{-1} = \Sigma_G$, the leaf-root functor $\Omega_G^0 \xrightarrow{\text{lr}} \Sigma_G$ makes Ω_G^\bullet into an augmented simplicial object and, furthermore, the maps $s_{-1}: \Omega_G^n \rightarrow \Omega_G^{n+1}$ sending $T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_n$ to $\text{lr}(T_0) \rightarrow T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_n$ equip it with extra degeneracies.

Remark 3.88. The identification $\Omega_G^{-1} = \Sigma_G$ can be understood by noting that a string (3.85) is equivalent to a string

$$T_{-1} \xrightarrow{\varphi_0} T_0 \xrightarrow{\varphi_1} T_1 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_n} T_n \quad (3.89)$$

STRINGOBJALT EQ

where $T_{-1} = \text{lr}(T_0) = \dots = \text{lr}(T_n)$.

Remark 3.90. Since for any planar n -string it is $r(T_i) = r(T_j)$ for any $1 \leq i, j \leq n$, one has a well defined functor $r: \Omega_G^n \rightarrow \mathcal{O}_G$, which is readily seen to be a split Grothendieck fibration. Furthermore, generalizing Remark 3.64, all operators d_i, s_j are maps of split Grothendieck fibrations.

VGDEF NOT

Notation 3.91. We extend the vertex functor to a functor $V_G: \Omega_G^{n+1} \rightarrow \mathbf{F}_s \wr \Omega_G^n$ by

$$V_G(T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_n) = (T_{1, v_{Ge}} \rightarrow \cdots \rightarrow T_{n, v_{Ge}})_{v_{Ge} \in V_G(T_0)} \quad (3.92) \quad \text{VGDEF EQ}$$

where we abuse notation by writing $T_{i, v_{Ge}}$ for $(T_{i, \bar{\varphi}_i(f)}^\dagger \leq \bar{\varphi}_i(f))_{f \in v_{Ge}}$, where $\bar{\varphi}_i = \varphi_i \circ \cdots \circ \varphi_1$.

Alternatively, regarding $T_0 \rightarrow \cdots \rightarrow T_n$ as a string of $n-1$ arrows in $T_0 \downarrow \Omega_G^{\text{pt}}$, the object $V_G(T_0 \rightarrow \cdots \rightarrow T_n)$ can be thought of as the image of the inverse functor in Corollary 3.79, written according to the reformulation in Remark 3.76 (where since we are in the planar case we need not distinguish between $U_{(-)}^l$ and $U_{(-)}$ notation (cf. Remark 3.81)). Note however that from this perspective functoriality needs to be checked separately.

We now obtain a key reinterpretation (and slight strengthening) of Corollary 3.79.

Proposition 3.93. For any $n \geq 0$ the commutative diagram

$$\begin{array}{ccc} \Omega_G^n & \xrightarrow{V_G} & \mathbf{F}_s \wr \Omega_G^{n-1} \\ d_{1, \dots, n} \downarrow & & \downarrow \text{Fid}_{0, \dots, n-1} \\ \Omega_G^0 & \xrightarrow{V_G} & \mathbf{F}_s \wr \Sigma_G \end{array} \quad (3.94) \quad \text{PTPULL EQ}$$

is a pullback diagram in \mathbf{Cat} .

Proof. Let us write $P = \Omega_G^0 \times_{\mathbf{F}_s \wr \Sigma_G} \mathbf{F}_s \wr \Omega_G^{n-1}$ for the pullback, so that our goal is to show that the canonical map $\Omega_G^n \rightarrow P$ is an isomorphism.

That $\Omega_G^n \rightarrow P$ is an isomorphism on objects follows by combining the alternative description of V_G in Notation 3.91 with the planar half of Corollary 3.79 (in fact, this yields isomorphisms of the fibers over Ω_G^0 , but we will not directly use this fact). We will hence write $T_0 \rightarrow \cdots \rightarrow T_n$ to denote an object of P as well.

An arrow in P from $T_0 \rightarrow \cdots \rightarrow T_n$ to $T'_0 \rightarrow \cdots \rightarrow T'_n$ then consists of a quotient $\pi_0: T_0 \rightarrow T'_0$ together with a $V_G(T_0)$ indexed tuple of quotients of strings (where we write $e' = \pi_0(e)$)

$$\begin{array}{ccccccc} T_{0, v_{Ge}} & \rightarrow & T_{1, v_{Ge}} & \rightarrow & \cdots & \rightarrow & T_{n, v_{Ge}} \\ \pi_{0, e} \downarrow & & \pi_{1, e} \downarrow & & & & \downarrow \pi_{n, e} \\ T'_{0, v_{Ge'}} & \rightarrow & T'_{1, v_{Ge'}} & \rightarrow & \cdots & \rightarrow & T'_{n, v_{Ge'}} \end{array} \quad (3.95) \quad \text{PTNARROWLOC EQ}$$

That $\Omega_G^n \rightarrow P$ is injective on arrows is then clear.

For surjectivity, note first that by Lemma 3.72 the composite $P \rightarrow \Omega_G^0 \rightarrow \mathbf{O}_G$ is a split Grothendieck fibration and $P \rightarrow \Omega_G^0$ is a map of split Grothendieck fibrations. Indeed, pullbacks in P can be built explicitly as those arrows such that π_0 and all $\pi_{i, e}$ in (3.95) are pullbacks (alternatively, an abstract argument also works). The alternative description of V_G in Notation 3.91 combined with (3.83) then show that $\Omega_G^n \rightarrow P$ preserves pullbacks, so that injectivity needs only be checked for maps in the fibers over \mathbf{O}_G , i.e. on rooted maps. Tautologically, a map in P is rooted iff $\pi_0: T_0 \rightarrow T'_0$ is. But since a quotient is an isomorphism iff it is so on roots, we further have that a map in P is rooted iff $\pi_0: T_0 \rightarrow T'_0$ is a rooted isomorphism and each $\pi_{i, e}$ in (3.95) is an isomorphism. But now rewriting 3.95 as a tuple of diagrams indexed by $f \in Ge$ one obtains a diagram in $\mathbf{Sub}(T_0)$ of the same shape which, after converted to a diagram in $T_0 \downarrow \Omega_G^{\text{rt}}$ using the rooted half of Corollary 3.79, yields the desired rooted map (3.86) in Ω_G^n lifting the rooted map in P . \square

INDVNG NOT

Notation 3.96. For $0 \leq k \leq n$ we will let

$$V_G^k: \Omega_G^n \rightarrow \mathbf{F}_s \wr \Omega_G^{n-k-1}$$

be inductively defined by $V_G^0 = V_G$ and $V_G^{n+1} = \sigma^0 \circ (\mathbf{F}_s \wr V_G^n) \circ V_G$.

VGN REM

Remark 3.97. When $n = 2$, V_G^2 is thus the composite

$$\Omega_G^2 \xrightarrow{V_G} F_s \wr \Omega_G^1 \xrightarrow{V_G} F_s \wr F_s \wr \Omega_G^0 \xrightarrow{V_G} F_s \wr F_s \wr F_s \wr \Sigma_G \xrightarrow{\sigma^0} F_s \wr F_s \wr \Sigma_G \xrightarrow{\sigma^0} F_s \wr \Sigma_G$$

while for $n = 4$, V_G^1 is the composite

$$\Omega_G^4 \xrightarrow{V_G} F_s \wr \Omega_G^3 \xrightarrow{V_G} F_s \wr F_s \wr \Omega_G^2 \xrightarrow{\sigma^0} F_s \wr \Omega_G^2.$$

In light of Remarks 3.52 and 3.66, $V_G^*(T_0 \rightarrow \cdots \rightarrow T_n)$ is identified with the tuple

$$(T_{k,v_{Ge}} \rightarrow \cdots \rightarrow T_{n,v_{Ge}})_{v_{Ge} \in V_G(T_k)}, \quad (3.98)$$

VGNISO EQ

where we note that strings are written in prepended notation as in (3.89), so that $T_{i,v_{Ge}}$ is superfluous unless $k = n$. Further, note that this requires changing the order of $V_G(T_k)$. Rather than using the order induced by T_k , one instead equips $V_G(T_k)$ with the order induced lexicographically from the maps $V_G(T_k) \rightarrow V_G(T_{k-1}) \rightarrow \cdots \rightarrow V_G(T_0)$ of Remark 3.52. I.e., for $v, w \in V_G(T_k)$ the condition $v < w$ is determined by the lowest l such that the images of $v, w \in V_G(T_l)$ are distinct.

Therefore, for each d_i with $i < k$ there are natural isomorphisms as on the left below which interchange the lexicographical order on the indexing set $V_G(T_k)$ induced by the string $V_G(T_k) \rightarrow V_G(T_{k-1}) \rightarrow \cdots \rightarrow V_G(T_0)$ with the one induced by the string that omits $V_G(T_i)$. For d_i with $i > k$ one has a commutative diagram as on the right below. Note that no such diagram is defined for d_k .

$$\begin{array}{ccc} \Omega_G^n & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-1} \\ d_i \downarrow & \swarrow \pi_i & \parallel \\ \Omega_G^{n-1} & \xrightarrow{V_G^{k-1}} & F_s \wr \Omega_G^{n-k-1} \end{array} \quad \begin{array}{ccc} \Omega_G^n & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-1} \\ d_i \downarrow & & \downarrow d_{i-k-1} \\ \Omega_G^{n-1} & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-2} \end{array} \quad (3.99)$$

PIIDEFDI EQ

Similarly, for s_j with $j < k$ (resp. $j \geq k$) one has commutative diagrams as on the left (resp. right) below. Note that for s_k one uses the extra degeneracy $s_{k-k-1} = s_{-1}$.

$$\begin{array}{ccc} \Omega_G^n & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-1} \\ s_j \downarrow & & \parallel \\ \Omega_G^{n+1} & \xrightarrow{V_G^{k+1}} & F_s \wr \Omega_G^{n-k-1} \end{array} \quad \begin{array}{ccc} \Omega_G^n & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-1} \\ s_j \downarrow & & \downarrow s_{j-k-1} \\ \Omega_G^{n+1} & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k} \end{array} \quad (3.100)$$

PIIDEFDI2 EQ

The functors V_G^k and isomorphisms π_i satisfy a number of useful conditions that we now catalog.

PIIPROP PROP

Proposition 3.101. (a) The composite

$$\Omega_G^n \xrightarrow{V_G^k} F_s \wr \Omega_G^{n-k-1} \xrightarrow{V_G^l} F_s^2 \wr \Omega_G^{n-k-l-2} \xrightarrow{\sigma^0} F_s \wr \Omega_G^{n-k-l-2}$$

equals the functor V_G^{k+l+1} .

- (b) The functors V_G^k send pullbacks for the split Grothendieck fibration $\Omega_G^k \rightarrow \mathcal{O}_G$ to pullbacks for $F_s \wr \Omega_G^{n-k-1} \rightarrow F_s$.
- (c) The isomorphisms $\pi_i(T_0 \rightarrow \cdots \rightarrow T_n)$ are pullbacks for the split Grothendieck fibration $F_s \wr \Omega_G^{n-k-1} \rightarrow F_s$. Moreover, the projection of $\pi_i(T_0 \rightarrow \cdots \rightarrow T_n)$ onto F_s depends only on $T_0 \rightarrow \cdots \rightarrow T_i$.

PIIDEFDI EQ PIIDEFDI2 EQ

- (d) The rightmost diagrams in both (3.99) and (3.100) are pullbacks diagrams in \mathbf{Cat} .

(e) For $i < k$ the composite natural transformation in the diagram below is π_i .

$$\begin{array}{ccccccc}
\Omega_G^n & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-1} & \xrightarrow{F_s \wr V_G^l} & F_s^{i2} \wr \Omega_G^{n-k-l-2} & \xrightarrow{\sigma^0} & F_s \wr \Omega_G^{n-k-l-2} \\
d_i \downarrow & \swarrow \pi_i & \parallel & & \parallel & & \parallel \\
\Omega_G^{n-1} & \xrightarrow{V_G^{k-1}} & F_s \wr \Omega_G^{n-k-1} & \xrightarrow{F_s \wr V_G^l} & F_s^{i2} \wr \Omega_G^{n-k-l-2} & \xrightarrow{\sigma^0} & F_s \wr \Omega_G^{n-k-l-2}
\end{array} \quad (3.102) \quad \boxed{\text{INDPI1 EQ}}$$

For $k < i < k+l+1$ the composite natural transformation in the diagram below is π_{i+1} .

$$\begin{array}{ccccccc}
\Omega_G^n & \xrightarrow{V_G^k} & F_s \wr \Omega_G^{n-k-1} & \xrightarrow{F_s \wr V_G^l} & F_s^{i2} \wr \Omega_G^{n-k-l-2} & \xrightarrow{\sigma^0} & F_s \wr \Omega_G^{n-k-l-2} \\
d_i \downarrow & & F_s \wr d_{i-k-1} \downarrow & \swarrow F_s \wr \pi_i & \parallel & & \parallel \\
\Omega_G^{n-1} & \xrightarrow{V_G^{k-1}} & F_s \wr \Omega_G^{n-k-2} & \xrightarrow{F_s \wr V_G^{l-1}} & F_s^{i2} \wr \Omega_G^{n-k-l-2} & \xrightarrow{\sigma^0} & F_s \wr \Omega_G^{n-k-l-2}
\end{array} \quad (3.103) \quad \boxed{\text{INDPI2 EQ}}$$

(f) Restricting to the case $k = n$, the pairs (d_i, π_i) and $(s_j, id_{V_G^n})$ satisfy all possible simplicial identities (i.e. those with $i \neq n$). Explicitly, for $0 \leq i' < i < n$ the composite natural transformations in the diagrams

$$\begin{array}{ccc}
\Omega_G^n \longrightarrow F_s \wr \Sigma_G & \Omega_G^n \longrightarrow F_s \wr \Sigma_G & \\
d_i \downarrow \swarrow \pi_i \parallel & d_{i'} \downarrow \swarrow \pi_{i'} \parallel & \\
\Omega_G^{n-1} \longrightarrow F_s \wr \Sigma_G & \Omega_G^{n-1} \longrightarrow F_s \wr \Sigma_G & (3.104) \quad \boxed{\text{SIMPPI EQ}} \\
d_{i'} \downarrow \swarrow \pi_{i'} \parallel & d_{i-1} \downarrow \swarrow \pi_{i-1} \parallel & \\
\Omega_G^{n-2} \longrightarrow F_s \wr \Sigma_G & \Omega_G^{n-2} \longrightarrow F_s \wr \Sigma_G &
\end{array}$$

coincide, and similarly for the face-degeneracy relations.

Proof. (a) follows by induction on k , with $k = 0$ being the definition. More generally (and writing F for F_s) one has

$$\begin{aligned}
\sigma^0(F \wr V_G^l) V_G^{k+1} &= \sigma^0(F \wr V_G^l) \sigma^0(F \wr V_G^k) V_G = \sigma^0 \sigma^0(F^{i2} \wr V_G^l)(F \wr V_G^k) V_G \\
&= \sigma^0 \sigma^1(F^{i2} \wr V_G^l)(F \wr V_G^k) V_G = \sigma^0(F \wr \sigma^0)(F^{i2} \wr V_G^l)(F \wr V_G^k) V_G \\
&= \sigma^0(F \wr (\sigma^0(F \wr V_G^l) V_G^k)) V_G = \sigma^0(F \wr V_G^{k+l-1}) V_G = V_G^{k+l}.
\end{aligned}$$

(b) generalizes Lemma [B.72](#), and follows by induction using that result, Lemma [B.20](#), [VGPULL LEM](#) [FWRGROTH LEM](#)

and the obvious claim that $F \wr F \wr A \xrightarrow{\sigma^0} F \wr A$ sends pullbacks over $F \wr F$ to pullbacks over F .

(c) is clear. Also, (e) and (f) are easy consequences of (b) and (c): since all natural transformations involved consist of pullbacks, one needs only check each claim after forgetting to the F_s coordinate, which is straightforward.

Lastly, (d) is argued by induction on k and n . The case $k = 0$ for the rightmost diagram in (3.99) follows by the diagram on the left below, combined with Proposition [3.93](#) applied to the bottom and total squares. The general case then follows from the right diagram, with the left square being in the case $k = 0$, the middle square being a pullback by induction (and since $F \wr (-)$ preserves pullback squares), and the rightmost square by direct verification.

$$\begin{array}{ccccc}
\Omega_G^n & \xrightarrow{V_G} & F_s \wr \Omega_G^{n-1} & \xrightarrow{V_G^k} & F_s^{i2} \wr \Omega_G^{n-k-2} & \xrightarrow{\sigma^0} & F_s \wr \Omega_G^{n-k-2} \\
d_i \downarrow & & d_{i-1} \downarrow & & F_s \wr d_{i-1} \downarrow & & F_s \wr d_{i-1} \downarrow \\
\Omega_G^{n-1} & \xrightarrow{V_G} & F_s \wr \Omega_G^{n-2} & \xrightarrow{V_G^k} & F_s^{i2} \wr \Omega_G^{n-k-3} & \xrightarrow{\sigma^0} & F_s \wr \Omega_G^{n-k-3} \\
d_{1,\dots,n} \downarrow & & d_{0,\dots,n-1} \downarrow & & & & \\
\Omega_G^0 & \xrightarrow{V_G} & F_s \wr \Sigma_G & & & &
\end{array} \quad (3.105)$$

The claim for the rightmost diagram in (3.100) follows by the analogous diagrams with the d_i (but not $d_{1,\dots,n}, d_{0,\dots,n-1}$) replaced by s_j . \square

4 Genuine equivariant operads

In this section we now build the category $\mathbf{Op}_G(\mathcal{V})$ of genuine equivariant operads. We will do so by building a monad \mathbb{F}_G on the category $\mathbf{Sym}_G(\mathcal{V}) = \mathbf{Fun}(\Sigma_G^{op}, \mathcal{V})$, that we refer to as the category of G -symmetric sequences on \mathcal{V} . The underlying endofunctor of \mathbb{F}_G is easy enough to describe. Given $X \in \mathbf{Sym}_G(\mathcal{V})$, $\mathbb{F}_G X$ is given by the left Kan extension diagram

$$\begin{array}{ccccc} (\Omega_G^0)^{op} & \xrightarrow{V_G^{op}} & (F \wr \Sigma_G)^{op} & \xrightarrow{(F \wr X)^{op}} & (F \wr \mathcal{V}^{op})^{op} & \xrightarrow{\Pi} & \mathcal{V} \\ \downarrow \text{lr} & \swarrow & & \searrow \mathbb{F}_G X & & & \\ \Sigma_G^{op} & & & & & & \end{array} \quad (4.1) \quad \text{FGXDEF EQ}$$

To describe the monad structure on \mathbb{F}_G , however, we will find it preferable to separate the left Kan extension step from the remaining construction. As such, we will first build a monad N on a larger category $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V})$ which we then transfer via the (\mathbf{Lan}, ι) adjunction in Remark 4.6.

4.1 A monad on spans

Definition 4.2. We will write $\mathbf{WSpan}^l(\mathcal{C}, \mathcal{D})$ (resp. $\mathbf{WSpan}^r(\mathcal{C}, \mathcal{D})$), which we call the category of *left weak spans* (resp. *right weak spans*), to denote the category with objects the spans

$$\mathcal{C} \xleftarrow{k} A \xrightarrow{X} \mathcal{D},$$

arrows the diagrams as on the left (resp. right) below

$$\begin{array}{ccc} & A_1 & \\ k_1 \swarrow & & \searrow X_1 \\ \mathcal{C} & & \mathcal{D} \\ k_2 \swarrow & & \searrow X_2 \\ & A_2 & \end{array} \quad \begin{array}{ccc} & A_1 & \\ k_1 \swarrow & & \searrow X_1 \\ \mathcal{C} & & \mathcal{D} \\ k_2 \swarrow & & \searrow X_2 \\ & A_2 & \end{array} \quad (4.3) \quad \text{TWISTEDARROWRIGHT EQ}$$

which we write as $(i, \varphi): (k_1, X_1) \rightarrow (k_2, X_2)$, and composition given in the obvious way.

Remark 4.4. There are canonical natural isomorphisms

$$\mathbf{WSpan}^r(\mathcal{C}, \mathcal{D}) \simeq \mathbf{WSpan}^l(\mathcal{C}^{op}, \mathcal{D}^{op}). \quad (4.5) \quad \text{LRSPANISO EQ}$$

Remark 4.6. The terms *left/right* are motivated by the existence of adjunctions (which are seen to be equivalent by using (4.5))

$$\mathbf{Lan}: \mathbf{WSpan}^l(\mathcal{C}, \mathcal{D}) \rightleftarrows \mathbf{Fun}(\mathcal{C}, \mathcal{D}): \iota$$

$$\iota: \mathbf{Fun}(\mathcal{C}, \mathcal{D}) \rightleftarrows \mathbf{WSpan}^r(\mathcal{C}, \mathcal{D})^{op}: \mathbf{Ran}$$

where the functors ι denote the obvious inclusions (note the need for the $(-)^{op}$ in the second adjunction) and $\mathbf{Lan}/\mathbf{Ran}$ denote the left/right Kan extension functors.

We will mainly be interested in the span categories $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V}) \simeq \mathbf{WSpan}^r(\Sigma_G, \mathcal{V}^{op})$.

OMEGAGNA NOT

Notation 4.7. Given a functor $\rho: A \rightarrow \Sigma_G$, $n \geq 0$, we let $\Omega_G^n \wr A$ denote the pullback in **Cat**

$$\begin{array}{ccc} \Omega_G^n \wr A & \xrightarrow{V_G^n} & F_s \wr A \\ \downarrow & & \downarrow \\ \Omega_G^n & \xrightarrow{V_G^n} & F_s \wr \Sigma_G \end{array} \quad (4.8)$$

OMGGNA

We will write the top V_G^n functor as $V_G^n \wr A$ whenever we need to distinguish such functors. Explicitly, by Remark 3.97 the objects of $\Omega_G^n \wr A$ are pairs

$$(T_0 \rightarrow \cdots \rightarrow T_n, (a_{v_{Ge}})_{v_{Ge} \in V_G(T_n)}) \quad (4.9)$$

OMEGAGNA EQ

such that $\rho(a_{v_{Ge}}) = T_{n,v_{Ge}}$, and where $V_G(T_n)$ is ordered lexicographically according to the string $T_0 \rightarrow \cdots \rightarrow T_n$.

Remark 4.10. Generalizing the notation $\Omega_G^{-1} = \Sigma_G$, we will also write $\Omega_G^{-1} \wr A = A$, in which case $V_G^{-1} \wr A: \Omega_G^{-1} \wr A \rightarrow F_s \wr A$ is the obvious “simpleton map” $\delta^0: A \rightarrow F_s \wr A$.

Remark 4.11. An alternative, and arguably more suggestive, notation for $\Omega_G^n \wr A$ would be $\Omega_G^n \wr_{\Sigma_G} A$, since we are really defining a “relative” analogue of the wreath product (so that in particular $\Omega_G^n \wr_{\Sigma_G} \Sigma_G \simeq \Omega_G^n$). However, we will prefer $\Omega_G^n \wr A$ due to space concerns.

Remark 4.12. The definition of $\Omega_G^n \wr A$ in (4.8) is unchanged by replacing F_s with F . As such, to avoid cluttering the diagrams in this section we will from now on abuse notation by writing simply F instead of F_s .

Expand on if necessary

Our primary interest here will be in the $\Omega_G^0 \wr (-)$ construction, which can be iterated thanks to the existence of the composite maps $\Omega_G^0 \wr A \rightarrow \Omega_G^0 \rightarrow \Sigma_G$. The role of the higher strings $\Omega_G^n \wr A$ will then be to provide more convenient models for iterated $\Omega_G^0 \wr (-)$ constructions. Indeed, Proposition 3.93 can be reinterpreted as providing a canonical identification $\Omega_G^0 \wr \Omega_G^n \simeq \Omega_G^{n+1}$ with the functor $V_G^0 \wr \Omega_G^n$ identified with the functor V_G as defined in Notation 3.91. Moreover, arguing by induction on n , the fact that the rightmost squares in (3.99) are pullbacks (Proposition 3.101) provides further identifications $\Omega_G^k \wr \Omega_G^n \simeq \Omega_G^{n+k+1}$ with $V_G^k \wr \Omega_G^n$ identified with V_G^k as defined by Notation 3.96.

Our first task is now to produce analogous identifications between $\Omega_G^k \wr \Omega_G^n \wr A = \Omega_G^k \wr (\Omega_G^n \wr A)$ and $\Omega_G^{n+k+1} \wr A$ (note that iterated wreath expressions should always be read as bracketed on the right, i.e. we do *not* define the expression $(\Omega_G^k \wr \Omega_G^n) \wr A$). We start by generalizing the key functors from §3.4.

Proposition 4.13. *There are functors*

$$\Omega_G^n \wr A \xrightarrow{V_G^k} F_s \wr \Omega_G^{n-k-1} \wr A \quad \Omega_G^n \wr A \xrightarrow{d_i} \Omega_G^{n-1} \wr A \quad \Omega_G^n \wr A \xrightarrow{s_j} \Omega_G^{n+1} \wr A$$

where $i < n$, and natural isomorphisms

$$\pi_i: V_G^k \Rightarrow V_G^{k-1} \circ d_i$$

for $i < k$. Further, all of these are natural in A and they satisfy all the analogues of the properties listed in Proposition 3.101.

Proof. While not hard to explicitly write formulas for V_G^k , d_i , s_j , π_i (which we list in Remark 4.16), and then verify the desired properties, we here instead argue that the desired properties themselves can be used to uniquely, and coherently, define those functors.

Firstly, the functors V_G are defined from the following diagram

$$\begin{array}{ccccccc} \Omega_G^{n+1} \wr A & \xrightarrow{V_G} & F \wr \Omega_G^n \wr A & \xrightarrow{F \wr V_G^n} & F^{\wr 2} \wr A & \xrightarrow{\sigma^0} & F \wr A \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_G^{n+1} & \xrightarrow{V_G} & F \wr \Omega_G^n & \xrightarrow{F \wr V_G^n} & F^{\wr 2} \wr \Sigma_G & \xrightarrow{\sigma^0} & F \wr \Sigma_G \end{array} \quad (4.14)$$

ALLSQUARES2 EQ

by noting that the center and right squares are pullbacks, and choosing V_G to be the unique functor such that the top composite is V_G^{n+1} . The higher functors V_G^k are defined exactly as in (3.92), and the analogue of Proposition 3.101(a) follows by the same proof.

The analogue of Proposition 3.101(b) is tautological, as pullback arrows for $\Omega_G^n \wr A \rightarrow \mathcal{O}_G$ are defined as compatible pairs of pullbacks in Ω_G^n and $\mathbf{F} \wr A$.

To define d_i we consider the diagram below (for some $i < k$).

$$\begin{array}{ccccc}
 \Omega_G^n \wr A & \xrightarrow{V_G^k} & \mathbf{F} \wr \Omega_G^{n-k-1} \wr A & & \\
 \downarrow & \searrow d_i & \swarrow \pi_i & \downarrow & \\
 \Omega_G^n & & \Omega_G^{n-1} \wr A & \xrightarrow{V_G^{k-1}} & \mathbf{F} \wr \Omega_G^{n-k-1} \\
 \downarrow & \searrow d_i & \swarrow \pi_i & \downarrow & \\
 \Omega_G^{n-1} & & \Omega_G^{n-1} & \xrightarrow{V_G^{k-1}} & \mathbf{F} \wr \Omega_G^{n-k-1}
 \end{array} \quad (4.15) \quad \boxed{\text{PICUBOIDAB EQ}}$$

The desiderata that the top π_i consist of pullback arrows lifting the lower π_i implies that it is uniquely defined by the top V_G^k functor, and hence so is the top composite $V_G^{k-1} d_i$. But since the front face is a pullback square (by arguing by induction on k), there is a unique choice for d_i . The fact that this definition of $d_i \wr A$ is not dependent on k is ensured by natural transformation in (3.102) is π_i . Similarly, that the analogues of the left diagrams in (3.100) hold follows by an identical argument from the fact that the composites of (3.103) are π_{i+1} .

The definitions of the s_j are similar, except easier since there are no π_i to contend with.

The analogue of Proposition 3.101(c),(e),(f) are then tautological, and the analogue of Proposition 3.101(d) follows by an identical argument. \square

VGDEFA REM

Remark 4.16. Explicitly, $V_G^k: \Omega_G^n \wr A \rightarrow \mathbf{F} \wr \Omega_G^{n-k-1} \wr A$ is defined by sending (4.9) to

$$\left(\left(T_{k,v_{Gf}} \rightarrow \cdots \rightarrow T_{n,v_{Gf}}, (a_{v_{Ge}})_{v_{Ge} \in V_G(T_{n,v_{Gf}})} \right) \right)_{v_{Gf} \in V_G(T_k)} \quad (4.17) \quad \boxed{\text{VGDEFA EQ}}$$

where both $V_G(T_k)$ and $T_{n,v_{Gf}}$ are ordered lexicographically according to the obvious strings.

Similarly, functors $d_i: \Omega_G^n \wr A \rightarrow \Omega_G^{n-1} \wr A$ for $0 \leq i < n$ and $s_j: \Omega_G^n \wr A \rightarrow \Omega_G^{n+1} \wr A$ for $-1 \leq j \leq n$ are defined on (4.9) by performing the corresponding operation on the $T_0 \rightarrow \cdots \rightarrow T_n$ coordinate and, in the d_i case suitably reordering $V_G(T_n)$.

Remark 4.18. One upshot of Proposition 4.13 is that formally applying the symbol $(-)\wr A$ to the diagrams in Proposition 3.101 yields sensible statements. As such, we will simply refer to the corresponding part of Proposition 3.101 when using one of the generalized claims.

IDEN COR

Corollary 4.19. One has identifications $\Omega_G^k \wr \Omega_G^n \wr A \simeq \Omega_G^{n+k+1} \wr A$ which identify $V_G^k \wr \Omega_G^n \wr A$ with $V_G^k \wr A$. Further, these are associative in the sense that the identifications

$$\Omega_G^k \wr \Omega_G^l \wr \Omega_G^n \wr A \simeq \Omega_G^{k+l+1} \wr \Omega_G^n \wr A \simeq \Omega_G^{k+l+n+2} \wr A$$

$$\Omega_G^k \wr \Omega_G^l \wr \Omega_G^n \wr A \simeq \Omega_G^k \wr \Omega_G^{l+n+1} \wr A \simeq \Omega_G^{k+l+n+2} \wr A$$

coincide. Lastly, one obtains identifications

$$d_i \wr \Omega_G^n \simeq d_i \quad \pi_i \wr \Omega_G^n \simeq \pi_i \quad s_j \wr \Omega_G^n \simeq s_j \quad \Omega_G^k \wr d_i \simeq d_{i+k+1} \quad \Omega_G^k \wr \pi_i \simeq \pi_{i+k+1} \quad \Omega_G^k \wr s_j \simeq s_{j+k+1}$$

Proof. The identification $\Omega_G^k \wr \Omega_G^n \wr A \simeq \Omega_G^{n+k+1} \wr A$ follows since by Proposition 3.101(a) both

expressions compute the limit of the solid part of the diagram below.

$$\begin{array}{ccccccc}
\bullet & \xrightarrow{\quad\quad\quad} & \bullet & \xrightarrow{\quad\quad\quad} & F^{i2} \wr A & \xrightarrow{\sigma^0} & F \wr A \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Omega_G^{n+k+1} & \xrightarrow{V_G^k} & F \wr \Omega_G^n & \xrightarrow{F \wr V_G^n} & F^{i2} \wr \Sigma_G & \xrightarrow{\sigma^0} & F \wr \Sigma_G \\
\downarrow & & \downarrow & & & & \\
\Omega_G^k & \xrightarrow{V_G^k} & F \wr \Sigma_G & & & &
\end{array}$$

Associativity follows similarly. The remaining identifications are obvious. \square

We now have all the necessary ingredients to define our monad on spans.

Definition 4.20. Suppose \mathcal{V} has finite products or, more generally, that it is a symmetric monoidal category with diagonals in the sense of Remark 2.18.

We define an endofunctor N of $\mathbf{Wspan}^r(\Sigma_G, \mathcal{V}^{op})$ by letting $N(\Sigma_G \leftarrow A \rightarrow \mathcal{V}^{op})$ be the span $\Sigma_G \leftarrow \Omega_G^0 \wr A \rightarrow \mathcal{V}^{op}$ given composition along the diagram

$$\begin{array}{ccccc}
\Omega_G^0 \wr A & \xrightarrow{V_G} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \xrightarrow{\Pi^{op}} \mathcal{V}^{op} \\
\downarrow & & \downarrow & & \\
\Omega_G^0 & \xrightarrow{V_G} & F \wr \Sigma_G & & \\
\downarrow & & & & \\
\Sigma_G & & & &
\end{array}$$

and defined on maps of spans in the obvious way.

One has a multiplication $\mu: N \circ N \Rightarrow N$ given by the natural isomorphism

$$\begin{array}{ccccccc}
\Sigma_G \longleftarrow \Omega_G^1 \wr A & \xrightarrow{V_G} & F \wr \Omega_G^0 \wr A & \xrightarrow{F \wr V_G} & F^{i2} \wr A & \longrightarrow & F^{i2} \wr \mathcal{V}^{op} \xrightarrow{\Pi^{op}} F \wr \mathcal{V}^{op} \xrightarrow{\Pi^{op}} \mathcal{V}^{op} \\
\parallel & \searrow d_0 & \nearrow \pi_0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & \nearrow \alpha & \parallel \\
\Sigma_G \longleftarrow \Omega_G^0 \wr A & \xrightarrow{V_G} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\Pi^{op}} & \mathcal{V}^{op}
\end{array} \quad (4.21)$$

where we note that the top right composite in the π_0 square is indeed V_G^1 using the inductive description in (the $(-) \wr A$ analogue of) Notation 3.96.

Lastly, there is a unit $\eta: id \Rightarrow N$ given by the strictly commutative diagrams (where to see that the second square commutes we recall that $V_G^{-1} = \delta^0$)

$$\begin{array}{ccccccc}
\Sigma_G \longleftarrow A & \xlongequal{\quad} & A & \longrightarrow & \mathcal{V}^{op} & \xlongequal{\quad} & \mathcal{V}^{op} \\
\parallel & \searrow s_{-1} & \downarrow \delta^0 & & \downarrow \delta^0 & & \parallel \\
\Sigma_G \longleftarrow \Omega_G^0 \wr A & \xrightarrow{V_G} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\Pi^{op}} & \mathcal{V}^{op}.
\end{array} \quad (4.22)$$

Proposition 4.23. (N, μ, η) form a monad on $\mathbf{Wspan}^r(\Sigma_G, \mathcal{V}^{op})$.

Proof. The natural transformation component of $\mu \circ (N\mu)$ is given by the composite diagram

$$\begin{array}{ccccccccccc}
\Omega_G^2 \wr A & \rightarrow & F \wr \Omega_G^1 \wr A & \rightarrow & F^{i2} \wr \Omega_G^0 \wr A & \rightarrow & F^{i3} \wr A & \rightarrow & F^{i3} \wr \mathcal{V}^{op} & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
d_1 \downarrow & & F \wr d_0 \downarrow & & \nearrow F \wr \pi_0 & & \downarrow \sigma^1 & & \downarrow \sigma^1 & & \nearrow F \wr \alpha & & \parallel & & \parallel \\
\Omega_G^1 \wr A & \rightarrow & F \wr \Omega_G^0 \wr A & \rightarrow & F^{i2} \wr A & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
d_0 \downarrow & & \nearrow \pi_0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel & & \parallel \\
\Omega_G^0 \wr A & \rightarrow & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op}
\end{array} \quad (4.24)$$

whereas the natural transformation component of $\mu \circ (\mu N)$ is given by

$$\begin{array}{ccccccccccc}
 \Omega_G^2 \wr A & \rightarrow & F \wr \Omega_G^1 \wr A & \rightarrow & F^2 \wr \Omega_G^0 \wr A & \rightarrow & F^3 \wr A & \rightarrow & F^3 \wr \mathcal{V}^{op} & \rightarrow & F^2 \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0 \downarrow & \nearrow \pi_0 & & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & & \downarrow \sigma^0 & \nearrow \alpha & & & & & \\
 \Omega_G^1 \wr A & \xrightarrow{\quad} & F \wr \Omega_G^0 \wr A & \rightarrow & F^2 \wr A & \rightarrow & F^2 \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & & & \\
 d_0 \downarrow & \nearrow \pi_0 & & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & & \downarrow \sigma^0 & \nearrow \alpha & & & & & \\
 \Omega_G^0 \wr A & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & & & & & & &
 \end{array}
 \tag{4.25}$$

ASSOCSPAN2 EQ

That the rightmost sections of (4.24) and (4.25) coincide follows from the associativity of the isomorphisms α in (2.15). On the other hand, the leftmost sections coincide since they are instances of the “simplicial relation” diagrams in (3.104), as is seen by using (3.102) and (3.103) to reinterpret the top left sections.

As for unit conditions, $\mu \circ (N\eta)$ is represented by

$$\begin{array}{ccccccccccc}
 \Omega_G^0 \wr A & \xrightarrow{\quad} & F \wr A & = & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & = & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 s_0 \downarrow & & s_{-1} \downarrow & & \downarrow \delta^1 & & \downarrow \delta^1 & & \parallel & & \parallel \\
 \Omega_G^1 \wr A & \rightarrow & F \wr \Omega_G^0 \wr A & \rightarrow & F^2 \wr A & \rightarrow & F^2 \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0 \downarrow & \nearrow \pi_0 & & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel \\
 \Omega_G^0 \wr A & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & & &
 \end{array}
 \tag{4.26}$$

UNITSPAN1 EQ

while $\mu \circ (\eta N)$ is represented by

$$\begin{array}{ccccccccccc}
 \Omega_G^0 \wr A & = & \Omega_G^0 \wr A & \rightarrow & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} & = & \mathcal{V}^{op} \\
 s_{-1} \downarrow & & \downarrow \delta^0 & & \downarrow \delta^0 & & \downarrow \delta^0 & & \downarrow \delta^0 & & \parallel \\
 \Omega_G^1 \wr A & \rightarrow & F \wr \Omega_G^0 \wr A & \rightarrow & F^2 \wr A & \rightarrow & F^2 \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0 \downarrow & \nearrow \pi_0 & & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel \\
 \Omega_G^0 \wr A & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & & &
 \end{array}
 \tag{4.27}$$

UNITSPAN2 EQ

That (4.26) and (4.27) coincide follows analogously by the unital condition for α and the face degeneracy relations in Proposition 3.101(f). \square

4.2 The genuine equivariant operad monad

Since $\mathbf{Wspan}^r(\Sigma_G, \mathcal{V}^{op}) \simeq \mathbf{Wspan}^l(\Sigma_G^{op}, \mathcal{V})$, Proposition 4.23 and Remark 4.6 give an adjunction

$$\mathbf{Lan} : \mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V}) \rightleftarrows \mathbf{Fun}(\Sigma_G^{op}, \mathcal{V}) : \iota
 \tag{4.28}$$

LANIOTAADJ EQ

together with a monad N in the leftmost category $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V})$.

We will now show that under reasonable conditions on \mathcal{V} this monad can be transferred by using Proposition 2.26, i.e. we will show that the natural transformations $\mathbf{Lan} \circ N \Rightarrow \mathbf{Lan} \circ N \circ \iota \circ \mathbf{Lan}$ and $\mathbf{Lan} \circ \iota \Rightarrow id$ are isomorphisms.

This will require us to introduce a slight modification of the category of spans. For motivation, note that iterations $N^{on+1} \circ \iota$ produce spans of the form $\Sigma_G \leftarrow \Omega_G^n \rightarrow \mathcal{V}^{op}$ (where we use the identification $\Omega_G^n \wr \Sigma_G \simeq \Omega_G^n$). As noted in Remark 3.90, the maps $\Omega_G^n \rightarrow \Sigma_G$ are maps of split fibrations over \mathbf{O}_G , as are all other simplicial operators d_i, s_j .

Definition 4.29. The category $\mathbf{Wspan}_r^l(\Sigma_G^{op}, \mathcal{V})$ of *rooted (left) spans* has as objects spans $\Sigma_G^{op} \leftarrow A^{op} \rightarrow \mathcal{V}$ together with a split Grothendieck fibration $r : A \rightarrow \mathbf{O}_G$ such that $A \rightarrow \Sigma_G$ is a map of split fibrations.

Similarly, arrows are maps of spans that induce maps of split fibrations.

We refer split fibrations $A \rightarrow \mathbf{O}_G$ as *root fibrations* and to maps between them as *root fibration maps*.

Remark 4.30. The condition that $A \rightarrow \mathbf{O}_G$ be a root fibration requires additional *choices* of root pullbacks. Therefore, the forgetful functor $\mathbf{Wspan}_i^l(\Sigma_G^{op}, \mathcal{V}) \rightarrow \mathbf{Wspan}^l(\Sigma_G^{op}, \mathcal{V})$ is not quite injective on objects.

The relevance of rooted spans is given by the following couple of lemmas.

Lemma 4.31. *If $A \rightarrow \Sigma_G$ is a root fibration map then so is $\Omega_G^0 \wr A \rightarrow \Omega_G^0$, naturally in A .*

Proof. The hypothesis that $A \rightarrow \Sigma_G$ is root fibration map implies that the rightmost map below in is a map of split fibrations over $\mathbf{F} \wr \mathbf{O}_G$.

$$\begin{array}{ccc} \Omega_G^0 \wr A & \xrightarrow{V_G} & \mathbf{F} \wr A \\ \downarrow & & \downarrow \\ \Omega_G^0 & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G \end{array} \quad (4.32)$$

Since by Lemma 3.72 the map V_G sends pullback arrows in Ω_G^0 (over \mathbf{O}_G) to pullback arrows of $\mathbf{F} \wr \Sigma_G$ (over $\mathbf{F} \wr \mathbf{O}_G$), the root pullback arrows in $\Omega_G^0 \wr A$ can be defined as compatible pairs of pullback arrows in Ω_G^0 and $\mathbf{F} \wr A$, and the result follows. \square

Remark 4.33. Explicitly, if $\psi: Y \rightarrow X$ is a map in \mathbf{O}_G , and $\tilde{T} = (T, (A_{v_{Ge}})_{v_{Ge}(T)}) \in \Omega_G^0 \wr A$, the pullback $\psi^* \tilde{T}$ is given by

$$(\psi^* T, (\bar{\psi}_{Ge}^* A_{v_{Ge}})_{v_{Ge}(\psi^* T)})$$

where $\bar{\psi}$ is the map $\bar{\psi}: \psi^* T \rightarrow T$ and $\bar{\psi}_{Ge}$ denote the restrictions $\bar{\psi}: Ge \rightarrow G\bar{\psi}(e)$, as in Remark 3.82.

Lemma 4.34. *Suppose that \mathcal{V} is complete and that $\rho: A \rightarrow \Sigma_G$ is a root fibration map. If the rightmost triangle in*

$$\begin{array}{ccccc} \Omega_G^0 \wr A & \xrightarrow{V_G} & \mathbf{F} \wr A & \xrightarrow{\quad} & \mathcal{V}^{op} \\ \downarrow & & \downarrow & \nearrow & \\ \Omega_G^0 & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G & & \end{array} \quad (4.35)$$

is a right Kan extension diagram then so is the composite diagram.

Proof. Unpacking definitions using the pointwise formula for right Kan extensions ([McL, X.3.1]), it suffices to check that for each $T \in \Omega_G^0$ the induced functor

$$T \downarrow \Omega_G^0 \wr A \xrightarrow{V_G} V_G(T) \downarrow \mathbf{F} \wr A \quad (4.36)$$

is initial. We will slightly abuse notation by writing $(T \rightarrow U, (A_{v_{Gf}})_{v_{Gf}(U)})$ for the objects of $T \downarrow \Omega_G^0$, as well as $((T_{v_{Ge}} \rightarrow U_{\phi(v_{Ge})})_{v_{Ge} \in V_G(T)}, (A_v)_{v \in V})$ for the objects of $V_G(T) \downarrow \mathbf{F} \wr A$, with the map $\phi: V_G(T) \rightarrow V$ and the condition $\rho(A_v) = U_v$ left implicit.

By Proposition 2.5, $T \downarrow \Omega_G^0 \wr A$ has an initial subcategory $T \downarrow_r \Omega_G^0 \wr A$ of those objects such that $T \rightarrow U$ is the identity on roots. Similarly, again by Proposition 2.5, $V_G(T) \downarrow \mathbf{F} \wr A$ has an initial subcategory

$$\prod_{v_{Ge} \in V_G(T)} T_{v_{Ge}} \downarrow_r A \quad (4.37)$$

of those objects inducing an identity on $\mathbf{F} \wr \mathbf{O}_G$. Moreover, (4.37) comes together with a right retraction r , i.e. a right adjoint to the inclusion i into $V_G(T) \downarrow \mathbf{F} \wr A$, which is built using pullbacks. We now compute the following composite (where we abbreviate expressions $T_{v_{Ge}}$

as $T_{v_{Ge}}$ and implicitly assume that tuples with index Ge (resp. Gf) run over $V_G(T)$ (resp. $V_G(U)$).

$$T \downarrow_r \Omega_G^0 \wr A \xrightarrow{V_G} V_G(T) \downarrow F \wr A \xrightarrow{r} \prod_{v_{Ge} \in V_G(T)} T_{v_{Ge}} \downarrow_r A$$

$$(T \xrightarrow{\psi} U, (A_{Gf})) \mapsto ((T_{Ge} \rightarrow U_{G\psi(e)}), (A_{Gf})) \mapsto ((T_{Ge} \rightarrow \psi_{Ge}^* U_{G\psi(e)}), (\psi_{Ge}^* A_{G\psi(e)}))$$

Since rooted quotients are isomorphisms, the ψ and ψ_{Ge} appearing above are isomorphisms, and hence the natural transformation $i \circ r \circ V_G \Rightarrow V_G$ is a natural isomorphism. Therefore, to check that V_G is initial it suffices to verify that $r \circ V_G$ is an isomorphism.

But now note that an arbitrary choice of rooted isomorphisms $T_{v_{Ge}} \rightarrow U_{v_{Ge}}^r$ uniquely determines a compatible planar structure on T , and thus a unique isomorphism $\psi: U$. Therefore, arbitrary choices of $\psi_{Ge}^* U_{G\psi(e)}$, $\psi_{Ge}^* A_{G\psi(e)}$ uniquely determine U , A_{Gf} , finishing the proof. \square

ROOTFIBPULL LEM

WSPAN_MONAD_DEFINITION

Lemmas 4.31 implies that copying Definition 4.20 yields a monad N_r on $\mathcal{Wspan}_r^!(\Sigma_G^{op}, \mathcal{V})$ lifting the monad N .

Corollary 4.38. *Suppose that finite products in \mathcal{V} commute with colimits in each variable or, more generally, that \mathcal{V} is a monoidal category with diagonals such that \otimes preserves colimits in each variable. Then the functors*

$$\mathbf{Lan} \circ N_r \Rightarrow \mathbf{Lan} \circ N_r \circ \iota \circ \mathbf{Lan}, \quad \mathbf{Lan} \circ \iota \Rightarrow id$$

are natural isomorphisms.

LANPULLCOMA LEM

FINWREATPRODLIM LEM

Proof. This follows by combining Lemma 4.34 with Lemma 2.21. \square

THEMONAD DEF

Definition 4.39. The *genuine equivariant operad monad* is the monad \mathbb{F}_G on $\mathbf{Fun}(\Sigma_G^{op}, \mathcal{V})$ given by

$$\mathbb{F}_G = \mathbf{Lan} \circ N_r \circ \iota$$

and with multiplication and unit given by the composites

$$\mathbf{Lan} \circ N_r \circ \iota \circ \mathbf{Lan} \circ N_r \circ \iota \xrightarrow{\sim} \mathbf{Lan} \circ N_r \circ N_r \circ \iota \Rightarrow \mathbf{Lan} \circ N_r \circ \iota$$

$$id \xleftarrow{\sim} \mathbf{Lan} \circ \iota \Rightarrow \mathbf{Lan} \circ N_r \circ \iota.$$

We will write $\mathbf{Op}_G(\mathcal{V})$ for the category $\mathbf{Alg}_{\mathbb{F}_G}(\mathcal{V})$ of *genuine equivariant operads*.

Remark 4.40. The functor $\mathbf{Lan} \circ N_r \circ \iota$ is isomorphic to $\mathbf{Lan} \circ N \circ \iota$, and this isomorphism is compatible with the multiplication and unit in Definition 4.39, and hence we will henceforth simply write N rather than N_r . **THEMONAD DEF**

From this point of view, the role of root fibrations is to guarantee that $\mathbf{Lan} \circ N \circ \iota$ is indeed a monad, though unnecessary to describe the monad structure itself.

Remark 4.41. Since a map

$$\mathbb{F}_G X = \mathbf{Lan} \circ N \circ \iota X \rightarrow X$$

is adjoint to a map

$$N \circ \iota X \rightarrow \iota X$$

one easily verifies that X is a genuine equivariant operad, i.e. a \mathbb{F}_G -algebra, iff ιX is a N -algebra.

Moreover, the bar resolution $\mathbb{F}_G^{on+1} X$ is isomorphic to $\mathbf{Lan}(N^{on+1} \iota X)$.

4.3 Comparison with (regular) equivariant operads

We start by noting that in the case $G = *$, genuine operads simply recover the usual notion of symmetric operads, i.e. $\text{Sym}_*(\mathcal{V}) \simeq \text{Sym}(\mathcal{V})$ and $\text{Op}_*(\mathcal{V}) \simeq \text{Op}(\mathcal{V})$, and in what follows we will adopt the notations $\text{Sym}^G(\mathcal{V})$ and $\text{Op}^G(\mathcal{V})$ for the corresponding categories of G -objects. Our goal will be to relate these to the categories $\text{Sym}_G(\mathcal{V})$ and $\text{Op}_G(\mathcal{V})$ of genuine equivariant sequences and genuine operads.

We will throughout this section fix a total order of G such that the identity e is the first element, though we note that the exact order is unimportant, as any other such choice would lead to unique isomorphisms between the constructions in this section.

We thus have an inclusion functor

$$\begin{aligned} \iota: G \times \Sigma &\hookrightarrow \Sigma_G \\ C &\longmapsto G \cdot C \end{aligned}$$

where $G \cdot C$ is the constant tuple $(C)_{g \in G}$, which we think of as $|G|$ copies of C , planarized according to C and the order on G . Moreover, letting $\Sigma_G^{\text{fr}} \hookrightarrow \Sigma_G$ denote the full subcategory of G -free corollas, there is an induced retraction $\rho: \Sigma_G^{\text{fr}} \rightarrow G \times \Sigma$ defined by $\rho((C_i)_{1 \leq i \leq |G|}) = G \cdot C_1$ together with isomorphisms $C \simeq \rho(C)$ uniquely determined by the condition that they are the identity on the first tree component C_1 .

We now consider the associated adjunctions.

$$\begin{array}{ccc} & \xleftarrow{\iota_!} & \\ \text{Sym}_G(\mathcal{V}) & \xrightarrow{\quad \quad} & \text{Sym}^G(\mathcal{V}) \\ & \xleftarrow{\iota_*} & \end{array} \quad (4.42) \quad \boxed{\text{TWOADJOINTS EQ}}$$

Explicitly, we have the formulas (where we write G -corollas as $(C_i)_I$ for $I \in \mathbf{O}_G$)

$$\iota_! Y((C_i)_I) = \begin{cases} Y(C_1), & (C_i)_I \in \Sigma_G^{\text{fr}} \\ \emptyset, & (C_i)_I \notin \Sigma_G^{\text{fr}} \end{cases}, \quad \iota_* X(C) = X(G \cdot C), \quad \iota_* Y((C_i)_I) = \left(\prod_I Y(C_i) \right)^G,$$

where in the formula for $\iota_*(-)$ the action of G interchanges factors according to the action on the indexing set I . As a side note, note that the formulas for $\iota_!$ and ι_* are independent of the chosen order of G .

Remark 4.43. $\iota_!$ essentially identifies $\text{Sym}^G(\mathcal{V})$ as the coreflexive subcategory of sequences $X \in \text{Sym}_G(\mathcal{V})$ such that $X(C) = \emptyset$ whenever C is not a free corolla.

By contrast, ι_* identifies $\text{Sym}^G(\mathcal{V})$ with a far more interesting reflexive subcategory of sequences $X \in \text{Sym}_G(\mathcal{V})$ such that $X(C)$ for each C not a free corolla must satisfy a fixed point condition. Concretely, letting $\varphi: G \rightarrow \mathbf{r}(C)$ denote the unique map preserving the minimal element, one has

$$X(C) \xrightarrow{\sim} X(\varphi^* C)^\Gamma$$

for $\Gamma \leq \text{Aut}(\varphi^* C)$ the subgroup preserving the quotient map $\varphi^* C \rightarrow C$ under precomposition (note that $\varphi^* C \in \Sigma_G^{\text{fr}}$).

There is an obvious natural transformation $\beta: \iota_! \Rightarrow \iota_*$ which for $(C_i)_I \in \Sigma_G^{\text{fr}}$ sends $Y(C_1)$ to the “ G -twisted diagonal” of $\prod_I Y(C_i)$. Moreover, letting $\eta_!, \epsilon_!$ (resp. η_*, ϵ_*) denote the unit and counit of the $(\iota_!, \iota^*)$ adjunction (resp. (ι^*, ι_*) adjunction) it is straightforward to check that the following diagram commutes.

$$\begin{array}{ccc} \iota_! \iota^* \iota_* & \xrightarrow{\epsilon_!} & \iota_* \\ \epsilon_* \downarrow \simeq & \nearrow \beta & \simeq \downarrow \eta_! \\ \iota_! & \xrightarrow{\eta_*} & \iota_* \iota^* \iota_! \end{array} \quad (4.44) \quad \boxed{\text{BETADEFSQUARE EQ}}$$

Remark 4.45. An exercise in adjunctions shows that any outer square as in (4.44) commutes provided at least one of the adjunctions in 4.42 is (co)reflexive, so that (4.44) can be regarded as an alternative definition of β . BETADEFSQUARE EQ

Proposition 4.46. *One has the following:*

- (i) the map $\iota^* \mathbb{F}_G \xrightarrow{\eta_*} \iota^* \mathbb{F}_{G\iota_*} \iota^*$ is an isomorphism, and thus (cf. Prop. 2.26) $\iota^* \mathbb{F}_{G\iota_*}$ is a monad; MONADADJ PROP
- (ii) the map $\iota^* \mathbb{F}_G \iota_! \xrightarrow{\beta} \iota^* \mathbb{F}_{G\iota_*}$ is an isomorphism of monads;
- (iii) the map $\iota_! \iota^* \mathbb{F}_G \iota_! \xrightarrow{\epsilon_!} \mathbb{F}_{G\iota_!}$ is an isomorphism;
- (iv) there is a natural isomorphism of monads $\alpha: \mathbb{F} \rightarrow \iota^* \mathbb{F}_{G\iota_!}$.

Proof. We first show (i), starting with some notation. In analogy with Σ_G^{fr} , we write $\Omega_G^{0,\text{fr}}$ for the subcategory of free trees and note that the leaf-root and vertex functors then restrict to functors $\text{lr}: \Omega_G^{0,\text{fr}} \rightarrow \Sigma_G^{\text{fr}}$, $V_G: \Omega_G^{0,\text{fr}} \rightarrow \mathbb{F} \wr \Sigma_G^{\text{fr}}$. Moreover, for each $C \in \Sigma_G^{\text{fr}}$ one has an equality of rooted undercategories between $C \downarrow_r \Omega_G^0$ and $C \downarrow_r \Omega_G^{0,\text{fr}}$, and thus $\iota^* \mathbb{F}_G X$ is computed by the Kan extension of the following diagram.

$$\begin{array}{ccccccc} \Omega_G^{0,\text{fr}} & \longrightarrow & \mathbb{F} \wr \Sigma_G^{\text{fr}} & \xrightarrow{\mathbb{F} \wr X} & \mathbb{F} \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\ \downarrow & & & & & & \\ \Sigma_G^{\text{fr}} & & & & & & \end{array} \quad (4.47) \quad \text{IFGI EQ}$$

(i) now follows by noting that $X \rightarrow \iota_* \iota^* X$ is an isomorphism when restricted to Σ_G^{fr} .

For (ii), to show that $\iota^* \mathbb{F}_{G\iota_!} \rightarrow \iota^* \mathbb{F}_{G\iota_*}$ is an isomorphism one just repeats the argument in the previous paragraph by noting that $\iota_! \rightarrow \iota_*$ is an isomorphism when restricted to Σ_G^{fr} . To check that this is a map of monads, we recall first that the monad structure on $\iota^* \mathbb{F}_{G\iota_*}$ is given as described in Proposition 2.26. Unpacking definitions, compatibility with multiplication reduces to showing that the composite $\iota_! \iota^* \xrightarrow{\epsilon_!} id \xrightarrow{\eta_*} \iota_* \iota^*$ coincides with $\beta \iota^*$ while compatibility with units reduces to showing that the composite $id \xrightarrow{\eta_!} \iota^* \iota_! \xrightarrow{\iota^* \beta} \iota^* \iota_* \xrightarrow{\epsilon_*} id$ is the identity. Both of these are a consequence of (4.44), following from the diagrams below (where the top composites are identities).

$$\begin{array}{ccc} \iota_! \iota^* & \xrightarrow{\iota_! \iota^* \eta_*} & \iota_! \iota^* \iota_* \iota^* \xrightarrow{\iota_! \epsilon_* \iota^*} \iota_! \iota^* \\ \epsilon_! \downarrow & & \downarrow \epsilon_! \iota_* \iota^* \\ id & \xrightarrow{\eta_*} & \iota_* \iota^* \end{array} \quad \begin{array}{ccc} \iota^* \iota_* & \xrightarrow{\eta_! \iota^* \iota_*} & \iota^* \iota_! \iota^* \iota_* \xrightarrow{\iota^* \epsilon_! \iota_*} \iota^* \iota_* \\ \epsilon_* \downarrow \simeq & & \downarrow \iota^* \iota_! \epsilon_* \simeq \\ id & \xrightarrow{\eta_!} & \iota^* \iota_! \end{array} \quad \begin{array}{ccc} & & \swarrow \beta \iota^* \\ & & \searrow \iota^* \beta \end{array} \quad (4.48)$$

(iii) amounts to showing that if $X(C) = \emptyset$ whenever $C \notin \Sigma_G^{\text{fr}}$ then it is also $\mathbb{F}_G X(C) = \emptyset$. But since for such $C \notin \Sigma_G^{\text{fr}}$ the undercategory $C \downarrow \Omega_G^0$ consists of trees with at least one non-free vertex (namely the root vertex), the composite

$$C \downarrow \Omega_G^0 \xrightarrow{V_G} \mathbb{F} \wr \Sigma_G \xrightarrow{\mathbb{F} \wr X} \mathbb{F} \wr \mathcal{V}^{op} \xrightarrow{\Pi} \mathcal{V}^{op}$$

is constant equal to \emptyset , and (iii) follows.

Finally, we show (iv). We will slightly abuse notation by writing $G \times \Sigma \hookrightarrow \Sigma_G$ for the image of ι and similarly $G \times \Omega^0 \hookrightarrow \Omega_G^0$ for the image of the obvious analogous functor $\iota: G \times \Omega^0 \rightarrow \Omega_G^0$. The map $\alpha: \mathbb{F} \rightarrow \iota^* \mathbb{F}_{G\iota_!}$ is the adjoint to the map $\tilde{\alpha}: \mathbb{F} \iota^* \rightarrow \iota^* \mathbb{F}_G$ encoded on spans by the

following diagram.

$$\begin{array}{ccccccc}
 G \times \Omega^0 & \longrightarrow & F \wr (G \times \Sigma) & \xrightarrow{\iota^* X} & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\
 \downarrow & \searrow & \downarrow & & \downarrow & \searrow & \downarrow \\
 & & \Omega_G^0 & \longrightarrow & F \wr \Sigma_G & \xrightarrow{X} & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 & \searrow & \downarrow & & \downarrow & & \\
 G \times \Sigma & & \Sigma_G & & & &
 \end{array} \quad (4.49) \quad \boxed{\text{MONADEQUIV DEF}}$$

That α is a natural isomorphism follows by the previous identifications $C \downarrow \Omega_G^0 \simeq C \downarrow \Omega_G^{0, \text{fr}}$ for $C \in G \times \Sigma$ together with the fact that the retraction $\rho: \Omega_G^{0, \text{fr}} \rightarrow G \times \Omega^0$ (built just as the retraction $\rho: \Sigma_G^{\text{fr}} \rightarrow G \times \Sigma$) retracts $C \downarrow \Omega_G^{0, \text{fr}}$ to the undercategory $C \downarrow G \times \Omega^0$, which is thus initial (as well as final).

Intuitively, the final claim that α is a map of monads follows from the fact that the composite $\mathbb{F}\mathbb{F} \rightarrow \iota^* \mathbb{F}_G \iota \iota^* \mathbb{F}_G \iota \rightarrow \iota^* \mathbb{F}_G \mathbb{F}_G \iota$ is encoded by the analogous natural transformation of diagrams for strings $G \times \Omega^1 \hookrightarrow \Omega_G^{1, \text{fr}}$. However, since the presence of left Kan extensions in the definitions of \mathbb{F} , \mathbb{F}_G can make a rigorous direct proof of this last claim fairly cumbersome, we sketch here a workaround argument. We first consider the adjunction $\iota: \mathbf{WSpan}^l((G \times \Sigma)^{op}, \mathcal{V}) \rightleftarrows \mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V}): \iota^*$ where ι is composition with ι and ι^* is the pullback of spans. Writing N, N_G for the monads on the span categories, mimicking (4.49) yields a map $\tilde{\alpha}: N \rightarrow \iota^* N_G \iota$ encoded by the diagram (where the front and back squares are pullbacks).

$$\begin{array}{ccccccc}
 (G \times \Omega^0) \wr \iota^* A & \longrightarrow & F \wr \iota^* A & \longrightarrow & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\
 \downarrow & \searrow & \downarrow & & \downarrow & \searrow & \downarrow \\
 & & \Omega_G^0 \wr A & \longrightarrow & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 & \searrow & \downarrow & & \downarrow & & \\
 G \times \Omega^0 & \longrightarrow & F \wr (G \times \Sigma) & \longrightarrow & F \wr \Sigma_G & \longrightarrow & \\
 \downarrow & \searrow & \downarrow & & \downarrow & & \\
 G \times \Sigma & & \Sigma_G & & & &
 \end{array}$$

The claim that $\tilde{\alpha}$ is a map of monads is then straightforward. Writing

$$\text{Lan}: \mathbf{WSpan}^l((G \times \Sigma)^{op}, \mathcal{V}) \rightleftarrows \mathbf{Fun}((G \times \Sigma)^{op}, \mathcal{V}): j \quad \text{Lan}_G: \mathbf{WSpan}^l(\Omega_G^{op}, \mathcal{V}) \rightleftarrows \mathbf{Fun}(\Omega_G^{op}, \mathcal{V}): j_G$$

for the span functor adjunctions, $\alpha: \mathbb{F} \rightarrow \iota^* \mathbb{F}_G \iota$ can then be written as the composite

$$\text{Lan} N j \rightarrow \text{Lan} \iota^* N_G \iota j \rightarrow \iota^* \text{Lan}_G N_G j_G \iota$$

where the first map is the isomorphism of monads induced by $\tilde{\alpha}$ and the second map can be shown directly to be a monad map by unpacking the monad structures in Propositions 2.25 and 2.26. \square

Combining the previous result with Propositions 2.25 and 2.26 now gives the following.

Corollary 4.50. *The adjunctions (4.42) extends to adjunctions*

$$\begin{array}{ccc}
 & \xleftarrow{\iota_!} & \\
 \text{Op}_G(\mathcal{V}) & \xrightarrow{\quad} & \text{Op}^G(\mathcal{V}). \\
 & \xleftarrow{\iota_*} &
 \end{array} \quad (4.51) \quad \boxed{\text{TWOADJOINTSOP EQ}}$$

In particular, ι_* identifies $\text{Op}^G(\mathcal{V})$ as a reflexive subcategory of $\text{Op}_G(\mathcal{V})$.

Remark 4.52. Remark [REFLCOREFL REM](#) 4.43 extends to operads mutatis mutandis.

Moreover, the isomorphism $\iota_! \iota^* \mathbb{F}_G \iota_! \xrightarrow{\epsilon_!} \mathbb{F}_G \iota_!$ then shows that \mathbb{F}_G essentially preserves the image of $\iota_!$, and can thus be identified with \mathbb{F} over it.

However, the analogous statement fails for ι_* , i.e., one does not always have that

$$\mathbb{F}_G \iota_* \xrightarrow{\eta_*} \iota_* \iota^* \mathbb{F}_G \iota_* \quad (4.53)$$


KEYNONISO EQ

is an isomorphism. In fact, showing that [KEYNONISO EQ](#) (4.53) *does* become an isomorphism when restricted to suitably cofibrant objects is one of the key technical ingredients for our proof of the Quillen equivalence between $\mathbf{Op}_G(\mathcal{V})$ and $\mathbf{Op}^G(\mathcal{V})$, and will be the subject of [§6](#). [COFIB SEC](#)


For now, we end this section with a minimal counterexample to the more general claim.

Let $G = \mathbb{Z}/2$ and $Y = * \in \mathbf{Sym}^G(\mathcal{V})$ be the singleton.

When evaluating $\mathbb{F}_G Y$ at the G -fixed stump corolla $G/G \cdot C_0$, the two G -trees T_1 and T_2 below encode two distinct points (since T_1, T_2 are not isomorphic as objects under $G/G \cdot T_0$).



However, when pulling these points back to the G -free stump corolla $G \cdot C_0$ one obtains the same point, namely that encoded by the G -tree T below.



Moreover, it is not hard to modify the example above to produce similar examples when evaluating $\mathbb{F}_G Y$ at non-empty corollas.

However, such counter-examples all require the use of trees with stumps. Indeed, it can be shown that [KEYNONISO EQ](#) (4.53) is an isomorphism whenever evaluated at a Y such that $Y(C_0) = \emptyset$.

4.4 Indexing systems and partial genuine operads

As discussed preceding Theorem [MAINEXIST2 THM](#) II, the Elmendorf-Piacenza equivalence [COFADJINT EQ](#) (II.7) has analogues

$$\mathbf{Top}^{\mathbf{Op}}_{\mathcal{F}} \xrightleftharpoons[\iota_*]{\iota^*} \mathbf{Top}^G_{\mathcal{F}} \quad (4.54)$$

for each family \mathcal{F} of subgroups of G . Here $\mathbf{Op}_{\mathcal{F}} \hookrightarrow \mathbf{Op}_G$ consists of those G/H such that $H \in \mathcal{F}$ so that $\mathbf{Top}^{\mathbf{Op}}_{\mathcal{F}}$ consists of partial coefficient systems. These specialized equivalences provide an alternative approach to universal $E\mathcal{F}$ spaces: rather than cofibrantly replacing the object $\delta_{\mathcal{F}} \in \mathbf{Top}^{\mathbf{Op}}_G$ discussed in the introduction, one builds an $E\mathcal{F}$ -space by $\iota^*(C^*) = (C^*)(G)$ where now $*$ $\in \mathbf{Top}^{\mathbf{Op}}_{\mathcal{F}}$ is the terminal object and C the cofibrant replacement in $\mathbf{Top}^{\mathbf{Op}}_{\mathcal{F}}$.

In keeping with the motivation that the Blumberg-Hill $N\mathcal{F}$ operads are the operadic analogues of universal $E\mathcal{F}$ spaces, we will now show that the closure conditions for indexing systems identified in [\[3, Def. 3.22\]](#) are (almost exactly) the necessary conditions to define categories $\mathbf{Op}_{\mathcal{F}}$ of partial genuine equivariant operads. [BH15](#)

We start by recalling that in the classic setting \mathcal{F} is a family of subgroups of G iff the associated subcategory $\mathbf{Op}_{\mathcal{F}} \hookrightarrow \mathbf{Op}_G$ is a sieve, defined as follows.

Definition 4.55. A *sieve* of a category \mathcal{D} is a subcategory \mathcal{S} such that for any arrow $f: d \rightarrow s$ of \mathcal{D} with $s \in \mathcal{S}$ then both d and f are also in \mathcal{S} . In particular, sieves are full subcategories.

ILY_COROLLAS_DEF

ILY_COROLLAS_REM

Definition 4.56. A family of G -corollas is a sieve $\Sigma_{\mathcal{F}} \hookrightarrow \Sigma_G$.

Remark 4.57. A family of corollas $\Sigma_{\mathcal{F}}$ can equivalently be encoded by a collection $\mathcal{F} = \{\mathcal{F}_n\}_{n \geq 0}$ of families \mathcal{F}_n of graph subgroups of $G \times \Sigma_n$, so that there is an equivalence of categories $\Sigma_{\mathcal{F}} \simeq \coprod \mathcal{O}_{\mathcal{F}_n}$. As such, we abuse notation and abbreviate either set of data as \mathcal{F} .

Writing $\gamma: \Sigma_{\mathcal{F}} \hookrightarrow \Sigma_G$ for the inclusion and $\text{Sym}_{\mathcal{F}}(\mathcal{V}) = \mathcal{V}^{\Sigma_{\mathcal{F}}^{op}}$, we thus have a pair of standard adjunctions

$$\begin{array}{ccc} & \xrightarrow{\gamma_!} & \\ \text{Sym}_{\mathcal{F}}(\mathcal{V}) & \xleftarrow{\gamma^*} & \text{Sym}_G(\mathcal{V}) \\ & \xleftarrow{\gamma_*} & \end{array} \quad (4.58)$$

F_TWADJOINTS_EQ

Our focus will be on the $(\gamma_!, \gamma^*)$ adjunction. The requirement that $\Sigma_{\mathcal{F}}$ be a sieve then implies that $\gamma_!$ simply extends presheaves by the initial object $\emptyset \in \mathcal{V}$, so that $\gamma_!$ identifies $\text{Op}_{\mathcal{F}}(\mathcal{V})$ with a (coreflexive) subcategory of $\text{Op}_G(\mathcal{V})$. One may then ask for conditions on the family of corollas \mathcal{F} such that the genuine operad monad \mathbb{F}_G preserves this subcategory and, as it turns out, the answer is almost exactly given by the Blumberg-Hill indexing systems.

FTREE_DEF

Definition 4.59. Let \mathcal{F} be a family of G -corollas.

We say that a G -tree T is a \mathcal{F} -tree if all of its G -vertices T_v , $v \in V_G(T)$ are in $\Sigma_{\mathcal{F}}$, and we denote by $\Omega_{\mathcal{F}} \hookrightarrow \Omega_G$, $\Omega_{\mathcal{F}}^0 \hookrightarrow \Omega_G^0$ the full subcategories spanned by the \mathcal{F} -trees.

Remark 4.60. By vacuousness the stick G -trees $G \cdot_H \eta \simeq (\eta)_{G/H}$ are always \mathcal{F} -trees.

VACUOUSNESS_REM

INDEXSYS_DEF

Definition 4.61. A family \mathcal{F} of G -corollas is called a *weak indexing system* if for any \mathcal{F} -tree $T \in \Omega_{\mathcal{F}}^0$ it is $\text{lr}(T) \in \Sigma_{\mathcal{F}}$, i.e. if the leaf-root functor restricts to a functor $\text{lr}: \Omega_{\mathcal{F}}^0 \rightarrow \Sigma_{\mathcal{F}}$. Moreover, \mathcal{F} is called simply an *indexing system* if all trivial corollas $(G/H) \cdot C_n \simeq (C_n)_{G/H}$ are in $\Sigma_{\mathcal{F}}$.

Remark 4.62. In light of Remark 4.60 any weak indexing system must contain the 1-corollas $(G/H) \cdot C_1 \simeq (C_1)_{G/H}$.

VACUOUSNESS_REM

Remark 4.63. The notion of indexing system was first introduced in [3, Def. 3.22], though packaged quite differently. Moreover, a third definition of (weak) indexing systems as the sieves $\Omega_{\mathcal{F}} \hookrightarrow \Omega_G$ was presented by the second author in [15, §9]. The equivalence between the definitions in [3] and [15] was addressed in [15, Bmk. 9.7], hence here we address only the easier equivalence between Definition 4.61 and the sieve definition in [15, §9].

The existence of canonical maps $\text{lr}(T) \rightarrow T$ shows that the sieve condition implies the lr condition in Definition 4.61. Conversely, as discussed immediately preceding [15, Def. 9.5], the sieve condition needs only be checked for inner faces and degeneracies, i.e. tall maps, and thus follows from Definition 4.61 since planar tall strings $\Omega_{\mathcal{F}}^1$ between \mathcal{F} -trees match the pullback $\Omega_{\mathcal{F}}^0 \rightarrow \mathbb{F} \wr \Sigma_{\mathcal{F}} \leftarrow \mathbb{F} \wr \Omega_{\mathcal{F}}^0$.

The connection between weak indexing systems and \mathbb{F}_G is given by the following, which generalizes Proposition 4.46.

MONAD_COMPARISON_PROP

_COMPARISON_PROP

Proposition 4.64. Let \mathcal{F} be a weak indexing system. Then:

- (i) the map $\gamma^* \mathbb{F}_G \xrightarrow{\eta_*} \gamma^* \mathbb{F}_G \gamma_* \gamma^*$ is an isomorphism, and thus (cf. Prop. 2.26) $\gamma^* \mathbb{F}_G \gamma_*$ is a monad;
- (ii) the map $\gamma^* \mathbb{F}_G \gamma_! \xrightarrow{\beta} \gamma^* \mathbb{F}_G \gamma_*$ is an isomorphism of monads;
- (iii) the map $\gamma_! \gamma^* \mathbb{F}_G \gamma_! \xrightarrow{\epsilon_!} \mathbb{F}_G \gamma_!$ is an isomorphism.

MONADADJ_PROP

Proof. This follows just like the analogous parts of Proposition 4.46 by replacing $\text{lr}: \Omega_G^{0, \text{fr}} \rightarrow \Sigma_G^{\text{fr}}$ with $\text{lr}: \Omega_{\mathcal{F}}^0 \rightarrow \Sigma_{\mathcal{F}}$. For (i), note that if $C \in \Sigma_{\mathcal{F}}$ there is an identification between $C \downarrow \Omega_G^0$ and $C \downarrow \Omega_{\mathcal{F}}^0$, so that $\mathbb{F}_G X(C)$ only depends on the values of X on $\Sigma_{\mathcal{F}}$. (ii) is immediate. Lastly, (iii) follows since if $C \notin \Sigma_{\mathcal{F}}$ then any tree in $C \downarrow \Omega_G^0$ must contain at least one G -vertex not in $\Sigma_{\mathcal{F}}$, so that indeed $\mathbb{F}_G \gamma_! Y(C) = \emptyset$. \square

MONAD_COMPARISON_PROP

Notation 4.65. We write $\mathbb{F}_{\mathcal{F}} = \gamma^* \mathbb{F}_G \gamma_!$ for the induced monad on $\text{Sym}_{\mathcal{F}}(\mathcal{V})$, and $\text{Op}_{\mathcal{F}}(\mathcal{V})$ for the corresponding algebras.

Corollary 4.66. *The adjunction (4.58) lifts to an adjunction on algebras*

$$\begin{array}{ccc} & \gamma_! & \\ \text{Op}_{\mathcal{F}}(\mathcal{V}) & \xleftarrow{\gamma^*} & \text{Op}_G(\mathcal{V}) \\ & \gamma_* & \end{array} \quad (4.67)$$

TWOADJOINTSOPF EQ

Remark 4.68. Part (iii) of Proposition 4.64 states that if \mathcal{F} is a weak indexing system then \mathbb{F}_G essentially preserves the image of $\gamma_!$ (moreover, the converse is easily seen to also hold). As such, we will sometimes find it conceptually convenient to regard $\mathbb{F}_{\mathcal{F}}$ as “restricting \mathbb{F}_G ”.

Remark 4.69. The free corollas of §4.3 form a weak indexing system $\Sigma_{\mathcal{F}}^{\text{fr}} = \Sigma_{\mathcal{F}}$, and moreover, there is an equivalence of categories $\text{Op}_{\mathcal{F}}^G \simeq \text{Op}_{\Sigma_{\mathcal{F}}}^G$, so that (4.51) is a particular case of (4.67). However, while our discussion of (4.51) focuses on the (ι^*, ι_*) -adjunction, due to the fact that the intended model structures on $\text{Op}_G^G(\mathcal{V})$ in Theorem I are defined via fixed point conditions, our discussion of (4.67) focuses on the $(\iota_!, \iota^*)$ -adjunction, due to the model structures in Theorem II being projective.

Remark 4.70. In most cases, the rightmost (ι^*, ι_*) -adjunction appearing in Theorem III is induced by an inclusion $\iota: \Sigma_G^{\text{fr}} \hookrightarrow \Sigma_{\mathcal{F}}$. However, it is possible for $\Sigma_G^{\text{fr}} \not\hookrightarrow \Sigma_{\mathcal{F}}$ (the most interesting case being that of $\Sigma_G^{\geq 1} \hookrightarrow \Sigma_G$ the corollas of arity ≥ 1 , which models non-unital operads), in which case (and compatibly with the $\Sigma_G^{\text{fr}} \hookrightarrow \Sigma_{\mathcal{F}}$ case), we instead use the composite adjunction

$$\text{Op}_{\mathcal{F}}(\mathcal{V}) \xleftarrow[\gamma^*]{\gamma_!} \text{Op}_G(\mathcal{V}) \xleftarrow[\iota_*]{\iota^*} \text{Op}^G(\mathcal{V}) \quad (4.71)$$

COMPADJ EQ

We note that the right adjoint $\gamma^* \iota_*$ is still defined by computing fixed points while the left adjoint $\iota^* \gamma_!$ is still essentially a forgetful functor, with those levels not present in \mathcal{F} declared to be \emptyset .

In practice, however, the use of the composite adjunction (4.71) is fairly benign, requiring only minor adjustments to the notation of the proofs in §6.4.

5 Free extensions and the existence of model structures

In order to prove all of our main theorems we will need to homotopically analyze free extensions of genuine equivariant operads, i.e. pushouts of the form

$$\begin{array}{ccc} \mathbb{F}_G X & \longrightarrow & \mathcal{P} \\ \mathbb{F}_G u \downarrow & & \downarrow \\ \mathbb{F}_G Y & \longrightarrow & \mathcal{P}[u] \end{array} \quad (5.1)$$

FREE_FG_EXT_EQ

in the category Op_G . As is common in the literature (e.g. [18, 19, 1, 22, 14]), the key technical ingredient will be the identification of a suitable filtration

$$\mathcal{P} = \mathcal{P}_0 \rightarrow \mathcal{P}_1 \rightarrow \mathcal{P}_2 \rightarrow \cdots \rightarrow \mathcal{P}_{\infty} = \mathcal{P}[u] \quad (5.2)$$

FILTR EQ

of the map $\mathcal{P} \rightarrow \mathcal{P}[u]$ in the underlying category Sym_G . To explain how this filtration is obtained, and abbreviating \mathbb{F}_G as \mathbb{F} , note first that $\mathcal{P}[u]$ is given by a coequalizer

$$\mathcal{P} \amalg \mathbb{F}X \amalg \mathbb{F}Y \xrightleftharpoons{\quad} \mathcal{P} \amalg \mathbb{F}Y \quad (5.3)$$

REFLCOEQ EQ

where $\tilde{\sqcup}$ denotes the algebraic coproduct, i.e. the coproduct in \mathbf{Op}_G , and, a priori, the coequalizer is also calculated in \mathbf{Op}_G . However, (5.3) is a so called *reflexive coequalizer*, meaning that the maps being coequalized have a common section, and standard arguments show that it is hence also an underlying coequalizer in \mathbf{Sym}_G .

In practice, we will need to enlarge (5.3) somewhat. Firstly, note that (5.3) corresponds to the two bottom levels of the bar construction $B_l(\mathcal{P}, \mathbb{F}X, \mathbb{F}Y) = \mathcal{P} \tilde{\sqcup} (\mathbb{F}X)^{\tilde{\sqcup} l} \tilde{\sqcup} \mathbb{F}Y$, whose colimit (over Δ^{op}) is again $\mathcal{P}[u]$. For technical reasons, we prefer the double bar construction

$$B_l(\mathcal{P}, \mathbb{F}X, \mathbb{F}X, \mathbb{F}X, \mathbb{F}Y) = \mathcal{P} \tilde{\sqcup} (\mathbb{F}X)^{\tilde{\sqcup} l} \tilde{\sqcup} \mathbb{F}X \tilde{\sqcup} (\mathbb{F}X)^{\tilde{\sqcup} l} \tilde{\sqcup} \mathbb{F}Y = \mathcal{P} \tilde{\sqcup} (\mathbb{F}X)^{\tilde{\sqcup} 2l+1} \tilde{\sqcup} \mathbb{F}Y. \quad (5.4)$$

To actually describe the individual levels of (5.4) one further resolves \mathcal{P} so as to obtain the bisimplicial object

$$B_l(\mathbb{F}^{n+1}\mathcal{P}, \mathbb{F}X, \mathbb{F}X, \mathbb{F}X, \mathbb{F}Y) = \mathbb{F}^{n+1}\mathcal{P} \tilde{\sqcup} (\mathbb{F}X)^{\tilde{\sqcup} 2l+1} \tilde{\sqcup} \mathbb{F}Y \simeq \mathbb{F}(\mathbb{F}^n\mathcal{P} \sqcup X^{\sqcup 2l+1} \sqcup Y), \quad (5.5)$$

where \sqcup denotes the coproduct in \mathbf{Sym}_G . As in Remark 4.41, each level of (5.5) can then be described as

$$\mathbf{Lan} N(N^n \iota \mathcal{P} \sqcup \iota X^{\sqcup 2l+1} \sqcup \iota Y), \quad (5.6)$$

for N the span monad (cf. Definition 4.20) and \sqcup now the coproduct of spans. In particular, each level of (5.5) is thus a left Kan extension over some category Ω_G^{n, λ_l} , which we explicitly identify in §5.1, giving the first identification below.

$$\mathcal{P} \tilde{\coprod}_{\mathbb{F}X} \mathbb{F}Y \simeq \operatorname{colim}_{(\Delta \times \Delta)^{op}} \left(\mathbf{Lan}_{(\Omega_G^{n, \lambda_l} \rightarrow \Sigma_G)^{op}} N_{n, l}^{(\mathcal{P}, X, Y)} \right) \simeq \mathbf{Lan}_{(\Omega_G^e \rightarrow \Sigma_G)^{op}} \tilde{N}^{(\mathcal{P}, X, Y)} \quad (5.7)$$

The second identification, which reduces the calculation to a single left Kan extension, is an instance of Proposition 5.40, a result whose proof is straightforward but lengthy, and thus postponed to the appendix. The category Ω_G^e of *extension trees* appearing on the right side is obtained as a categorical realization $\Omega_G^e = |\Omega_G^{n, \lambda_l}|$, which we explicitly describe and analyze in §5.2. In particular, we identify a smaller and more convenient subcategory $\widehat{\Omega}_G^e \hookrightarrow \Omega_G^e$ that is suitably initial, so that Ω_G^e can be replaced with $\widehat{\Omega}_G^e$ in (5.7).

The desired filtration (5.2) then follows from a filtration of the category $\widehat{\Omega}_G^e$ itself, and this is the subject of §5.3.

HERE

5.1 Labeled planar strings

In this section we explicitly identify the categories underlying the left Kan extensions in (5.6).


In the notation of Remark 2.30, letting $\langle\langle l \rangle\rangle = \{-\infty, -l, \dots, -1, 0, 1, \dots, l, \infty\}$ and writing λ_l for the partition $\lambda_{l, a} = \{-\infty\}$, $\lambda_{l, i} = \langle\langle l \rangle\rangle - \{-\infty\}$, (5.6) can be repackaged as an instance of the functor $\mathbf{Lan} \circ N \circ \coprod \circ (N^{\times \lambda_l})^{on} \circ \iota^{\times \langle\langle l \rangle\rangle}$. Our goal is thus to understand the underlying categories of the spans in the image of the functor $N \circ \coprod \circ (N^{\times \lambda_l})^{on}$, though we will find it preferable and no harder to tackle the more general case of the functors $N^{s+1} \circ \coprod \circ (N^{\times \lambda})^{on-s}$.

Definition 5.8. A l -node labeled G -tree (or just l -labeled G -tree) is a pair $(T, V_G(T) \rightarrow \{1, \dots, l\})$ with $T \in \Omega_G$, which we think of as a G -tree together with G -vertices labels in $1, \dots, l$.

Further, a tall map $\varphi: T \rightarrow S$ between l -labeled trees is called a *label map* if for each G -vertex v_{Ge} of T with label j , the vertices of the subtree $S_{v_{Ge}}$ are all labeled by j .

Lastly, given a subset $J \subset l$, a planar label map $\varphi: T \rightarrow S$ is said to be J -inert if for every G -vertex v_{Ge} of T with label $j \in J$ it is $S_{v_{Ge}} = T_{v_{Ge}}$.

Example 5.9. Consider the 2-labeled trees below (for $G = *$ the trivial group), with black nodes (\bullet) denoting labels by the number 1 and white nodes (\circ) labels by the number 2. The planar map φ (sending $a_i \mapsto a$, $b \mapsto b$, $c \mapsto c$, $d \mapsto d$, $e \mapsto e$) is a label map which is $\{1\}$ -inert.



(5.10) SUBSDATUM TREES LAB EQ

Definition 5.11. Let $-1 \leq s \leq n$ and $\lambda = \lambda_a \sqcup \lambda_i$ a partition of $\{1, 2, \dots, l\}$.

We define $\Omega_G^{n,s,\lambda}$ to have as objects n -planar strings (where $T_{-1} = \text{lr}(T_0)$ as in (3.89))

$$T_{-1} \xrightarrow{\varphi_0} T_0 \xrightarrow{\varphi_1} T_1 \xrightarrow{\varphi_2} \dots \xrightarrow{\varphi_s} T_s \xrightarrow{\varphi_{s+1}} T_{s+1} \xrightarrow{\varphi_{s+2}} \dots \xrightarrow{\varphi_n} T_n \quad (5.12)$$

NSTRING LAB EQ

together with l -labelings of T_s, T_{s+1}, \dots, T_n such that the $\varphi_r, r > s$ are λ_i -inert label maps.

Arrows in $\Omega_G^{n,s,\lambda}$ are quotients of strings $(\pi_r: T_r \rightarrow T'_r)$ such that $\pi_r, r \geq s$ are label maps.

Further, for any $s < 0$ or $n < s'$ we write

$$\Omega_G^{n,s,\lambda} = \Omega_G^{n,-1,\lambda}, \quad \Omega_G^{n,s',\lambda} = \Omega_G^n. \quad (5.13)$$

EXTRACASES EQ

Intuitively, $\Omega_G^{n,s,\lambda}$ consists of strings that are labeled in the range $s \leq r \leq n$, with the extra cases (5.13) interpreted by infinitely prepending and postpending copies of T_{-1} and T_n to (5.12).

The main case of interest is that of $s = 0$, which we abbreviate as $\Omega_G^{n,\lambda} = \Omega_G^{n,0,\lambda}$, with the remaining $\Omega_G^{n,s,\lambda}$ playing an auxiliary role. The $s = -1$ case also deserves special attention.

Remark 5.14. For $s < 0$ there are identifications

$$\Omega_G^{n,s,\lambda} = \Omega_G^{n,-1,\lambda} \simeq \coprod_{\lambda_a} \Omega_G^n \sqcup \coprod_{\lambda_i} \Sigma_G. \quad (5.15)$$

OMEGAN MINUS ONE EQ

Indeed, since T_{-1} is a G -corolla, the label of its unique G -vertex determines all other labels.

Notation 5.16. We will write $(\Omega_G^n)^{\times \lambda}$ to denote the l -tuple with $(\Omega_G^n)_j^{\times \lambda} = \Omega_G^n$ if $j \in \lambda_a$ and $(\Omega_G^n)_j^{\times \lambda} = \Sigma_G$ if $j \in \lambda_i$. As such, (5.15) is abbreviated as $\Omega_G^{n,-1,\lambda} = \coprod (\Omega_G^n)^{\times \lambda}$.

The $\Omega_G^{n,s,\lambda}$ categories are related by a number of obvious functors, which we now catalog.

Firstly, if $s \leq s'$ there are forgetful functors

$$\Omega_G^{n,s,\lambda} \rightarrow \Omega_G^{n,s',\lambda} \quad (5.17)$$

NKNFGT EQ

and the simplicial operators in Notation 3.87 generalize to operators (for $0 \leq i \leq n, -1 \leq j \leq n$)

$$\begin{aligned} d_i: \Omega_G^{n,s,\lambda} &\rightarrow \Omega_G^{n-1,s-1,\lambda} & i < s & & s_j: \Omega_G^{n,s,\lambda} &\rightarrow \Omega_G^{n+1,s+1,\lambda} & j < s \\ d_i: \Omega_G^{n,s,\lambda} &\rightarrow \Omega_G^{n-1,s,\lambda} & s \leq i & & s_j: \Omega_G^{n,s,\lambda} &\rightarrow \Omega_G^{n+1,s,\lambda} & s \leq j \end{aligned} \quad (5.18)$$

LABSTSIM EQ

which are compatible with the forgetful functors in the obvious way.

We will prefer to reorganize (5.17) and (5.18) somewhat. Defining functions $d_i: \mathbb{Z} \rightarrow \mathbb{Z}$ and $s_j: \mathbb{Z} \rightarrow \mathbb{Z}$ by

$$d_i(s) = \begin{cases} s-1, & i < s \\ s, & s \leq i \end{cases} \quad s_j(s) = \begin{cases} s+1, & j < s \\ s, & s \leq j \end{cases} \quad (5.19)$$

INTERMAPDEF EQ

LABSTSIM EQ
(5.18) is rewritten as maps $d_i: \Omega_G^{n,s,\lambda} \rightarrow \Omega_G^{n-1,d_i(s),\lambda}$ and $s_j: \Omega_G^{n,s,\lambda} \rightarrow \Omega_G^{n+1,s_j(s),\lambda}$. Therefore, we henceforth write simply $\Omega_G^{n,\bullet,\lambda}$ to denote the string of categories $\Omega_G^{n,s,\lambda}$ and forgetful functors, and abbreviate (5.18) as

$$d_i: \Omega_G^{n,\bullet,\lambda} \rightarrow \Omega_G^{n-1,\bullet,\lambda} \quad s_j: \Omega_G^{n,\bullet,\lambda} \rightarrow \Omega_G^{n+1,\bullet,\lambda} \quad (5.20)$$

LABSTSIM2 EQ

ORDLABEL REM

INTERMAPDEF EQ
Remark 5.21. Considering the ordered sets $\langle n \rangle = \{0 < 1 < \dots < n < +\infty\}$, the formulas (5.19) define functions $d_i: \langle n \rangle \rightarrow \langle n-1 \rangle$, $s_j: \langle n \rangle \rightarrow \langle n+1 \rangle$ which preserve 0 and $+\infty$, except for s_{-1} which preserves only $+\infty$. This recovers the description of Δ^{op} as the category of intervals (i.e. ordered finite sets with a minimum and maximum and maps preserving them).

Next, the vertex functors V_G^k of (3.98) generalize to functors $V_G^k: \Omega_G^{n,s,\lambda} \rightarrow \mathbf{F}_s \wr \Omega_G^{n-k-1,s-k-1,\lambda}$ given by the same formula

$$(T_k, v_{G_e} \rightarrow \dots \rightarrow T_n, v_{G_e})_{v_{G_e} \in V_G(T_k)}, \quad (5.22)$$

as in (3.98), except now with the T_m, v_{G_e} inheriting the node labels from T_m (if any).

The diagrams in (3.99) for $i < k$ and $i > k$ now generalize to diagrams

$$\begin{array}{ccc} \Omega_G^{n,\bullet,\lambda} & \xrightarrow{V_G^k} & \mathbf{F}_s \wr \Omega_G^{n-k-1,\bullet,\lambda} \\ d_i \downarrow & \swarrow \pi_i & \parallel \\ \Omega_G^{n-1,\bullet,\lambda} & \xrightarrow{V_G^{k-1}} & \mathbf{F}_s \wr \Omega_G^{n-k-1,\bullet,\lambda} \end{array} \quad \begin{array}{ccc} \Omega_G^{n,\bullet,\lambda} & \xrightarrow{V_G^k} & \mathbf{F}_s \wr \Omega_G^{n-k-1,\bullet,\lambda} \\ d_i \downarrow & & \downarrow d_{i-k-1} \\ \Omega_G^{n-1,\bullet,\lambda} & \xrightarrow{V_G^k} & \mathbf{F}_s \wr \Omega_G^{n-k-2,\bullet,\lambda} \end{array} \quad (5.23)$$

PIIDEFDILAB EQ

while the diagrams in (3.100) for $j < k$ and $j > k$ generalize to diagrams

$$\begin{array}{ccc} \Omega_G^{n,\bullet,\lambda} & \xrightarrow{V_G^k} & \mathbf{F}_s \wr \Omega_G^{n-k-1,\bullet,\lambda} \\ s_j \downarrow & & \parallel \\ \Omega_G^{n+1,\bullet,\lambda} & \xrightarrow{V_G^{k+1}} & \mathbf{F}_s \wr \Omega_G^{n-k-1,\bullet,\lambda} \end{array} \quad \begin{array}{ccc} \Omega_G^{n,\bullet,\lambda} & \xrightarrow{V_G^k} & \mathbf{F}_s \wr \Omega_G^{n-k-1,\bullet,\lambda} \\ s_j \downarrow & & \downarrow s_{j-k-1} \\ \Omega_G^{n+1,\bullet,\lambda} & \xrightarrow{V_G^k} & \mathbf{F}_s \wr \Omega_G^{n-k,\bullet,\lambda} \end{array} \quad (5.24)$$

PIIDEFDI2LAB EQ

where we note that in all cases the s -index \bullet varies according to (5.18).

Lastly, the $\Omega_G^{n,s,\lambda}$ are also functorial in λ . Explicitly, given $\alpha: \{1, \dots, l\} \rightarrow \{1, \dots, m\}$ and partitions such that $\lambda' \leq \alpha^* \lambda$ one has forgetful functors

$$\Omega_G^{n,s,\lambda'} \rightarrow \Omega_G^{n,s,\lambda} \quad (5.25)$$

LAMBINC EQ

compatible with the forgetful functors (5.17), the simplicial operators d_i, s_j and the isomorphisms π_i .

Remark 5.26. When α is the identity and $\lambda' \leq \lambda$ the forgetful functors (5.25) are fully faithful inclusions. However, this is not the case for the (5.17) forgetful functors. Indeed, regarding the map $T \rightarrow U$ in (5.10) as an object in $\Omega_G^{1,0,\lambda}$ for $\lambda = \lambda_a \sqcup \lambda_i = \{2\} \sqcup \{1\} = \{\bullet\} \sqcup \{\circ\}$, changing the label of $a_1 \leq a_2$ to a \bullet -label produces a non isomorphic object $\bar{T} \rightarrow U$ of $\Omega_G^{1,0,\lambda}$ that forgets to the same object of $\Omega_G^{1,1,\lambda}$.

We now extend Notation 4.7.

Notation 5.27. Let $(A_j) = (A_j \rightarrow \Sigma_G)_{1 \leq j \leq l}$ be a l -tuple of maps over Σ_G . We define $\Omega_G^{n,s,\lambda} \wr (A_j)$ as the pullback

$$\begin{array}{ccc} \Omega_G^{n,s,\lambda} \wr (A_j) & \xrightarrow{V_G^n} & \mathbf{F} \wr \coprod A_j \\ \downarrow & & \downarrow \\ & & \mathbf{F} \wr \coprod \Sigma_G \\ \downarrow & & \downarrow \\ \Omega_G^{n,s,\lambda} & \xrightarrow{V_G^n} & \mathbf{F} \wr \Omega_G^{-1,s-n-1,\lambda} \end{array} \quad (5.28)$$

OMEGAWRTUP EQ

Remark 5.29. To unpack $\Omega_G^{n,s,\lambda} \wr (A_j)$, note first that by (5.13) $\Omega_G^{n,s,\lambda}$ is simply either Σ_G^{ul} if $r < 0$ or Σ_G for $r \geq 0$. We can thus break down (5.28) into the three cases $s < 0$, $0 \leq s \leq n$ and $n < s$, depicted below.

$$\begin{array}{ccccc}
\Omega_G^{n,s,\lambda} \wr (A_j) & \xrightarrow{V_G^n} & F \wr \coprod_j A_j & & \Omega_G^{n,s,\lambda} \wr (A_j) & \xrightarrow{V_G^n} & F \wr \coprod_j A_j & & \Omega_G^{n,s,\lambda} \wr (A_j) & \xrightarrow{V_G^n} & F \wr \coprod_j A_j \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
\coprod (\Omega_G^n)^{\times \lambda} & \xrightarrow{V_G^n} & F \wr \coprod_l \Sigma_G & & \Omega_G^{n,s,\lambda} & \xrightarrow{V_G^n} & F \wr \coprod_l \Sigma_G & & \Omega_G^n & \xrightarrow{V_G^n} & F \wr \Sigma_G
\end{array}$$

Therefore, for $s > n$ (5.28) coincides with $\Omega_G^n \wr (\coprod_j A_j)$ as defined in Notation 4.7. Moreover, for $s < 0$ both squares in the diagram below are pullbacks and the bottom composite is V_G^n ,

$$\begin{array}{ccccc}
\coprod (\Omega_G^n)^{\times \lambda} \wr (A_j) & \xrightarrow{\coprod (V_G^n)^{\times \lambda}} & \coprod F \wr A_j & \longrightarrow & F \wr \coprod_j A_j \\
\downarrow & & \downarrow & & \downarrow \\
\coprod (\Omega_G^n)^{\times \lambda} & \xrightarrow{\coprod (V_G^n)^{\times \lambda}} & \coprod_l F \wr \Sigma_G & \longrightarrow & F \wr \coprod_l \Sigma_G
\end{array} \tag{5.31}$$

BOTTOM EQ

so that there is an identification $\Omega_G^{n,s,\lambda} \wr (A_j) \simeq \coprod (\Omega_G^n)^{\times \lambda} \wr (A_j)$, where in the right side $(-) \wr (-)$ is computed entry-wise.

ORDLABEL REM

Remark 5.32. The naturality of the $\Omega_G^{n,s,\lambda} \wr (A_j)$ constructions with regards to λ interacts with the tuple (A_j) in the obvious way, i.e., given $\alpha: \{1, \dots, l\} \rightarrow \{1, \dots, m\}$, $\lambda' \leq \alpha^* \lambda$ and a map $(B_k) \rightarrow \alpha^*(A_j)$ one obtains a natural map

$$\Omega_G^{n,s,\lambda'} \wr (B_k) \rightarrow \Omega_G^{n,s,\lambda} \wr (A_j).$$

Proposition 5.33. The analogue statements of Proposition 3.101 hold for the $\Omega_G^{n,s,\lambda}$ and the $\Omega_G^{n,s,\lambda} \wr (A_j)$ constructions, where in the latter case we exclude the statements involving d_n .

Additionally, the natural squares (for $n \geq -1$)

$$\begin{array}{ccc}
\Omega_G^{n,n,\lambda} & \xrightarrow{V_G^n} & F \wr \coprod_l \Sigma_G \\
\downarrow & & \downarrow \\
\Omega_G^n & \xrightarrow{V_G^n} & F \wr \Sigma_G
\end{array} \tag{5.34}$$

ADDSQUARE EQ

are also pullback squares.

Proof. Firstly, we note that the $\Omega_G^{n,s,\lambda}$ analogues, as well as the claim for (5.34), all follow by keeping track of the labels on the strings, with the only part worthy of note being the analogue of (d), stating that the right squares in (5.23) and (5.24) are pullbacks. Since in these diagrams the s -coordinate \bullet is determined by the top left corner, a direct analysis shows that compatible choices of labels for strings on the top right and bottom left corners do assemble to the correct labels on the top left corner, so that the claim follows by the unlabeled one.

For the more general $\Omega_G^{n,s,\lambda} \wr (A_j)$ constructions, one can either build the general V_G^k , d_i , s_j , π_i explicitly, or simply mimic the argument in Proposition 4.13, thereby reducing to the $\Omega_G^{n,s,\lambda}$ case. \square

LABIDEN COR

Corollary 5.35. For $-1 \leq s \leq n$ there are natural identifications

$$\Omega_G^k \wr \Omega_G^{n,s,\lambda} \wr (A_j) \simeq \Omega_G^{n+k+1,s+k+1,\lambda} \wr (A_j) \quad \Omega_G^{n,s,\lambda} \wr (\Omega_G^k)^{\times \lambda} \wr (A_j) \simeq \Omega_G^{n+k+1,s,\lambda} \wr (A_j)$$

which identify $V_G^k \wr \Omega_G^{n,s,\lambda} \wr (A_j)$ with $V_G^k \wr (A_j)$ and $V_G^n \wr (\Omega_G^k)^{\times \lambda} \wr (A_j)$ with $V_G^n \wr (A_j)$.

Further, these identifications are compatible with each other and associative in the obvious ways, and they induce identifications

$$\begin{aligned} d_i \wr (\Omega_G^n)^{\times \lambda} &\simeq d_i & \pi_i \wr (\Omega_G^n)^{\times \lambda} &\simeq \pi_i & s_j \wr (\Omega_G^n)^{\times \lambda} &\simeq s_j \\ \Omega_G^k \wr d_i &\simeq d_{i+k+1} & \Omega_G^k \wr \pi_i &\simeq \pi_{i+k+1} & \Omega_G^k \wr s_j &\simeq s_{j+k+1} \end{aligned}$$

as well as obvious identifications of forgetful functors.

Proof. This is analogous to Corollary 4.19. ^{**IDEN COR**} For the first identification, the case $s \geq 0$ follows from the diagram below, where we note that the bottom arrow is $V_G^k: \Omega_G^k \rightarrow F \wr \Sigma_G$.

$$\begin{array}{ccccccc} \bullet & \xrightarrow{\quad \quad \quad} & \bullet & \xrightarrow{\quad \quad \quad} & F^{i2} \wr \coprod (A_j) & \xrightarrow{\sigma^0} & F \wr \coprod (A_j) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_G^{n+k+1, s+k+1, \lambda} & \xrightarrow{V_G^k} & F \wr \Omega_G^{n, s, \lambda} & \xrightarrow{F \wr V_G^n} & F^{i2} \wr \coprod_l \Sigma_G & \xrightarrow{\sigma^0} & F \wr \coprod_l \Sigma_G \\ \downarrow & & \downarrow & & & & \\ \Omega_G^{k, k+1, \lambda} & \xrightarrow{V_G^k} & F \wr \Omega_G^{-1, 0, \lambda} & & & & \end{array}$$

In the $s = -1$ case, the bottom arrow is instead $V_G^k: \Omega_G^{k, k, \lambda} \rightarrow F \wr \Omega_G^{-1, -1, \lambda} = F \wr \coprod_l \Sigma_G$, in which case one further attaches ^{**ADDSQUARE EQ**} (5.34) to the diagram.

The second identification is analogous, using the pullback diagram below, with the composite of the central horizontal arrows reinterpreted using ^{**BOTTOM EQ**} (5.31).

$$\begin{array}{ccccccc} \bullet & \xrightarrow{\quad \quad \quad} & \bullet & \xrightarrow{\quad \quad \quad} & F \wr \coprod F \wr A_j & \xrightarrow{\quad \quad \quad} & F^{i2} \wr \coprod A_j \xrightarrow{\sigma^0} F \wr \coprod A_j \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_G^{n+k+1, s, \lambda} & \xrightarrow{V_G^n} & F \wr \coprod (\Omega_G^k)^{\times \lambda} & \xrightarrow{F \wr \coprod (V_G^n)^{\times \lambda}} & F \wr \coprod_l F \wr \Sigma_G & \xrightarrow{\quad \quad \quad} & F^{i2} \wr \coprod \Sigma_G \xrightarrow{\sigma^0} F \wr \coprod_l \Sigma_G \\ \downarrow & & \downarrow & & & & \\ \Omega_G^{n, s, \lambda} & \xrightarrow{V_G^n} & F \wr \coprod_l \Sigma_G & & & & \end{array}$$

The additional claims are straightforward. \square

Remark 5.36. The identifications in Corollary 5.35 ^{**LABIDEN COR**} allow for the case $n = -1$, which is non-trivial due to the existence of $\Omega_G^{-1, -1, \lambda} = \coprod_l \Sigma_G$, in which case $\Omega_G^{-1, -1, \lambda} \wr (A_j) \simeq \coprod A_j$. For $-1 \leq s \leq n$ the identifications

$$\Omega_G^{n, s, \lambda} = \Omega_G^s \wr \Omega_G^{-1, -1} \wr (\Omega_G^{n-s-1})^{\times \lambda}$$

then show that $\Omega_G^{n, s, \lambda} \wr (-)$ encodes (the underlying category of) the functor $N^{\circ s+1} \coprod (N^{\times \lambda})^{\circ n-s}$. Furthermore, the left commutative square below, where vertical arrows are forgetful functors (and the right diagram merely unpacks notation)

$$\begin{array}{ccc} \Omega_G^{0, -1, \lambda} & \xrightarrow{\coprod (V_G^0)^{\times \lambda}} & \coprod F \wr (\Omega_G^{-1})^{\times \lambda} \longrightarrow F \wr \Omega_G^{-1, -2, \lambda} \\ \downarrow & & \parallel \\ \Omega_G^{0, 0, \lambda} & \xrightarrow{V_G^0} & F \wr \Omega_G^{-1, -1, \lambda} \\ \downarrow & & \downarrow \\ \Omega_G^{0, 1, \lambda} & \xrightarrow{V_G^0} & F \wr \Omega_G^{-1, 0, \lambda} \end{array} \quad \begin{array}{ccc} \coprod (\Omega_G^0)^{\times \lambda} & \longrightarrow & \coprod F \wr \Sigma_G \\ \downarrow & & \downarrow \\ \Omega_G^{0, 0, \lambda} & \longrightarrow & F \wr \coprod \Sigma_G \\ \downarrow & & \downarrow \\ \Omega_G^0 & \longrightarrow & F \wr \Sigma_G \end{array} \quad (5.37)$$

NATCOP EQ

shows that the forgetful functor $\Omega_G^{0, -1, \lambda} \wr (A_j) \rightarrow \Omega_G^{0, 0, \lambda} \wr (A_j)$ encodes the natural map $\coprod \circ N \Rightarrow N \circ \coprod$.

5.2 The category of extension trees

The purpose of this section is to make (5.7) explicit. We start by discussing realizations of simplicial objects in \mathbf{Cat} .

Recalling the standard cosimplicial object $[\bullet] \in \mathbf{Cat}^\Delta$ given by $[n] = (0 \rightarrow 1 \rightarrow \dots \rightarrow n)$ yields the following definition.

Definition 5.38. The left adjoint below is called the *realization* functor.

$$|-| : \mathbf{Cat}^{\Delta^{op}} \rightleftarrows \mathbf{Cat} : (-)^{[\bullet]}$$

Remark 5.39. Suppose that $\mathcal{C} \in \mathbf{Cat}$ contains subcategories $\mathcal{C}_h, \mathcal{C}^v$ whose arrows span those of \mathcal{C} . Defining $\mathcal{C}_{h,\bullet}^v \in \mathbf{Cat}^{\Delta^{op}}$ so that the objects of $\mathcal{C}_{h,n}^v$ are n -strings in \mathcal{C}_h and the arrows are compatible n -tuples of arrows in \mathcal{C}^v , it is straightforward to show that it is $|\mathcal{C}_{h,\bullet}^v| = \mathcal{C}$.

An immediate example is given by the planar strings in Definition 3.84. Writing $\mathcal{C} = \Omega_G^t$ the category of tall maps, $\mathcal{C}_h = \Omega_G^{pt}$ the category of planar tall maps and $\mathcal{C}^v = \Omega_G^0$ the category of quotients, one has $\mathcal{C}_{h,\bullet}^v = \Omega_G^n$ and thus $|\Omega_G^n| = \Omega_G^t$.

Similarly, noting that the $\Omega_G^{n,\lambda} = \Omega_G^{n,0,\lambda}$ form a simplicial object, we have that the $|\Omega_G^{n,\lambda}| = \Omega_G^{t,\lambda}$ is the category of tall maps between l -labeled trees that induce quotients on nodes with λ -inert labels.

In the following statement, we note that it is shown in Lemma A.3 that $ob(|A_\bullet|) \simeq ob(A_0)$ and that arrows in $|A_\bullet|$ are generated by the arrows in A_0 together with arrows $d_1(a) \rightarrow d_0(a)$ for each $a \in A_1$.

Proposition 5.40. Given a simplicial object $\Sigma_G \leftarrow A_\bullet \xrightarrow{N} \mathcal{V}^{op}$ in $\mathbf{WSpan}^r(\Sigma_G, \mathcal{V}^{op})$ such that the natural transformation components of the differential operators d_i , $0 \leq i < n$ and s_j , $0 \leq j \leq n$ are isomorphisms, there is an identification

$$\lim_{\Delta} (\mathbf{Ran}_{A_n \rightarrow \Sigma_G} N_n) \simeq \mathbf{Ran}_{|A_\bullet| \rightarrow \Sigma_G} \tilde{N}$$

where $\tilde{N} : |A_\bullet| \rightarrow \mathcal{V}^{op}$ is given by N_0 on objects and generating arrows in A_0 , and on generating arrows $d_1(a) \rightarrow d_0(a)$ for $a \in A_1$ as the composite

$$\begin{array}{ccccc} A_0 & & \xleftarrow{d_1} & A_1 & \xrightarrow{d_0} & A_0 \\ & \searrow & & \downarrow & \swarrow & \\ & & & \mathcal{V}^{op} & & \end{array}$$

Proposition 5.40 applies to both directions of the bisimplicial object $N(N^{on} \iota \mathcal{P} \sqcup \iota X^{\sqcup 2l+1}) \sqcup Y$ in (5.6). Indeed, in the n direction all d_i with $0 < i < n$ are induced by the multiplication $NN \rightarrow N$ defined in (4.21) while d_0 is induced by the composite $N \circ \sqcup \circ N \rightarrow NN \circ \sqcup \rightarrow N \circ \sqcup$, with the second map again given by composition and the first induced by the natural map $\sqcup \circ N \rightarrow N \circ \sqcup$, which is encoded by a strictly commutative diagram of spans, as seen using the top part of (5.37) (or, more abstractly, it also suffices to note that N preserves arrows in $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V})$ given by strictly commutative diagrams). Degeneracies are similar.

As for the l direction, we note that our convention on the double bar construction $B_l(\mathcal{P}, \mathbb{F}X, \mathbb{F}X, \mathbb{F}Y)$, is symmetric, with d_l encoding both of the maps $\mathbb{F}X \rightarrow \mathbb{F}Y$ and $\mathbb{F}X \rightarrow \mathbb{F}\mathcal{P}$ and the remaining differentials given by fold maps. Or, more precisely, the action of the differential operators on the sets of labels $\langle \langle l \rangle \rangle = \{-\infty, -l, \dots, -1, 0, 1, \dots, l, +\infty\}$ is given by extending the functions in Remark 5.32 anti-symmetrically. But then the differential operators d_i , s_j for $0 \leq i < l$ and $0 \leq j \leq l$ correspond to instances of the naturality in Remark 5.32 when $(B_k) = \alpha^*(A_j)$, and are hence given by strictly commutative maps of spans.

Our next task is thus that of identifying the category of extension trees Ω_G^e appearing in (5.7), i.e. to produce an explicit model for the double realization $|\Omega_G^{n,\lambda_l}|$. By Remark

REALEX REM


5.39 we can first perform the realization in the n direction, so as to obtain $|\Omega_G^{n,\lambda_l}| = |\Omega_G^{t,\lambda_l}|$, where we recall that Ω_G^{t,λ_l} consists of $\langle l \rangle$ -labelled trees together with tall maps that induce quotients on all nodes not labeled by $-\infty$.

We now identify Ω_G^e directly.

Definition 5.41. The *extension tree category* Ω_G^e is the category whose objects are $\{\mathcal{P}, X, Y\}$ -labeled trees and whose maps $\varphi: T \rightarrow S$ are tall maps of trees such that

- (i) if $T_{v_{Ge}}$ has a X -label, then $S_{v_{Ge}} = T_{v_{Ge}}$ and $S_{v_{Ge}}$ has a X -label;
- (ii) if $T_{v_{Ge}}$ has a Y -label, then $S_{v_{Ge}} = T_{v_{Ge}}$ and $S_{v_{Ge}}$ has either an X -label or a Y -label;
- (iii) if $T_{v_{Ge}}$ has a \mathcal{P} -label, then $S_{v_{Ge}}$ has only X and \mathcal{P} -labels.

Example 5.42. The following is an example of a planar map in Ω_G^e for $G = *$, where black nodes represent \mathcal{P} -labeled nodes, grey nodes represent Y -labeled nodes and white nodes represent X -labeled nodes.



REGALTERNMAP EQ

Remark 5.44. By changing any X -labels in $S_{v_{Ge}}$ into Y -labels (resp. \mathcal{P} -labels) whenever $T_{v_{Ge}}$ has a Y -label (resp. \mathcal{P} -label), one obtains a factorization

$$T \rightarrow \bar{S} \rightarrow S \quad (5.45)$$

LABRE EQ

such that $T \rightarrow \bar{S}$ is a label map (cf. Definition 5.8) and $\bar{S} \rightarrow S$ is an underlying identity of trees that merely changes some of the Y and \mathcal{P} labels into X labels. We refer to the latter kind of map as a *relabel map*. It is clear that the label-relabel factorization (5.45) is unique.

Proposition 5.46. *There is an identification*

$$\Omega_G^e \simeq |\Omega_G^{t,\lambda_l}|.$$

Proof. We will show that Remark 5.39 applies to $\mathcal{C} = \Omega_G^e$, with \mathcal{C}_h and \mathcal{C}^v the categories of relabel and label maps. More precisely, we claim that there is an isomorphism $\mathcal{C}_{h,l}^v \simeq \Omega_G^{t,\lambda_l}$ of objects in $\text{Cat}^{\Delta^{op}}$. Unpacking notation, one must first show that strings

$$T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_l \quad (5.47)$$

RELABSTR EQ

of relabel arrows in Ω_G^e are in bijection with objects of Ω_G^{t,λ_l} , i.e. with trees labeled by $\langle l \rangle = \{-\infty, -l, \dots, -1, 0, 1, \dots, l, +\infty\}$. Noting that the maps in (5.47) are simply underlying identities on some fixed tree T that convert some of the \mathcal{P}, Y labels into X labels, we label a vertex $T_{v_{Ge}}$ by (i) $0 < j \leq +\infty$ if the last j labels of $T_{v_{Ge}}$ in (5.47) are Y labels (where $+\infty = l + 1$); (ii) $-\infty \leq -j < 0$ if the last j labels of $T_{v_{Ge}}$ in (5.47) are \mathcal{P} labels; (iii) $j = 0$ if all labels in (5.47) are X labels. This process clearly established the desired bijection on objects.

The compatibilities with arrows and with the simplicial structure are straightforward. \square

Our next task will be that of identifying a convenient initial subcategory $\widehat{\Omega}_G^e \hookrightarrow \Omega_G^e$. We first introduce the auxiliary notion of alternating trees. Recall the notion of input path (Notation 3.5) $I(e) = \{f \in T : e \leq_d f\}$ for an edge $e \in T$, which naturally extends to T in any of $\in \Omega, \Phi, \Omega_G, \Phi_G$.

OMEGAA DEF

Definition 5.48. A G -tree $T \in \Omega_G$ is called *alternating* if, for all leaves $l \in T$ one has that the input path $I(l)$ has an even number of elements.


Further, a vertex $e^\dagger \leq e$ is called *active* if $|I(e)|$ is odd and *inert* otherwise.

Finally, a tall map $T \xrightarrow{\varphi} S$ between alternating G -trees is called a *tall alternating map* if for any inert vertex $e^\dagger \leq e$ of T one has that $S_{e^\dagger \leq e}$ is an inert vertex of S .

We will denote the category of alternating G -trees and tall alternating maps by Ω_G^a .

Remark 5.49. A G -tree (resp. map of G -trees) is alternating (resp. an alternating map) iff each component is.

Example 5.50. Two alternating trees (for $G = *$ the trivial group) and a planar tall alternating map between them follow, with active nodes in black (\bullet) and white nodes in white (\circ).



REGALTERNMAPLR EQ

The term “alternating” reflects the fact that adjacent nodes have different colors, though there is an additional restriction: the “outer vertices”, i.e. those immediately below a leaf or above the root, are necessarily black/active (this does not, however, apply to stumps).

ALTSUB REM

Remark 5.52. If $T \in \Omega$ is alternating, it follows from Remark 5.54 that a tall map $\varphi: T \rightarrow U$ is an alternating map iff the corresponding substitution datum under Proposition 3.47 is given by the identity $U_{e^\dagger \leq e} = T_{e^\dagger \leq e}$ when $e^\dagger \leq e$ is inert and by an alternating tree $U_{e^\dagger \leq e}$ when $e^\dagger \leq e$ is active.

Definition 5.53. $\widehat{\Omega}_G^e \hookrightarrow \Omega_G^e$ is the full subcategory of (\mathcal{P}, X, Y) -labeled trees whose underlying tree is alternating, active nodes are labeled by \mathcal{P} , and passive nodes are labeled by X or Y .

Note that conditions (i) and (ii) in Definition 5.41 imply that for any map in $\widehat{\Omega}_G^e$ the underlying map is an alternating map.

The following is the key to establishing the desired initiality of $\widehat{\Omega}_G^e$ in Ω_G^e .

LXP PROP

Proposition 5.54. For each $U \in \Omega_G^e$ there exists a unique $\text{lr}_{\mathcal{P}}(U) \in \widehat{\Omega}_G^e$ together with a unique planar label map in Ω_G^e

$$\text{lr}_{\mathcal{P}}(U) \rightarrow U.$$

Furthermore, $\text{lr}_{\mathcal{P}}$ extends to a right retraction $\text{lr}_{\mathcal{P}}: \Omega_G^e \rightarrow \widehat{\Omega}_G^e$.

Proof. We first address the non-equivariant case $U \in \Omega^e$.

To build $\text{lr}_{\mathcal{P}}(U)$, consider the collection of outer faces $\{U_i^X\} \sqcup \{U_j^Y\} \sqcup \{U_k^{\mathcal{P}}\}$ where the U_i^X, U_j^Y are simply the X, Y -labeled nodes and the $\{U_k^{\mathcal{P}}\}$ are the maximal outer subtrees whose nodes have only \mathcal{P} -labels (these may possibly be sticks). Lemma 3.57 guarantees that each edge and each \mathcal{P} -labeled node belong to exactly one of the $V_G(U_k^X)$, and applying Proposition 3.55(iii) yields a planar tall map

$$T = \text{lr}_{\mathcal{P}}(U) \rightarrow U \tag{5.55}$$

LRXDEF EQ

such that $\{U_{e^\dagger \leq e}\}_{(e^\dagger \leq e) \in V(T)} = \{U_i^X\} \sqcup \{U_j^Y\} \sqcup \{U_k^P\}$. T has an obvious (\mathcal{P}, X, Y) -labeling making (5.55) into a label map, but we must still check $T \in \widehat{\Omega}_G^e$, i.e. that T is alternating with active vertices precisely those labeled by \mathcal{P} . But since the image of each $e \in T$ belongs to precisely one U_k^P , e belongs to precisely one of the \mathcal{P} -labeled nodes of T , so that any leaf input path $I(l) = (l = e_n \leq e_{n-1} \leq \dots \leq e_1 \leq e_0)$ must start with, end with, and alternate between \mathcal{P} -nodes, and thus have even length.

To check uniqueness, note that for any other planar label map $S \rightarrow U$ with S alternating and $e^\dagger \leq e$ a \mathcal{P} vertex of S the outer face $U_{e^\dagger \leq e}$ must be a maximal \mathcal{P} -labeled outer face since the vertices adjacent to its root and leaves are labeled by either X or Y . The condition $V(U) = \coprod_{V(S)} V(U_{e^\dagger \leq e})$ thus guarantees that the collection of outer faces determined by S matches that determined by T except perhaps in the number of stick faces, so that the degeneracy-face factorizations $S \rightarrow F \rightarrow U$, $T \rightarrow F \rightarrow U$ factor through the same planar inner face F , with the unique labeling that makes the inclusion a label map. S, T are thus both trees in $\widehat{\Omega}_G^e$ obtained from F by adding degenerate \mathcal{P} vertices, and since this can be done in at most one way, we conclude $S = T$.


To check functoriality, consider the diagram below, where $T \rightarrow U$ is the map defined above and $\varphi: U \rightarrow V$ any map in Ω_G^e .

$$\begin{array}{ccc} T & \longrightarrow & U \\ \downarrow & & \downarrow \varphi \\ S & \dashrightarrow & V \end{array} \quad (5.56) \quad \boxed{\text{LRPFUN EQ}}$$

The composite $T \rightarrow V$ is encoded by a substitution datum $\{T_{e^\dagger \leq e} \rightarrow V_{e^\dagger \leq e}\}$ which is given by an isomorphism if $e^\dagger \leq e$ has label X or Y (possibly changing a Y label to a X label), and by some X, \mathcal{P} -labeled tree $V_{e^\dagger \leq e}$ if $e^\dagger \leq e$ has a \mathcal{P} -label. We now consider the factorization problem in (5.56), where we want $S \in \widehat{\Omega}_G^e$ and for the map $S \rightarrow V$ to be a planar label map. Combining Remark 5.52 with the uniqueness of the $\text{lr}_{\mathcal{P}}(V_{e^\dagger \leq e})$, the only possibility is for S to be defined using the T substitution datum that replaces $T_{e^\dagger \leq e} \rightarrow V_{e^\dagger \leq e}$ with $T_{e^\dagger \leq e} \rightarrow \text{lr}_{\mathcal{P}}(V_{e^\dagger \leq e})$ whenever $e^\dagger \leq e$ has a \mathcal{P} -label. Uniqueness of $\text{lr}_{\mathcal{P}}(V)$ then implies $S = \text{lr}_{\mathcal{P}}(V)$, and one sets $\text{lr}_{\mathcal{P}}(\varphi)$ to be the map $T \rightarrow S$. Associativity and unitality are automatic from the uniqueness of the factorization of (5.56).

In the case of $T = (T_x)_{x \in X}$ in Ω_G^e for a general group G , one sets $\text{lr}_{\mathcal{P}}(T) = (\text{lr}_{\mathcal{P}}(T_x))_{x \in X}$. \square

Example 5.57. The following illustrates the $\text{lr}_{\mathcal{P}}$ construction when applied to the map φ in (5.43). Intuitively, the functor $\text{lr}_{\mathcal{P}}$ replaces each of the maximal *mathcal{P}*-labeled subtrees T_k^P, S_k^P with the corresponding leaf-root $\text{lr}(T_k^P), \text{lr}(S_k^P)$, which is then \mathcal{P} -labeled.



(5.58)

KANRED COR

Corollary 5.59. The inclusion $\widehat{\Omega}_G^e \hookrightarrow \Omega_G^e$ is *Ran*-initial over Σ_G . I.e., for \mathcal{C} any a complete category and functor $N: \Omega_G^e \rightarrow \mathcal{C}$ it is

$$\text{Ran}_{\Omega_G^e \rightarrow \Sigma_G} N \simeq \text{Ran}_{\widehat{\Omega}_G^e \rightarrow \Sigma_G} N.$$

Proof. Since $\text{lr}_{\mathcal{P}}$ is a right retraction over Σ_G , the undercategories $C \downarrow \widehat{\Omega}_G^e$ are right retractions of $C \downarrow \Omega_G^e$ for any $C \in \Sigma_G$. \square

5.3 Filtrations of free extensions

Summarizing the previous section, the discussion following Proposition 5.40 establishes (5.7), and hence Corollary 5.59 gives the alternate formula

$$\mathcal{P}[u] \simeq \mathcal{P} \coprod_{\mathbb{F}X} \mathbb{F}Y \simeq \text{Lan}_{(\widehat{\Omega}_G^e \rightarrow \Sigma_G)^{op}} \tilde{N}^{(\mathcal{P}, X, Y)}, \quad (5.60)$$

which we will now use to filter the map $\mathcal{P} \rightarrow \mathcal{P}[u]$ in the underlying category $\text{Sym}_G(\mathcal{V})$.

First, given $T = (T_i)_{i \in I} \in \Omega_G^e$, we write $V^X(T_i)$ (resp. $V^Y(T_i)$) to denote the set of X -labeled (Y -labeled) vertices of T_i . We define *degrees* of T by

$$|T|_X = |V^X(T_i)|, \quad |T|_Y = |V^Y(T_i)|, \quad |T| = |T|_X + |T|_Y,$$

which we note do not depend on the choice of $i \in I$.

Similarly, for $T = (T_i)_{i \in I} \in \Omega_G^a$ we write $V^{in}(T_i)$ for the inert vertices and $|T| = |V^{in}(T_i)|$.

Remark 5.61. One key property of the degrees $|T|$, $|T|_X$, $|T|_Y$ is that they are invariant under root pullbacks, which are defined by generalizing Definition 3.24 in the obvious way.

Definition 5.62. We specify some rooted (i.e. closed under root pullbacks) full subcategories of $\widehat{\Omega}_G^e$:

- (i) $\widehat{\Omega}_G^e[\leq k]$ (resp. $\widehat{\Omega}_G^e[k]$) is the subcategory of T with $|T| \leq k$ ($|T| = k$);
- (ii) $\widehat{\Omega}_G^e[\leq k \setminus Y]$ (resp. $\widehat{\Omega}_G^e[k \setminus Y]$) is the subcategory of $\widehat{\Omega}_G^e[\leq k]$ ($\widehat{\Omega}_G^e[k]$) of T with $|T|_Y \neq k$.

Similarly, we define subcategories $\Omega_G^a[\leq k]$, $\Omega_G^a[k]$ of Ω_G^a by the conditions $|T| \leq k$, $|T| = k$.

Remark 5.63. The categories $\widehat{\Omega}_G^e[k]$, $\widehat{\Omega}_G^e[k \setminus Y]$ and $\Omega_G^a[k]$ have rather limited morphisms.

Indeed, it is clear from Definitions 5.41 and 5.48 that maps never lower degree, and Remark 5.52 further ensures that degree is preserved iff \mathcal{P} -vertices are substituted by \mathcal{P} -vertices (rather than larger trees which would necessarily have inert vertices, thus increasing degree). Therefore, all maps in $\Omega_G^a[k]$ are quotients while maps in $\widehat{\Omega}_G^e[k]$, $\widehat{\Omega}_G^e[k \setminus Y]$ are underlying quotients of G -trees that relabel some Y -vertices to X -vertices. Moreover, this can be repackaged as saying that the diagonal forgetful functors in

$$\begin{array}{ccc} \widehat{\Omega}_G^e[k \setminus Y] & \hookrightarrow & \widehat{\Omega}_G^e[k] \\ & \searrow & \swarrow \\ & \Omega_G^a[k] & \end{array}$$

are Grothendieck fibrations whose fibers over $T \in \Omega_G^a[k]$ are the punctured cube and cube categories

$$(Y \rightarrow X)^{\times V_G^{in}(T)} - Y^{\times V_G^{in}(T)}, \quad (Y \rightarrow X)^{\times V_G^{in}(T)}$$

for $V_G^{in}(T)$ the set of inert G -vertices.


Note that though $|V^{in}(T_i)| = k$ for each T_i composing $T = (T_i)_{i \in I}$, one can only guarantee $|V_G^{in}(T)| \leq k$.

Lemma 5.64. $\widehat{\Omega}_G^e[\leq k - 1]$ is *Ran-initial* in $\widehat{\Omega}_G^e[\leq k \setminus Y]$ over Σ_G .

The proof will make use of an additional construction on Ω_G^e : given $T \in \Omega_G^e$ let $T_{\mathcal{P}}$ denote the result of replacing all X -labeled nodes of T with \mathcal{P} -labeled nodes.

Remark 5.65. In contrast to the functor $\text{lr}_{\mathcal{P}}: \Omega_G^e \rightarrow \widehat{\Omega}_G^e$ of Proposition 5.54, the $(-)^{\mathcal{P}}$ construction does not define a full functor $\Omega_G^e \rightarrow \Omega_G^e$, instead being functorial, and the obvious maps $T_{\mathcal{P}} \rightarrow T$ natural, only with respect to the Y -inert maps of Ω_G^e .

Example 5.66. Combining the $(-)_\mathcal{P}$ and $\text{lr}_\mathcal{P}$ constructions one obtains a construction sending trees in $\widehat{\Omega}_G^e$ to trees in $\widehat{\Omega}_G^e$. We illustrate this for the tree $T \in \widehat{\Omega}^e$ below (so that $G = *$), where black nodes are \mathcal{P} -labeled, white nodes are X -labeled, and grey nodes are Y -labeled.



Proof of Lemma 5.64. By Proposition 2.5 it suffices to show that for each $C \in \Sigma_G$ the map of rooted undercategories

$$C \downarrow_r \widehat{\Omega}_G^e[\leq k-1] \rightarrow C \downarrow_r \widehat{\Omega}_G^e[\leq k \setminus Y]$$

is initial, i.e. (cf. (McL, X.3.1)) that for each $(S, \pi: C \rightarrow \text{lr}(S))$ in $C \downarrow_r \widehat{\Omega}_G^e[\leq k \setminus Y]$ the overcategory

$$(C \downarrow_r \widehat{\Omega}_G^e[\leq k-1]) \downarrow (S, \pi) \quad (5.67)$$

UNDERCATPR EQ

is non-empty and connected. By definition of rooted undercategory, π is the identity on roots and thus an isomorphism on Σ_G , so that objects of (5.67) correspond to maps $T \rightarrow S$ that induce a rooted isomorphism on lr , i.e. rooted tall maps.

The case $S \in \widehat{\Omega}_G^e[\leq k-1]$ is immediate, since then the identity $S = S$ is terminal in (5.67). Otherwise, since $|S|_k \neq k$ we have $|\text{lr}_\mathcal{P}(S_\mathcal{P})| < k$ and the map $\text{lr}_\mathcal{P}(S_\mathcal{P}) \rightarrow S$, which is a rooted tall, shows that (5.67) is indeed non-empty.

Now, consider a rooted tall map $T \rightarrow S$ with $T \in \widehat{\Omega}_G^e[\leq k-1]$. One can form a diagram

$$\begin{array}{ccccc} & & S & \longleftarrow & \text{lr}_\mathcal{P}(S_\mathcal{P}) \\ & \nearrow & \uparrow Y\text{-inert} & & \uparrow \\ T & \longrightarrow & T' & \longleftarrow & \text{lr}_\mathcal{P}(T'_\mathcal{P}) \end{array} \quad (5.68)$$

K-1LANFINAL EQ

where $T \rightarrow T' \rightarrow S$ is the natural factorization such that $T' \rightarrow S$ is Y -inert, i.e., T' is obtained from T by simply relabeling to X those Y -labeled vertices of T that become X -vertices in S . Note that by Remark 5.65 the existence of the right square relies on $T' \rightarrow S$ being Y -inert. Since all maps in (5.68) are rooted tall, this produces the necessary zigzag connecting the objects $T \rightarrow S$ and $\text{lr}_\mathcal{P}(S_\mathcal{P}) \rightarrow S$ in the category (5.67), finishing the proof. \square

In what follows we write $\tilde{N}: \widehat{\Omega}_G^{e,op} \rightarrow \mathcal{V}$ for the functor in (5.60) and any of its restrictions. We are now in a position to produce the filtration (5.2) of the map $\mathcal{P} \rightarrow \mathcal{P}[u]$ in (5.1).

PK_DEFN

Definition 5.69. \mathcal{P}_k is the left Kan extension

$$\begin{array}{ccc} \widehat{\Omega}_G^e[\leq k]^{op} & \xrightarrow{\tilde{N}} & \mathcal{V} \\ \downarrow \text{lr} & \Downarrow & \uparrow \mathcal{P}_k \\ \Sigma_G^{op} & & \end{array}$$

Noting that $\widehat{\Omega}_G^e[\leq 0] \simeq \Sigma_G$ (since $|T| = 0$ only if T is a G -corolla with \mathcal{P} -labeled vertex) and that $\widehat{\Omega}_G^e$ is the union of (the nerves of) the $\widehat{\Omega}_G^e[\leq k]$, we obtain the desired filtration

$$\mathcal{P} = \mathcal{P}_0 \rightarrow \mathcal{P}_1 \rightarrow \mathcal{P}_2 \rightarrow \cdots \rightarrow \operatorname{colim}_k \mathcal{P}_k = \mathcal{P}[u]. \quad (5.70) \quad \boxed{\text{FILT EQ}}$$

To analyze $\boxed{\text{FILT EQ}}$ homotopically we will further need a pushout description of each map $\mathcal{P}_{k-1} \rightarrow \mathcal{P}_k$. To do so, note that the diagram of inclusions

$$\begin{array}{ccc} \widehat{\Omega}_G^e[k \setminus Y] & \longrightarrow & \widehat{\Omega}_G^e[\leq k \setminus Y] \\ \downarrow & & \downarrow \\ \widehat{\Omega}_G^e[k] & \longrightarrow & \widehat{\Omega}_G^e[\leq k] \end{array} \quad (5.71) \quad \boxed{\text{INCDIAG EQ}}$$

is a pushout of at the level of nerves. Indeed, this follows since

$$\widehat{\Omega}_G^e[k] \cap \widehat{\Omega}_G^e[\leq k \setminus Y] = \widehat{\Omega}_G^e[k \setminus Y], \quad \widehat{\Omega}_G^e[k] \cup \widehat{\Omega}_G^e[\leq k \setminus Y] = \widehat{\Omega}_G^e[\leq k],$$

and since a map $T \rightarrow S$ in $\widehat{\Omega}_G^e[\leq k]$ is in one of subcategories in $\boxed{\text{INCDIAG EQ}}$.

Since Lemma 5.64 provides an identification $\operatorname{Lan}_{\widehat{\Omega}_G^e[k \setminus Y]^{op}} \tilde{N} \simeq \operatorname{Lan}_{\widehat{\Omega}_G^e[\leq k-1]^{op}} \tilde{N} = \mathcal{P}_{k-1}$, applying left Kan extensions to $\boxed{\text{INCDIAG EQ}}$ yields the pushout diagram below.

$$\begin{array}{ccc} \operatorname{Lan}_{\widehat{\Omega}_G^e[k \setminus Y]^{op}} \tilde{N} & \longrightarrow & \mathcal{P}_{k-1} \\ \downarrow & & \downarrow \\ \operatorname{Lan}_{\widehat{\Omega}_G^e[k]^{op}} \tilde{N} & \longrightarrow & \mathcal{P}_k \end{array} \quad (5.72) \quad \boxed{\text{FILTRATION_LAN_SQUARE}}$$

We will also make use of an explicit levelwise description of $\boxed{\text{FILTRATION_LAN_SQUARE_DIAGRAM}}$.

Proposition 5.73. *For each G -corolla $C \in \Sigma_G$, $\boxed{\text{FILTRATION_LAN_SQUARE_DIAGRAM}}$ (5.72) is given by the following pushout in $\mathcal{V}^{\operatorname{Aut}(C)}$*

$$\begin{array}{ccc} \coprod_{[T] \in \operatorname{Iso}(C \downarrow_r \Omega_G^a[k])} \left(\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) \otimes Q_T^{in}[u] \right)_{\operatorname{Aut}(T)} \otimes \operatorname{Aut}(C) & \longrightarrow & \mathcal{P}_{k-1}(C) \\ \downarrow & & \downarrow \\ \coprod_{[T] \in \operatorname{Iso}(C \downarrow_r \Omega_G^a[k])} \left(\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) \otimes \bigotimes_{v \in V_G^{in}(T)} Y(T_v) \right)_{\operatorname{Aut}(T)} \otimes \operatorname{Aut}(C) & \longrightarrow & \mathcal{P}_k(C) \end{array} \quad (5.74) \quad \boxed{\text{FILTRATION_LAN_LEVEL}}$$

where $V_G^{ac}(T)$, $V_G^{in}(T)$ denote the active and inert vertices of $T \in \Omega_G^a[k]$, and $Q_T^{in}[u]$ is the domain of the iterated pushout product

$$\square_{v \in V_G^{in}(T)} u(T_v): Q_T^{in}[u] \rightarrow \bigotimes_{v \in V_G^{in}(T)} Y(T_v).$$

Proof. This is a consequence of Remark 5.63. Iteratively computing left Kan extensions by first left Kan extending to $\Omega_G^a[k]$, we can rewrite the leftmost map in $\boxed{\text{FILTRATION_LAN_SQUARE_DIAGRAM}}$ as

$$\operatorname{Lan}_{\Omega_G^a[k]^{op} \rightarrow \Sigma_G^{op}} \left(\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) \otimes \square_{v \in V_G^{in}(T)} u(T_v) \right). \quad (5.75) \quad \boxed{\text{FILTINTALT EQ}}$$

The desired description of the leftmost map given in $\boxed{\text{FILTRATION_LAN_LEVEL}}$ (5.74) now follows by noting that the rooted undercategories $C \downarrow_r \Omega_G^a[k]$ are groupoids. \square

5.4 Proof of Theorems I and II

In this section we use the filtrations just developed to prove our first two main results, Theorems I and II, concerning the existence of model structures on $\mathbf{Op}^G(\mathcal{V})$ and $\mathbf{Op}_G(\mathcal{V})$.

Recall that given a group Σ , the genuine model structure (if it exists) on \mathcal{V}^Σ , which we denote $\mathcal{V}_{\text{gen}}^\Sigma$, has weak equivalences (resp. fibrations) those maps $X \rightarrow Y$ such that $X^H \rightarrow Y^H$ is a weak equivalence (fibration) for all $H \leq \Sigma$.

Our main proof will require some auxiliary results concerning genuine model structures. However, since these results are particular instances of subtler results in §6, which require a far more careful analysis, we defer the harder proofs to those of the results in §6.

Remark 5.76. The genuine model structure $\mathcal{V}_{\text{gen}}^\Sigma$ exists whenever \mathcal{V} has *cellular fixed points*. The exact condition, originally from [20], can be found in Definition 6.2. Moreover, note that this is condition (iii) in our main theorems. For our immediate purposes, however, we will only need to know that $\mathcal{V}_{\text{gen}}^\Sigma$ is then cofibrantly generated with generating (trivial) cofibrations the maps $\Sigma/H \cdot i$ for $H \leq \Sigma$ and i a generating (trivial) cofibration of \mathcal{V} .

Moreover given a family \mathcal{F} (or even collection of subgroups) of Σ , there exists a model structure $\mathcal{V}_{\mathcal{F}}^\Sigma$ with weak equivalences, fibrations and generating (trivial) cofibrations all described by restricting H to \mathcal{F} .

Remark 5.77. A skeletal filtration argument shows that all objects in $\mathbf{sSet}_{\text{gen}}^\Sigma$, $\mathbf{sSet}_{*,\text{gen}}^\Sigma$ are cofibrant.

Remark 5.78. Suppose \mathcal{V} has cellular fixed points and is a closed monoidal model category.

- (i) Propositions 6.5 and 6.7 imply that for a group homomorphism $\phi: \Sigma \rightarrow \bar{\Sigma}$ the functors

$$\bar{\Sigma} \cdot \Sigma (-): \mathcal{V}_{\text{gen}}^\Sigma \longrightarrow \mathcal{V}_{\text{gen}}^{\bar{\Sigma}} \quad \text{res}_{\bar{\Sigma}}: \mathcal{V}_{\text{gen}}^{\bar{\Sigma}} \longrightarrow \mathcal{V}_{\text{gen}}^\Sigma$$

are left Quillen functors.

- (ii) (6.19) says that the monoidal product on \mathcal{V} lifts to a left Quillen bifunctor

$$\mathcal{V}_{\text{gen}}^\Sigma \times \mathcal{V}_{\text{gen}}^{\bar{\Sigma}} \xrightarrow{\otimes} \mathcal{V}_{\text{gen}}^{\Sigma \times \bar{\Sigma}}.$$

The following lemma is the key to our main proof. Here, a map f in $\mathbf{Sym}_G(\mathcal{V})$ is called a level genuine (trivial) cofibration if each of the maps $f(C)$ for $C \in \Sigma_G$ are genuine trivial cofibrations in $\mathcal{V}_{\text{gen}}^{\text{Aut}(C)}$.

Lemma 5.79. Suppose \mathcal{V} is a cofibrantly generated closed monoidal model category with cellular fixed points and with cofibrant symmetric pushout powers (cf. Proposition 6.30).

Suppose that $\mathcal{P} \in \mathbf{Sym}_G(\mathcal{V})$ is level genuine cofibrant and that $u: X \rightarrow Y$ in $\mathbf{Sym}_G(\mathcal{V})$ is a level genuine cofibration. Then for each $T \in \Omega_G^a[k]$ and writing $C = \text{lr}(T)$ the map

$$\left(\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) \otimes \bigotimes_{v \in V_G^{in}(T)} u(T_v) \right) \otimes_{\text{Aut}(T)} \text{Aut}(C). \quad (5.80)$$

is a genuine cofibration in $\mathcal{V}_{\text{gen}}^{\text{Aut}(C)}$, which is trivial if $k \geq 1$ and u is trivial.

Proof. Combining the homomorphism $\text{Aut}(T) \rightarrow \text{Aut}(C)$ with the leftmost left Quillen functor in Remark 5.78(i), it suffices to check that the parenthesized expression in (5.80) is a (trivial) genuine $\text{Aut}(T)$ -cofibration.

Furthermore, the homomorphism $\text{Aut}(T) \rightarrow \text{Aut}\left(\bigotimes_{v \in V_G^{ac}(T)} (T_v) \otimes \bigotimes_{v \in V_G^{in}(T)} (T_v)\right)$ combined with the rightmost left Quillen functor in Remark 5.78(i) and Remark 5.78(ii) then yield that it suffices to check that

$$\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) = \bigotimes_{v \in V_G^{ac}(T)} (\emptyset \rightarrow \mathcal{P})(T_v), \quad \bigotimes_{v \in V_G^{in}(T)} u(T_v)$$

are, respectively, $\text{Aut}\left((T_v)_{v \in V_G^{gc}(T)}\right)$ and $\text{Aut}\left((T_v)_{v \in V_G^{in}(T)}\right)$ genuine cofibrations, with the latter trivial if u is. Here, the automorphism groups are taken in the category in $\mathbf{F} \wr \Sigma_G$, and thus admit a product description of the form $\Sigma_{|\lambda_1|} \wr \text{Aut}(T_{v_1}) \times \cdots \times \Sigma_{|\lambda_k|} \wr \text{Aut}(T_{v_k})$ as in Remark 2.9. A further application of Remark 5.78(ii) yields that the required conditions need only be checked for the partial pushout product indexed by each λ_i , thus reducing to Proposition 6.30 (when \mathcal{F} is the family of all subgroups). \square

Remark 5.81. When $T \in \Omega^a[k]$ is a non-equivariant alternating tree, \mathcal{P} is cofibrant in $\text{Sym}^G u: X \rightarrow Y$ is a (trivial) cofibration in Sym^G , and writing $C_n = \text{lr}(T)$, the previous result applied to $G \cdot T = (T)_{g \in G} \cdot \iota_! \mathcal{P}$, $\iota_! u$, yields that the analogue of the map (5.80) is a $\text{Aut}(G \cdot C_n) \simeq G \times \Sigma_n$ genuine cofibration.

proof of Theorems I and II. We first build a seemingly unrelated model structure. Consider the composite adjunction below, with right adjoints on the bottom, and where the rightmost right adjoint simply forgets structure and the leftmost right adjoint is given by evaluation.

$$\prod_{C \in \Sigma_G} \mathcal{V}_{\text{gen}}^{\text{Aut}(C)} \xrightleftharpoons[(\text{ev}_C(-))]{\quad} \text{Sym}_G(\mathcal{V}) \xrightleftharpoons[\mathbb{F}_G]{\quad} \text{Op}_G(\mathcal{V}) \quad (5.82)$$

We claim that $\text{Op}_G(\mathcal{V})$ admits a (semi-)model structure with weak equivalences and fibrations defined by the composite right adjoint in (5.82). Noting that the left adjoint to $(\text{ev}_C(-))$ is given by $(X_D) \mapsto \coprod_{D \in \Sigma_G} \text{Hom}_{\Sigma_G}(-, D) \cdot_{\text{Aut}(D)} X_D$ and using either [10, Thm. 11.3.2] in the model structure case $\mathcal{V} = \mathbf{sSet}, \mathbf{sSet}_*$ or [23, Thm. 2.2.2] in the semi-model category structure case, one must analyze free \mathbb{F}_G -extension diagrams of the form

$$\begin{array}{ccc} \mathbb{F}_G(\text{Hom}_{\Sigma_G}(-, D)/H \cdot A) & \longrightarrow & \mathcal{P} \\ u \downarrow & & \downarrow \\ \mathbb{F}_G(\text{Hom}_{\Sigma_G}(-, D)/H \cdot B) & \longrightarrow & \mathcal{P}[u] \end{array}$$

where $D \in \Sigma_G$, $H \leq \text{Aut}(D)$, and $u: A \rightarrow B$ is a generating (trivial) cofibration in \mathcal{V} .

The map $\mathcal{P} \rightarrow \mathcal{P}[u]$ is then filtered as in (5.70), and since $\text{Hom}_{\Sigma_G}(C, D)/H \cdot u$ is a (trivial) cofibration in $\mathcal{V}_{\text{gen}}^{\text{Aut}(C)}$ for all $C \in \Sigma_G$ (cf. Remark 5.76), combining the inductive description of the filtration in (5.74) with Lemma 5.79 shows that if \mathcal{P} is level genuine cofibrant then $\mathcal{P} \rightarrow \mathcal{P}[u]$ is a level genuine cofibration, trivial whenever u is.

In the model structure case $\mathcal{V} = \mathbf{sSet}, \mathbf{sSet}_*$, Remark 5.77 guarantees that any \mathcal{P} is level genuine cofibrant, and thus the conditions in [10, Thm. 11.3.2] are met, showing the existence of the model structure. In the semi-model structure case, the condition that \mathcal{P} is level genuine cofibrant does not quite coincide with the cell complex condition in [23, Thm. 2.2.2]. However, the regular (i.e. not trivial) cofibration case in the previous paragraph together with a routine induction argument over the cell decomposition of cellular \mathcal{P} shows that cellular \mathcal{P} are indeed level genuine cofibrant. Thus, the semi-model structure case also follows.

We now turn to showing the existence of the (semi-)model structures appearing in Theorems I and II, which are essentially corollaries of the existence of that defined by (5.82).

Firstly, consider the projective (semi-)model structure on $\text{Op}_G(\mathcal{V})$. This model structure is transferred from the exact same adjunction (5.82), except equipping the leftmost $\mathcal{V}_{\text{gen}}^{\text{Aut}(C)}$ with their naive model structure, where weak equivalences and fibrations are defined by forgetting the $\text{Aut}(C)$ -action, and ignoring fixed point conditions. The desired projective model structure thus has both less generating (trivial) cofibrations and more weak equivalences than the “genuine projective” model structure defined by (5.82). Therefore, transfinite composites of pushouts of generating projective trivial cofibrations are genuine projective equivalences and hence also projective equivalences, showing that the condition

in [18, Lemma 2.3(1)] holds, establishing the existence of the projective model structure¹. In the semi-model structure case, one replaces [18, Lemma 2.3(1)] with the obvious analogue (unfortunately, we know of no direct reference for this analogue, but its proof is identical).

To address the remaining cases in Theorems I and II, note first that by replacing the model structure in the leftmost category of (5.82) with $\prod_{C \in \Sigma_G} \mathcal{V}_{\mathcal{F}_C}^{\text{Aut}(C)}$ for an arbitrary choice of collections of subgroups \mathcal{F}_C of $\text{Aut}(C)$ for $C \in \Sigma_G$, the exact same argument as in the previous paragraph yields a transferred $\{\mathcal{F}_C\}$ model structure in $\text{Op}_G(\mathcal{V})$.

Letting \mathcal{F} now denote a weak indexing system as in Theorem II and (4.67), one concludes in particular that there exists a “ \mathcal{F} -projective” (semi-)model structure on $\text{Op}_G(\mathcal{V})$, with weak equivalences and fibrations determined by evaluation at C for $C \in \Sigma_{\mathcal{F}}$. This does not quite coincide with the \mathcal{F} -projective model structure on $\text{Op}_{\mathcal{F}}(\mathcal{V})$ appearing in Theorem II, since $\text{Op}_G(\mathcal{V})$ is a larger category. But, since the inclusions $\gamma_! : \text{Sym}_{\mathcal{F}}(\mathcal{V}) \rightarrow \text{Sym}_G(\mathcal{V})$, $\gamma_! : \text{Op}_{\mathcal{F}}(\mathcal{V}) \rightarrow \text{Op}_G(\mathcal{V})$ in (4.58), (4.67) preserve colimits and the monad $\mathbb{F}_{\mathcal{F}}$ defining $\text{Op}_{\mathcal{F}}(\mathcal{V})$ can be regarded as a restriction of \mathbb{F}_G , the desired condition in [18, Lemma 2.3(1)] when applied to the intended model structure on $\text{Op}_{\mathcal{F}}(\mathcal{V})$ turns out to coincide with the corresponding condition for the \mathcal{F} -projective model structure on $\text{Op}_G(\mathcal{V})$. The existence of the projective (semi-)model structures on $\text{Op}_{\mathcal{F}}(\mathcal{V})$ follows, finishing the proof of Theorem II.

We now turn for Theorem I. Should it be the case that (\mathcal{V}, \otimes) has diagonals (which is not a requirement of Theorem I), one can simply use the inclusion $\iota_! : \text{Op}^G(\mathcal{V}) \rightarrow \text{Op}_G(\mathcal{V})$ of (4.42) and repeat the argument in the previous paragraph, except now for an arbitrary collection $\{\mathcal{F}_C\}$. Otherwise, one instead adapts the entire proof, starting with the obvious $\text{Op}^G(\mathcal{V})$ analogue of (5.82) and using Remark 5.81 instead of Lemma 5.79. \square

6 Cofibrancy and Quillen equivalences

In this final section we prove our main result Theorem III, i.e. we show that there are Quillen equivalences

$$\text{Op}_G(\mathcal{V}) \xrightleftharpoons[\iota_*]{\iota^*} \text{Op}^G(\mathcal{V}) \quad \text{Op}_{\mathcal{F}}(\mathcal{V}) \xrightleftharpoons[\iota_*]{\iota^*} \text{Op}_{\mathcal{F}}^G(\mathcal{V})$$

In contrast to the existence of model structure results shown in §5.4, this will require a far more careful analysis of the genuine model structures $\mathcal{V}_{\mathcal{F}}^G$ mentioned in Remark 5.76. This analysis is the subject of §7.7 and §6.2, the results of which are converted to the setup of G -trees in §6.3 and culminate in the characterization of cofibrant objects in Op_G , $\text{Op}_{\mathcal{F}}$ given by Lemma 6.72 in §6.4, with this final lemma tantamount to Theorem III.

6.1 Families of subgroups

Throughout \mathcal{F} denotes a family of subgroups of a finite group G , i.e. a collection of subgroups closed under conjugation and inclusion or, equivalently (cf. §4.4), a sieve $\mathcal{O}_{\mathcal{F}} \hookrightarrow \mathcal{O}_G$.

Remark 6.1. For fixed G families form a lattice, ordered by inclusion, with meet and join given by intersection and union.

As mentioned in Remark 5.76, whenever \mathcal{V} is cofibrantly generated and has cellular fixed points, [20, Prop. 2.6] shows the existence of a model structure $\mathcal{V}_{\mathcal{F}}^G$ on the G -object category \mathcal{V}^G whose fibrations and weak equivalences are determined by fixed points $(-)^H$ for $H \in \mathcal{F}$.

Our analysis will require an explicit understanding of this cellularity condition, which we now recall.

¹ The condition in [18, Lemma 2.3(1)] is clearly also necessary for the existence of the model structure therein, and hence a necessary and sufficient condition. By contrast, the condition in [10, Thm. 11.3.2], while more convenient to state, is merely a sufficient condition.

CELL DEF

Definition 6.2. A model category \mathcal{V} is said to have *cellular fixed points* if for all finite groups G and subgroups $H, K \leq G$ one has that:

- (i) fixed points $(-)^H: \mathcal{V}^G \rightarrow \mathcal{V}$ preserve direct colimits;
- (ii) fixed points $(-)^H$ preserve pushouts where one leg is $(G/K) \cdot f$, for f a cofibration;
- (iii) for each object $X \in \mathcal{V}$, the natural map $(G/K)^H \cdot X \rightarrow ((G/K) \cdot X)^H$ is an isomorphism.

This section will establish some simple useful properties of the \mathcal{V}^G model structures. We start by strengthening the cellularity conditions in Definition 6.2.

CELL DEF

Proposition 6.3. Let \mathcal{V} be a cofibrantly generated model category with cellular fixed points. Then:

- (i) $(-)^H: \mathcal{V}^G \rightarrow \mathcal{V}$ preserves cofibrations and pushouts where one leg is a genuine cofibration;
- (ii) if X is G -genuine cofibrant the map $(G/K)^H \cdot X^H \rightarrow (G \cdot_K X)^H$ is an isomorphism.

Proof. Since both conditions are compatible with retracts, we are free to assume each cofibration $f: X \rightarrow Y$ (or, for Y cofibrant, the map $\emptyset \rightarrow Y$) is a transfinite composition

$$X_0 \xrightarrow{f_0} X_1 \xrightarrow{f_1} X_2 \xrightarrow{f_2} X_3 \xrightarrow{f_3} \dots \rightarrow Y = X_\beta = \text{colim}_{\alpha < \beta} X_\alpha \quad (6.4)$$

TRANSFCOMP EQ

where each $f_\alpha: X_\alpha \rightarrow X_{\alpha+1}$ is the pushout of a generating cofibration $(G/H) \cdot i_\alpha$. Both (i) and (ii) now follow by transfinite induction on α in the partial composite map $X_0 \rightarrow X_\alpha$, with the successor ordinal case following by Def. 6.2 (ii), (iii) and the limit ordinal case by Def. 6.2 (i). We note that (ii) also includes an obvious base case $X_0 = \emptyset$. \square

CELL DEF

CELL DEF

Proposition 6.5. Let $\phi: G \rightarrow \bar{G}$ be a homomorphism and \mathcal{V} be cofibrantly generated with cellular fixed points. Then the adjunction

$$\phi_! = \bar{G} \cdot_G (-): \mathcal{V}_{\mathcal{F}}^G \rightleftarrows \mathcal{V}_{\bar{\mathcal{F}}}^{\bar{G}}: \text{res}_{\bar{G}}^{\bar{G}} = \phi^* \quad (6.6)$$

is a Quillen adjunction provided that for any $H \in \mathcal{F}$ it is $\phi(H) \in \bar{\mathcal{F}}$.

Proof. Since one has a canonical isomorphism of fixed points $(\text{res}(X))^H \simeq X^{\phi(H)}$, it is immediate that the right adjoint preserves fibrations and trivial fibrations. \square

Proposition 6.7. Let $\phi: G \rightarrow \bar{G}$ be a homomorphism and \mathcal{V} be cofibrantly generated with cellular fixed points. Then the adjunction

$$\phi^* = \text{res}_{\bar{G}}^{\bar{G}}: \mathcal{V}_{\bar{\mathcal{F}}}^{\bar{G}} \longrightarrow \mathcal{V}_{\mathcal{F}}^G: \text{Hom}_G(\bar{G}, -) = \phi_* \quad (6.8)$$

is a Quillen adjunction provided that for any $H \in \bar{\mathcal{F}}$ it is $\phi^{-1}(H) \in \mathcal{F}$.

Proof. Since the double coset formula yields that

$$\text{res}(\bar{G}/H \cdot f) \simeq \text{res}(\bar{G}/H) \cdot f \simeq \left(\coprod_{[a] \in \phi(G) \backslash \bar{G}/H} G/\phi^{-1}(H^a) \right) \cdot f$$

it follows that the left adjoint res preserves generating (trivial) cofibrations. \square

FGTRIGHTFGLEFT PROP

Propositions 6.5 and 6.7 motivate the following definition.

Definition 6.9. Let $\phi: G \rightarrow \bar{G}$ be a homomorphism and \mathcal{F} and $\bar{\mathcal{F}}$ families in G and \bar{G} . We define

$$\phi^*(\bar{\mathcal{F}}) = \{H \leq G : \phi(H) \in \bar{\mathcal{F}}\} \quad (6.10)$$

PHISTARDEF EQ

$$\phi_!(\mathcal{F}) = \{\phi(H)^{\bar{g}} \leq \bar{G} : \bar{g} \in \bar{G}, H \in \mathcal{F}\} \quad (6.11)$$

$$\phi_*(\mathcal{F}) = \{\bar{H} \leq \bar{G} : \forall \bar{g} \in \bar{G} (\phi^{-1}(\bar{H}^{\bar{g}}) \in \mathcal{F})\} \quad (6.12)$$

PHISTARDEF3 EQ

Lemma 6.13. The $\phi^*(\bar{\mathcal{F}})$, $\phi_!(\mathcal{F})$, $\phi_*(\mathcal{F})$ just defined are themselves families. Furthermore

- (i) The “provided that” condition in Proposition 6.5 holds iff $\mathcal{F} \subset \phi^*(\bar{\mathcal{F}})$ iff $\phi_!(\mathcal{F}) \subset \bar{\mathcal{F}}$. FGTRIGHT PROP
- (ii) The “if” condition in Proposition 6.7 holds iff $\phi^*(\bar{\mathcal{F}}) \subset \mathcal{F}$ iff $\bar{\mathcal{F}} \subset \phi_*(\mathcal{F})$. FGTLEFT PROP

Proof. Since the result is elementary, we include only the proof of the second iff in (ii), which is the hardest step and illustrates the necessary arguments. This follows by the following equivalences.

$$\phi^*(\bar{\mathcal{F}}) \subset \mathcal{F} \Leftrightarrow \left(\bigvee_{\substack{H \leq G \\ \phi(H) \in \bar{\mathcal{F}}}} H \in \mathcal{F} \right) \Leftrightarrow \left(\bigvee_{\bar{H} \in \bar{\mathcal{F}}} \phi^{-1}(\bar{H}) \in \mathcal{F} \right) \Leftrightarrow \left(\bigvee_{\substack{\bar{H} \in \bar{\mathcal{F}} \\ \bar{g} \in G}} \phi^{-1}(\bar{H}^{\bar{g}}) \in \mathcal{F} \right) \Leftrightarrow \bar{\mathcal{F}} \subset \phi_*(\mathcal{F})$$

Here the second equivalence follows since $H \leq \phi^{-1}(\phi(H))$ and \mathcal{F} is closed under subgroups while the third equivalence follows since $\bar{\mathcal{F}}$ is closed under conjugation. \square

Proposition 6.14. Suppose that \mathcal{V} is cofibrantly generated, has cellular fixed points, and is also a closed monoidal model category. Then the bifunctor

$$\mathcal{V}_{\mathcal{F}}^G \times \mathcal{V}_{\bar{\mathcal{F}}}^G \xrightarrow{\otimes} \mathcal{V}_{\mathcal{F} \cap \bar{\mathcal{F}}}^G \quad (6.15)$$

is a left Quillen bifunctor.

Proof. The double coset formula yields

$$(G/H \cdot f) \square (G/\bar{H} \cdot g) \simeq (G/H \times G/\bar{H}) \cdot (f \square g) \simeq \left(\coprod_{[a] \in H \backslash G / \bar{H}} G/H \cap \bar{H}^a \cdot (f \square g) \right) \quad (6.16)$$

and hence the result follows since families are closed under conjugation and subgroups. \square

Definition 6.17. Let \mathcal{F} and $\bar{\mathcal{F}}$ be families of G and \bar{G} , respectively.

We define their *external intersection* to be the family of $G \times \bar{G}$ given by

$$\mathcal{F} \sqcap \bar{\mathcal{F}} = (\pi_G)^*(\mathcal{F}) \cap (\pi_{\bar{G}})^*(\bar{\mathcal{F}})$$

for $\pi_G: G \times \bar{G} \rightarrow G$, $\pi_{\bar{G}}: G \times \bar{G} \rightarrow \bar{G}$ the projections.

Remark 6.18. Combining Proposition 6.7 with Proposition 6.14 yields that the following composite is a left Quillen bifunctor. BIQUILLENQ PROP

$$\mathcal{V}_{\mathcal{F}}^G \times \mathcal{V}_{\bar{\mathcal{F}}}^{\bar{G}} \xrightarrow{\text{res}} \mathcal{V}_{(\pi_G)^*(\mathcal{F})}^{G \times \bar{G}} \times \mathcal{V}_{(\pi_{\bar{G}})^*(\bar{\mathcal{F}})}^{G \times \bar{G}} \xrightarrow{\otimes} \mathcal{V}_{\mathcal{F} \sqcap \bar{\mathcal{F}}}^{G \times \bar{G}} \quad (6.19)$$

6.2 Pushout powers

That (6.19) is a left Quillen bifunctor (and its obvious higher order analogues) is one of the key properties of pushout products of \mathcal{F} cofibrations when those cofibrations (and the group) are allowed to change. However, when those cofibrations (and hence G) coincide there is an additional symmetric group action that we will need to consider.

To handle these actions we will need two new axioms, each concerning cofibrancy and fixed point properties. We start by discussing the cofibrancy axiom.

HERE

Definition 6.20. We say that a symmetric monoidal model category \mathcal{V} has *cofibrant symmetric pushout powers* if for each (trivial) cofibration f the pushout product power $f^{\square n}$ is a Σ_n -genuine (trivial) cofibration.

Remark 6.21. When \mathcal{V} is cofibrantly generated the condition in Definition 6.20 needs only be checked for generating cofibrations. However, the argument needed is somewhat harder than usual due to $(-)^{\square n}$ not preserving composition of maps: one instead follows the argument in the proof of Proposition 6.30 below when $G = *$. COFSYMPUSHPOW

EXTERINT DEF 6.17.

We now turn to describing the symmetric power analogue of Definition 6.17. We start with some notation. Letting λ be a partition $E = \lambda_1 \sqcup \dots \sqcup \lambda_k$ of a finite set E , we write $\Sigma_\lambda = \Sigma_{\lambda_1} \times \dots \times \Sigma_{\lambda_k} \leq \Sigma_E$ for the subgroup of permutations preserving λ . In addition, given any $e \in E$ we write λ_e for the partition $E = \{e\} \sqcup (E - e)$, so that Σ_{λ_e} is then the isotropy of e .

Definition 6.22. Let \mathcal{F} be a family of G , E a finite set and $e \in E$ any fixed element.

We define the n -th semidirect power of \mathcal{F} to be the family of $\Sigma_E \wr G = \Sigma_E \ltimes G^{\times E}$ given by

$$\mathcal{F}^{\times E} = (\iota_{\Sigma_{\lambda_e} \wr G})_* ((\pi_G)^*(\mathcal{F})), \quad (6.23)$$

FLTIMESN EQ

where ι is the inclusion $\Sigma_{\lambda_e} \wr G \rightarrow \Sigma_E \wr G$ and π the projection $\Sigma_{\lambda_e} \wr G = \Sigma_{\{e\}} \times G \times \Sigma_{E-e} \wr G \rightarrow G$.

More explicitly, since in (6.12) one needs only consider conjugates by coset representatives of $\bar{G}/\phi(G)$, when computing $(\iota_{\Sigma_{\lambda_e} \wr G})_*$ one needs only conjugate by coset representatives of $\Sigma_E \wr G / \Sigma_{\lambda_e} \wr G \simeq \Sigma_E / \Sigma_{\lambda_e}$, so that

$$K \in \mathcal{F}^{\times E} \text{ iff } \forall_{e \in E} \pi_G(K \cap (\Sigma_{\lambda_e} \wr G)) \in \mathcal{F}, \quad (6.24)$$

FLTIMESN2 EQ

showing that in particular (6.23) is independent of the choice of $e \in E$.

Remark 6.25. The previous definition is likely to seem mysterious at first sight. Ultimately, the origin of this definition is best understood by working through this section backwards: the study of the interactions between equivariant trees and graph families, namely Lemma 6.57, requires the study of the families $\mathcal{F}^{\times G^n}$ in Notation 6.41, which are variants of the $\mathcal{F}^{\times n}$ construction for graph families. It then suffices, and is notationally far more convenient, to establish the required results first for the $\mathcal{F}^{\times n}$ families and then translate them to the $\mathcal{F}^{\times G^n}$ families.

Proposition 6.26. Writing $\iota: \Sigma_E \times \Sigma_{\bar{E}} \rightarrow \Sigma_{E \sqcup \bar{E}}$ for the inclusion, one has

$$\mathcal{F}^{\times E} \sqcap \mathcal{F}^{\times \bar{E}} \subset \iota^*(\mathcal{F}^{\times E \sqcup \bar{E}}). \quad (6.27)$$

LTIMESPRODINC EQ

Hence, the following is a left Quillen bifunctor.

$$\Sigma_{E \sqcup \bar{E}} \wr \Sigma_{E \times \Sigma_{\bar{E}}} \cdot (- \otimes -): \mathcal{V}^{\Sigma_E \wr G} \times \mathcal{V}^{\Sigma_{\bar{E}} \wr G} \rightarrow \mathcal{V}^{\Sigma_{E \sqcup \bar{E}} \wr G} \quad (6.28)$$

LTIMESPRODQUI EQ

Proof. Let $K \in \mathcal{F}^{\times E} \sqcap \mathcal{F}^{\times \bar{E}}$ and $e \in E$. We write λ_e for the partition of $E \sqcup \bar{E}$ and λ_e^E for the partition of E . One then has

$$\pi_G(K \cap (\Sigma_{\lambda_e} \wr G)) = \pi_G(\pi_{\Sigma_E \wr G}(K) \cap (\Sigma_{\lambda_e^E} \wr G)), \quad (6.29)$$

where on the right we write $\pi_{\Sigma_E \wr G}: \Sigma_E \wr G \times \Sigma_{\bar{E}} \wr G \rightarrow \Sigma_E \wr G$ and $\pi_G: \Sigma_{\lambda_e^E} \wr G = \Sigma_{\{e\}} \times G \times \Sigma_{E-e} \wr G \rightarrow G$. Therefore K satisfies (6.24) for $\mathcal{F}^{\times E \sqcup \bar{E}}$ since $\pi_{\Sigma_E \wr G}(K)$ does so for $\mathcal{F}^{\times E}$. The case of $e \in \bar{E}$ is identical.

(6.28) simply combines the left Quillen bifunctor (6.19) with Proposition 6.5. \square

POWERF PROP

Proposition 6.30. Suppose that \mathcal{V} is a cofibrantly generated closed monoidal model category with cellular fixed points and with cofibrant symmetric pushout powers.

Then, for all n and cofibration (resp. trivial cofibration) f of $\mathcal{V}_{\mathcal{F}}^G$ one has that $f^{\square n}$ is a cofibration (resp. trivial cofibration) of $\mathcal{V}_{\mathcal{F}^{\times n}}^{\Sigma_n \wr G}$.

Our proof of Proposition 6.30 will essentially repeat the main argument in the proof of [Pe16, Thm. 1.2]. However, both for the sake of completeness and to stress that the argument is independent of the (fairly technical) model structures in [Pe16], we include an abridged version of the proof below, the key ingredient of which is that (6.28) is a left Quillen bifunctor.

Proof. We first note that in the case of $i = (G/H) \cdot \bar{i}$, $H \in \mathcal{F}$, a generating (trivial) cofibration it is

$$i^{\square n} = (G/H)^{\times n} \cdot \bar{i}^{\square n} \simeq \Sigma_n \wr G_{\Sigma_n \wr H} \cdot \bar{i}^{\square n}.$$

$\bar{i}^{\square n}$ is thus a $\Sigma_n \wr H$ -genuine (trivial) cofibration by the cofibrant symmetric pushout powers hypothesis combined with Proposition 6.7 and hence $i^{\square n}$ is a $\mathcal{F}^{\times n}$ (trivial) cofibration by Proposition 6.5 since $\Sigma_n \wr H \in \mathcal{F}^{\times n}$.

For the general case, we start by making the key observation that for composable arrows $\bullet \xrightarrow{g} \bullet \xrightarrow{h} \bullet$ the n -fold pushout product $(hg)^{\square n}$ has a factorization

$$\bullet \xrightarrow{k_0} \bullet \xrightarrow{k_1} \dots \xrightarrow{k_n} \bullet \quad (6.31) \quad \text{COMPINFOLDFACT EQ}$$

where each k_i , $0 \leq i \leq n$, fits into a pushout product

$$\begin{array}{ccc} \bullet & \xrightarrow{\quad} & \bullet \\ \Sigma_n \wr_{\Sigma_{n-i} \times \Sigma_i} (g^{\square n-i} \square h^{\square i}) \downarrow & \scriptstyle r & \downarrow k_i \\ \bullet & \xrightarrow{\quad} & \bullet \end{array} \quad (6.32) \quad \text{COMPINFOLDFACTPUSH EQ}$$

Briefly, (6.31) follows from suitable Σ_n -symmetric convex subposets $P_0 \subset P_1 \subset \dots \subset P_n$ of the poset $P_n = (0 \rightarrow 1 \rightarrow 2)^{\times n}$ where P_0 consists of “tuples with at least one 0-coordinate” and P_i is obtained from P_{i-1} by adding the “tuples with $n-i$ 1-coordinates and i 2-coordinates”. Additional details concerning this filtration appear in the proof of [Pe16, Lemma 4.8].

The general proof now follows by writing f as a retract of a transfinite composition of pushouts of generating (trivial) cofibrations as in (6.4). As usual, retracts can be ignored, and we can hence assume that there is an ordinal κ and $X_\bullet: \kappa \rightarrow \mathcal{V}^G$ such that (i) $f_\beta: X_\beta \rightarrow X_{\beta+1}$ is the pushout of a (trivial) cofibration i_β ; (ii) $\text{colim}_{\alpha < \beta} X_\alpha \xrightarrow{\simeq} X_\beta$ for limit ordinals $\beta < \kappa$; (iii) setting $X_\kappa = \text{colim}_{\beta < \kappa} X_\beta$, f equals the transfinite composite $X_0 \rightarrow X_\kappa$.

We argue by transfinite induction on κ . Writing $\bar{f}_\beta: X_0 \rightarrow X_\beta$ for the partial composites, it suffices to check that the natural transformation of κ -diagrams (rightmost map not included)

$$\begin{array}{ccccccc} Q^n(\bar{f}_1) & \longrightarrow & Q^n(\bar{f}_2) & \longrightarrow & Q^n(\bar{f}_3) & \longrightarrow & Q^n(\bar{f}_4) \longrightarrow \dots \longrightarrow Q^n(\bar{f}_\kappa) \\ \bar{f}_1^{\square n} \downarrow & & \bar{f}_2^{\square n} \downarrow & & \bar{f}_3^{\square n} \downarrow & & \bar{f}_4^{\square n} \downarrow & & \downarrow \bar{f}_\kappa^{\square n} = \text{colim}_{\beta < \kappa} \bar{f}_\beta^{\square n} \\ X_1^{\otimes n} & \longrightarrow & X_2^{\otimes n} & \longrightarrow & X_3^{\otimes n} & \longrightarrow & X_4^{\otimes n} & \longrightarrow & \dots \longrightarrow X_\kappa^{\otimes n}, \end{array}$$

is κ -cofibrant, i.e. that the maps $Q^n(\bar{f}_\beta) \sqcup_{\text{colim}_{\alpha < \beta} Q^n(\bar{f}_\alpha)} \text{colim}_{\alpha < \beta} X_\alpha^{\otimes n} \rightarrow X_\beta^{\otimes n}$ are cofibrations in $\mathcal{V}_{\mathcal{F}^{\times n}}^{\Sigma_n \wr G}$. Condition (ii) above implies that this map is an isomorphism for β a limit ordinal while for $\beta + 1$ a successor ordinal it is the map $Q^n(\bar{f}_{\beta+1}) \sqcup_{Q^n(\bar{f}_\beta)} X_\beta^{\otimes n} \rightarrow X_{\beta+1}^{\otimes n}$. But since $Q^n(\bar{f}_{\beta+1}) \rightarrow Q^n(\bar{f}_{\beta+1}) \sqcup_{Q^n(\bar{f}_\beta)} X_\beta^{\otimes n}$ is precisely the map k_0 of (6.31) for $g = \bar{f}_\beta$, this last map is the composite $k_n k_{n-1} \dots k_1$ so that the result now follows from (6.32) combined with (6.28), the induction hypothesis applied to \bar{f}_β , the fact that $\bar{f}_\beta^{\square k}$ is a pushout of $i_\beta^{\square k}$ (cf. [Pe16, Lemma 4.11]) and the cofibrancy of $i_\beta^{\square k}$ proven at the beginning. \square

We will also need to understand the fixed points of $f^{\square n}$ for general subgroups $K \leq \Sigma_n \wr G$.

To do so recall first that $f^{\square n}$ can be built from the composite


$$f^{\otimes n}: (0 \rightarrow 1)^{\times n} \xrightarrow{f^{\times n}} \mathcal{V}^{\times n} \xrightarrow{\otimes} \mathcal{V}$$

as the map

$$\text{colim}_{(0 \rightarrow 1)^{\times n} - (1, \dots, 1)} f^{\otimes n} \rightarrow Y^{\otimes n},$$

where Y is the target of f . Any $K \leq \Sigma_n \wr G$ acts on the poset $(0 \rightarrow 1)^{\times n}$ itself (via $K \rightarrow \Sigma_n \wr G \rightarrow \Sigma_n$). Moreover, the fixed subposet $((0 \rightarrow 1)^{\times n})^K$ then consists of those tuples in $\{0, 1\}^{\times n}$ whose coordinates coincide if their indexes are in the same coset of n/K , i.e. there is an identification $((0 \rightarrow 1)^{\times n})^K \simeq (0 \rightarrow 1)^{\times n/K}$.

Example 6.33. When $n = 3$ and $n/K = \{\{1, 2\}, \{3\}\}$ the fixed subposet $(0 \rightarrow 1)^{n/K}$ is displayed on the right below.



It will be key for our purposes to know that fixed points $(f^{\square n})^K$ can be computed by first restricting to the smaller cube $(0 \rightarrow 1)^{n/K}$, resulting in a cube of objects with K -actions, and then computing a pushout over that smaller cube. The formal result follows.

Proposition 6.34. Suppose that \mathcal{V} is as in Proposition 6.30 and \mathcal{V} is strongly cellular. Let $K \leq \Sigma_n \wr G$ be a subgroup, $f: X \rightarrow Y$ a map in \mathcal{V}^G and consider the natural maps (in the arrow category)

$$\coprod_{[i] \in n/K} (f^{\otimes [i]})^K \rightarrow (f^{\square n})^K. \quad (6.35)$$

If f is a cofibration between cofibrant objects then (6.35) is an isomorphism.

Proof. The result will follow by induction on n . The base case $n = 1$ is obvious.

Moreover, it is obvious that (6.35), which is a map of arrows, is an isomorphism on the target objects, hence the real claim is that this map is also an isomorphism on sources.

We now note that by considering (6.31) for $g = \emptyset \rightarrow X$, $h = f$ and removing the last map k , one obtains a filtration of the source of $f^{\square n}$. Applying $(-)^K$ to the leftmost map in (6.32) one has isomorphisms

$$\begin{aligned} \left(\Sigma_n \cdot \prod_{\Sigma_{n-i} \times \Sigma_i} X^{\otimes n-i} \otimes f^{\square i} \right)^K &\simeq \coprod_{\substack{n/K=A/K \sqcup B/K \\ |A|=n-i, |B|=i}} (X^{\otimes A} \otimes f^{\square B})^K \simeq \coprod_{\substack{n/K=A/K \sqcup B/K \\ |A|=n-i, |B|=i}} (X^{\otimes A})^K \otimes (f^{\square B})^K \\ &\simeq \coprod_{\substack{n/K=A/K \sqcup B/K \\ |A|=n-i, |B|=i}} \left(\bigotimes_{[j] \in A/K} (X^{\otimes [j]})^K \right) \otimes \left(\coprod_{[k] \in B/K} (f^{\otimes [k]})^K \right) \end{aligned}$$

where the first step is an instance of Prop. 6.3(ii), the second step follows from Prop. 6.3(iii) (with the required cofibrancy of the objects following from Propositions 6.14 and 6.30), and the last step follows by Prop. 6.3(iii) together with the induction hypothesis (which applies since $|B| \leq i < n$).

We have thus shown that the leftmost map in the pushouts (6.32) for $(f^{\square n})^K$ is isomorphic to the leftmost map in the pushouts for the corresponding filtration of $\coprod_{[i] \in n/K} (f^{\otimes [i]})^K$, and since $(-)^K$ preserves such pushouts (cf. Prop. 6.3(i)), the result now follows. \square

Corollary 6.36. Let \mathcal{V} be strongly cellular. Given a partition λ given by $\{1, 2, \dots, n\} = \lambda_1 \sqcup \dots \sqcup \lambda_k$, cofibrations between cofibrant objects f_i in \mathcal{V}^{G_i} , $1 \leq i \leq k$ and a subgroup $K \leq \Sigma_{\lambda_1} \wr G_1 \times \dots \times \Sigma_{\lambda_k} \wr G_k$, the natural map

$$\coprod_{1 \leq i \leq k} \coprod_{[j] \in \lambda_i/K} (f_i^{\otimes [j]})^K \rightarrow \left(\coprod_{1 \leq i \leq k} f_i^{\square \lambda_i} \right)^K. \quad (6.37)$$

is an isomorphism.

Proof. This simply combines Propositions 6.3(iii) and 6.34. \square

6.3 G -graph families and G -trees

We will now convert the results in the previous sections to the context of the type of families we are mainly interested in: graph families. We note that in this section we use Σ to denote a general group (usually meant to be some type of permutation group).

Definition 6.38. A subgroup $\Gamma \leq G \times \Sigma$ is called a G -graph subgroup if $\Gamma \cap \Sigma = *$.

Further, a family \mathcal{F} of $G \times \Sigma$ is called a G -graph family if it consists of G -graph subgroups.

Remark 6.39. Γ is a G -graph subgroup iff it can be written as

$$\Gamma = \{(k, \varphi(k)) : k \in K \leq G\}$$

for some partial homomorphism $G \geq K \xrightarrow{\varphi} \Sigma$, thus motivating the terminology.

Remark 6.40. The collection of all G -graph subgroups is itself a family. Indeed, it coincides with $(\iota_\Sigma)_*(\{*\})$ for the inclusion homomorphism $\iota_\Sigma: \Sigma \rightarrow G \times \Sigma$.

Notation 6.41. Letting $\mathcal{F}, \bar{\mathcal{F}}$ be G -graph families of $G \times \Sigma$ and $G \times \bar{\Sigma}$ we will write

$$\mathcal{F} \sqcap_G \bar{\mathcal{F}} = \Delta^*(\mathcal{F} \sqcap \bar{\mathcal{F}}) \quad \mathcal{F}^{\kappa_{G^n}} = \Delta^*(\mathcal{F}^{\kappa_n})$$

where Δ denotes either of the diagonal inclusions $\Delta: G \times \Sigma \times \bar{\Sigma} \rightarrow G \times \Sigma \times G \times \bar{\Sigma}$ or $\Delta: G \times \Sigma_n \wr \Sigma \rightarrow \Sigma_n \wr (G \times \Sigma)$.

Remark 6.42. Unpacking Definition 6.17 one has that $\Gamma \in \mathcal{F} \sqcap_G \bar{\mathcal{F}}$ iff $\pi_{G \times \Sigma}(\Gamma) \in \mathcal{F}$, $\pi_{G \times \bar{\Sigma}}(\Gamma) \in \bar{\mathcal{F}}$.

Remark 6.43. Unpacking (6.24) and noting that

$$(G \times \Sigma_E \wr \Sigma) \cap (\Sigma_{\lambda_e} \wr (G \times \Sigma)) = G \times \Sigma_{\lambda_e} \wr \Sigma$$

one has

$$K \in \mathcal{F}^{\kappa_{G^E}} \text{ iff } \forall_{e \in E} \pi_{G \times \Sigma}(K \cap (G \times \Sigma_{\lambda_e} \wr \Sigma)) \in \mathcal{F}. \quad (6.44)$$

Combining either the left Quillen bifunctor (6.19) or Proposition 6.30 with Proposition 6.7 yields the following results.

Proposition 6.45. Suppose that \mathcal{V} is a cofibrantly generated closed monoidal model category with cellular fixed points. Let $\mathcal{F}, \bar{\mathcal{F}}$ be G -graph families of $G \times \Sigma$ and $G \times \bar{\Sigma}$. Then the following (with diagonal G -action on the images) is a left Quillen bifunctor.

$$\mathcal{V}_{\mathcal{F}}^{G \times \Sigma} \times \mathcal{V}_{\bar{\mathcal{F}}}^{G \times \bar{\Sigma}} \xrightarrow{\otimes} \mathcal{V}_{\mathcal{F} \sqcap_G \bar{\mathcal{F}}}^{G \times \Sigma \times \bar{\Sigma}} \quad (6.46)$$

Proposition 6.47. Suppose that \mathcal{V} is a cofibrantly generated closed monoidal model category with cellular fixed points and with cofibrant symmetric pushout powers.

Let \mathcal{F} be a G -graph family of $G \times \Sigma$. If f is a cofibration (resp. trivial cofibration) in $\mathcal{V}_{\mathcal{F}}^{G \times \Sigma}$ then so is f^{\square_n} a cofibration (resp. trivial cofibration) in $\mathcal{V}_{\mathcal{F}^{\kappa_{G^n}}}^{G \times \Sigma_n \wr \Sigma}$.

Remark 6.48. It is straightforward to check that $\mathcal{F} \sqcap_G \bar{\mathcal{F}}$ is in fact also a G -graph family of $G \times \Sigma \times \bar{\Sigma}$. However, $\mathcal{F}^{\kappa_{G^n}}$ is not a G -graph family of $G \times \Sigma_n \wr \Sigma$, due to the need to consider the power Σ_n -action.

The G -graph families we will be interested in encode certain families of G -trees. We start with the case of families of corollas (see Definition 4.56).

Lemma 6.49. The set of families of corollas $\Sigma_{\mathcal{F}}$ is in bijection with the set of collections $\mathcal{F} = \{\mathcal{F}_n\}_{n \geq 0}$ of G -graph families \mathcal{F}_n of $G \times \Sigma_n$. Moreover, we have an equivalence of categories $\coprod \mathcal{O}_{\mathcal{F}_n} \xrightarrow{\cong} \Sigma_{\mathcal{F}}$.

Proof. We define \mathcal{F}_n so that any G -corolla C with n leaves per component is in $\Sigma_{\mathcal{F}}$ iff

$$C \simeq (C_{\Gamma})_{G/H}$$

for some $C_{\Gamma} \in \Sigma^H$ encoded by a partial homomorphism $G \leq H \rightarrow \Sigma_n$ whose graph Γ is in \mathcal{F}_n ; conversely, any \mathcal{F} dually defines a full subcategory $\Sigma_{\mathcal{F}}$ of Σ_G , and these operations are clearly inverse.

Further, Γ_1 is subconjugate to Γ_2 iff we have a quotient map

$$(C_{\Gamma_1})_{G/H_1} \rightarrow (C_{\Gamma_2})_{G/H_2}$$

where $H_i = \pi_G(\Gamma_i)$, and hence \mathcal{F} is a collection of *families* iff $\Sigma_{\mathcal{F}}$ is a sieve of Σ_G .

Finally, the equivalence of categories just sends $\Gamma \leq H \times \Sigma_n$ to $(C_{\Gamma})_{G/H}$. \square

As in §4.4, we will abuse notation and use \mathcal{F} and $\Sigma_{\mathcal{F}}$ interchangably².

Proposition 6.50. *Let \mathcal{F} be a family of G -corollas and $T \in \Omega$ a tree with automorphism group Σ_T . Write \mathcal{F}_T for the collection of G -graph subgroups of $G \times \Sigma_T$ encoded by partial homomorphisms $G \geq H \rightarrow \Sigma_T$ such that the associated G -tree $G \cdot_H T$ is a \mathcal{F} -tree.*

Then \mathcal{F}_T is a G -graph family.

Proof. Closure under conjugation follows since conjugate graph subgroups produce isomorphic G -trees. As for subgroups, they correspond to restrictions $K \leq H \rightarrow \Sigma_T$, which induce quotient maps $G \cdot_K T \rightarrow G \cdot_H T$. \mathcal{F}_T is thus closed under subgroups since the G -vertices of $G \cdot_H T$ are quotients of those of $G \cdot_K T$. \square

Remark 6.51. The closure condition required of weak indexing systems from Definition 4.61 can be translated in terms of families as saying that for any tree $T \in \Omega$ and letting $\phi: \Sigma_T \rightarrow \Sigma_{\text{lr}(T)}$ be the natural homomorphism, one has $(G \times \phi)(\Gamma) \in \mathcal{F}_{\text{lr}(T)}$ for any $\Gamma \in \mathcal{F}_T$. Proposition 6.5 then says that

$$\phi!: \mathcal{V}_{\mathcal{F}_T}^{G \times \Sigma_T} \rightarrow \mathcal{V}_{\mathcal{F}_{\text{lr}(T)}}^{G \times \text{lr}(T)} \quad (6.52)$$

LRLEFTQUILLEN EQ

is a left Quillen functor.

Remark 6.53. Unpacking definitions, a partial homomorphism $G \geq H \rightarrow \Sigma_T$ encodes a subgroup in \mathcal{F}_T iff, for each vertex $v = e^{\uparrow} \leq e$ of T with $H_e \leq H$ the H -isotropy of the edge e , the induced homomorphism

$$H_e \rightarrow \Sigma_{T_v} \simeq \Sigma_{|v|} \quad (6.54)$$

PARTIALHOMEDGE EQ

encodes a subgroup in $\mathcal{F}_{|v|}$, where $|v| = |e^{\uparrow}|$.

Remark 6.55. Recall that any tree $T \in \Omega$ other than the stick η has an essentially unique grafting decomposition $T = C_n \sqcup_{n \cdot \eta} (T_1 \sqcup \dots \sqcup T_n)$ where C_n is the root corolla and the leaves of C_n are identified with the roots of the T_i . We now let λ be the partition $\{1, \dots, n\} = \lambda_1 \sqcup \dots \sqcup \lambda_k$ such that $1 \leq i_1, i_2 \leq n$ are in the same class iff $T_{i_1}, T_{i_2} \in \Omega$ are isomorphic.

Writing $\Sigma_{\lambda} = \Sigma_{\lambda_1} \times \dots \times \Sigma_{\lambda_k}$ and picking representatives $i_j \in \lambda_j$ one then has isomorphisms

$$\Sigma_T \simeq \Sigma_{\lambda} \wr \prod_i \Sigma_{T_i} \simeq \Sigma_{|\lambda_1|} \wr \Sigma_{T_{i_1}} \times \dots \times \Sigma_{|\lambda_k|} \wr \Sigma_{T_{i_k}} \quad (6.56)$$

TREEISOT EQ

where the second isomorphism, while not canonical (it depends on choices of isomorphisms $T_{i_j} \simeq T_l$ for each $i_j \neq l \in \lambda_j$) is nonetheless well defined up to conjugation.

The following, which is the core result in this section, is a reinterpretation of Remark 6.53 in light of the inductive description of trees in Remarks 6.55.

Lemma 6.57. *Let $\Sigma_{\mathcal{F}}$ be a family of G -corollas and $T \in \Omega$ a tree other than η . Then*

$$\mathcal{F}_T = (\pi_{G \times \Sigma_n})^* (\mathcal{F}_n) \cap \left(\mathcal{F}_{T_{i_1}}^{\kappa_G |\lambda_1|} \sqcap_G \dots \sqcap_G \mathcal{F}_{T_{i_k}}^{\kappa_G |\lambda_k|} \right), \quad (6.58)$$

KEYLEMMA GECO EQ

where $\pi_{G \times \Sigma_n}$ denotes the composite $G \times \Sigma_T \rightarrow G \times \Sigma_{\lambda} \rightarrow G \times \Sigma_n$.

² Alternatively, the reader can think of \mathcal{F} as a “family in the groupoid Σ of finite sets”.

Proof. The argument is by induction on the decomposition $T = C_n \sqcup_{n,\eta} (T_1 \sqcup \dots \sqcup T_n)$ with the base case, that of a corolla, being immediate.

Consider now a partial homomorphism $G \geq H \rightarrow \Sigma_T$ encoding a G -graph subgroup $\Gamma \leq G \times \Sigma_T$. The condition that $\Gamma \in (\pi_{G \times \Sigma_n})^*(\mathcal{F}_n)$ states that the composite $H \rightarrow \Sigma_T \rightarrow \Sigma_n$ is in \mathcal{F}_n , and this is precisely the condition (6.54) in Remark 6.53 for $e = r$ the root of T .

As for the condition $\Gamma \in (\mathcal{F}_{T_i}^{\kappa_G|\lambda_1|} \sqcap_G \dots \sqcap_G \mathcal{F}_{T_{i_k}}^{\kappa_G|\lambda_k|})$, by unpacking it by combining Remarks 6.42 and 6.43, this translates to the condition that, for each $i \in \{1, \dots, n\}$, one has

$$\pi_{G \times \Sigma_{T_i}} \left(\Gamma \cap \left(G \times \Sigma_{\{i\}} \times \Sigma_{T_i} \times \Sigma_{\lambda - \{i\}} \wr \prod_{j \neq i} \Sigma_{T_j} \right) \right) \in \mathcal{F}_{T_i} \quad (6.59)$$

KEYLEMMAECOR EQ

where $\lambda - \{i\}$ denotes the induced partition of $\{1, \dots, n\} - \{i\}$. Noting that the intersection subgroup inside $\pi_{G \times \Sigma_{T_i}}$ in (6.59) can be rewritten as $\Gamma \cap \pi_{\Sigma_n}^{-1}(\Sigma_{\{i\}} \times \Sigma_{\{1, \dots, n\} - \{i\}})$, we see that this is the graph subgroup encoded by the restriction $H_i \leq H \rightarrow \Sigma_T$, where H_i is the isotropy subgroup of the root r_i of T_i (equivalently, this is also the subgroup sending T_i to itself). But since for any edge $e \in T_i$ its isotropy H_e (cf. 6.54) is a subgroup of H_i , the induction hypothesis implies that (6.59) is equivalent to condition (6.54) across all vertices other than the root vertex.

The previous paragraphs show that (6.58) indeed holds when restricted to G -graph subgroups. However, it still remains to show that any group Γ in the right family in (6.58) is indeed a G -graph subgroup, i.e. $\Gamma \cap \Sigma_T = *$ or, in other words, that any element $\gamma \in \Gamma \cap \Sigma_T = G \times \Sigma_\lambda \wr \prod_i \Sigma_{T_i}$ with G -coordinate $\gamma_G = e$ is indeed the identity. But the condition $\pi_{G \times \Sigma_n}(\Gamma) \in \mathcal{F}_n$ now implies that for such γ the Σ_λ -coordinate is $\gamma_{\Sigma_\lambda} = e$ and thus (6.59) in turn implies that the Σ_{T_i} -coordinates are $\gamma_{\Sigma_{T_i}} = e$, finishing the proof. \square

The results just developed will allow us to analyze cofibrancy properties of the left maps in the key pushouts (5.74). The first part of the analysis concerns the maps

$$\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) \otimes \bigsqcup_{v \in V_G^{in}(T)} u(T_v) \quad (6.60)$$

COFIBMAPSTREE EQ

that constitute the inner part of (5.75), where we recall that $T \in \Omega^a$ is an alternating tree. We will in turn subdivide the cofibrancy analysis of (6.60) itself into two parts: (i) showing a \mathcal{F}_{T_e} -cofibrancy claim when $T = G \cdot T_e$ is free and; (ii) showing a fixed point claim for non free trees, as in Remark 4.43.

It will find it convenient to slightly reinterpret (6.60): writing $p(T_v): \emptyset \rightarrow \mathcal{P}(T_v)$ for the unique map, we can rewrite (6.60) as

$$\bigsqcup_{v \in V_G^{ac}(T)} p(T_v) \bigsqcup \bigsqcup_{v \in V_G^{in}(T)} u(T_v).$$

For both the sake of generality and to simplify notation in the proof, we extend the context of the following results to the l -labeled trees Ω_G^l of §5.1.

Remark 6.61. l -labeled \mathcal{F} -trees can be defined exactly as in Definition 4.59. Moreover, it is then clear that a l -labeled G -tree T is an \mathcal{F} -tree iff the underlying G -tree is.

Remarks 6.53, 6.55 and Lemma 6.57 then extend to the l -labeled context, by now writing Σ_T for the group of label isomorphisms (a subgroup of the isomorphisms of the underlying tree) and defining the partition λ in Remark 6.55 by using label isomorphism classes.

Proposition 6.62. Suppose that \mathcal{V} is a cofibrantly generated closed monoidal model category with cellular fixed points and with cofibrant symmetric pushout powers.

Let \mathcal{F} be a family of corollas and suppose that $f_i: A \rightarrow B$, $1 \leq i \leq l$ are level \mathcal{F} -cofibrations (resp. trivial cofibrations) in $\text{Sym}^G(\mathcal{V})$, i.e. that $f_i(r): A(r) \rightarrow B(r)$ are cofibrations (resp. trivial cofibrations) in $\mathcal{V}_{\mathcal{F}_n}^{G \times \Sigma_n}$. Then for any l -labeled tree $T \in \Omega^l$ the map

$$f^{\square V(T)} = \bigsqcup_{1 \leq i \leq l, v \in V_i(T)} f_i(v)$$

(where $V_i(T)$ denotes vertices with label i) is a cofibration (resp. trivial cofibration) in $\mathcal{V}_{\mathcal{F}_T}^{G \times \Sigma_T}$.

Proof. This follows by induction on the decomposition $T = C_n \sqcup_{n,\eta} (T_1 \sqcup \dots \sqcup T_n)$ with the base cases of corollas and η being immediate. The description of \mathcal{F}_T in (6.58) combined with the left Quillen functors in Propositions 6.45, 6.14 and 6.7 then yield that

$$\mathcal{V}_{\mathcal{F}_n}^{G \times \Sigma_n} \times \mathcal{V}_{\mathcal{F}_{T_{i_1}}}^{G \times \Sigma_{|\lambda_1|} \wr \Sigma_{T_{i_1}}} \times \dots \times \mathcal{V}_{\mathcal{F}_{T_{i_k}}}^{G \times \Sigma_{|\lambda_k|} \wr \Sigma_{T_{i_k}}} \xrightarrow{\otimes} \mathcal{V}_{\mathcal{F}_T}^{G \times \Sigma_T}$$

is a left Quillen multifunctor. The result now follows by Proposition 6.47 together with the induction hypothesis. \square

Remark 6.63. When $G = *$, Proposition 6.62 coincides with [2, Lemma 5.9]. Moreover, it is not hard to adapt the proof of that non-equivariant result to provide a proof of Proposition 6.62 in the case of the universal family Σ_G of all G -corollas. However, the reader of [2] may note that the proof therein is technically less involved, using no analogue of the rather subtle $\mathcal{F}^{\kappa_{G^n}}$ families. Indeed, this is reflected on the last paragraph of our proof of Lemma 6.57, which effectively uses the subtle condition (6.59) to deduce the much simpler condition $\Gamma \cap \prod \Sigma_{T_i} = *$ that would suffice for the direct generalization of [2, Lemma 5.9] mentioned above. One may thus wonder if the $\mathcal{F}^{\kappa_{G^n}}$ families are indeed required to prove Proposition 6.62, or whether a more direct adaptation of the proof of [2, Lemma 5.9] is possible. Reverse engineering our proofs, the most natural “simplification” would be to replace the condition (6.59) with the condition

$$\pi_{G \times \Sigma_{T_i}}(\Gamma \cap \prod \Sigma_{T_i}) \in \mathcal{F}, \quad (6.64)$$

thus replacing the families \mathcal{F}^{κ_n} of (6.23) with the families $(\iota_{G \times n})_*(\mathcal{F} \sqcap \dots \sqcap \mathcal{F})$. However, it is not hard to build indexing systems \mathcal{F} (other than the universal one) for which these simpler families do not satisfy the analogue of Lemma 6.57, and thus for which (6.52) fails.

Proposition 6.65. *Let \mathcal{V} be as in Proposition 6.62, and suppose additionally that \mathcal{V} is strongly cofibrantly generated (so \mathcal{V} is strongly cellular) and that \mathcal{V} is Cartesian monoidal.*

Let $f_i: A \rightarrow B$, $1 \leq i \leq l$ be genuine cofibrations between genuine cofibrant objects in $\text{Sym}^G(\mathcal{V})$. For each $T \in \Omega_{G,0}^l$ define

$$f^{\square V_G(T)} = \bigsqcup_{1 \leq i \leq l, v \in V_{G,i}(T)} \iota_* f_i(v). \quad (6.66)$$

Then the canonical map

$$f^{\square V_G(-)} \rightarrow \iota_* \iota^* f^{\square V_G(-)} \quad (6.67)$$

is an isomorphism in $\mathcal{V}_{\Omega_G^{0,l,op}}$ (with $G \times \Omega_0^l \xrightarrow{\iota} \Omega_G^{0,l}$ the inclusion).

Proof. Note first that there is a coproduct decomposition

$$\Omega_{G,0}^l \simeq \coprod_{U \in \text{Iso}(\Omega_0^l)} \Omega_{G,0}^l[U]$$

where $\Omega_{G,0}^l[U]$ is the full subcategory formed by the quotients of $G \cdot U$. It suffices to establish (6.67) for each subcategory $\Omega_{G,0}^l[U]$.

Moreover, writing T as $T \simeq G \cdot_H T_e$ for $T_e \in \Omega^H$, we are free by induction on $|G|$ to assume $H = G$. Indeed, otherwise there are identifications $V_G(T) \simeq V_H(T_e)$ and $f^{\square V_G(T)} \simeq (\text{res}_H^G f)^{\square V_H(T_e)}$ from which the desired isomorphism follows.

We thus reduce to the case where there is a quotient map $G \cdot U \rightarrow U_G$ where U_G denotes the underlying tree U together a G -action. Moreover, the automorphisms of $G \cdot U$ compatible

with the quotient map $G \cdot U \rightarrow U_G$ are the subgroup $K \leq G \times \Sigma_U$ encoding the action $G \rightarrow \Sigma_U$ of G on U_G . One then has identifications

$$\left(\coprod_{[v] \in V_G(G \cdot U)} \iota_* f_\bullet([v]) \right)^K \simeq \left(\coprod_{v \in V(U)} f_\bullet(v) \right)^G \simeq \left(\coprod_{[v] \in V(U)/G} \prod_{v \in [v]} f_\bullet(v) \right)^G \simeq \left(\coprod_{[v] \in V_G(U_G)} \iota_* f_\bullet([v]) \right)$$

where the middle step is Corollary 6.36 establishing the desired isomorphism (6.67). \square

6.4 Cofibrancy and the proof of Theorem III

Propositions 6.62 and 6.65 from the previous section provide most of the backbone of the main technical step in the proof of our Elmendorf-type result Theorem III. The last remaining step is the following refinement of the key argument in the proof of [20, Thm. 2.10].

Proposition 6.68. *Let \mathcal{V} be a cofibrantly generated model category with cellular fixed points, \mathcal{F} a non-empty family of subgroups of G , and consider the reflexive adjunction*

$$\mathcal{V}_{\mathcal{F}}^{\text{Opp}} \xrightleftharpoons[\iota_*]{\iota^*} \mathcal{V}_{\mathcal{F}}^G. \quad (6.69)$$

Then the cofibrant objects of $\mathcal{V}_{\mathcal{F}}^{\text{Opp}}$ are precisely the essential image under ι_* of the cofibrant objects of $\mathcal{V}_{\mathcal{F}}^G$. Moreover, the analogous statement for cofibrations between cofibrant objects also holds.

Proof. Note first that since ι_* identifies $\mathcal{V}^G(\mathcal{F})$ as a reflexive subcategory of $\mathcal{V}_{\mathcal{F}}^{\text{Opp}}$, it is $X \simeq \iota_* Y$ for some $Y \in \mathcal{V}^G(\mathcal{F})$ (i.e. $X \in \mathcal{V}_{\mathcal{F}}^{\text{Opp}}$ is in the essential image of ι_*) iff both $\iota^* X \simeq Y$ and the unit map $X \xrightarrow{\simeq} \iota_* \iota^* X$ is an isomorphism.

Letting $C_{\mathcal{F}}$ (resp. $C^{\mathcal{F}}$) denote the classes of cofibrant objects in $\mathcal{V}_{\mathcal{F}}^{\text{Opp}}$ (resp. $\mathcal{V}_{\mathcal{F}}^G$) we need to show $C_{\mathcal{F}} = \iota_*(C^{\mathcal{F}})$, where we slightly abuse notation by writing $\iota_*(-)$ for the essential image rather than the image. Since $C_{\mathcal{F}}$ is characterized as being the smallest class closed under retracts and transfinite composition of cellular extensions that contains the initial presheaf \emptyset , it suffices to show that $\iota_*(C^{\mathcal{F}})$ satisfies this same characterization.

It is immediate that $\iota_*(\emptyset) = \emptyset$. Further, the characterization in the first paragraph yields that $X \in \iota_*(C^{\mathcal{F}})$ iff $\iota^*(X) \in C^{\mathcal{F}}$ and $X \xrightarrow{\simeq} \iota_* \iota^* X$ is an isomorphism, showing that $\iota_*(C^{\mathcal{F}})$ is closed under retracts.

The crux of the proof will be to compare cellular extensions in $C_{\mathcal{F}}$ with the images under ι_* of the cellular extensions in $C^{\mathcal{F}}$. Firstly, note that the generating cofibrations in $\mathcal{V}_{\mathcal{F}}^{\text{Opp}}$ have the form $\text{Hom}(-, G/H) \cdot f$, and that by the cellularity axiom (iii) in Definition 6.2 this map is isomorphic to the map $\iota_*(G/H \cdot f)$. We now claim that the cellular extensions of objects in $\iota_*(C^{\mathcal{F}})$, i.e. pushforward diagrams as on the left below

$$\begin{array}{ccc} \iota_* X & \longrightarrow & \iota_* V \\ \iota_* u \downarrow & & \downarrow \\ \iota_* Y & \dashrightarrow & \tilde{W} \end{array} \quad \begin{array}{ccc} X & \longrightarrow & V \\ u \downarrow & & \downarrow \\ Y & \dashrightarrow & W \end{array} \quad (6.70)$$

are precisely the essential image under ι_* of the cellular extensions of objects in $C^{\mathcal{F}}$, i.e., pushforward diagrams as on the right above. That the solid subdiagrams in either side of (6.70) are indeed in bijection up to isomorphism is simply the claim that ι^* is fully faithful, hence the real claim is that $\tilde{W} \simeq \iota_* W$. But this follows since by the cellularity axiom (ii) in Definition 6.2 the map ι_* preserves the rightmost pushforward in (6.70) (note that $u: X \rightarrow Y$ is assumed to be a generating cofibration of $\mathcal{V}_{\mathcal{F}}^G$).

Noting that the cellularity axiom (i) in Definition 6.2 implies that ι_* preserves filtered colimits finishes the proof that $C_{\mathcal{F}} = \iota_*(C^{\mathcal{F}})$.

The additional claim concerning cofibrations between cofibrant objects follows by the same argument. \square

FINALCOR COR

Corollary 6.71. *Let \mathcal{V} be as above, $\phi: G \rightarrow \bar{G}$ a homomorphism, and $\mathcal{F}, \bar{\mathcal{F}}$ families of G, \bar{G} such that $\phi_! \mathcal{F} \subset \bar{\mathcal{F}}$. Then the diagram*

$$\begin{array}{ccc} \mathcal{V}_{\mathcal{F}}^{Op} & \xleftarrow{\iota_*} & \mathcal{V}_{\mathcal{F}}^G \\ \phi_! \downarrow & & \downarrow \phi_! \\ \mathcal{V}_{\bar{\mathcal{F}}}^{Op} & \xleftarrow{\iota_*} & \mathcal{V}_{\bar{\mathcal{F}}}^G \end{array}$$

commutes up to isomorphism when restricted to cofibrant objects of $\mathcal{V}_{\mathcal{F}}^G$.

Proof. It is straightforward to check that the left adjoints commute, i.e. that there is a natural isomorphism $\iota^* \phi_! \simeq \phi_! \iota^*$ which by adjunction induces a natural transformation $\phi_! \iota_* \rightarrow \iota_* \phi_!$. More explicitly, this natural transformation is the composite

$$\phi_! \iota_* \rightarrow \iota_* \iota^* \phi_! \iota_* \xrightarrow{\simeq} \iota_* \phi_! \iota^* \iota_* \xrightarrow{\simeq} \iota_* \phi_!$$

where the last two maps are always isomorphisms. But when restricting to cofibrant objects the previous result guarantees both that $\phi_! \iota_*$ lands in cofibrant objects and that cofibrant objects are in the essential image of ι_* . The result follows. \square

We now possess all the technical ingredients needed to prove Theorem MAINQUILLENEQUIV THM

MAINLEM LEM

Lemma 6.72. *Let \mathcal{V} be strongly cellular with diagonals, and let \mathcal{F} be a weak indexing system.*

Then in both of the adjunctions

$$\mathrm{Op}_{\mathcal{F}}(\mathcal{V}) \xrightleftharpoons[\iota_*]{\iota^*} \mathrm{Op}_{\mathcal{F}}^G(\mathcal{V}) \qquad \mathrm{Sym}_{\mathcal{F}}(\mathcal{V}) \xrightleftharpoons[\iota_*]{\iota^*} \mathrm{Sym}_{\mathcal{F}}^G(\mathcal{V}) \quad (6.73) \quad \text{COFADJ2 EQ}$$

the cofibrant objects in the leftmost category are the essential image under ι_ of the cofibrant objects in the rightmost category.*

Moreover, both forgetful functors

$$\mathrm{Op}_{\mathcal{F}}(\mathcal{V}) \xrightarrow{\mathrm{fgt}} \mathrm{Sym}_{\mathcal{F}}(\mathcal{V}) \qquad \mathrm{Op}_{\mathcal{F}}^G(\mathcal{V}) \xrightarrow{\mathrm{fgt}} \mathrm{Sym}_{\mathcal{F}}^G(\mathcal{V}) \quad (6.74) \quad \text{FGTFUNC EQ}$$

preserve cofibrant objects.

Proof. We first observe that the claim concerning the symmetric sequence adjunction of (6.73) is not really new. Indeed, as there are equivalences of categories $\mathrm{Sym}_{\mathcal{F}}(\mathcal{V}) \simeq \prod_{n \geq 0} \mathcal{V}_{\mathcal{F}_n}^{Op}$, $\mathrm{Sym}_{\mathcal{F}}^G(\mathcal{V}) \simeq \prod_{n \geq 0} \mathcal{V}_{\mathcal{F}_n}^{G \times \Sigma_n}$, compatible with both the model structures and the (ι^*, ι_*) adjunctions, the symmetric sequence statement merely repackages Proposition 6.68. COFADJ2 EQ

For the operad adjunction in (6.73), most of the argument in the proof of Proposition 6.68 applies mutatis mutandis except for the claim that $\mathbb{F}_{\mathcal{F}}(\emptyset) \simeq \iota_* \mathbb{F}(\emptyset)$, which is readily checked directly, and the comparison of cellular extensions, which is the key claim. COFESSIM PROP

Now, borrowing the notation from Proposition 6.68, let $C^{\mathcal{F}}$ and $C_{\mathcal{F}}$ denote the classes of cofibrant objects in $\mathrm{Op}_{\mathcal{F}}^G(\mathcal{V})$ and $\mathrm{Op}_{\mathcal{F}}$. We note that the generating cofibrations for $\mathrm{Op}_{\mathcal{F}}(\mathcal{V})$ are precisely those for $\mathrm{Op}_G(\mathcal{V})$ which are generated from those in $\mathrm{Sym}_{\mathcal{F}}^G(\mathcal{V})$, as $\gamma_!$ includes $\mathrm{Op}_{\mathcal{F}}(\mathcal{V})$ as a reflective subcategory and moreover detects cofibrations, since Remark 6.7 implies F MODEL REM

$$\gamma_!(\gamma^* \mathbb{F}_G \gamma_! \iota^*(f)) = \mathbb{F}_G \iota^*(f) \quad (6.75) \quad \text{UPGAMMA_COFIB_EQ}$$

for f a generating cofibration (with cofibrant source by assumption) in $\mathrm{Sym}_{\mathcal{F}}^G(\mathcal{V})$.

Thus, recalling that ι_* in the statement of the lemma abusively denotes the composite $\gamma^* \iota_*$, to prove the desired equality $\gamma^* \iota_*(C^{\mathcal{F}}) = C_{\mathcal{F}}$, we may equivalently prove $\iota_*(C^{\mathcal{F}}) =$

$\gamma_!(C_{\mathcal{F}})$, for which in turn it suffices by [\(6.75\)](#) to show that cellular extensions of objects in $\iota_*(C^{\mathcal{F}})$, such as on the left below

$$\begin{array}{ccc} \mathbb{F}_G \iota_* X & \longrightarrow & \iota_* \mathcal{O} \\ \iota_* u \downarrow & & \downarrow \\ \mathbb{F}_G \iota_* Y & \dashrightarrow & (\iota_* \mathcal{O})[\iota_* u] \end{array} \quad \begin{array}{ccc} \mathbb{F} X & \longrightarrow & \mathcal{O} \\ u \downarrow & & \downarrow \\ \mathbb{F} Y & \dashrightarrow & \mathcal{O}[u] \end{array} \quad (6.76) \quad \boxed{\text{TWOCELLEXT EQ}}$$

are precisely the essential image under ι_* of cellular extensions of objects in $C^{\mathcal{F}}$, as on the right above, with $u : X \rightarrow Y$ a cofibration in $\text{Sym}_{\mathcal{F}}^G(\mathcal{V})$. Moreover, by induction on the minimum number of cellular extensions needed to generate \mathcal{O} , we may assume $\text{fgt}(\mathcal{O})$ is cofibrant in $\text{Sym}_{\mathcal{F}}^G(\mathcal{V})$.

Now, recalling that there are natural isomorphisms

$$\iota^* \mathbb{F}_G \iota_* \simeq \mathbb{F} \iota^* \iota_* \simeq \mathbb{F} \quad \gamma^* \mathbb{F}_G \gamma_* \simeq \gamma^* \mathbb{F}_G \gamma_!,$$

we see that the two solid subdiagrams in [\(6.76\)](#) are in fact adjoint up to isomorphism, so that there is a bijection between such data. We claim that it now suffices to check that all four objects in the leftmost diagram of [\(6.76\)](#) are in the essential image of ι_* . Indeed, if that is the case then

$$\mathbb{F}_G \iota_* Z \simeq \iota_* \iota^* \mathbb{F}_G \iota_* Z \simeq \iota_* \mathbb{F} Z$$

for $Z = X, Y$ and since ι_* reflects colimits³ it must then indeed be that $(\iota_* \mathcal{O})[\iota_* u] \simeq \iota_*(\mathcal{O}[u])$.

To establish the remaining claim that the objects in the leftmost diagram in [\(6.76\)](#) are in the essential image of ι_* , we claim it suffices to show this for the bottom right corner $(\iota_* \mathcal{O})[\iota_* u]$ when $u : X \rightarrow Y$ is a general cofibration between cofibrant objects in $\text{Sym}_{\mathcal{F}}^G(\mathcal{V})$. Indeed, setting $X = \emptyset$ and $\mathcal{O} = \mathbb{F}(\emptyset)$, one has $(\iota_* \mathcal{O})[\iota_* u] = \mathbb{F}_G \iota_* Y$, and similarly for $\mathbb{F}_G \iota_* X$.

Now, writing $\mathcal{P} = \iota_* \mathcal{O}$, so that $(\iota_* \mathcal{O})[\iota_* u] = \mathcal{P}[\iota_* u]$, the required condition that $\mathcal{P}[\iota_* u] \rightarrow \iota_* \iota^* \mathcal{P}[\iota_* u]$ is an isomorphism can be checked by forgetting to $\text{Sym}_G(\mathcal{V})$, and we can thus appeal to the filtration [\(5.70\)](#) of $\mathcal{P} \rightarrow \mathcal{P}[\iota_* u]$. It thus suffices to verify by induction on k that each \mathcal{P}_k is in the essential image of $\iota_* : \text{Sym}_{\mathcal{F}}^G(\mathcal{V}) \rightarrow \text{Sym}_G(\mathcal{V})$. Using the iterative description of the \mathcal{P}_k in [\(5.74\)](#) it suffices by [Proposition 6.68](#) to check that the leftmost map in [\(5.74\)](#) is a cofibration between cofibrant objects in $\text{Sym}_G(\mathcal{V})$. We now recall that that map can also be described (cf. [\(5.75\)](#)) as

$$\text{Lan}_{\Omega_{\mathcal{F}}^a[k]^{op} \rightarrow \Sigma_G^{op}} \left(\bigotimes_{v \in V_G^{ac}(T)} \mathcal{P}(T_v) \otimes \bigoplus_{v \in V_G^{in}(T)} \iota_* u(T_v) \right). \quad (6.77) \quad \boxed{\text{FILTINTALTAG EQ}}$$

where we have been able to replace $\Omega_G^a[k]$ with $\Omega_{\mathcal{F}}^a[k]$ in the left Kan extension, as both \mathcal{P} and $\iota_* u$ are in the image of $\gamma_!$ (cf. proof of [Proposition 4.64](#)). Noting that there is an equivalence $\mathcal{V}_{\mathcal{F}}^{\Omega_{\mathcal{F}}^a[k]^{op}} \simeq \prod_{T \in \text{Iso}(\Omega_{\mathcal{F}}^a[k])} \mathcal{V}_{\mathcal{F}}^{\mathcal{O}_{\mathcal{F}}^{op}}$, [Propositions 6.62](#) and [6.65](#) show that the inner map inside the left Kan extension in [\(6.77\)](#) is in the essential image of the cofibrations between cofibrant objects of $\mathcal{V}_{\mathcal{F}}^{G \times \Omega^a[k]^{op}}$ under the functor $\iota_* : \mathcal{V}_{\mathcal{F}}^{G \times \Omega^a[k]^{op}} \rightarrow \mathcal{V}_{\mathcal{F}}^{\Omega_{\mathcal{F}}^a[k]^{op}}$. Since the Lan in [\(6.77\)](#) can now be identified with an instance of the functor $\phi_!$ in [Corollary 6.71](#) one has (using the further identification $\mathcal{V}_{\mathcal{F}}^{G \times \Omega^a[k]^{op}} \simeq \prod_{T \in \text{Iso}(\Omega_{\mathcal{F}}^a[k])} \mathcal{V}_{\mathcal{F}}^{G \times \Sigma_T}$) that the main claim now follows by [Corollary 6.71](#) with [Propositions 6.68](#) and [6.71](#).

As for the additional claim concerning the forgetful functors in [\(6.74\)](#), that $\text{fgt} : \text{Op}_{\mathcal{F}}(\mathcal{V}) \rightarrow \text{Sym}_{\mathcal{F}}(\mathcal{V})$ preserves cofibrant objects is precisely what was argued in the previous paragraph (and independently is immediate from the main result and the fact that $\iota^* \iota_* = \text{id}$). Moreover, since the forgetful functors commute with ι_* , the claim that $\text{fgt} : \text{Op}_{\mathcal{F}}(\mathcal{V}) \rightarrow \text{Sym}_{\mathcal{F}}(\mathcal{V})$ also preserves cofibrant objects follows from the “essential image characterization” of cofibrant objects in [\(6.73\)](#).

□

³I.e. any diagram that becomes a colimit upon applying $\gamma^* \iota_*$ must have already been a colimit diagram.

Remark 6.78. A slightly more careful analysis of the argument in the previous proof shows that we have in fact shown the slightly more general claim that operads (in either $\mathbf{Op}_{\mathcal{F}}(\mathcal{V})$ or $\mathbf{Op}_{\mathcal{F}}^G(\mathcal{V})$) that forget to cofibrant symmetric sequences (in either $\mathbf{Sym}_{\mathcal{F}}(\mathcal{V})$ or $\mathbf{Sym}_{\mathcal{F}}^G(\mathcal{V})$) are closed under cellular extensions of operads.

MAINQUILLENEQUIV THM
proof of Theorem III. It suffices to show that both the derived unit and derived counit for the adjunction are given by weak equivalences.

For the counit, it is immediate from Lemma 6.72 that if $X \in \mathbf{Op}^G(\mathcal{V})$ is bifibrant the functor $\iota^* \iota_* X$ is already derived, and hence the derived counit is identified with the counit isomorphism $\iota^* \iota_* X \xrightarrow{\sim} X$.

For the unit, note first that it is immediate from the definitions that $\iota_*: \mathbf{Op}_{\mathcal{F}}^G(\mathcal{V}) \rightarrow \mathbf{Op}_{\mathcal{F}}(\mathcal{V})$ detects fibrations (as well as weak equivalences), and thus by Lemma 6.72 $Y \in \mathbf{Op}_{\mathcal{F}}(\mathcal{V})$ is bifibrant iff $Y \simeq \iota_* X$ for $X \in \mathbf{Op}_{\mathcal{F}}^G(\mathcal{V})$ bifibrant. But then the functor $\iota_* \iota^* Y$ is also already derived (since $\iota^* Y \simeq \iota^* \iota_* X \simeq X$ is fibrant) and the derived unit is thus the isomorphism $Y \xrightarrow{\sim} \iota_* \iota^* Y$. \square

6.5 N_{∞} -operads

HERE

Combining the existence of \mathcal{F} -model structures and the above cofibrancy result, we give our first proof of the following conjecture of Blumberg-Hill in [3].

NINFTY-REAL-COR-MAIN
Proof of Corollary IV, version one. Recall that $\mathbf{Comm}(n) = *$ for all n . Consider the functorial factorization

$$\emptyset \twoheadrightarrow N\mathcal{F} \xrightarrow{\sim} \mathbf{Comm}$$

in $\mathbf{Op}_{\mathcal{F}}^G(\mathbf{sSet})$ of the unique map into a cofibration and trivial fibration. Since the initial operad is cofibrant, Theorem 6.72 implies that $\emptyset \rightarrow N\mathcal{F}$ is a level \mathcal{F} -cofibration, and hence each $N\mathcal{F}(n)$ is cofibrant in $\mathbf{sSet}_{\mathcal{F}_n}^{G \times \Sigma_n}$; thus, for all $\Gamma \notin \mathcal{F}_n$, $N\mathcal{F}(n)^{\Gamma} = \emptyset$. Further, since $N\mathcal{F}$ is \mathcal{F} -equivalent to $*$, $N\mathcal{F}(n)^{\Gamma} \simeq *$ for all $\Gamma \in \mathcal{F}_n$. Hence, each $N\mathcal{F}(n)$ is a universal space for \mathcal{F}_n , as desired. \square

come back

Theorem 6.79. $B(\mathbb{F}_G, \mathbb{F}_G, \delta_{\mathcal{F}})$ is

- in the image of i_*
- weakly equivalent to $\delta_{\mathcal{F}}$.

Hence, $i^*(B(\mathbb{F}_G, \mathbb{F}_G, \delta_{\mathcal{F}}))$ is an $N\mathcal{F}$ -operad.

A Transferring Kan extensions

The purpose of this appendix is to provide the somewhat long proof of Proposition 5.40, which is needed when repackaging free extensions of genuine equivariant operads in (5.7).

We start with a more detailed discussion of the realization functor $|-|$ defined by the adjunction

$$|-|: \mathbf{Cat}^{\Delta^{op}} \rightleftarrows \mathbf{Cat}: (-)^{[\bullet]}$$

in Definition 5.38. More explicitly, one has

$$|\mathcal{I}_{\bullet}| = \operatorname{coeq} \left(\coprod_{[n] \rightarrow [m]} [n] \times \mathcal{I}_m \rightrightarrows \coprod_{[n]} [n] \times \mathcal{I}_n \right). \quad (\text{A.1})$$

Example A.2. Any $\mathcal{I} \in \mathbf{Cat}$ induces objects $\mathcal{I}, \mathcal{I}_{\bullet}, \mathcal{I}^{[\bullet]} \in \mathbf{Cat}^{\Delta^{op}}$ where \mathcal{I} is the constant simplicial object and \mathcal{I}_{\bullet} is the nerve $N\mathcal{I}$ with each level regarded as a discrete category. It is straightforward to check that $|\mathcal{I}| \simeq |\mathcal{I}_{\bullet}| \simeq |\mathcal{I}^{[\bullet]}| \simeq \mathcal{I}$.

Lemma A.3. Given $\mathcal{I}_\bullet \in \text{Cat}^{\Delta^{op}}$ one has an identification $ob(|\mathcal{I}_\bullet|) \simeq ob(\mathcal{I}_0)$. Furthermore, the arrows of $|\mathcal{I}_\bullet|$ are generated by the image of the arrows in $\mathcal{I}_0 \simeq \mathcal{I}_0 \times [0]$ and the image of the arrows in $[1] \times ob(\mathcal{I}_1)$.

For each $i_1 \in \mathcal{I}_1$, we will denote the arrow of $|\mathcal{I}_\bullet|$ induced by the arrow in $[1] \times \{i_1\}$ by

$$d_1(i_1) \xrightarrow{i_1} d_0(i_1).$$

Proof. We write $d_{\hat{k}}, d_{\hat{k}, \hat{l}}$ for the simplicial operators induced by the maps $[0] \xrightarrow{0 \mapsto k} [n]$, $[1] \xrightarrow{0 \mapsto k, 1 \mapsto l} [n]$ which can informally be thought of as the “composite of all faces other than d_k, d_l ”. Using (A.1) one has equivalence relations of objects

$$[n] \times \mathcal{I}_n \ni (k, i_n) \sim (0, d_{\hat{k}}(i_n)) \in [0] \times \mathcal{I}_0$$

and since for any generating relation $(k, i_n) \sim (l, i'_n)$ it is $d_{\hat{k}}(i_n) = d_{\hat{l}}(i'_n)$ the identification $ob(|\mathcal{I}_\bullet|) \simeq ob(\mathcal{I}_0)$ follows.

To verify the claim about generating arrows, note that any arrow of $[n] \times \mathcal{I}_n$ factors as

$$(k, i_n) \rightarrow (l, i_n) \xrightarrow{I_n} (l, i'_n) \quad (\text{A.4})$$

for $I_n: i_n \rightarrow i'_n$ an arrow of \mathcal{I}_n . The $d_{\hat{l}}$ relation identifies the right arrow in (A.4) with

$(0, d_{\hat{l}}(i_n)) \xrightarrow{d_{\hat{l}}(I_n)} (0, d_{\hat{l}}(i'_n))$ in $[0] \times \mathcal{I}_0$ while (if $k < l$) the $d_{\hat{k}, \hat{l}}$ relation identifies the left arrow with $(0, d_{\hat{k}, \hat{l}}(i_n)) \rightarrow (1, d_{\hat{k}, \hat{l}}(i_n))$ in $[1] \times \mathcal{I}_1$. The result follows. \square

Remark A.5. Given $\mathcal{I}_\bullet \in \text{Cat}^{\Delta^{op}}$, $\mathcal{C} \in \text{Cat}$, the isomorphisms

$$\text{Hom}_{\text{Cat}}(|\mathcal{I}_\bullet|, \mathcal{C}) \simeq \text{Hom}_{\text{Cat}^{\Delta^{op}}}(\mathcal{I}_\bullet, \mathcal{C}^{[\bullet]})$$

together with the fact that $\mathcal{C}^{[\bullet]}$ is always 2-coskeletal show that $|\mathcal{I}_\bullet|$ is determined by the categories $\mathcal{I}_0, \mathcal{I}_1, \mathcal{I}_2$ and maps between them, i.e. by the truncated version of formula (A.1) with $n, m \leq 2$.

Indeed, one can show that a sufficient set of generating relations in $|\mathcal{I}_\bullet|$ is given by (i) the relations in \mathcal{I}_0 (including relations stating that identities of \mathcal{I}_0 are identities of $|\mathcal{I}_\bullet|$); (ii) relations stating that for each $i_0 \in \mathcal{I}_0$ the arrow $i_0 = d_1(s_0(i_0)) \xrightarrow{s_0(i_0)} d_1(s_0(i_0)) = i_0$ is an identity; (iii) for each arrow $I_1: i_1 \rightarrow i'_1$ in \mathcal{I}_1 the relation that the square below commutes

$$\begin{array}{ccc} d_1(i_1) & \xrightarrow{i_1} & d_0(i_1) \\ d_1(I_1) \downarrow & & \downarrow d_0(I_1) \\ d_1(i'_1) & \xrightarrow{i'_1} & d_0(i'_1) \end{array}$$

and (iv) for each object $i_2 \in \mathcal{I}_2$ the relation that the following triangle commutes.

$$\begin{array}{ccc} d_{1,2}(i_2) & \xrightarrow{d_1(i_2)} & d_{0,1}(i_2) \\ & \searrow d_2(i_2) & \nearrow d_0(i_2) \\ & d_{0,2}(i_2) & \end{array}$$

We now relate diagrams in the span categories with the Grothendieck constructions in Definition 2.2.

Lemma A.6. Functors $F: \mathcal{D} \times \mathcal{I}_\bullet \rightarrow \mathcal{C}$ are in bijection with lifts

$$\begin{array}{ccc} & & \text{WSpan}^l(*, \mathcal{C}) \\ & \nearrow \mathcal{I}_\bullet^F & \downarrow \text{fgt} \\ \mathcal{D} & \xrightarrow{\mathcal{I}_\bullet} & \text{Cat}. \end{array}$$

where fgt is the functor forgetting the maps to $*$ and \mathcal{C} .

Proof. This is a matter of unpacking notation. The restrictions $F|_{\mathcal{I}_d}$ to the fibers $\mathcal{I}_d \subset \mathcal{D} \ltimes \mathcal{I}_\bullet$ are precisely the functors $\mathcal{I}_d^F: \mathcal{I}_d \rightarrow \mathcal{C}$ describing $\mathcal{I}_\bullet^F(d)$.

Furthermore, the images $F((d, i) \rightarrow (d', f_*(i)))$ of the pushout arrows over a fixed arrow $f: d \rightarrow d'$ of \mathcal{D} assemble to a natural transformation

$$\begin{array}{ccc} \mathcal{I}_d & \xrightarrow{I_d^F} & \mathcal{C} \\ f_* \downarrow & \searrow & \uparrow I_{d'}^F \\ \mathcal{I}_{d'} & \xrightarrow{I_{d'}^F} & \mathcal{C} \end{array} \quad (\text{A.7})$$

which describes $\mathcal{I}_\bullet^F(f)$. One readily checks that the associativity and unitality conditions coincide. \square

In the cases of interest we have $\mathcal{D} = \Delta^{op}$. The following is the key result in this section.

Proposition A.8. *Let $\mathcal{I}_\bullet \in \text{Cat}^{\Delta^{op}}$. Then there is a natural functor*

$$\Delta^{op} \ltimes \mathcal{I}_\bullet \xrightarrow{s} |\mathcal{I}_\bullet|. \quad (\text{A.9})$$

Further, s is final.

Remark A.10. The s in the result above stands for *source*. This is because, for $\mathcal{I} \in \text{Cat}$, the map $\Delta^{op} \ltimes \mathcal{I}^{[\bullet]} \rightarrow |\mathcal{I}^{[\bullet]}| \simeq \mathcal{I}$ is given by $s(i_0 \rightarrow \dots \rightarrow i_n) = i_0$.

Proof. Recall that $|\mathcal{I}_\bullet|$ is the coequalizer REALDEF EQ (A.1). Given $(k, g_m) \in [n] \times \mathcal{I}_m$, we write $[k, g_m]$ for the corresponding object in $|\mathcal{I}_\bullet|$. To simplify notation, we write objects of \mathcal{I}_n as i_n and implicitly assume that $[k, i_n]$ refers to the class of the object $(k, i_n) \in [n] \times \mathcal{I}_n$.

We define s on objects by $s([n], i_n) = [0, i_n]$ and on an arrow $(\phi, I_m): (n, i_n) \rightarrow (m, i'_m)$ as the composite (note that $\phi: [m] \rightarrow [n]$ and $I_m: \phi^* i_n \rightarrow i'_m$)

$$[0, i_n] \rightarrow [\phi(0), i_n] = [0, \phi^* i_n] \xrightarrow{I_m} [0, i'_m]. \quad (\text{A.11})$$

TARGETDEFINITION EQ

To check compatibility with composition, the cases of a pair of either two fiber arrows (i.e. arrows where ϕ is the identity) or two pushforward arrows (i.e. arrows where I_m is the identity) are immediate from (A.11), hence we are left with the case $([n], i_n) \xrightarrow{I_n} ([n], i'_n) \rightarrow ([m], \phi^* i'_n)$ of a fiber arrow followed by a pushforward arrow. Noting that in $\Delta^{op} \ltimes \mathcal{I}_\bullet$ this composite can be rewritten as $([n], i_n) \rightarrow ([m], \phi^* i_n) \xrightarrow{\phi^* I_n} ([m], \phi^* i'_n)$ this amounts to checking that

$$\begin{array}{ccccc} [0, i_n] & \longrightarrow & [\phi(0), i_n] & = & [0, \phi^* i_n] \\ I_n \downarrow & & I_n \downarrow & & \downarrow \phi^* I_n \\ [0, i'_n] & \longrightarrow & [\phi(0), i'_n] & = & [0, \phi^* i_n] \end{array} \quad (\text{A.12})$$

commutes in $|\mathcal{I}_\bullet|$, which is the case since the left square is encoded by a square in $[n] \times \mathcal{I}_n$ and the right square is encoded by an arrow in $[m] \times \mathcal{I}_n$.

We now show that s is final. Fix $h \in \mathcal{I}_0$. We must check that $[0, h] \downarrow \Delta^{op} \ltimes \mathcal{I}_\bullet$ is connected. By Lemma A.3 any object in this undercategory has a description (not necessarily unique) as a pair

$$\left(([n], i_n), [0, h] \xrightarrow{f_1} \dots \xrightarrow{f_r} s([n], i_n) \right) \quad (\text{A.13})$$

UNDERCATOBJ EQ

where each f_i is a generating arrow of $|\mathcal{I}_\bullet|$ induced by either an arrow I_0 of \mathcal{I}_0 or object $i_1 \in \mathcal{I}_1$. We will connect (A.13) to the canonical object $(([0], h), [0, h] = [0, h])$, arguing by induction on r . If $n \neq 0$, the map $d_0^*: ([n], i_n) \rightarrow ([0], d_0^*(i_n))$ and the fact that $s(d_0^*) = id_{[0, d_0^*(i_n)]}$ provides an arrow to an object with $n = 0$ without changing r . If $n = 0$, one can apply the induction hypothesis by lifting f_r to $\Delta^{op} \ltimes \mathcal{I}_\bullet$ according to one of two cases: (i) if f_r is induced by an arrow I_0 of \mathcal{I}_0 , the lift of f_r is simply $([0], i'_0) \xrightarrow{I_0} ([0], i_0)$; (ii) if f_r is induced by $i_1 \in \mathcal{I}_1$ the lift is provided by the map $([1], i_1) \rightarrow ([0], d_0(i_1))$. \square

Remark A.14. The involution

$$\Delta \xrightarrow{\tau} \Delta$$

which sends $[n]$ to itself and d_i, s_i to d_{n-i}, s_{n-i} induces vertical isomorphisms

$$\begin{array}{ccc} \Delta^{op} \ltimes (\mathcal{I}_\bullet \circ \tau) & \xrightarrow{s} & |\mathcal{I}_\bullet \circ \tau| \\ \downarrow \simeq & & \downarrow \simeq \\ \Delta^{op} \ltimes \mathcal{I}_\bullet & \xrightarrow{t} & |\mathcal{I}_\bullet^{op}|^{op} \end{array}$$

which reinterpret the “source” functor as what one might call the “target” functor, with $t([n], i_n) = [n, i_n]$ rather than $s([n], i_n) = [0, i_n]$. The target functor is thus also final.

Moreover, the source/target formulations of all the results that follow are equivalent.

In practice, we will need to know that the source s and target t satisfy a stronger finality condition with respect to left Kan extensions.

Lemma A.15. *Let $\mathcal{J} \in \mathbf{Cat}$ be a small category and $j \in \mathcal{J}$. Then the under and over category functors*

$$\mathbf{Cat} \downarrow \mathcal{J} \xrightarrow{(-) \downarrow j} \mathbf{Cat}, \quad \mathbf{Cat} \downarrow \mathcal{J} \xrightarrow{j \downarrow (-)} \mathbf{Cat} \quad (\text{A.16})$$

UNDEROVER EQ

preserve colimits.

Proof. The result can easily be shown directly, so here we note instead that one can in fact write explicit formulas for the right adjoints of $(-) \downarrow j$, $j \downarrow (-)$. Moreover, since $j \downarrow \mathcal{I} = (\mathcal{I}^{op} \downarrow j)^{op}$ it suffices to do so for $(-) \downarrow j$. The right adjoint $(-) \downarrow j: \mathbf{Cat} \rightarrow \mathbf{Cat} \downarrow \mathcal{J}$ is then defined on objects by the Grothendieck constructions $\mathcal{C}^{\downarrow j} = \mathcal{J} \ltimes \mathcal{C}^{\mathcal{J}^{(-,j)}}$ for the functors

$$\begin{aligned} \mathcal{J} &\longrightarrow \mathbf{Cat} \\ i &\longmapsto \mathcal{C}^{\mathcal{J}^{(i,j)}}. \end{aligned}$$

□

Corollary A.17. *Consider a map $\mathcal{I}_\bullet \rightarrow \mathcal{J}$ between $\mathcal{I}_\bullet \in \mathbf{Cat}^{\Delta^{op}}$ and a constant object $\mathcal{J} = \mathcal{J}_\bullet \in \mathbf{Cat}^{\Delta^{op}}$. Then the source and target maps*

$$\begin{array}{ccc} \Delta^{op} \ltimes \mathcal{I}_\bullet & \xrightarrow{s} & |\mathcal{I}_\bullet| \\ & \searrow & \swarrow \\ & \mathcal{J} & \end{array} \quad \begin{array}{ccc} \Delta^{op} \ltimes \mathcal{I}_\bullet & \xrightarrow{t} & |\mathcal{I}_\bullet^{op}|^{op} \\ & \searrow & \swarrow \\ & \mathcal{J} & \end{array}$$

are Lan-final over \mathcal{J} , i.e. the functors $s \downarrow j: (\Delta^{op} \ltimes \mathcal{I}_\bullet) \downarrow j \rightarrow |\mathcal{I}_\bullet| \downarrow j$ are final for all $j \in \mathcal{J}$, and similarly for t .

Proof. It is clear that $(\Delta^{op} \ltimes \mathcal{I}_\bullet) \downarrow j \simeq \Delta^{op} \ltimes (\mathcal{I}_\bullet \downarrow j)$ while Lemma A.15 guarantees that since $(-) \downarrow j$ is a left adjoint, $|\mathcal{I}_\bullet| \downarrow j \simeq |\mathcal{I}_\bullet \downarrow j|$. One thus reduces to Proposition A.8. □

We will require two additional straightforward lemmas.

Lemma A.18. *Let $\mathcal{I}_\bullet^F \in \mathbf{Span}(*, \mathcal{C})^{\Delta^{op}}$ be such that the diagrams*

$$\begin{array}{ccc} \mathcal{I}_n & & \\ d_i \downarrow & \swarrow F_n & \searrow \\ \mathcal{I}_{n-1} & \delta_i & \mathcal{C} \end{array} \quad \begin{array}{ccc} \mathcal{I}_n & & \\ s_j \downarrow & \swarrow F_n & \searrow \\ \mathcal{I}_{n+1} & \sigma_j & \mathcal{C} \end{array} \quad (\text{A.19})$$

IDENTSIMPRELSISO EQ

are given by natural isomorphisms for $0 < i \leq n$, $0 \leq j \leq n$. Then the functors $\tilde{F}_n: \mathcal{I}_n \rightarrow \mathcal{C}$ given by the composites

$$\mathcal{I}_n \xrightarrow{d_{1,\dots,n}} \mathcal{I}_0 \xrightarrow{F_0} \mathcal{C}$$

assemble to an object $\mathcal{I}_\bullet^{\tilde{F}} \in \text{Span}(*, \mathcal{C})^{\Delta^{op}}$ which is isomorphic to \mathcal{I}_\bullet^F and such that the corresponding diagrams (A.19) for $0 < i \leq n$, $0 \leq j \leq n$ are strictly commutative.

Dually, if (A.19) are natural isomorphisms for $0 \leq i < n$ and $0 \leq j \leq n$ one can form $\mathcal{I}_\bullet^{\tilde{F}} \in \text{Span}(*, \mathcal{C})^{\Delta^{op}}$ such that the corresponding diagrams are strictly commutative.

Proof. This follows by a straightforward verification. \square

Lemma A.20. A (necessarily unique) factorization

$$\Delta^{op} \ltimes \mathcal{I}_\bullet \xrightarrow{F_\bullet} \mathcal{C} \quad \begin{array}{c} \searrow s \\ \text{---} \mathcal{I}_\bullet \end{array} \quad \begin{array}{c} \xrightarrow{F} \\ \text{---} \end{array} \quad \mathcal{C} \quad (\text{A.21}) \quad \text{SOURCEFACT EQ}$$

exists iff for the associated object $\mathcal{I}_\bullet \in \text{Span}(*, \mathcal{C})^{\Delta^{op}}$ (cf. Lemma A.6) all faces d_i for $0 < i \leq n$ and degeneracies s_j for $0 \leq j \leq n$ are strictly commutative, i.e. they are given by diagrams

$$\begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ d_0 \downarrow & \searrow \varphi_n & \nearrow F_{n-1} \\ \mathcal{I}_{n-1} & \xrightarrow{F_{n-1}} & \mathcal{C} \end{array} \quad \begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ d_i \downarrow & \searrow & \nearrow F_{n-1} \\ \mathcal{I}_{n-1} & \xrightarrow{F_{n-1}} & \mathcal{C} \end{array} \quad \begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ s_j \downarrow & \searrow & \nearrow F_{n+1} \\ \mathcal{I}_{n+1} & \xrightarrow{F_{n+1}} & \mathcal{C} \end{array} \quad (\text{A.22}) \quad \text{IDENTSIMPRELS EQ}$$

Dually, a factorization through the target $t: \Delta^{op} \ltimes \mathcal{I}_\bullet \rightarrow |\mathcal{I}_\bullet|^{op}$ exists iff the faces d_i and degeneracies s_j are strictly commutative for $0 \leq i < n$, $0 \leq j \leq n$.

Proof. For the “only if” direction, it suffices to note that s sends all pushout arrows of $\Delta^{op} \ltimes \mathcal{I}_\bullet$ for faces d_i , $0 < i \leq n$ and degeneracies s_j , $0 \leq j \leq n$ to identities, yielding the required commutative diagrams in (A.22).

For the “if” direction, this will follow by building a functor $\mathcal{I}_\bullet \xrightarrow{\tilde{F}_\bullet} \mathcal{C}^{[\bullet]}$ together with the naturality of the source map s (recall that $|\mathcal{C}^{[\bullet]}| \simeq \mathcal{C}$). We define $\tilde{F}_n|_{k \rightarrow k+1}$ as the map

$$F_{n-k}d_{0,\dots,k-1} \xrightarrow{\varphi_{n-k}d_{0,\dots,k-1}} F_{n-k-1}d_{0,\dots,k}. \quad (\text{A.23}) \quad \text{EQUIVALENCEDEF EQ}$$

The claim that $s \circ (\Delta^{op} \ltimes \tilde{F})$ recovers the horizontal map in (A.21) is straightforward, hence the real task is to prove that (A.23) defines a map of simplicial objects. First, functoriality of the original F_\bullet yields identities

$$\varphi_{n-1}d_i = \varphi_n, \quad 1 < i \quad \varphi_{n-1}d_1 = (\varphi_{n-1}d_0) \circ \varphi_n, \quad \varphi_{n+1}s_i = \varphi_n, \quad 0 < i, \quad \varphi_{n+1}s_0 = id_{F_n} \quad (\text{A.24})$$

Next, note that there is no ambiguity in writing simply $\varphi_{n-k}d_{0,\dots,k-1}$ to denote the map (A.23). We now check that $\tilde{F}_{n-1}d_i = d_i\tilde{F}_n$, $0 \leq i \leq n$, which must be verified after restricting to each $k \rightarrow k+1$, $0 \leq k \leq n-2$. There are three cases, depending on i and k :

$$\begin{aligned} (i < k+1) \quad & \varphi_{n-k-1}d_{0,\dots,k-1}d_i = \varphi_{n-k-1}d_{0,\dots,k}; \\ (i = k+1) \quad & \varphi_{n-k-1}d_{0,\dots,k-1}d_i = \varphi_{n-k-1}d_1d_{0,\dots,k-1} = (\varphi_{n-k-1}d_0 \circ \varphi_{n-k})d_{0,\dots,k-1} = (\varphi_{n-k-1}d_{0,\dots,k}) \circ (\varphi_{n-k}d_{0,\dots,k-1}); \\ (i > k+1) \quad & \varphi_{n-k-1}d_{0,\dots,k-1}d_i = \varphi_{n-k-1}d_{i-k}d_{0,\dots,k-1} = \varphi_{n-k}d_{0,\dots,k-1}. \end{aligned}$$

The case of degeneracies is similar. \square

RANTRANS PROP
proof of Proposition 5.40. The result follows from the following string of identifications.

$$\begin{aligned} \lim_{\Delta} (\text{Ran}_{A_n \rightarrow \Sigma_G} N_n) & \simeq \text{Ran}_{\Delta \times \Sigma_G \rightarrow \Sigma_G} (\text{Ran}_{A_n \rightarrow \Sigma_G} N_n) \simeq \\ & \simeq \text{Ran}_{\Delta \times \Sigma_G \rightarrow \Sigma_G} (\text{Ran}_{(\Delta^{op} \ltimes A_\bullet^{op})^{op} \rightarrow \Delta \times \Sigma_G} N_\bullet) \simeq \\ & \simeq \text{Ran}_{(\Delta^{op} \ltimes A_\bullet^{op})^{op} \rightarrow \Sigma_G} N_\bullet \simeq \text{Ran}_{(\Delta^{op} \ltimes A_\bullet^{op})^{op} \rightarrow \Sigma_G} \tilde{N}_\bullet \simeq \text{Ran}_{|A_\bullet| \rightarrow \Sigma_G} \tilde{N} \end{aligned}$$

The first step simply rewrites \lim_{Δ} . The second step follows from Proposition [FIBERKANMAP_PROP](#) applied to the map $(\Delta^{op} \ltimes A_{\bullet}^{op})^{op} \rightarrow \Delta \times \Sigma_G$ of Grothendieck fibrations over Δ . The third step follows since iterated Kan extensions are again Kan extensions. The fourth step twists N_{\bullet} as in Lemma [TWISTING_LEMMA](#) to obtain \tilde{N}_{\bullet} such that the d_i, s_j are given by strictly commutative diagrams for $0 \leq i < n, 0 \leq j \leq n$. Lastly, the final step uses Lemma [SOURCEFACT_LEM](#) to conclude that \tilde{N} factors through the target functor t , obtaining \tilde{N} , and then uses Corollary [SOURCELANFINAL_COR](#) to conclude that the Kan extensions indeed coincide. \square

References

- [BM03](#) [1] C. Berger and I. Moerdijk. Axiomatic homotopy theory for operads. *Commentarii Mathematici Helvetici*, 78:805–831, 2003.
- [BM08](#) [2] C. Berger and I. Moerdijk. On an extension of the notion of Reedy category. *Math. Z.*, 269(3-4):977–1004, 2011.
- [BH15](#) [3] A. J. Blumberg and M. A. Hill. Operadic multiplications in equivariant spectra, norms, and transfers. *Adv. Math.*, 285:658–708, 2015.
- [BV73](#) [4] M. Boardman and R. Vogt. *Homotopy invariant algebraic structures on topological spaces*, volume 347 of *Lecture Notes in Mathematics*. Springer-Verlag, 1973.
- [CM13b](#) [5] D.-C. Cisinski and I. Moerdijk. Dendroidal sets and simplicial operads. *J. Topol.*, 6(3):705–756, 2013.
- [CW91](#) [6] S. R. Costenoble and S. Waner. Fixed set systems of equivariant infinite loop spaces. *Trans. Amer. Math. Soc.*, 326(2):485–505, 1991.
- [Elm83](#) [7] A. D. Elmendorf. Systems of fixed point sets. *Transactions of the American Mathematical Society*, 277:275–284, 1983.
- [Hatcher](#) [8] A. Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [HHR](#) [9] M. A. Hill, M. J. Hopkins, and D. C. Ravenel. On the non-existence of elements of Kervaire invariant one. *Annals of Mathematics*, 184:1–262, 2016.
- [Hi03](#) [10] P. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [McL](#) [11] S. Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [May72](#) [12] J. P. May. *The geometry of iterated loop spaces*. Springer-Verlag, Berlin-New York, 1972. Lectures Notes in Mathematics, Vol. 271.
- [MW07](#) [13] I. Moerdijk and I. Weiss. Dendroidal sets. *Algebr. Geom. Topol.*, 7:1441–1470, 2007.
- [Pe16](#) [14] L. A. Pereira. Cofibrancy of operadic constructions in positive symmetric spectra. *Homology Homotopy Appl.*, 18(2):133–168, 2016.
- [Pe17](#) [15] L. A. Pereira. Equivariant dendroidal sets. arXiv preprint: 1702.08119, 2017.
- [Pia91](#) [16] R. J. Piacenza. Homotopy theory of diagrams and CW-complexes over a category. *Canadian Journal of Mathematics*, 43:814–824, 1991.
- [Rub17](#) [17] J. Rubin. On the realization problem for N_{∞} operads. arXiv preprint: 1705.03585, 2017.
- [SS00](#) [18] S. Schwede and B. E. Shipley. Algebras and modules in monoidal model categories. *Proc. London Math. Soc. (3)*, 80(2):491–511, 2000.
- [Spitz01](#) [19] M. Spitzweck. Operads, algebras and modules in general model categories. arXiv preprint: 0101102, 2001.
- [Ste16](#) [20] M. Stephan. On equivariant homotopy theory for model categories. *Homology Homotopy Appl.*, 18(2):183–208, 2016.

We12

[21] I. Weiss. Broad posets, trees, and the dendroidal category. Available at: <https://arxiv.org/abs/1201.3987>, 2012.

Whi14

[22] D. White. Monoidal Bousfield localizations and algebras over operads. arXiv preprint: 1404.5197v1, 2014.

WY15

[23] D. White and D. Yau. Bousfield localization and algebras over colored operads. arXiv preprint: 1503.06720v2, 2015.