

Genuine equivariant operads

Peter Bonventre, Luís A. Pereira

April 14, 2017

Abstract

We build new algebraic structures, which we call genuine equivariant operads, which can be thought of as a hybrid between equivariant operads and coefficient systems. We then prove an Elmendorf type theorem stating that equivariant operads, with their graph model structure, are equivalent to genuine equivariant operads with their projective model structure.

As an application, we build explicit models for the N_∞ -operads of Blumberg and Hill.

Contents

1	Introduction	1
2	Planar and tall maps	2
2.1	Planar structures	2
2.2	Outer faces and tall maps	6
2.3	Substitution	7
3	The genuine equivariant operad monad	9
3.1	Wreath product over finite sets	9
3.2	Equivariant leaf-root and vertex functors	10
3.3	Planar strings	13
3.4	A monad on spans	14
4	Filtration of Cellular Extensions	18
4.1	Stuff Which Probably Will Be Included in an Above Section	18
4.2	Filtration Lemmas	20
4.3	Notation	22
5	Appendix	24
5.1	Grothendieck fibrations	24
5.2	Kan Extensions - May Also be Part of the Above	24
5.2.1	Naturality	24
5.2.2	Left Kan Extensions and Pushouts	26

1 Introduction

No content yet.

2 Planar and tall maps

2.1 Planar structures


Throughout we will work with trees possessing *planar structures* or, more intuitively, trees embedded into the plane.

Our preferred model for trees will be that of broad posets first introduced by Weiss in [3] and further worked out by the second author in [2]. We now define planar structures in this context.

Definition 2.1. Let $T \in \Omega$ be a tree. A *planar structure* of T is an extension of the descendanty partial order \leq_d to a total order \leq_p such that:

- *Planar*: if $e \leq_p f$ and $e \not\leq_d f$ then $g \leq_d f$ implies $e \leq_p g$.

Example 2.2. An example of a planar structure on a tree T follows, with \leq_r encoded by the number labels.



(2.3)

PLANAREX EQ

Intuitively, given a planar depiction of a tree T , $e \leq_d f$ holds when the downward path from e passes through f and $e \leq_p f$ holds if either $e \leq_d f$ or if the downward path from e is to the left of the downward path from f (as measured at the node where the paths intersect).

Intuitively, a planar depiction of a tree amounts to choosing a total order for each of the sets of *input edges* of each node (i.e. those edges immediately above that node).

While we will not need to make this last statement precise, we will nonetheless find it convenient to show that Definition 2.1 is equivalent to such choosing total orders for each of the sets of input edges. To do so, we first introduce some notation.

Notation 2.4. Let $T \in \Omega$ be a tree and $e \in T$ and edge. We will denote

$$I(e) = \{f \in T : e \leq_d f\}$$

and refer to this poset as the *input path* of e .

We will repeatedly use the following, which is a consequence of [2, Cor. 5.26].

Lemma 2.5. If $e \leq_d f$, $e \leq_d f'$, then f, f' are \leq_d -comparable.

Proposition 2.6. Let $T \in \Omega$ be a tree. Then

- (a) for any $e \in T$ the finite poset $I(e)$ is totally ordered;
- (b) the poset (T, \leq_d) has all joins, denoted \vee . In fact, $\vee_i e_i = \min(\cap_i I(e_i))$.

Proof. (a) is immediate from Lemma 2.5. To prove (b) we note that $\min(\cap_i I(e_i))$ exists by (a), and that this is clearly the join $\vee e_i$. \square

Notation 2.7. Let $T \in \Omega$ be a tree and suppose that $e <_d b$. We will denote by $b_e^\dagger \in T$ the predecessor of b in $I(e)$.

Proposition 2.8. Suppose e, f are \leq_d -incomparable edges of T and write $b = e \vee f$. Then

- (a) $e <_d b$, $f <_d b$ and $b_e^\dagger \neq b_f^\dagger$;
- (b) $b_e^\dagger, b_f^\dagger \in b^\dagger$. In fact $\{b_e^\dagger\} = I(e) \cap b^\dagger$, $\{b_f^\dagger\} = I(f) \cap b^\dagger$;

(c) if $e' \leq_d e$, $f' \leq_d f$ then $b = e' \vee f'$ and $b_{e'}^\dagger = b_e^\dagger$, $b_{f'}^\dagger = b_f^\dagger$.

Proof. (a) is immediate: the condition $e = g$ (resp. $f = g$) would imply $f \leq_d e$ (resp. $e \leq_d f$) while the condition $b_e^\dagger = b_f^\dagger$ would provide a predecessor of b in $I(e) \cap I(f)$.

For (b), note that any relation $a <_d b$ factors as $a \leq_d b_a^* <_d b$ for some unique $b_a^* \in b^\dagger$, where uniqueness follows from Lemma 2.5. Choosing $a = e$ implies $I(e) \cap b^\dagger = \{b_e^*\}$ and letting a range over edges such that $e \leq_d a <_d b$ shows that b_e^* is in fact the predecessor of b .

To prove (c) one reduces to the case $e' = e$, in which case it suffices to check $I(e) \cap I(f') = I(e) \cap I(f)$. But if it were otherwise there would exist an edge a satisfying $f' \leq_d a <_d f$ and $e \leq_d a$, and this would imply $e \leq_d f$, contradicting our hypothesis. \square

Proposition 2.9. Let $c = e_1 \vee e_2 \vee e_3$. Then $c = e_i \vee e_j$ iff $c_{e_i}^\dagger \neq c_{e_j}^\dagger$.
Therefore, all ternary joins in (T, \leq_d) are binary, i.e.


$$c = e_1 \vee e_2 \vee e_3 = e_i \vee e_j \quad (2.10)$$

for some $1 \leq i < j \leq 3$, and (2.10) fails for at most one choice of $1 \leq i < j \leq 3$.

Proof. If $c_{e_i}^\dagger \neq c_{e_j}^\dagger$, then $c = \min(I(e_i) \cap I(e_j)) = e_i \vee e_j$, whereas the converse follows from Proposition 2.8(a).

The “therefore” part follows by noting that $c_{e_1}^\dagger, c_{e_2}^\dagger, c_{e_3}^\dagger$ can not all coincide, or else c would not be the minimum of $I(e_1) \cap I(e_2) \cap I(e_3)$. \square

Example 2.11. In the following example $b = e \vee f$, $c = e \vee f \vee g$, $c_e^\dagger = c_f^\dagger = b$.



Notation 2.12. Given a set S of size n we write $\text{Ord}(S) \simeq \text{Iso}(S, \{1, \dots, n\})$. We will usually abuse notation by regarding its objects as pairs (S, \leq) where \leq is a total order in S .

Proposition 2.13. Let $T \in \Omega$ be a tree. There is a bijection

$$\{\text{planar structures } (T, \leq_p)\} \longrightarrow \prod_{(a^\dagger \leq a) \in V(T)} \text{Ord}(a^\dagger) \quad (2.14)$$

$$\leq_p \longmapsto (\leq_p \upharpoonright_{a^\dagger})$$

Proof. We will keep the setup of Proposition 2.8 throughout: e, f are \leq_d -incomparable edges and we write $b = e \vee f$.

We first show that (2.14) is injective, i.e. that the restrictions $\leq_p \upharpoonright_{a^\dagger}$ determine if $e <_p f$ holds or not. If $b_e^\dagger <_p b_f^\dagger$, the relations $e \leq_d b_e^\dagger <_p b_f^\dagger \geq_d f$ and Definition 2.1 imply it must be $e <_p f$. Dually, if $b_f^\dagger <_p b_e^\dagger$ then $f <_p e$. Thus $b_e^\dagger <_p b_f^\dagger \Leftrightarrow e <_p f$ and hence (2.14) is indeed injective.

To check that (2.14) is surjective, it suffices (recall that e, f are assumed \leq_d -incomparable) to check that defining $e \leq_p f$ to hold iff $b_e^\dagger < b_f^\dagger$ holds in b^\dagger yields a planar structure.

Antisymmetry and the total order conditions are immediate, and it thus remains to check the transitivity and planar conditions. Transitivity of \leq_p in the case $e' \leq_p e <_p f$ and the planar condition, which is the case $e <_p f \geq_d f'$, follow from Proposition 2.8(c). Transitivity of \leq_p in the case $e <_p f \leq_d f'$ follows since either $e \leq_d f'$ or else e, f' are \leq_d -incomparable, in which case one can apply 2.8(c) with the roles of f, f' reversed.


It remains to check transitivity in the hardest case, that of $e <_p f <_p g$ with e, f, g incomparable. We write $c = e \vee f \vee g$. By the “therefore” part of Proposition 2.9, either (i) $e \vee f <_d c$, in which case Proposition 2.9 implies $c_e^\dagger = c_f^\dagger$ and transitivity follows; (ii) $f \vee g <_d c$, which follows just as (i); (iii) $e \vee f = f \vee g = c$, in which case $c_e^\dagger < c_f^\dagger < c_g^\dagger$ in c^\dagger so that $c_e^\dagger \neq c_g^\dagger$ and by Proposition 2.9 it is also $e \vee g = c$ and transitivity follows. \square

Remark 2.15. Definition 2.1 readily extends to forests $F \in \Phi$. The analogue of Proposition 2.13 then states that the data of a planar structure is equivalent to total orderings of the nodes of F together with a total ordering of its set of roots. Indeed, this follows by either adapting the proof above or by noting that planar structures on F are clearly in bijection with planar structures on the join tree $F \star \eta$ (cf. [2, Def. 7.44]), which adds a single edge η to F , serving as the (unique) root of $F \star \eta$.

When discussing the substitution procedure in §2.3 we will find it convenient to work with a model for the category Ω that possesses exactly one representative of each possible planar structure on each tree or, more precisely, such that the only isomorphisms preserving the planar structures are the identities. On the other hand, using such a model for Ω throughout would, among other issues, make the discussion of faces in §2.2 rather awkward. We now outline our conventions to address such issues.

Let Ω^p , the category of *planarized trees*, denote the category with objects pairs $T_{\leq_p} = (T, \leq_p)$ of trees together with a planar structure and morphisms the *underlying* maps of trees (so that the planar structures are ignored). There is a full subcategory $\Omega^s \hookrightarrow \Omega^p$, whose objects we call *standard models*, of those T_{\leq_p} whose underlying set is one of the sets $\underline{n} = \{1, 2, \dots, n\}$ and for which \leq_p coincides with the canonical order.

Example 2.16. Some examples of standard models, i.e. objects of Ω^s , follow (further, (2.3) can also be interpreted as such an example).



Here T_1 and T_2 are isomorphic to each other but not isomorphic to any other standard model in Ω^s while both C and U are the unique objects in their isomorphism classes.

Given $T_{\leq_p} \in \Omega^p$ there is an obvious standard model $T_{\leq_p}^s \in \Omega^s$ given by replacing each edge by its order following \leq_p . Indeed, this defines a retraction $(-)^s: \Omega^p \rightarrow \Omega^s$ and a natural transformation $\sigma: id \Rightarrow (-)^s$ given by isomorphisms preserving the planar structure (in fact, the pair $((-)^s, \sigma)$ is clearly unique).

Convention 2.18. From now on, we will write simply Ω, Ω_G to denote the categories Ω^s, Ω_G^s of standard models (where planar structures are defined in the underlying forest as in Remark 2.15). Similarly \mathbf{O}_G will denote the model \mathbf{O}_G^s for the orbital category whose objects are the orbital G -sets whose underlying set is one of the sets $\underline{n} = \{1, 2, \dots, n\}$.

Therefore, whenever one of our constructions produces an object/diagram in $\Omega^p, \Omega_G^p, \mathbf{O}_G^p$ (of trees, G -trees, orbital G -sets with a planarization/total order) we will hence implicitly reinterpret it by using the standardization functor $(-)^s$.

Example 2.19. To illustrate our convention, we consider the trees in Example 2.16.

One has subfaces $F_1 \subset F_2 \subset U$ where F_1 is the subtree with edge set $\{1, 2, 6, 7\}$ and F_2 is the subtree with edge set $\{1, 2, 3, 6, 7\}$, both with inherited tree and planar structures. Applying $(-)^s$ to the inclusion diagram on the left below then yields a diagram as on the right.




Similarly, let $\leq_{(12)}$ and $\leq_{(45)}$ denote alternate planar structures for U exchanging the orders of the pairs 1, 2 and 4, 5, so that one has objects $U_{\leq_{(12)}}$, $U_{\leq_{(45)}}$ in Ω^p . Applying $(-)^s$ to the diagram of underlying identities on the left yields the permutation diagram on the right.

$$\begin{array}{ccc} U & \xrightarrow{id} & U_{\leq_{(45)}} \\ id \searrow & & \nearrow id \\ & U_{\leq_{(12)}} & \end{array} \qquad \begin{array}{ccc} U & \xrightarrow{(45)} & U \\ (12) \searrow & & \nearrow (12)(45) \\ & U & \end{array}$$

Example 2.20. An additional reason to leave the use of $(-)^s$ implicit is that when depicting G -trees it is preferable to choose edge labels that describe the action rather than the planarization (which is already implicit anyway).

For example, when $G = \mathbb{Z}/4$, in both diagrams below the orbital representation on the left represents the isomorphism class consisting of the two trees $T_1, T_2 \in \Omega_G$ on the right.



Definition 2.21. A morphism $S \xrightarrow{\varphi} T$ in Ω that is compatible with the planar structures \leq_p is called a *planar map*.

More generally, a morphism $F \rightarrow G$ in the categories Φ , Φ^G , Ω^G of forests, G -forests, G -trees is called a *planar map* if it is an independent map (cf. [2, Def. 5.28]) compatible with the planar structures \leq_p .

Remark 2.22. The need for the independence condition is justified by [2, Lemma 5.33] and its converse, since non independent maps do not reflect \leq_d inequalities.

We note that in the Ω_G case a map φ is independent iff φ does not factor through a non trivial quotient iff φ is injective on each edge orbit.

Proposition 2.23. Let $F \xrightarrow{\varphi} G$ be an independent map in Φ (or Ω , Ω_G , Φ_G). Then there is a unique factorization

$$F \xrightarrow{\cong} \bar{F} \rightarrow G$$

such that $F \xrightarrow{\cong} \bar{F}$ is an isomorphism and $\bar{F} \rightarrow G$ is planar.

Proof. We need to show that there is a unique planar structure $\leq_{\bar{F}}$ on the underlying forest of F making the underlying map a planar map. Simplicity of G ensures that for any vertex $e^\dagger \leq e$ of F the edges in $\varphi(e^\dagger)$ are all distinct while independence of φ likewise ensures that the edges in $\varphi(e^\dagger)$ are distinct. The result now follows from (the forest version of) Proposition 2.13: one simply orders each set e^\dagger and \underline{r}_F according to its image.

not quite complete... maybe that \leq_p is the closure of \leq_d and the vertex relations under transitivity and the planar condition \square

Remark 2.24. Proposition 2.23 says that planar structures can be pulled back along independent maps. However, they can not always be pushed forward. As an example, in the notation of (2.17), consider the map $C \rightarrow T_1$ defined by $1 \mapsto 1$, $2 \mapsto 4$, $3 \mapsto 2$, $4 \mapsto 5$.

Remark 2.25. Given any tree $T \in \Omega$ there is a unique corolla $\text{lr}(T) \in \Sigma$ and planar tall map $\text{lr}(T) \rightarrow T$. Explicitly, the number of leaves of $\text{lr}(T)$ matches that of T , together with the inherited order.

2.2 Outer faces and tall maps

In preparation for our discussion of the substitution operation in §2.3, we now recall some basic notions and results concerning outer subtrees and tree grafting, as in [2, §5].

Definition 2.26. Let $T \in \Omega$ be a tree and $e_1 \cdots e_n = \underline{e} \leq e$ a broad relation in T .

We define the *planar outer face* $T_{\underline{e} \leq e}$ to be the subtree with underlying set those edges $f \in T$ such that

$$f \leq_d e, \quad \forall_i e_i \not\leq_d f, \quad (2.27)$$

generating broad relations the relations $f^\dagger \leq f$ for f satisfying (2.27) and $\forall_i f \neq e_i$, and planar structure pulled back from T .

Remark 2.28. If one forgoes the requirement that $T_{\underline{e} \leq e}$ be equipped with the pullback planar structure, the inclusion $T_{\underline{e} \leq e} \rightarrow T$ is usually called simply an *outer face*.

We now recap some basic results.

Proposition 2.29. Let $T \in \Omega$ be a tree.

- (a) $T_{\underline{e} \leq e}$ is a tree with root e and edge tuple \underline{e} ;
- (b) there is a bijection

$$\{\text{planar outer faces of } T\} \leftrightarrow \{\text{broad relations of } T\};$$

- (c) if $R \rightarrow S$ and $S \rightarrow T$ are outer face maps then so is $R \rightarrow T$;
- (d) any pair of broad relations $\underline{g} \leq v$, $\underline{f}v \leq e$ induces a grafting pushout diagram

$$\begin{array}{ccc} \eta & \xrightarrow{v} & T_{\underline{g} \leq v} \\ v \downarrow & & \downarrow \\ T_{\underline{f}v \leq e} & \longrightarrow & T_{\underline{f}g \leq v} \end{array} \quad (2.30)$$

Proof. We first show (a). That $T_{\underline{e} \leq e}$ is indeed a tree is the content of [2, Prop. 5.20]; more precisely, $T_{\underline{e} \leq e} = (T^{\leq e})_{< \underline{e}}$ in the notation therein. That the root of $T_{\underline{e} \leq e}$ is e is clear and that the root tuple is \underline{e} follows from [2, Remark 5.23].

(b) follows from (a), which shows that $\underline{e} \leq e$ can be recovered from $T_{\underline{e} \leq e}$.

(c) follows from the definition of outer face together with [2, Lemma 5.33], which states that the \leq_d relations on S, T coincide.

Since by (c) both $T_{\underline{g} \leq v}$ and $T_{\underline{f}v \leq e}$ are outer faces of $T_{\underline{f}g \leq v}$, (d) is a restatement of [2, Prop. 5.15]. \square

Definition 2.31. A map $S \xrightarrow{\varphi} T$ in Ω is called a *tall map* if

$$\varphi(l_S) = l_T, \quad \varphi(r_S) = r_T,$$

where $l_{(-)}$ denotes the leaf tuple and $r_{(-)}$ the root.

The following is a restatement of [2, Cor. 5.24]

Proposition 2.32. Any map $S \xrightarrow{\varphi} T$ in Ω has a factorization, unique up to unique isomorphism,

$$S \xrightarrow{\varphi^t} U \xrightarrow{\varphi^u} T$$

as a tall map followed by an outer face (in fact, $U = T_{\varphi(l_S) \leq r_S}$).

We recall that a face $F \rightarrow T$ is called inner if is obtained by iteratively removing inner edges, i.e. edges other than the root or the leaves. In particular, it follows that a face is inner iff it is tall. The usual face-degeneracy decomposition thus combines with Corollary 2.32 to give the following.

Corollary 2.33. Any map $S \xrightarrow{\varphi} T$ in Ω has a factorization, unique up to unique isomorphisms,

$$S \xrightarrow{\varphi^-} U \xrightarrow{\varphi^i} V \xrightarrow{\varphi^u} T \quad (2.34) \quad \text{TRIPLEFACT EQ}$$

as a degeneracy followed by an inner face followed by an outer face.


Proof. The factorization (2.34) can be built by first performing the degeneracy-face decomposition and then performing the tall-outer decomposition on the face map. \square

2.3 Substitution

One of the key ideas needed to describe operads is that of substitution of tree nodes, a process that we will prefer to repackage in terms of maps of trees. We start by discussing an example, focusing on the related notion of iterated graftings of trees (as described in (2.30)).

Example 2.35. The trees U_1, U_2, \dots, U_6 on the left below can be grafted into the tree U in the middle. More precisely (among other possible grafting orders), one has

$$U = (((((U_6 \sqcup_a U_2)) \sqcup_a U_1) \sqcup_b U_3) \sqcup_d U_5) \sqcup_c U_4) \quad (2.36) \quad \text{UFORMULA EQ}$$



We now consider the tree T , which is built by converting each U_i into the corolla $\text{Irr}(U_i)$ (cf. Remark 2.25), and then performing the same grafting operations as in (2.36). T can then be regarded as encoding the combinatorics of the iterated grafting in (2.36), with alternative ways to reorder operations in (2.36) in bijection with ways to assemble T out of its nodes.

One can now therefore think of the iterated grafting (2.36) as being instead encoded by

the tree T together with the (unique) planar tall maps φ_i below.

(2.38)

SUBSDATUMTREES2 EQ

From this perspective, U can now be thought as obtained from T by substituting each of its nodes with the corresponding U_i . Moreover, the φ_i assemble to a planar tall map $\varphi: T \rightarrow U$ (such that $a_i \mapsto a, b \mapsto b, \dots, e \mapsto e$), which likewise encodes the same information.

Our perspective will then be that data for substitution of tree nodes such as in (2.38) can equivalently be repackaged using planar tall maps.

SUBSDATUMTREES2 EQ

UBSTITUTIONDATUM

Definition 2.39. Let $T \in \Omega$ be a tree.

A T -substitution datum is a tuple $\{U_{e \uparrow \leq e}\}_{(e \uparrow \leq e) \in V(T)}$ such that $\text{lr}(U_{e \uparrow \leq e}) = T_{e \uparrow \leq e}$.

Further, a map of T -substitution data $\{U_{e \uparrow \leq e}\} \rightarrow \{V_{e \uparrow \leq e}\}$ is a tuple of planar tall maps $\{U_{e \uparrow \leq e} \rightarrow V_{e \uparrow \leq e}\}$.

Definition 2.40. Let $T \in \Omega$.

The *Segal core poset* $\text{Sc}(T)$ is the poset with objects the edge subtrees η_e and vertex subtrees $T_{e \uparrow \leq e}$. The order relation is given by inclusion.

Remark 2.41. Note that the only maps in $\text{Sc}(T)$ are inclusions of the form $\eta_a \subset T_{e \uparrow \leq e}$. In particular, there are no pairs of composable non-identity relations in $\text{Sc}(T)$.

Given a T -substitution datum $\{U_{e \uparrow \leq e}\}$ we abuse notation by writing

$$U_{(-)}: \text{Sc}(T) \rightarrow \Omega$$

for the functor $\eta_a \mapsto \eta, T_{e \uparrow \leq e} \mapsto U_{e \uparrow \leq e}$ and sending the inclusions $\eta_a \subset T_{e \uparrow \leq e}$ to the composites

$$\eta \xrightarrow{a} T_{e \uparrow \leq e} = \text{lr}(U_{e \uparrow \leq e}) \rightarrow U_{e \uparrow \leq e}.$$

Proposition 2.42. Let $T \in \Omega$ be a tree. There is an isomorphism of categories

$$\begin{aligned} \text{Sub}(T) &\xleftrightarrow{\quad} \Omega_{T/}^{\text{pt}} \\ \{U_{e \uparrow \leq e}\} &\longmapsto (T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}) \\ \{U_{\varphi(e \uparrow) \leq \varphi(e)}\} &\longleftarrow (T \xrightarrow{\varphi} U) \end{aligned} \tag{2.43}$$

SUBDATAUNDERPLAN EQ

Where $\text{Sub}(T)$ denotes the category of T -substitution data and $\Omega_{T/}^{\text{pt}}$ the category of planar tall maps under T .

Proof. We first claim that (i) the $\text{colim}_{\text{Sc}(T)} U_{(-)}$ indeed exists; (ii) for the canonical datum $\{T_{e \uparrow \leq e}\}$, it is $T = \text{colim}_{\text{Sc}(T)} T_{(-)}$; (iii) the induced map $T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}$ is planar tall.

The argument is by induction on the number of vertices of T , with the base cases of T with 0 or 1 vertices being immediate, since then T is the terminal object of $\text{Sc}(T)$. Otherwise, one can choose a non trivial grafting decomposition so as to write $T = R \sqcup_e S$,

resulting in identifications $\text{Sc}(R) \subset \text{Sc}(T)$, $\text{Sc}(S) \subset \text{Sc}(T)$ so that $\text{Sc}(R) \cup \text{Sc}(S) = \text{Sc}(T)$ and $\text{Sc}(R) \cap \text{Sc}(S) = \{\eta_e\}$. The existence of $\text{colim}_{\text{Sc}(T)} U_{(-)}$ is thus equivalent to the existence of the pushout below.

$$\begin{array}{ccc} \eta & \longrightarrow & \text{colim}_{\text{Sc}(R)} U_{(-)} \\ \downarrow & & \downarrow \\ \text{colim}_{\text{Sc}(S)} U_{(-)} & \dashrightarrow & \text{colim}_{\text{Sc}(T)} U_{(-)} \end{array} \quad (2.44)$$

ASSEMBLYGRAFT EQ

By induction, the top right and bottom left colimits exist for any $U_{(-)}$, equal R and S in the case $U_{(-)} = T_{(-)}$, and the maps $R \rightarrow \text{colim}_{\text{Sc}(R)} U_{(-)}$, $S \rightarrow \text{colim}_{\text{Sc}(S)} U_{(-)}$ are planar tall. But is now follows that (2.44) is a grafting pushout diagram, so that the pushout indeed exists. The conditions that $T = \text{colim}_{\text{Sc}(T)} T_{(-)}$ and $T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}$ is planar tall follow.

The fact that the two functors in (2.43) are inverse to each other is clear by the same inductive argument. \square

Remark 2.45. It follows from the previous proof that, writing $U = \text{colim}_{\text{Sc}(T)} U_{(-)}$, one has

$$V(U) = \coprod_{(e^\dagger \leq e) \in V(T)} V(U_{e^\dagger \leq e}). \quad (2.46)$$

VERTEXDECOMP EQ

Alternatively, (2.46) can be regarded as a map $f^*: V(U) \rightarrow V(T)$ induced by the planar tall map $f: T \rightarrow U$. Explicitly, $f^*(U_{u^\dagger \leq u})$ is the unique $T_{t^\dagger \leq t}$ such that $U_{u^\dagger \leq u} \subset U_{t^\dagger \leq t}$. We note that f^* is indeed contravariant in the tall planar map f .

3 The genuine equivariant operad monad

We now turn to the task of building the monad encoding genuine equivariant operads.

3.1 Wreath product over finite sets

In what follows we will let \mathbf{F} denote the usual skeleton of the category of finite sets and all set maps. Explicitly, its objects are the finite sets $\{1, 2, \dots, n\}$ for $n \geq 0$. However, much as in the discussion in Convention 2.18 we will often find it more convenient to regard the elements of \mathbf{F} as equivalence classes of finite sets equipped with total orders.

Definition 3.1. For a category \mathcal{C} , we let $\mathbf{F} \wr \mathcal{C}$ denote the opposite of the Grothendieck construction for the functor

$$\begin{array}{ccc} \mathbf{F}^{op} & \longrightarrow & \mathbf{Cat} \\ I & \longmapsto & \mathcal{C}^I \end{array}$$

Explicitly, the objects of $\mathbf{F} \wr \mathcal{C}$ are tuples $(c_i)_{i \in I}$ and a map $(c_i)_{i \in I} \rightarrow (d_j)_{j \in J}$ consists of a pair

$$(\phi: I \rightarrow J, (f_i: c_i \rightarrow d_{\phi(i)})_{i \in I}),$$

henceforth abbreviated as $(\phi, (f_i))$.

The following is immediate.

Proposition 3.2. Suppose \mathcal{C} has all finite coproducts. One then has a functor as on the left below. Dually, if \mathcal{C} has all finite products, one has a functor as on the right below.

$$\begin{array}{ccc} \mathbf{F} \wr \mathcal{C} & \xrightarrow{\Pi} & \mathcal{C} \\ (c_i)_{i \in I} & \longmapsto & \coprod_{i \in I} c_i \end{array} \quad \begin{array}{ccc} (\mathbf{F} \wr \mathcal{C}^{op})^{op} & \xrightarrow{\Pi} & \mathcal{C} \\ (c_i)_{i \in I} & \longmapsto & \prod_{i \in I} c_i \end{array}$$

Lemma 3.3. Suppose that \mathcal{E} is a bicomplete category such that coproducts commute with limits in each variable. If the leftmost diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{E} \\ k \downarrow & \nearrow \eta & \\ \mathcal{D} & & \end{array} \quad \begin{array}{ccccc} \mathbf{F} \wr \mathcal{C} & \xrightarrow{\mathbf{F} \wr F} & \mathbf{F} \wr \mathcal{E} & \xrightarrow{\Pi} & \mathcal{E} \\ \mathbf{F} \wr k \downarrow & \nearrow \mathbf{F} \wr \eta & \nearrow \mathbf{F} \wr G & \searrow \Pi \circ \mathbf{F} \wr G & \\ \mathbf{F} \wr \mathcal{D} & & & & \end{array} \quad (3.4) \quad \boxed{\text{WRRAN EQ}}$$

is a right Kan extension diagram then so is the composite of the rightmost diagram.

Dually, if in \mathcal{E} products commute with colimits in each variable, and the leftmost diagram

$$\begin{array}{ccc} \mathcal{C}^{op} & \xrightarrow{F} & \mathcal{E} \\ k \downarrow & \nearrow \epsilon & \\ \mathcal{D}^{op} & & \end{array} \quad \begin{array}{ccccc} (\mathbf{F} \wr \mathcal{C})^{op} & \xrightarrow{(\mathbf{F} \wr F)^{op}} & (\mathbf{F} \wr \mathcal{E}^{op})^{op} & \xrightarrow{\Pi} & \mathcal{E} \\ (\mathbf{F} \wr k)^{op} \downarrow & \nearrow & \nearrow (\mathbf{F} \wr G)^{op} & \searrow \Pi \circ (\mathbf{F} \wr G)^{op} & \\ (\mathbf{F} \wr \mathcal{D})^{op} & & & & \end{array} \quad (3.5) \quad \boxed{\text{WRLAN EQ}}$$

is a left Kan extension diagram then so is the composite of the rightmost diagram.

Proof. Unpacking definitions using the pointwise formula for Kan extensions ($\frac{\text{McL}}{\text{I, X.3.1}}$), the claim concerning (3.4) amounts to showing that for each $(d_i) \in \mathbf{F} \wr \mathcal{D}$ one has natural isomorphisms

$$\lim_{((d_i) \rightarrow (kc_j)) \in ((d_i) \downarrow \mathbf{F} \wr \mathcal{C})} \left(\coprod_j F(c_j) \right) \simeq \coprod_i \lim_{(d_i \rightarrow kc_i) \in d_i \downarrow \mathcal{C}} (F(c_i)). \quad (3.6) \quad \boxed{\text{POINTKAN EQ}}$$

Noting that the canonical factorizations of each $(\varphi, (f_i)): (d_i)_{i \in I} \rightarrow (kc_j)_{j \in J}$ as

$$(d_i)_{i \in I} \rightarrow (c_{\phi(i)})_{i \in I} \rightarrow (kc_j)_{j \in J}$$

exhibit $\prod_i (d_i \downarrow \mathcal{C})$ as a coreflexive subcategory of $(d_i) \downarrow \mathbf{F} \wr \mathcal{C}$, we see that it is an initial subcategory. Therefore

$$\lim_{((d_i) \rightarrow (kc_j)) \in ((d_i) \downarrow \mathbf{F} \wr \mathcal{C})} \left(\coprod_j F(c_j) \right) \simeq \lim_{((d_i) \rightarrow (kc_i)) \in \prod_i (d_i \downarrow \mathcal{C})} \left(\coprod_i F(c_i) \right)$$

and hence (3.6) now follows from the assumption that coproducts commute with limits in each variable. \square

Notation 3.7. Using the coproduct functor $\mathbf{F}^{\wr 2} = \mathbf{F}^{\wr \{0,1\}} = \mathbf{F} \wr \mathbf{F} \xrightarrow{\Pi} \mathbf{F}$ (where $\coprod_{i \in I} J_i$ is ordered lexicographically) and the simpleton $\{1\} \in \mathbf{F}$ one can regard the collection of categories $\mathbf{F}^{\wr \{0, \dots, n\}} \wr \mathcal{C} = \mathbf{F}^{\wr n} \wr \mathcal{C}$ as a coaugmented cosimplicial object in \mathbf{Cat} . As such, we will denote by

$$\delta^i: \mathbf{F}^{\wr n-1} \wr \mathcal{C} \rightarrow \mathbf{F}^{\wr n} \wr \mathcal{C}, \quad 0 \leq i \leq n$$

the cofaces obtained by inserting simpletons $\{1\} \in \mathbf{F}$ and by

$$\sigma^i: \mathbf{F}^{\wr n+1} \wr \mathcal{C} \rightarrow \mathbf{F}^{\wr n} \wr \mathcal{C}, \quad 0 \leq i \leq n$$

the codegeneracies obtained by applying the coproduct $\mathbf{F}^{\wr 2} \xrightarrow{\Pi} \mathbf{F}$ to adjacent \mathbf{F} coordinates.

3.2 Equivariant leaf-root and vertex functors

Definition 3.8. A morphism $T \xrightarrow{\varphi} S$ in Ω_G is called a *quotient* if the underlying morphism of forests

$$\coprod_{[g] \in G/H} T_{[g]} \rightarrow \coprod_{[h] \in G/K} S_{[h]}$$

maps each tree component (or, equivalently, some tree component) isomorphically onto its image component.

We denote the subcategory of G -trees and quotients by Ω_G^q .


Definition 3.9. The G -symmetric category, which we will also call the *category of G -corollas*, is the full subcategory $\Sigma_G \subset \Omega_G^q$ of those G -trees that are corollas, i.e. G -trees such that each edge is either a root or a leaf (but not both).

Definition 3.10. The *leaf-root functor* is the functor $\Omega_G^q \xrightarrow{\text{lr}} \Sigma_G$ defined by

$$\text{lr}(T) = \{\text{leaves of } T\} \sqcup \{\text{roots of } T\}$$

with a broad relation $l_1 \cdots l_n \leq_r$ holding in $\text{lr}(T)$ iff its image holds in T and similarly for the planar structure \leq_p .

Remark 3.11. Generalizing Remark 2.25, $\text{lr}(T)$ can alternatively be characterized as being the *unique* G -corolla which admits an also unique (tree-wise) tall planar map $\text{lr}(T) \rightarrow T$. Moreover, $\text{lr}(T)$ can usually be regarded as the “smallest inner face” of T , obtained by removing all the inner edges, although this characterization fails when $T = G \cdot_H \eta$ is a stick G -tree. Some examples with $G = \mathbb{Z}_{/4}$ follow.




Remark 3.12. One consequence of the fact that planarizations can not be pushed forward along tree maps (cf. Remark 2.24) is that $\text{lr}: \Omega_G^q \rightarrow \Sigma_G$ is not a categorical fibration. **maybe add to this.**

Definition 3.13. Given $T \in \Omega_G$ we define the set $V_G(T)$ of G -vertices of T to be the orbit set $V(T)/G$, i.e. the quotient of the vertex set $V(T)$ by its G -action.

Furthermore, we will regard $V_G(T)$ as an object in \mathbf{F} by equipping it with its lexicographic order: i.e. vertex equivalence classes $[e^\dagger \leq e]$ are ordered according to the planar order \leq_p of the smallest representative ge , $g \in G$.

Remark 3.14. Following Remark 2.45, a planar tall map $f: T \rightarrow U$ of G -trees induces a G -equivariant map $f^*: V(U) \rightarrow V(T)$ and thus also a map of orbits $f^*: V_G(U) \rightarrow V_G(T)$. We note, however, that f^* is not in general compatible with the order on V_G , as is indeed the case even in the non-equivariant case.

A minimal example follows.



In $V(T)$ the vertices are ordered as $a < c < d$ while in $V(U)$ they are ordered as $a < e < c < d$ but the map $f^*: V(U) \rightarrow V(T)$ is given by $a \mapsto a, c \mapsto c, d \mapsto d, e \mapsto d$.


Note that each element of $V_G(T)$ corresponds to an unique edge orbit Ge for e not a leaf. As such, we will represent the corresponding G -vertex by $v_{Ge} = (Ge)^\dagger \leq Ge$ (which we interpret as the concatenation of the relations $f^\dagger \leq f$ for $f \in Ge$) and write

$$T_{v_{Ge}} = T_{(Ge)^\dagger \leq Ge} = \coprod_{f \in Ge} T_{f^\dagger \leq f}.$$

We note that $T_{v_{Ge}}$ is always a G -corolla. Indeed, noting that a quotient map $\varphi: T \rightarrow S$ induces quotient maps $T_{v_{Ge}} \rightarrow S_{v_{G\varphi(e)}}$ one obtains a functor

$$\begin{aligned} \Omega_G^q &\xrightarrow{V_G} \mathbf{F} \wr \Sigma_G \\ T &\longmapsto (T_{v_{Ge}})_{v_{Ge} \in V_G(T)}. \end{aligned} \tag{3.15} \quad \boxed{\text{VFUNCTOR EQ}}$$

Remark 3.16. The need to introduce the $\mathbf{F} \wr \mathcal{C}$ categories comes from the fact that general quotient maps do not preserve the number of G -vertices. For a simple example, let $G = \mathbb{Z}/4$ and consider the quotient map



sending edges labeled a, b, c to the edges with the same name and the edges a^{+1}, b^{+1} to the edges $a+1, b+1$. We note that T has three G -vertices v_{Gc}, v_{Gb}, v_{Gb+1} while S has only two G -vertices v_{Gc} and v_{Gb} . $V(\phi)$ then maps the two corollas $T_{v_{Gb}}$ and $T_{v_{Gb+1}}$ isomorphically onto $T_{S_{Gb}}$ and the corolla $T_{v_{Gc}}$ non-isomorphically onto $S_{v_{Gc}}$.

Definition 2.39 now immediately generalizes.

Definition 3.17. Let $T \in \Omega_G$ be a G -tree.

A T -substitution datum is a tuple $\{U_{v_{Ge}}\}_{v_{Ge} \in V_G(T)}$ such that $\text{lr}(U_{v_{Ge}}) = T_{v_{Ge}}$.

Further, a map of T -substitution data $\{U_{v_{Ge}}\} \rightarrow \{V_{v_{Ge}}\}$ is a tuple of planar tall maps $\{U_{v_{Ge}} \rightarrow V_{v_{Ge}}\}$.

Remark 3.18. To establish the equivariant analogue of Proposition 2.42 we will prefer to repackage equivariant substitution data in terms of non-equivariant terms.

Noting that there are decompositions $U_{v_{Ge}} = \coprod_{ge \in Ge} U_{ge^\dagger \leq ge}$ and letting $G \ltimes V(T)$ denote the Grothendieck construction for the action of G on the non-equivariant vertices $V(T)$ (often called the action groupoid), it is immediate that an equivariant T -substitution datum is the same as a functor $G \ltimes V(T) \rightarrow \Omega$ whose restriction to $V(T) \subset G \ltimes V(T)$ is a (non-equivariant) substitution datum.

Proposition 3.19. Let $T \in \Omega_G$ be a G -tree. There is an isomorphism of categories

$$\begin{aligned} \text{Sub}(T) &\xrightleftharpoons{\quad} \Omega_{G,T}^{\text{pt}} \\ \{U_{v_{Ge}}\} &\longmapsto (T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}) \end{aligned} \tag{3.20} \quad \boxed{\text{SUBDATAUNDERPLANG EQ}}$$

Proof. This is a minor adaptation of the non-equivariant analogue Proposition 3.19. Since $\text{Sc}(T)$ inherits a G -action, one can form the Grothendieck construction $G \ltimes \text{Sc}(T)$ and by Remark 3.18 equivariant substitution data $\{U_{v_{Ge}}\}$ therefore induce functors $U_{(-)}: G \ltimes \text{Sc}(T) \rightarrow \Omega$. It is then immediate that $\text{colim}_{\text{Sc}(T)} U_{(-)}$ inherits a G -action, provided it exists. The key observation is then that, since $\text{Sc}(T)$ is now a disconnected poset, this colimit is to be interpreted as taken in the category Φ of forests rather than in Ω . \square

Remark 3.21. We will need to know that each of the maps

$$U_{v_{Ge}} \rightarrow U = \text{colim}_{\text{Sc}(T)} U_{(-)}$$

induced by the previous proof is a planar map of G -trees. This requires two observations: (i) the restrictions to each of the constituent non-equivariant trees $U_{ge \uparrow \leq ge}$ is planar by Proposition 3.19; (ii) the restriction to the roots of $U_{v_{Ge}}$ is injective and order preserving since it matches the inclusion of the roots of $T_{v_{Ge}}$, and the map $T \rightarrow U$ is a planar map of G -trees.

3.3 Planar strings

The leaf-root and vertex functors will allow us to reinterpret our results concerning substitution.

Definition 3.22. The category $\Omega_{G,n}$ of *substitution n -strings* is the category whose objects are strings

$$T_0 \xrightarrow{f_1} T_1 \xrightarrow{f_2} \dots \xrightarrow{f_n} T_n$$

where $T_i \in \Omega_G$ and the f_i are tall planar maps, and arrows are commutative diagrams

$$\begin{array}{ccccccc} T_0 & \xrightarrow{f_1} & T_1 & \xrightarrow{f_2} & \dots & \xrightarrow{f_n} & T_n \\ q_0 \downarrow & & q_1 \downarrow & & & & q_n \downarrow \\ T'_0 & \xrightarrow{f'_1} & T'_1 & \xrightarrow{f'_2} & \dots & \xrightarrow{f'_n} & T'_n \end{array} \quad (3.23)$$

PTNARROW EQ

where each q_i is a quotient map.

Notation 3.24. Since compositions of planar tall arrows are planar tall and identity arrows are planar tall it follows that $\Omega_{G,\bullet}$ forms a simplicial object in \mathbf{Cat} , with faces given by composing and degeneracies by inserting identities.

Noting that $\Omega_{G,0} = \Omega_G^q$ and setting $\Omega_{G,-1} = \Sigma_G$, the leaf-root functor $\Omega_G^q \xrightarrow{\text{lr}} \Sigma_G$ makes $\Omega_{G,\bullet}^q$ into an augmented simplicial object and, furthermore, the maps $s_{-1}: \Omega_{G,n}^q \rightarrow \Omega_{G,n+1}^q$ sending $T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_n$ to $\text{lr}(T_0) \rightarrow T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_n$ equip it with extra degeneracies.

Notation 3.25. We extend the vertex functor to a functor $V_G: \Omega_{G,n+1} \rightarrow \mathbf{F} \wr \Omega_{G,n}$ by

$$V_G(T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_n) = (V_G(T_0), (T_{1,v_{Ge}} \rightarrow \dots \rightarrow T_{n,v_{Ge}})_{v_{Ge} \in V_G(T_0)})$$

where we abuse notation by writing $T_{i,v_{Ge}}$ for $T_{i,(f_1 \circ \dots \circ f_i)(v_{Ge})}$.

The following is a reinterpretation of Proposition 3.19.

Proposition 3.26. *The diagram*

$$\begin{array}{ccc} \Omega_{G,n+1} & \xrightarrow{V_G} & \mathbf{F} \wr \Omega_{G,n} \\ d_{1,\dots,n+1} \downarrow & & \downarrow \text{Fid}_{0,\dots,n} \\ \Omega_{G,0} & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G \end{array} \quad (3.27)$$

PTPULL EQ

is a pullback diagram in \mathbf{Cat} .

Proof. An object in the pullback (3.27) over $T_0 \in \Omega_{G,0} = \Omega_G^q$ is precisely the same as a n -string in $\text{Sub}(T)$, and thus by Proposition 3.19 equivalent to a $n+1$ planar tall string starting at T .

The case of arrows is slightly more subtle. A quotient map $\pi: T \rightarrow T'$ induces a G -equivariant poset map $\pi_*: \mathbf{Sc}(T) \rightarrow \mathbf{Sc}(T')$ (or equivalently, a map of Grothendieck constructions $G \ltimes \mathbf{Sc}(T) \rightarrow G \ltimes \mathbf{Sc}(T')$) and diagrams as on the left below (where v_{Ge} ranges over $V_G(T)$ and $e' = \varphi(e)$) induce diagrams (of functors $\mathbf{Sc}(T) \rightarrow \Omega$) as on the right below.

$$\begin{array}{ccc}
T_{v_{Ge}} & \longrightarrow & T_{1,v_{Ge}} \longrightarrow \cdots \longrightarrow T_{n,v_{Ge}} \\
\downarrow & & \downarrow \\
T'_{v_{Ge'}} & \longrightarrow & T'_{1,v_{Ge'}} \longrightarrow \cdots \longrightarrow T'_{n,v_{Ge'}}
\end{array}
\quad
\begin{array}{ccc}
T_{(-)} & \longrightarrow & T_{1,(-)} \longrightarrow \cdots \longrightarrow T_{n,(-)} \\
\downarrow & & \downarrow \\
T'_{(-)} \circ \pi_* & \longrightarrow & T'_{1,(-)} \circ \pi_* \longrightarrow \cdots \longrightarrow T'_{n,(-)} \circ \pi_*
\end{array}
\quad (3.28)$$

Passing to colimits then gives the desired commutative diagram (3.23). Moreover, diagrams of the form (3.23) clearly induce diagrams as in (3.28) and it is straightforward to check that these are inverse processes. \square

DSCOM REM

Remark 3.29. The diagrams (with back and lower slanted faces instances of (3.27))

$$\begin{array}{ccc}
\Omega_{G,n+2} & \xrightarrow{\quad} & F \wr \Omega_{G,n+1} \\
\downarrow & \searrow d_{i+1} & \downarrow \\
\Omega_{G,n+1} & \xrightarrow{\quad} & F \wr \Omega_{G,n} \\
\downarrow & \searrow & \downarrow \\
\Omega_{G,0} & \xrightarrow{\quad} & F \wr \Sigma_G
\end{array}
\quad
\begin{array}{ccc}
\Omega_{G,n+1} & \xrightarrow{\quad} & F \wr \Omega_{G,n} \\
\downarrow & \searrow s_i & \downarrow \\
\Omega_{G,n+2} & \xrightarrow{\quad} & F \wr \Omega_{G,n+1} \\
\downarrow & \searrow & \downarrow \\
\Omega_{G,0} & \xrightarrow{\quad} & F \wr \Sigma_G
\end{array}$$

commute whenever defined (i.e. $0 \leq i \leq n+1$).

HERE

Notation 3.30. We will let

$$V_{G,n}: \Omega_{G,n} \rightarrow F \wr \Sigma_G$$

be inductively defined by $V_{G,n} = \sigma_0 \circ V_{G,n-1} \circ V_G$.

Remark 3.31. When $n = 2$, $V_{G,2}$ is thus the composite

$$\Omega_{G,2} \xrightarrow{V_G} F \wr \Omega_{G,1} \xrightarrow{V_G} F \wr F \wr \Omega_{G,0} \xrightarrow{V_G} F \wr F \wr F \wr \Sigma_G \xrightarrow{\sigma^0} F \wr F \wr \Sigma_G \xrightarrow{\sigma^0} F \wr \Sigma_G$$

In light of Remarks 2.45 and 3.14, $V_{G,n}(T_0 \rightarrow \cdots \rightarrow T_n)$ is identified with the tuple

$$(T_n, v_{Ge})_{v_{Ge} \in V_G(T_n)}, \quad (3.32)$$

VGNISO EQ

though this requires changing the total order in $V_G(T_n)$. Rather than using the order induced by T_n , one instead equips $V_G(T_n)$ with the order induced lexicographically from the maps $V_G(T_n) \rightarrow V_G(T_{n-1}) \rightarrow \cdots \rightarrow V_G(T_0)$, i.e., for $v, w \in V_G(T_n)$ the condition $v < w$ is determined by the lowest i such that the images of $v, w \in V_G(T_i)$ are distinct.

3.4 A monad on spans

Definition 3.33. We will write $\mathbf{WSpan}^l(\mathcal{C}, \mathcal{D})$ (resp. $\mathbf{WSpan}^r(\mathcal{C}, \mathcal{D})$), which we call the category of *left weak spans* (resp. *right weak spans*), to denote the category with objects the spans

$$\mathcal{C} \xleftarrow{k} A \xrightarrow{F} \mathcal{D},$$

arrows the diagrams as on the left (resp. right) below

$$\begin{array}{ccc}
& A_1 & \\
k_1 \swarrow & & \searrow F_1 \\
\mathcal{C} & & \mathcal{D} \\
k_2 \swarrow & & \searrow F_2 \\
& A_2 &
\end{array}
\quad
\begin{array}{ccc}
& A_1 & \\
k_1 \swarrow & & \searrow F_1 \\
\mathcal{C} & & \mathcal{D} \\
k_2 \swarrow & & \searrow F_2 \\
& A_2 &
\end{array}
\quad (3.34)$$

TWISTEDARROWRIGHT EQ

which we write as $(i, \varphi): (k_1, F_1) \rightarrow (k_2, F_2)$, and composition given in the obvious way.

Remark 3.35. There are natural isomorphisms

$$\mathbf{WSpan}^r(\mathcal{C}, \mathcal{D}) \simeq \mathbf{WSpan}^l(\mathcal{C}^{op}, \mathcal{D}^{op}). \quad (3.36)$$

LRSPANISO EQ

Remark 3.37. The terms *left/right* are motivated by the existence of adjunctions (which are seen to be equivalent by using (3.36))

$$\mathbf{Lan}: \mathbf{WSpan}^l(\mathcal{C}, \mathcal{D}) \rightleftarrows \mathbf{Fun}(\mathcal{C}, \mathcal{D}): \iota$$

$$\iota: \mathbf{Fun}(\mathcal{C}, \mathcal{D}) \rightleftarrows \mathbf{WSpan}^r(\mathcal{C}, \mathcal{D})^{op}: \mathbf{Ran}$$

where the functors ι denote the obvious inclusions (note the need for the $(-)^{op}$ in the second adjunction) and $\mathbf{Lan}/\mathbf{Ran}$ denote the left/right Kan extension functors.

We will mainly be interested in the span categories $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V}) \simeq \mathbf{WSpan}^r(\Sigma_G, \mathcal{V}^{op})$.

Notation 3.38. Given a functor $\pi: A \rightarrow \Sigma_G$, we let $\Omega_{G,n}^{(A)}$ denote the pullback (in \mathbf{Cat})

$$\begin{array}{ccc} \Omega_{G,n}^{(A)} & \xrightarrow{V_{G,n}^{(A)}} & \mathbf{F} \wr A \\ \downarrow & & \downarrow \\ \Omega_{G,n} & \xrightarrow{V_{G,n}} & \mathbf{F} \wr \Sigma_G \end{array}$$

Explicitly, the objects of $\Omega_{G,n}^{(A)}$ are pairs

$$(T_0 \rightarrow \cdots \rightarrow T_n, (a_{e^\dagger \leq e})_{(e^\dagger \leq e) \in V_G(T_n)})$$

such that $\pi(a_{e^\dagger \leq e}) = T_{n, e^\dagger \leq e}$.

Remark 3.39. Our primary interest here will be in the $\Omega_{G,0}^{(A)}$ construction. Importantly, the composite maps $\Omega_{G,0}^{(A)} \rightarrow \Omega_{G,0} \rightarrow \Sigma_G$ allow us to iterate the $\Omega_{G,0}^{(-)}$ construction. In practice, the role of higher strings $\Omega_{G,n}^{(A)}$ will then be to provide more convenient models for iterated $\Omega_{G,0}^{(-)}$ constructions.

SUBSASPULL PROP

Indeed, the content of Proposition 3.26 is then that there are compatible identifications $\Omega_{G,0}^{(\Omega_{G,n})} \simeq \Omega_{G,n+1}$ which identify $V_G^{(\Omega_{G,n})}$ with V_G .

Moreover, since all squares in the diagram

$$\begin{array}{ccccccc} \Omega_{G,n+1}^{(A)} & \xrightarrow{V_G^{(A)}} & \mathbf{F} \wr \Omega_{G,n}^{(A)} & \xrightarrow{\mathbf{F} \wr V_{G,n}^{(A)}} & \mathbf{F} \wr \mathbf{F} \wr A & \xrightarrow{\sigma^0} & \mathbf{F} \wr A \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_{G,n+1} & \xrightarrow{V_G} & \mathbf{F} \wr \Omega_{G,n} & \xrightarrow{\mathbf{F} \wr V_{G,n}} & \mathbf{F} \wr \mathbf{F} \wr \Sigma_G & \xrightarrow{\sigma^0} & \mathbf{F} \wr \Sigma_G \\ \downarrow & & \downarrow & & & & \\ \Omega_{G,0} & \longrightarrow & \mathbf{F} \wr \Sigma_G & & & & \end{array}$$

are pullback squares (the top center square is so by induction, the top right square by direct verification, the total top square by definition of $\Omega_{G,n+1}^{(A)}$ and the bottom left square by

Proposition 3.26), we likewise obtain identifications $\Omega_G^{(\Omega_{G,n}^{(A)})} \simeq \Omega_{G,n+1}^{(A)}$.

HERE

Proposition 3.40. For any $A \rightarrow \Sigma_G$ there are functors $d_0^{(A)}: \Omega_{G,1}^{(A)} \rightarrow \Omega_G^{(A)}$ and natural isomorphisms

$$\begin{array}{ccccc}
 \Omega_{G,1}^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A \\
 d_0^{(A)} \downarrow & & \nearrow \pi^{(A)} & & \downarrow \sigma^0 \\
 \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr A, & &
 \end{array} \quad (3.41) \quad \boxed{\text{SHUFFLEPERMA EQ}}$$

both natural in $A \rightarrow \Sigma$. Here naturality of $\pi^{(-)}$ means that for a functor $H: A \rightarrow B$ with corresponding diagram

$$\begin{array}{ccccccc}
 \Omega_{G,1}^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A & \xrightarrow{\sigma^0} & F \wr A \\
 \Omega_{G,1}^{(B)} & \xrightarrow{V_G} & F \wr \Omega_G^{(B)} & \xrightarrow{F \wr V_G} & F \wr F \wr B & \xrightarrow{\sigma^0} & F \wr B \\
 \downarrow d_0^{(A)} & & \downarrow d_0^{(B)} & & \downarrow d_0^{(A)} & & \downarrow d_0^{(B)} \\
 \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A & \xrightarrow{\sigma^0} & F \wr A \\
 \downarrow d_0^{(A)} & & \downarrow d_0^{(B)} & & \downarrow d_0^{(A)} & & \downarrow d_0^{(B)} \\
 \Omega_G^{(B)} & \xrightarrow{V_G} & F \wr \Omega_G^{(B)} & \xrightarrow{F \wr V_G} & F \wr F \wr B & \xrightarrow{\sigma^0} & F \wr B
 \end{array} \quad (3.42) \quad \boxed{\text{PICUBOIDAB EQ}}$$

one has an equality

$$(F \wr H)\pi^{(A)} = \pi^{(B)}\Omega_{G,1}^{(H)}$$

(i.e. the two natural isomorphisms between the two distinct functors $\Omega_{G,1}^{(A)} \Rightarrow F \wr B$ coincide).

Proof. Since d_0 has already been defined when $B = \Sigma_G$, one can generally define $d_0^{(A)}$ by using diagram (3.42) when $B = \Sigma$ and recalling that the front, back and left faces are all known to be pullbacks.

Fill in the definition of $\pi^{(A)}$. □

Definition 3.43. Suppose \mathcal{V} has finite products.

We define an endofunctor N of $\mathbf{Wspan}^r(\Sigma_G, \mathcal{V}^{op})$ by letting $N(\Sigma_G \leftarrow A \rightarrow \mathcal{V}^{op})$ be the span $\Sigma_G \leftarrow \Omega_G^{(A)} \rightarrow \mathcal{V}^{op}$ given by the diagram

$$\begin{array}{ccccc}
 \Omega_G^{(A)} & \longrightarrow & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \xrightarrow{\Pi^{op}} \mathcal{V}^{op} \\
 \downarrow & & \downarrow & & \\
 \Omega_G & \longrightarrow & F \wr \Sigma_G & & \\
 \downarrow & & & & \\
 \Sigma_G & & & &
 \end{array}$$

and defined on arrows in the obvious way.

One has a multiplication $\mu: N \circ N \Rightarrow N$ given by the natural isomorphisms

$$\begin{array}{ccccccc}
 \Sigma \leftarrow \Omega_{G,1}^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A & \longrightarrow & F \wr F \wr \mathcal{V}^{op} \longrightarrow F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 \parallel & & \downarrow d_0^{(A)} & \nearrow \pi^{(A)} & \downarrow \sigma^0 & & \downarrow \sigma^0 \\
 \Sigma \leftarrow \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A & \longrightarrow & F \wr F \wr \mathcal{V}^{op} \longrightarrow F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 & & \downarrow d_0^{(A)} & \nearrow \pi^{(A)} & \downarrow \sigma^0 & & \downarrow \sigma^0 \\
 \Sigma \leftarrow \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A & \longrightarrow & F \wr F \wr \mathcal{V}^{op} \longrightarrow F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op}
 \end{array} \quad (3.44) \quad \boxed{\text{MULTDEFSPAN EQ}}$$

where we note that naturality follows from the commutativity of (3.42).

Lastly, there is a unit $\eta: id \Rightarrow N$ given by the strictly commutative diagrams

$$\begin{array}{ccccccc}
 \Sigma & \longleftarrow & A & \xlongequal{\quad} & A & \longrightarrow & \mathcal{V}^{op} \xlongequal{\quad} \mathcal{V}^{op} \\
 \parallel & & \downarrow s_{-1}^{(A)} & & \downarrow & & \downarrow & \parallel \\
 \Sigma & \longleftarrow & \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op}.
 \end{array} \tag{3.45}$$

UNITSPAN EQ

MONSPAN PROP

Proposition 3.46. (N, μ, η) form a monad on $\mathcal{Wspan}^r(\Sigma_G, \mathcal{V}^{op})$.

Proof. The natural transformation component of $\mu \circ (N\mu)$ is given by the composite diagram

$$\begin{array}{ccccccccccc}
 \Omega_{G,2}^{(A)} & \rightarrow & F \wr \Omega_{G,1}^{(A)} & \rightarrow & F^{i2} \wr \Omega_G^{(A)} & \rightarrow & F^{i3} \wr A & \rightarrow & F^{i3} \wr \mathcal{V}^{op} & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_1^{(A)} \downarrow & & \downarrow & & \swarrow F \wr \pi^{(A)} & & \downarrow \sigma^1 & & \downarrow \sigma^1 & & \swarrow F \wr \alpha & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \rightarrow & F \wr \Omega_G^{(A)} & \xrightarrow{\quad} & F^{i2} \wr A & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \xrightarrow{\quad} & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} & & \parallel & & \parallel \\
 d_0^{(A)} \downarrow & & \swarrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \swarrow \alpha & & \parallel & & \parallel & & \parallel \\
 \Omega_G^{(A)} & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op}
 \end{array} \tag{3.47}$$

ASSOCSPAN1 EQ

whereas the natural transformation component of $\mu \circ (\mu N)$ is given by

$$\begin{array}{ccccccccccc}
 \Omega_{G,2}^{(A)} & \rightarrow & F \wr \Omega_{G,1}^{(A)} & \rightarrow & F^{i2} \wr \Omega_G^{(A)} & \rightarrow & F^{i3} \wr A & \rightarrow & F^{i3} \wr \mathcal{V}^{op} & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_1^{(A)} \downarrow & & \swarrow \pi(\Omega_G^{(A)}) & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \swarrow \alpha & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \xrightarrow{\quad} & F \wr \Omega_G^{(A)} & \xrightarrow{\quad} & F^{i2} \wr A & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} & & \parallel & & \parallel \\
 d_0^{(A)} \downarrow & & \swarrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \swarrow \alpha & & \parallel & & \parallel & & \parallel \\
 \Omega_G^{(A)} & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op}
 \end{array} \tag{3.48}$$

ASSOCSPAN2 EQ

Add argument that these commute

As for $\mu \circ (N\eta)$, it is represented by

$$\begin{array}{ccccccc}
 \Omega_G^{(A)} & \xrightarrow{\quad} & F \wr A & \xlongequal{\quad} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \xlongequal{\quad} F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 s_0^{(A)} \downarrow & & \downarrow & & \downarrow \delta^1 & & \downarrow \delta^1 & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \rightarrow & F \wr \Omega_G^{(A)} & \rightarrow & F^{i2} \wr A & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & & \swarrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \swarrow \alpha & & \parallel \\
 \Omega_G^{(A)} & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op}
 \end{array} \tag{3.49}$$

UNITSPAN1 EQ

while $\mu \circ (\eta N)$ is represented by

$$\begin{array}{ccccccc}
 \Omega_G^{(A)} & \xlongequal{\quad} & \Omega_G^{(A)} & \longrightarrow & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \xlongequal{\quad} \mathcal{V}^{op} \\
 s_{-1}^{(A)} \downarrow & & \downarrow & & \downarrow \delta^0 & & \downarrow \delta^0 & & \downarrow & & \parallel \\
 \Omega_{G,1}^{(A)} & \rightarrow & F \wr \Omega_G^{(A)} & \rightarrow & F \wr F \wr A & \rightarrow & F \wr F \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & & \swarrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \swarrow \alpha & & \parallel \\
 \Omega_G^{(A)} & \xrightarrow{\quad} & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \xrightarrow{\quad} & \mathcal{V}^{op} & & \mathcal{V}^{op} & & \mathcal{V}^{op}
 \end{array} \tag{3.50}$$

UNITSPAN2 EQ

Add argument that these commute

□

4 Filtration of Cellular Extensions

We now focus on a particular construction in the category $\mathcal{V}\mathbf{Op}$ of \mathcal{V} -enriched operads. In particular, given any $\mathcal{P} \in \mathcal{V}\mathbf{Op}$, and $X, Y \in \mathcal{V}\mathbf{Sym}$, consider the following pushout.

$$\begin{array}{ccc} \mathbb{F}X & \xrightarrow{h} & \mathcal{P} \\ u \downarrow & & \downarrow \\ \mathbb{F}Y & \longrightarrow & \mathcal{P}[u] \end{array}$$

Definition 4.1. We call both the map $\mathcal{P} \rightarrow \mathcal{P}[u]$ and the operad $\mathcal{P}[u]$ itself a *cellular extension* of \mathcal{P} .

Cellular extensions in various forms have been extensively studied, as they are of paramount importance to lifting model structures across free-forgetful adjunctions (as we will exploit in the next section). As has been done previously (see, for example, [SS00, Spitz201, BM03, White14a, White14b, WY15, Per16, ?]), we will construct a filtration of the map $\mathcal{P} \rightarrow \mathcal{P}[u]$ in the underlying category of sequences.

As we showed above in Example , we have that $\mathcal{P}[u] \simeq \mathrm{Lan}_{\Omega_{G,e}^{op}} N^e$. Thus, we are able to decompose $\mathcal{P}[u]$ by a filtration of $\Omega_{G,e}$.

doesn't
exist yet

4.1 Stuff Which Probably Will Be Included in an Above Section

We recall a generator-relation description of the category Ω_G^e . Objects are $(X, Y; \mathcal{P})$ -alternating G -trees: odd or *active* nodes are \mathcal{P} -labeled, and even or *passive* nodes are either X - or Y -labeled, such that G acts by labeled-preserving isomorphisms of the underlying G -tree. Maps come in three generating flavors:


1. quotient maps which preserve the labelings of the nodes;
2. “passive relabeling” maps $\partial_Y = \partial_Y^{(v_i)}$, which are the identity on the underlying tree, but change the labelings on the vertices v_1, \dots, v_n from Y to X ;
3. “active substitution” maps $\partial_{\mathcal{P}} = \partial_{\mathcal{P}}^{(v_i, S^{v_i})}$, which are underlying inner face maps, that take an active (\mathcal{P} -labeled) nodes T_{v_i} and substitute them with $(X; \mathcal{P})$ -alternating G -trees S^{v_i} of height 3 such that we have a (unique) planar-tall maps $T_{v_i} \rightarrow S^{v_i}$ on the underlying G -trees.

Relations are generated by the following:

1. any two passive relabeling maps commute;
2. compositions of active substitutions are identified if they induce the same underlying map in Ω_G after forgetting labels; and
3. if ∂_Y and $\partial_{\mathcal{P}}$ affect different nodes of T , then the composites $\partial_Y \partial_{\mathcal{P}}$ and $\partial_{\mathcal{P}} \partial_Y$ exist and are equal.

Example 4.2. The following is a composite of passive relabelings followed by some active

substitutions:



OMEGA_E_REMARKS

Remark 4.3. Some remarks:

1. Maps in Ω_G^e have unique decompositions of the form $f = q\partial_Y\partial_P$; in particular, for any composite $f = \partial q$, we have $f = q'\partial'$, where ∂ and ∂' are composites of passive relabeling and active substitutions, and q and q' are quotients.
2. Equivalently (and as in Ω_G), for any cartesian map $q : S \rightarrow T$ and any non-cartesian map $f : T \rightarrow T'$, there must exist a factorization

$$\begin{array}{ccc} S & \xrightarrow{\quad f' \quad} & S' \\ q \downarrow & & \downarrow q' \\ T & \xrightarrow{\quad f \quad} & T' \end{array}$$

where q' is also Cartesian, and f' is non-Cartesian.

3. $\Omega_{G,-1}^{(2,1)}$ naturally embeds into Ω_G^e as a wide subcategory.

Definition 4.4. The nerve-evaluation map $N_{(X,Y),P} : \Omega_{G,-1}^{(2,1)} \rightarrow \mathcal{V}$ extends to Ω_G^e . It sends passive relabeling maps ∂_Y^v to an application of the map $u(T_v) : X(T_v) \rightarrow Y(T_v)$ in the v -th component of the indexed tensor product; on active substitution maps ∂_P^{v,S^v} , it is given by first an application of $h(T_v) : X(T_v) \rightarrow \mathcal{P}(T_v)$, followed by the operad structure map 'collapsing' newly adjacent active nodes.

Definition 4.5. Given $T \in \Omega_G^e$, let $V_P(T)$, $V_X(T)$, and $V_Y(T)$ be the G -sets of vertices which are \mathcal{P} -, X -, and Y -labeled, respectively. Similarly, let $V_{G,P}(T) = V_P(T)/G$, $V_{G,X}(T) = V_X(T)/G$, and $V_{G,Y}(T) = V_Y(T)/G$ be the sets of G -vertices which are correctly labeled.

Further, let $|T| = |T|_X + |T|_Y$, where $|T|_X$ is the "number of X -labeled nodes in any single component of T ", and similarly for $|T|_Y$. Rigorously, we have

$$|T|_X = \frac{|V_X(T)|}{|G.r|} = \sum_{G.v \in V_{G,X}(T)} \frac{|G.v|}{|G.r|}$$


where $G.r$ is the root G -set of T ; and similarly for $|T|_Y$.

Definition 4.6. Given a $(\mathcal{P}; X, Y)$ -alternating G -tree T , let T_P^\wedge be the alternating tree with $|T|_X = 0$ (so T_P^\wedge is $(\mathcal{P}; Y)$ -alternating) created from T by

1. first relabeling all X -nodes \mathcal{P} (yielding a tree in $\lambda^3\mathbb{T}_0$); then
2. collapsing all connected components of \mathcal{P} -labeled nodes.

There is a unique planar-tall map $\partial_{\mathcal{P}} : T_{\mathcal{P}}^{\wedge} \rightarrow T$, and in fact this factors through all maps $\partial_{\mathcal{P}} : S \rightarrow T$.

Example 4.7. We demonstrate this functor on an example element in $\lambda_1^2 \mathbb{T}_G$ below. The source object T is on the left; the (non-alternating) tree $T_{\mathcal{P}}^{\wedge}$ which has relabeled all X -nodes as \mathcal{P} -nodes is in the middle; and the right tree has collapsed all the connected active components. As before, black nodes are active, while white are passive, and we have labeled the interior of the node to denote which kind of passive it is.



4.2 Filtration Lemmas

Definition 4.8. We define subcategories of Ω_G^e .

1. Let $\Omega_G^e[\leq k]$ (respectively $\Omega_G^e[k]$) be the full subcategory of Ω_G^e spanned by trees T with $|T| \leq k$ (respectively, $|T| = k$).
2. Let $\Omega_G^e[\leq k, -]$ (respectively $\Omega_G^e[k, -]$) be the full subcategory of $\Omega_G^e[\leq k]$ (respectively $\Omega_G^e[k]$) spanned by trees T with $|T|_Y \neq k$.
3. Let $\Omega_G^e[\leq k, 0]$ (respectively $\Omega_G^e[k, 0]$) be the full subcategory of $\Omega_G^e[\leq k]$ (respectively $\Omega_G^e[k]$) spanned by trees T with $|T|_X = 0$ (equivalently, $|T|_Y = k$).
4. If Ξ is any of the above categories, and $C \in \Sigma_G$, let $\Xi(C)$ denote the full subcategory of Ξ spanned by those trees T with $\text{val}(T) \simeq C$.

Remark 4.9. The categories $\Omega_G^e[k]$ and $\Omega_G^e[k, -]$ have only very limited morphisms, as there cannot be any “active substitutions”. Thus, any map $S \rightarrow T$ in these categories just changes some Y -labelings into X -labelings, while the underlying $(1, 1)$ -alternating tree remains fixed (where here the one passive colour encompasses both Y - and X -labels).

Lemma 4.10. $\Omega_G^e[\leq k-1]^{op}$ is Lan-final in $\Omega_G^e[\leq k, -]^{op}$ over Σ_G^{op} .

Proof. Fix an arbitrary $C \in \Sigma_G$, and consider an element $q_S : \text{val}(S) \leftarrow C$ in $\Omega_G^e[\leq k, -]^{op} \downarrow C$ (so in particular $S \in \Omega_G^e[\leq k, -]$). We must show that the overcategory

$$(\Omega_G^e[\leq k-1]^{op} \downarrow C) \downarrow (\text{val}(S) \leftarrow C)$$

is non-empty and connected. If in fact $S \in \Omega_G^e[\leq k-1]$, the result is immediate. Otherwise, consider the map

$$S_{\mathcal{P}}^{\wedge} \xrightarrow{\partial_{\mathcal{P}}} S.$$

Since $|S|_Y \neq k$, $|S_{\mathcal{P}}^{\wedge}| \leq k-1$, and hence we have a diagram

$$\begin{array}{ccc} \text{val}(S) & \xleftarrow{\partial_{\mathcal{P}}} & \text{val}(S_{\mathcal{P}}^{\wedge}) \\ & \nwarrow q_S & \nearrow q_S \\ & C & \end{array}$$

(4.11) K-1_LAN_FINAL_EQ1

showing that the desired overcategory is inhabited. Further, given any other element

$$\begin{array}{ccc} \text{val}(S) & \xleftarrow{f} & \text{val}(T) \\ & \nwarrow q_T \quad \nearrow q_S & \\ & C & \end{array} \quad (4.12) \quad \boxed{\text{K-1_LAN_FINAL_EQ2}}$$

in the overcategory, consider the following zig-zag of maps connecting the objects $(\text{K-1_LAN_FINAL_EQ1})$ and (H.12) :

$$\begin{array}{ccccc} & & S & & \\ & \nearrow \partial_{\mathcal{P}} & \uparrow \tilde{q}_T & \nwarrow f & \\ & & \tilde{q}_T^*(S) & & \\ & \nwarrow \partial_{\mathcal{P}} & \nwarrow \partial_{\mathcal{P}} & \nearrow \partial_{\mathcal{P}} & \\ S_{\mathcal{P}}^{\wedge} & \xleftarrow{\tilde{q}_T} & \tilde{q}_T^*(S_{\mathcal{P}}^{\wedge}) & \xrightarrow{\partial_{\mathcal{P}}} & T' & \xleftarrow{\partial_Y} & T \\ & \nwarrow q_S & \nwarrow q_T & \nearrow q_T & \nearrow q_T & \\ & & C & & \end{array} \quad (4.13) \quad \boxed{\text{K-1_LAN_FINAL_DIAGRAM}}$$

Here, I have omitted the notation “*val*” from the top three rows. To understand this diagram, we first record that we have a factorization:

$$q_S = \tilde{q}_T q_T,$$

Then, if we let $C_S = \text{val}(S) = \text{val}(S_{\mathcal{P}}^{\wedge})$ and $C_T = \text{val}(T)$, we have

$$C \xrightarrow{q_T} C_T \xrightarrow{\tilde{q}_T} C_S$$

and hence, via Remark $\boxed{\text{OMEGA_E_REMARKS}}$ 4.3 (2), a factorization

$$\begin{array}{ccc} C_T & \cdots \rightarrow & \tilde{q}_T^*(S_{\mathcal{P}}^{\wedge}) \\ \tilde{q}_T \downarrow & & \downarrow \tilde{q}_T \\ C_S & \longrightarrow & S_{\mathcal{P}}^{\wedge} \end{array}$$

(where we are recording $C \rightarrow \text{val}(S)$ as a planar-tall map $C \rightarrow S$). A similar analysis shows that the top left trapezoid commutes.

The other regions also commute by a straightforward analysis. Indeed, the top right trapezoid commutes by unique factorization, and finally the middle triangle of $\partial_{\mathcal{P}}$ maps commutes since $(\tilde{q}_T^* S)_{\mathcal{P}}^{\wedge} = \tilde{q}_T^*(S_{\mathcal{P}}^{\wedge})$.

Lastly, we must check that the middle two maps are in fact elements of the appropriate overcategory. This follows from the fact that $S_{\mathcal{P}}^{\wedge}$ and T have $|-|_Y < k$. Thus, the overcategory in question is connected, as desired. \square

come back: define S_Y^{\wedge} .

$\boxed{\text{N_FINALITY_LEMMA}}$

Lemma 4.14. $\Omega_G^e[k, 0]^{op}$ is Lan-final in $\Omega_G^e[k]$ over Σ_G^{op} .

Definition 4.17 ^{BM03} Given a map $u : Y_0 \rightarrow Y_1$ of sequences and $(A, D) \in \mathbf{F}_0 \wr \Sigma$, we borrow notation from [7] and define the functor $[u]^D : (0 \rightarrow 1)^A \rightarrow \mathcal{V}$ as the composite

$$(0 \rightarrow 1)^A \rightarrow \mathbf{F}_0 \wr V \xrightarrow{\otimes} \mathcal{V}$$

where the first map is defined on $\epsilon : A \rightarrow \{0, 1\}$ by

$$(\epsilon(a))_a \mapsto (A, (Y_{\epsilon(a)}(D(a)))_a).$$

We recall that, in a general category \mathcal{C} , a subcategory $\mathcal{C}' \subseteq \mathcal{C}$ is called *convex* if whenever $c' \in \mathcal{C}'$ and $c \mapsto c'$ is an arrow in \mathcal{C} , then both c' and the map are in \mathcal{C}' .

Q_DEFINITION

Definition 4.18. Let \mathcal{C} be a convex subcategory of $(0 \rightarrow 1)^A$. We define $Q_{\mathcal{C}}^A[u]^D := \text{colim}_{\mathcal{C}}[u]^D$; moreover, given nested convex subcategories $\mathcal{C}' \subseteq \mathcal{C}$, let

$$[u]^D \square_{\mathcal{C}'}^{\mathcal{C}} : Q_{\mathcal{C}'}^A[u]^D \rightarrow Q_{\mathcal{C}}^A[u]^D$$

denote the unique natural map.

In particular, if \mathcal{C} is the full “punctured cube” subcategory $(0 \rightarrow 1)^A \setminus \{(1)_a\}$, we simplify the notation as follows:

$$\begin{aligned} Q[u]^D &:= Q_{\mathcal{C}}[u]^D \\ [u]^{\square D} &:= [u]^D \square_{\mathcal{C}}^{(0 \rightarrow 1)^A} : Q[u]^D \rightarrow \bigotimes_{a \in A} Y_1(D(a)). \end{aligned}$$

5 Appendix


5.1 Grothendieck fibrations

This is missing a bunch of expository content

Definition 5.1. Suppose $(\mathcal{D}, \mathcal{D}_c)$ is a pair in \mathbf{Cat} such that \mathcal{D}_c is a wide subcategory of \mathcal{D} .

A functor $\pi: \mathcal{B} \rightarrow \mathcal{D}$ is called a partial Grothendieck construction if there are cartesian lifts for every arrow in \mathcal{D}_c .

Lemma 5.2. *bla bla*



5.2 Kan Extensions - May Also be Part of the Above

We collect some technical results about the naturality of Kan extensions on their input data, and their preservation under certain categorical constructions.

Remark 5.3. For all of the results below, their formal dual result is true of right Kan extensions.

We begin with an easy result about “stacking” Kan extensions.

Lemma 5.4. *Suppose we have functors*

$$\begin{array}{ccc} \mathcal{C}' & & \\ \downarrow i & \searrow X & \\ \mathcal{C} & \xrightarrow{Y} & \mathcal{V} \\ \downarrow j & & \\ \mathcal{D} & & \end{array}$$

such that $Y = \text{Lan}_i X$. Then $\text{Lan}_{ji} X \simeq \text{Lan}_j Y$.

Proof. This follows from the Yoneda Lemma by directly unpacking the universal properties of the two functors. \square

5.2.1 Naturality

We formulate precisely what data Kan extensions are natural over.

Definition 5.5. Let $\mathbf{WSpan}(\mathcal{D}, \mathcal{V})$ be the following category. Objects are spans of functors

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \mathcal{V} \\ \downarrow & & \\ \mathcal{D} & & \end{array}$$

while morphisms are pairs (F, Φ)

$$\begin{array}{ccc} & \mathcal{C}' & \\ \downarrow F & \searrow \Phi & \downarrow \\ \mathcal{C} & \xrightarrow{\quad} & \mathcal{V} \\ \downarrow & & \\ \Sigma & & \end{array}$$

with F a functor such that the left triangle commutes, and Φ a natural transformation.

NATURALITY_PROP

Proposition 5.6. *The left Kan extension operation is a functor $\mathbf{WSpan}(\mathcal{D}, \mathcal{V}) \rightarrow \mathcal{V}^{\mathcal{D}}$.*

Proof. This is a straightforward diagram chase via the universal property of the left Kan extension. Indeed, suppose we are given the following data:

$$\begin{array}{ccc} \mathcal{C} & & \\ \downarrow i & \searrow X & \\ \mathcal{C}' & \xrightarrow{\quad} & \mathcal{V} \\ \downarrow j & \searrow Y & \\ \mathcal{D} & & \end{array}$$

Then, since we always have a natural transformation $\text{Lan}_i Y \circ i \Rightarrow Y$, we have the following chain of bijections and maps:

$$\begin{aligned} \mathcal{V}^{\mathcal{D}}(\text{Lan}_j Y, \text{Lan}_j Y) &= \mathcal{V}^{\mathcal{C}'}(Y, \text{Lan}_j Y \circ j) \xrightarrow{(\tilde{id})^*} \mathcal{V}^{\mathcal{C}'}(\text{Lan}_i(Yi), \text{Lan}_j Y \circ j) = \mathcal{V}^{\mathcal{C}}(Yi, \text{Lan}_j Y \circ ji) \\ &\xrightarrow{\Phi^*} \mathcal{V}^{\mathcal{C}}(X, \text{Lan}_j Y \circ ji) = \mathcal{V}^{\mathcal{D}}(\text{Lan}_{ji} X, \text{Lan}_j Y). \end{aligned}$$

The image of the identity, denoted Φ_* , is the desired natural transformation. It can similarly be shown that this process preserves compositions. \square

Diagrammatically, $\Phi_* : \text{Lan}_{ji} X \rightarrow \text{Lan}_j Y$ is the unique map such that the diagram below commutes:

$$\begin{array}{ccccc} X & \xrightarrow{\Phi} & Y \circ i & \xrightarrow{\alpha_Y \circ i} & \text{Lan}_j Y \circ ji \\ & \searrow \alpha_X & & \nearrow \Phi_* \circ ji & \\ & & \text{Lan}_{ji} X \circ ji & & \end{array}$$

(5.7)

LAN_MAPS_COROLLARY_DI.

We highlight a special case:

RECOMP_LAN_LEMMA

Corollary 5.8. *Suppose we have functors $\mathcal{C} \xrightarrow{i} \mathcal{C}' \xrightarrow{j} \mathcal{D}$ and $X : \mathcal{C}' \rightarrow \mathcal{V}$. Then we have a natural transformation $\Phi : \text{Lan}_{ji} Xi \rightarrow \text{Lan}_j X$.*

$$\begin{array}{ccc} \mathcal{C} & & \\ \downarrow i & \searrow Xi & \\ \mathcal{C}' & \xrightarrow{\quad} & \mathcal{V} \\ \downarrow j & \searrow X & \\ \mathcal{D} & & \end{array}$$

\square

We can also consider “higher-dimensional” naturality, such as the following.

Lemma 5.9. *Suppose we have the following commutative diagram of natural transformations*

$$\begin{array}{ccccc}
 & & \mathcal{C}' & \xrightarrow{\quad} & \mathcal{V} \\
 & \nearrow & \downarrow & \nearrow & \parallel \\
 \mathcal{E}' & \xrightarrow{\quad} & \mathcal{V} & & \parallel \\
 \downarrow & & \downarrow & & \parallel \\
 & \nearrow & \mathcal{C} & \xrightarrow{\quad} & \mathcal{V} \\
 \mathcal{E} & \xrightarrow{\quad} & \mathcal{V} & & \parallel \\
 \downarrow & & \downarrow & & \parallel \\
 & & \mathcal{D} & &
 \end{array}$$

such that the left and right faces commute, the front and back faces are some natural transformations Φ and Ψ , and the top and bottom faces are left Kan extensions. Then the maps induced by Lemma 5.6 from Φ and Ψ are isomorphic.

Proof. We know the sources and targets are isomorphic by Lemma 5.4. The result then follows as in the proof of Lemma 5.6. \square

5.2.2 Left Kan Extensions and Pushouts

While dealing with general pushouts of categories requires solving a “word problem” on morphisms, there is a stronger notion which is much easier to understand. We recall that, given a square of categories

$$\begin{array}{ccc}
 \mathcal{A} & \longrightarrow & \mathcal{C} \\
 \downarrow & & \downarrow \\
 \mathcal{B} & \longrightarrow & \mathcal{D}
 \end{array}$$

if the nerve of this square is a pushout in \mathbf{sSet} , then this is a pushout of categories (since the nerve is the inclusion of a reflective subcategory). The pushouts that most concern us in this paper will be of this form.

Definition 5.10. We call such squares *nervous pushouts* of categories.

If we further assume that the span of functors is built out of fully-faithful inclusions, these pushouts behave as nicely as possible with left Kan extensions.

Lemma 5.11. *Given any diagram in categories of the form*

$$\begin{array}{ccccc}
 \mathcal{A} & \xrightarrow{f} & \mathcal{C} & & \\
 \downarrow & & \downarrow i & & \\
 \mathcal{B} & \xrightarrow{g} & \mathcal{D} & \xrightarrow{Y} & \mathcal{V} \\
 & & \downarrow j & & \\
 & & \mathcal{D} & &
 \end{array}$$

such that the square is a nervous pushout of fully-faithful functors, then $\text{Lan}_j Y$ is the pushout of the induced span

$$\begin{array}{ccc}
 \text{Lan}_{jif}(Yif) & \longrightarrow & \text{Lan}_{ji}(Yi) \\
 \downarrow & & \\
 \text{Lan}_{jg}(Yg) & &
 \end{array}$$

Proof. By the universal property of left Kan extensions, it suffices to show that, for any functor $Z : \mathcal{V} \rightarrow \mathcal{D}$, the natural map

$$\mathcal{V}^{\mathcal{D}}(Y, Zj) \longrightarrow \mathcal{V}^{\mathcal{B}}(Yg, Zjg) \prod_{\mathcal{V}^{\mathcal{A}}(Yif, Zjif)} \mathcal{V}^{\mathcal{C}}(Yi, Zji)$$

is a bijection. These two sets give the same data: a collection of maps $\Phi_b : Y(b) \rightarrow Z(b)$ and $\Phi_c : Y(c) \rightarrow Z(c)$ for all $b \in \mathcal{B}$ and $c \in \mathcal{C}$, such that $\Phi_b = \Phi_c$ whenever $b = c \in \mathcal{A}$. In general, the compatibilites required on the right are less demanding. However, with the above assumptions, a map $d \rightarrow d'$ in \mathcal{D} is *uniquely* a map in \mathcal{A} , $\mathcal{B} \setminus \mathcal{A}$, or $\mathcal{C} \setminus \mathcal{A}$, and thus all the necessary compatibilities are covered by (at least) one of the $\{\Phi_b\}$ or $\{\Phi_c\}$. \square

References

McL

- [1] S. Mac Lane. Categories for the working mathematician, volume 5 of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1998.

Pe16b

- [2] L. A. Pereira. Equivariant dendroidal sets. Available at: <http://www.faculty.virginia.edu/luisalex/>, 2016.

We12

- [3] I. Weiss. Broad posets, trees, and the dendroidal category. Available at: <https://arxiv.org/abs/1201.3987>, 2012.