

Genuine equivariant operads

Peter Bonventre, Luís A. Pereira

April 21, 2017

Abstract

We build new algebraic structures, which we call genuine equivariant operads, which can be thought of as a hybrid between equivariant operads and coefficient systems. We then prove an Elmendorf type theorem stating that equivariant operads, with their graph model structure, are equivalent to genuine equivariant operads with their projective model structure.

As an application, we build explicit models for the N_∞ -operads of Blumberg and Hill.

Contents

1	Introduction	1
2	Basic definitions	2
3	Planar and tall maps	2
3.1	Planar structures	2
3.2	Outer faces and tall maps	6
3.3	Substitution	8
4	The genuine equivariant operad monad	10
4.1	Wreath product over finite sets	10
4.2	Equivariant leaf-root and vertex functors	12
4.3	Planar strings	14
4.4	A monad on spans	16
4.5	The free genuine operad monad	19
5	Free extensions	22
5.1	Extensions over general monads	22
5.2	Labeled planar strings	23
5.3	Bar constructions on spans	26
5.4	Transferring simplicial colimits of left Kan extensions	27
5.5	The category of extension trees	32
6	Filtration of Cellular Extensions	35
6.1	Filtration Pieces	35
6.2	Notation	39
6.3	Filtration Result	40

1 Introduction

No content yet.

2 Basic definitions

In this section we recall some definitions that will be used throughout.

Recall that for a diagram category \mathcal{D} and functor \mathcal{I}_\bullet

$$\begin{aligned} \mathcal{D} &\xrightarrow{\mathcal{I}_\bullet} \mathbf{Cat} \\ d &\longmapsto \mathcal{I}_d \end{aligned} \tag{2.1}$$

the (covariant) Grothendieck construction $\mathcal{D} \ltimes \mathcal{I}_\bullet$ has objects pairs (d, i) with $d \in \mathcal{D}$, $i \in \mathcal{I}_d$ and arrows $(d, i) \rightarrow (d', i')$ given by pairs

$$(f: d \rightarrow d', g: f_*(i) \rightarrow i'),$$

where $f_*: \mathcal{I}_d \rightarrow \mathcal{I}_{d'}$ is a shorthand for the functor $\mathcal{I}_\bullet(f)$.

We now discuss a basic property of over and under categories that will be used in §5.4.

Given $\mathcal{J}, \mathcal{C} \in \mathbf{Cat}$ and $j \in \mathcal{J}$ we will let $\mathcal{C}^{\downarrow j}$ denote the Grothendieck construction for the functor

$$\begin{aligned} \mathcal{J} &\longrightarrow \mathbf{Cat} \\ i &\longmapsto \mathcal{C}^{\mathcal{J}(i, j)} \end{aligned}$$

Explicitly, an object of $\mathcal{C}^{\downarrow j}$ is a pair $(i, \mathcal{J}(i, j) \xrightarrow{\varphi} \mathcal{C})$ and an arrow $(i, \varphi) \rightarrow (i', \varphi')$ is a pair $(I: i \rightarrow i', \gamma: \varphi \circ I^* \rightarrow \varphi')$.

Lemma 2.2. *Let $\mathcal{J} \in \mathbf{Cat}$ be a small category and $j \in \mathcal{J}$. One then has adjunctions*

$$(- \downarrow j): \mathbf{Cat}_{/\mathcal{J}} \rightleftarrows \mathbf{Cat}: (-)^{\downarrow j}, \quad (j \downarrow -): \mathbf{Cat}_{/\mathcal{J}} \rightleftarrows \mathbf{Cat}: (-)^{j \downarrow}.$$

Proof. Since $j \downarrow \mathcal{I} = (\mathcal{I}^{op} \downarrow j)^{op}$ by defining $(\mathcal{C}^{j \downarrow}) = ((\mathcal{C}^{op})^{\downarrow j})^{op}$ one reduces to the leftmost adjunction.

Given $\mathcal{I} \xrightarrow{\pi} \mathcal{J}$ and \mathcal{C} we will show that functors $\mathcal{I} \downarrow j \xrightarrow{F} \mathcal{C}$ correspond to functors $\mathcal{I} \xrightarrow{G} \mathcal{C}^{\downarrow j}$ over \mathcal{J} .

On objects, F associates to each pair $(i, J: \pi(i) \rightarrow j)$ an object $F(i, J) \in \mathcal{C}$. One thus sets $G(i) = (\pi(i), F(i, -))$ and these are clearly inverse processes.

On arrows F associates to $(i, J' \circ \pi(I)) \xrightarrow{I} (i', J')$ an arrow $F(i, J' \circ \pi(I)) \xrightarrow{F(I)} F(i', J')$. One thus defines

$$G(I) = \left(\pi(i) \xrightarrow{\pi(I)} \pi(i'), F(i, (-) \circ \pi(i)) \xrightarrow{F(I)} F(i', -) \right)$$

and again it is clear that these are inverse processes. Finally, the fact that the associativity and unit conditions for F, G coincide is likewise clear. \square

3 Planar and tall maps

3.1 Planar structures


Throughout we will work with trees possessing *planar structures* or, more intuitively, trees embedded into the plane.

Our preferred model for trees will be that of broad posets first introduced by Weiss in [We12] and further worked out by the second author in [Pef16]. We now define planar structures in this context.

Definition 3.1. Let $T \in \Omega$ be a tree. A *planar structure* of T is an extension of the descandancy partial order \leq_d to a total order \leq_p such that:

- *Planar*: if $e \leq_p f$ and $e \not\leq_d f$ then $g \leq_d f$ implies $e \leq_p g$.

Example 3.2. An example of a planar structure on a tree T follows, with \leq_r encoded by the number labels.



(3.3)

PLANAREX EQ

Intuitively, given a planar depiction of a tree T , $e \leq_d f$ holds when the downward path from e passes through f and $e \leq_p f$ holds if either $e \leq_d f$ or if the downward path from e is to the left of the downward path from f (as measured at the node where the paths intersect).

Intuitively, a planar depiction of a tree amounts to choosing a total order for each of the sets of *input edges* of each node (i.e. those edges immediately above that node).

While we will not need to make this last statement precise, we will nonetheless find it convenient to show that Definition 3.1 is equivalent to such choosing total orders for each of the sets of input edges. To do so, we first introduce some notation.

PLANARIZE DEF

Notation 3.4. Let $T \in \Omega$ be a tree and $e \in T$ and edge. We will denote

$$I(e) = \{f \in T : e \leq_d f\}$$

and refer to this poset as the *input path* of e .

We will repeatedly use the following, which is a consequence of [Pe16b, Cor. 5.26].

Lemma 3.5. If $e \leq_d f$, $e \leq_d f'$, then f, f' are \leq_d -comparable.

Proposition 3.6. Let $T \in \Omega$ be a tree. Then

- (a) for any $e \in T$ the finite poset $I(e)$ is totally ordered;
- (b) the poset (T, \leq_d) has all joins, denoted \vee . In fact, $\vee_i e_i = \min(\cap_i I(e_i))$.

Proof. (a) is immediate from Lemma 3.5. To prove (b) we note that $\min(\cap_i I(e_i))$ exists by (a), and that this is clearly the join $\vee e_i$. \square

Notation 3.7. Let $T \in \Omega$ be a tree and suppose that $e <_d b$. We will denote by $b_e^\uparrow \in T$ the predecessor of b in $I(e)$.

Proposition 3.8. Suppose e, f are \leq_d -incomparable edges of T and write $b = e \vee f$. Then

- (a) $e <_d b$, $f <_d b$ and $b_e^\uparrow \neq b_f^\uparrow$;
- (b) $b_e^\uparrow, b_f^\uparrow \in b^\uparrow$. In fact $\{b_e^\uparrow\} = I(e) \cap b^\uparrow$, $\{b_f^\uparrow\} = I(f) \cap b^\uparrow$;
- (c) if $e' \leq_d e$, $f' \leq_d f$ then $b = e' \vee f'$ and $b_{e'}^\uparrow = b_e^\uparrow$, $b_{f'}^\uparrow = b_f^\uparrow$.

Proof. (a) is immediate: the condition $e = g$ (resp. $f = g$) would imply $f \leq_d e$ (resp. $e \leq_d f$) while the condition $b_e^\uparrow = b_f^\uparrow$ would provide a predecessor of b in $I(e) \cap I(f)$.

For (b), note that any relation $a <_d b$ factors as $a \leq_d b_a^* <_d b$ for some unique $b_a^* \in b^\uparrow$, where uniqueness follows from Lemma 3.5. Choosing $a = e$ implies $I(e) \cap b^\uparrow = \{b_e^*\}$ and letting a range over edges such that $e \leq_d a <_d b$ shows that b_e^* is in fact the predecessor of b .

To prove (c) one reduces to the case $e' = e$, in which case it suffices to check $I(e) \cap I(f') = I(e) \cap I(f)$. But if it were otherwise there would exist an edge a satisfying $f' \leq_d a <_d f$ and $e \leq_d a$, and this would imply $e \leq_d f$, contradicting our hypothesis. \square

TERNARYJOIN PROP

Proposition 3.9. Let $c = e_1 \vee e_2 \vee e_3$. Then $c = e_i \vee e_j$ iff $c_{e_i}^\dagger \neq c_{e_j}^\dagger$.
Therefore, all ternary joins in (T, \leq_d) are binary, i.e.

$$c = e_1 \vee e_2 \vee e_3 = e_i \vee e_j \quad (3.10)$$


TERNJOIN EQ

for some $1 \leq i < j \leq 3$, and (3.10) fails for at most one choice of $1 \leq i < j \leq 3$.

Proof. If $c_{e_i}^\dagger \neq c_{e_j}^\dagger$, then $c = \min(I(e_i) \cap I(e_j)) = e_i \vee e_j$, whereas the converse follows from Proposition 3.8(a).

The “therefore” part follows by noting that $c_{e_1}^\dagger, c_{e_2}^\dagger, c_{e_3}^\dagger$ can not all coincide, or else c would not be the minimum of $I(e_1) \cap I(e_2) \cap I(e_3)$. \square

Example 3.11. In the following example $b = e \vee f$, $c = e \vee f \vee g$, $c_e^\dagger = c_f^\dagger = b$.



Notation 3.12. Given a set S of size n we write $\text{Ord}(S) \simeq \text{Iso}(S, \{1, \dots, n\})$. We will usually abuse notation by regarding its objects as pairs (S, \leq) where \leq is a total order in S .

Proposition 3.13. Let $T \in \Omega$ be a tree. There is a bijection

$$\{\text{planar structures } (T, \leq_p)\} \longrightarrow \prod_{(a^\dagger \leq a) \in V(T)} \text{Ord}(a^\dagger) \quad (3.14)$$

$$\leq_p \longmapsto (\leq_p |_{a^\dagger})$$

PLANAR EQ

Proof. We will keep the setup of Proposition 3.8 throughout: e, f are \leq_d -incomparable edges and we write $b = e \vee f$.

We first show that (3.14) is injective, i.e. that the restrictions $\leq_p |_{a^\dagger}$ determine if $e <_p f$ holds or not. If $b_e^\dagger <_p b_f^\dagger$, the relations $e \leq_d b_e^\dagger <_p b_f^\dagger \leq_d f$ and Definition 3.1 imply it must be $e <_p f$. Dually, if $b_f^\dagger <_p b_e^\dagger$ then $f <_p e$. Thus $b_e^\dagger <_p b_f^\dagger \Leftrightarrow e <_p f$ and hence (3.14) is indeed injective.

To check that (3.14) is surjective, it suffices (recall that e, f are assumed \leq_d -incomparable) to check that defining $e \leq_p f$ to hold iff $b_e^\dagger < b_f^\dagger$ holds in b^\dagger yields a planar structure.

Antisymmetry and the total order conditions are immediate, and it thus remains to check the transitivity and planar conditions. Transitivity of \leq_p in the case $e' <_p e <_p f$ and the planar condition, which is the case $e <_p f \geq_d f'$, follow from Proposition 3.8(c). Transitivity of \leq_p in the case $e <_p f \leq_d f'$ follows since either $e \leq_d f'$ or else e, f' are \leq_d -incomparable, in which case one can apply 3.8(c) with the roles of f, f' reversed.


It remains to check transitivity in the hardest case, that of $e <_p f <_p g$ with e, f incomparable f, g . We write $c = e \vee f \vee g$. By the “therefore” part of Proposition 3.9, either (i) $e \vee f <_d c$, in which case Proposition 3.9 implies $c_e^\dagger = c_f^\dagger$ and transitivity follows; (ii) $f \vee g <_d c$, which follows just as (i); (iii) $e \vee f = f \vee g = c$, in which case $c_e^\dagger < c_f^\dagger < c_g^\dagger$ in c^\dagger so that $c_e^\dagger \neq c_g^\dagger$ and by Proposition 3.9 it is also $e \vee g = c$ and transitivity follows. \square

Remark 3.15. Definition 3.1 readily extends to forests $F \in \Phi$. The analogue of Proposition 3.13 then states that the data of a planar structure is equivalent to total orderings of the nodes of F together with a total ordering of its set of roots. Indeed, this follows by either adapting the proof above or by noting that planar structures on F are clearly in bijection with planar structures on the join tree $F \star \eta$ (cf. [3, Def. 7.44]), which adds a single edge η to F , serving as the (unique) root of $F \star \eta$.

When discussing the substitution procedure in §3.3 we will find it convenient to work with a model for the category Ω that possesses exactly one representative of each possible planar structure on each tree or, more precisely, such that the only isomorphisms preserving the planar structures are the identities. On the other hand, using such a model for Ω throughout would, among other issues, make the discussion of faces in §3.2 rather awkward. We now outline our conventions to address such issues.

Let Ω^p , the category of *planarized trees*, denote the category with objects pairs $T_{\leq p} = (T, \leq_p)$ of trees together with a planar structure and morphisms the *underlying* maps of trees (so that the planar structures are ignored). There is a full subcategory $\Omega^s \hookrightarrow \Omega^p$, whose objects we call *standard models*, of those $T_{\leq p}$ whose underlying set is one of the sets $\underline{n} = \{1, 2, \dots, n\}$ and for which \leq_p coincides with the canonical order.

Example 3.16. Some examples of standard models, i.e. objects of Ω^s , follow (further, (3.3) can also be interpreted as such an example).



(3.17)

PLANAROMEGAEX1 EQ

Here T_1 and T_2 are isomorphic to each other but not isomorphic to any other standard model in Ω^s while both C and U are the unique objects in their isomorphism classes.

Given $T_{\leq p} \in \Omega^p$ there is an obvious standard model $T_{\leq p}^s \in \Omega^s$ given by replacing each edge by its order following \leq_p . Indeed, this defines a retraction $(-)^s: \Omega^p \rightarrow \Omega^s$ and a natural transformation $\sigma: id \Rightarrow (-)^s$ given by isomorphisms preserving the planar structure (in fact, the pair $((-)^s, \sigma)$ is clearly unique).

Convention 3.18. From now on, we will write simply Ω , Ω_G to denote the categories Ω^s , Ω_G^s of standard models (where planar structures are defined in the underlying forest as in Remark 3.15). Similarly \mathbf{O}_G will denote the model \mathbf{O}_G^s for the orbital category whose objects are the orbital G -sets whose underlying set is one of the sets $\underline{n} = \{1, 2, \dots, n\}$.


Therefore, whenever one of our constructions produces an object/diagram in Ω^p , Ω_G^p , \mathbf{O}_G^p (of trees, G -trees, orbital G -sets with a planarization/total order) we will hence implicitly reinterpret it by using the standardization functor $(-)^s$.

Example 3.19. To illustrate our convention, we consider the trees in Example 3.16.

One has subfaces $F_1 \subset F_2 \subset U$ where F_1 is the subtree with edge set $\{1, 2, 6, 7\}$ and F_2 is the subtree with edge set $\{1, 2, 3, 6, 7\}$, both with inherited tree and planar structures. Applying $(-)^s$ to the inclusion diagram on the left below then yields a diagram as on the right.




Similarly, let $\leq_{(12)}$ and $\leq_{(45)}$ denote alternate planar structures for U exchanging the orders of the pairs 1, 2 and 4, 5, so that one has objects $U_{\leq_{(12)}}$, $U_{\leq_{(45)}}$ in Ω^p . Applying $(-)^s$ to the diagram of underlying identities on the left yields the permutation diagram on the right.



Example 3.20. An additional reason to leave the use of $(-)^s$ implicit is that when depicting G -trees it is preferable to choose edge labels that describe the action rather than the planarization (which is already implicit anyway).

For example, when $G = \mathbb{Z}/4$, in both diagrams below the orbital representation on the left represents the isomorphism class consisting of the two trees $T_1, T_2 \in \Omega_G$ on the right.



Definition 3.21. A morphism $S \xrightarrow{\varphi} T$ in Ω that is compatible with the planar structures \leq_p is called a *planar map*.

More generally, a morphism $F \rightarrow G$ in the categories Φ, Φ^G, Ω^G of forests, G -forests, G -trees is called a *planar map* if it is an independent map (cf. [3, Def. 5.28]) compatible with the planar structures \leq_p .

Remark 3.22. The need for the independence condition is justified by [3, Lemma 5.33] and its converse, since non independent maps do not reflect \leq_d inequalities.

We note that in the Ω_G case a map φ is independent iff φ does not factor through a non trivial quotient iff φ is injective on each edge orbit.

Proposition 3.23. Let $F \xrightarrow{\varphi} G$ be an independent map in Φ (or Ω, Ω_G, Φ_G). Then there is a unique factorization

$$F \xrightarrow{\sim} \bar{F} \rightarrow G$$

such that $F \xrightarrow{\sim} \bar{F}$ is an isomorphism and $\bar{F} \rightarrow G$ is planar.

Proof. We need to show that there is a unique planar structure $\leq_p^{\bar{F}}$ on the underlying forest of F making the underlying map a planar map. Simplicity of G ensures that for any vertex $e^\dagger \leq e$ of F the edges in $\varphi(e^\dagger)$ are all distinct while independence of φ likewise ensures that the edges in $\varphi(e^\dagger)$ are distinct. The result now follows from (the forest version of) Proposition 5.13: one simply orders each set e^\dagger and \bar{r}_F according to its image.

not quite complete... maybe that \leq_p is the closure of \leq_d and the vertex relations under transitivity and the planar condition \square

Remark 3.24. Proposition 3.23 says that planar structures can be pulled back along independent maps. However, they can not always be pushed forward. As an example, in the notation of (3.17), consider the map $C \rightarrow T_1$ defined by $1 \mapsto 1, 2 \mapsto 4, 3 \mapsto 2, 4 \mapsto 5$.

Remark 3.25. Given any tree $T \in \Omega$ there is a unique corolla $\text{lr}(T) \in \Sigma$ and planar tall map $\text{lr}(T) \rightarrow T$. Explicitly, the number of leaves of $\text{lr}(T)$ matches that of T , together with the inherited order.

3.2 Outer faces and tall maps

In preparation for our discussion of the substitution operation in §3.3, we now recall some basic notions and results concerning outer subtrees and tree grafting, as in [3, §5].

Definition 3.26. Let $T \in \Omega$ be a tree and $e_1 \cdots e_n = \underline{e} \leq e$ a broad relation in T .

We define the *planar outer face* $T_{\underline{e} \leq e}$ to be the subtree with underlying set those edges $f \in T$ such that

$$f \leq_d e, \quad \forall_i e_i \not\leq_d f, \quad (3.27)$$

generating broad relations the relations $f^\dagger \leq f$ for f satisfying (3.27) and $\forall_i f \neq e_i$, and planar structure pulled back from T .

Remark 3.28. If one forgoes the requirement that $T_{\underline{e} \leq e}$ be equipped with the pullback planar structure, the inclusion $T_{\underline{e} \leq e} \rightarrow T$ is usually called simply an *outer face*.

We now recap some basic results.

Proposition 3.29. *Let $T \in \Omega$ be a tree.*

- (a) $T_{\underline{e} \leq e}$ is a tree with root e and edge tuple \underline{e} ;
- (b) there is a bijection

$$\{\text{planar outer faces of } T\} \leftrightarrow \{\text{broad relations of } T\};$$

- (c) if $R \rightarrow S$ and $S \rightarrow T$ are outer face maps then so is $R \rightarrow T$;
- (d) any pair of broad relations $\underline{g} \leq v$, $\underline{f}v \leq e$ induces a grafting pushout diagram

$$\begin{array}{ccc} \eta & \xrightarrow{v} & T_{\underline{g} \leq v} \\ v \downarrow & & \downarrow \\ T_{\underline{f}v \leq e} & \longrightarrow & T_{\underline{f}g \leq v} \end{array} \quad (3.30) \quad \boxed{\text{GRATPUSH EQ}}$$

Proof. We first show (a). That $T_{\underline{e} \leq e}$ is indeed a tree is the content of [Pe16b, Prop. 5.20]: more precisely, $T_{\underline{e} \leq e} = (T^{\leq e})_{< \underline{e}}$ in the notation therein. That the root of $T_{\underline{e} \leq e}$ is e is clear and that the root tuple is \underline{e} follows from [Pe16b, Remark 5.23].

(b) follows from (a), which shows that $\underline{e} \leq e$ can be recovered from $T_{\underline{e} \leq e}$.

(c) follows from the definition of outer face together with [Pe16b, Lemma 5.33], which states that the \leq_d relations on S, T coincide.

Since by (c) both $T_{\underline{g} \leq v}$ and $T_{\underline{f}v \leq e}$ are outer faces of $T_{\underline{f}g \leq v}$, (d) is a restatement of [Pe16b, Prop. 5.15]. \square

Definition 3.31. A map $S \xrightarrow{\varphi} T$ in Ω is called a *tall map* if

$$\varphi(l_S) = l_T, \quad \varphi(r_S) = r_T,$$

where $l_{(-)}$ denotes the leaf tuple and $r_{(-)}$ the root.

The following is a restatement of [Pe16b, Cor. 5.24]

Proposition 3.32. *Any map $S \xrightarrow{\varphi} T$ in Ω has a factorization, unique up to unique isomorphism,*

$$S \xrightarrow{\varphi^t} U \xrightarrow{\varphi^u} T$$

as a tall map followed by an outer face (in fact, $U = T_{\varphi(l_S) \leq r_S}$).

We recall that a face $F \rightarrow T$ is called inner if it is obtained by iteratively removing inner edges, i.e. edges other than the root or the leaves. In particular, it follows that a face is inner iff it is tall. The usual face-degeneracy decomposition thus combines with Corollary 3.32 to give the following.

Corollary 3.33. *Any map $S \xrightarrow{\varphi} T$ in Ω has a factorization, unique up to unique isomorphisms,*

$$S \xrightarrow{\varphi^-} U \xrightarrow{\varphi^i} V \xrightarrow{\varphi^u} T \quad (3.34) \quad \boxed{\text{TRIPLEFACT EQ}}$$

as a degeneracy followed by an inner face followed by an outer face.

Proof. The factorization (3.34) can be built by first performing the degeneracy-face decomposition and then performing the tall-outer decomposition on the face map. \square


3.3 Substitution

One of the key ideas needed to describe operads is that of substitution of tree nodes, a process that we will prefer to repackage in terms of maps of trees. We start by discussing an example, focusing on the related notion of iterated graftings of trees (as described in (3.30)).

Example 3.35. The trees U_1, U_2, \dots, U_6 on the left below can be grafted into the tree U in the middle. More precisely (among other possible grafting orders), one has

$$U = (((((U_6 \sqcup_a U_2)) \sqcup_a U_1) \sqcup_b U_3) \sqcup_d U_5) \sqcup_c U_4) \quad (3.36)$$

UFORMULA EQ




(3.37)

SUBSDATUMTREES EQ

We now consider the tree T , which is built by converting each U_i into the corolla $\text{Irr}(U_i)$ (cf. Remark 3.25), and then performing the same grafting operations as in (3.36). T can then be regarded as encoding the combinatorics of the iterated grafting in (3.36), with alternative ways to reorder operations in (3.36) in bijection with ways to assemble T out of its nodes.

One can now therefore think of the iterated grafting (3.36) as being instead encoded by the tree T together with the (unique) planar tall maps φ_i below.



(3.38)

SUBSDATUMTREES2 EQ

From this perspective, U can now be thought as obtained from T by substituting each of its nodes with the corresponding U_i . Moreover, the φ_i assemble to a planar tall map $\varphi: T \rightarrow U$ (such that $a_i \mapsto a, b \mapsto b, \dots, e \mapsto e$), which likewise encodes the same information.

Our perspective will then be that data for substitution of tree nodes such as in (3.38) can equivalently be repackaged using planar tall maps.

Definition 3.39. Let $T \in \Omega$ be a tree.

A T -substitution datum is a tuple $\{U_{e^\dagger \leq e}\}_{(e^\dagger \leq e) \in V(T)}$ such that $\text{lr}(U_{e^\dagger \leq e}) = T_{e^\dagger \leq e}$.

Further, a map of T -substitution data $\{U_{e^\dagger \leq e}\} \rightarrow \{V_{e^\dagger \leq e}\}$ is a tuple of planar tall maps $\{U_{e^\dagger \leq e} \rightarrow V_{e^\dagger \leq e}\}$.

Definition 3.40. Let $T \in \Omega$.

The *Segal core poset* $\mathbf{Sc}(T)$ is the poset with objects the edge subtrees η_e and vertex subtrees $T_{e^\dagger \leq e}$. The order relation is given by inclusion.

Remark 3.41. Note that the only maps in $\mathbf{Sc}(T)$ are inclusions of the form $\eta_a \subset T_{e^\dagger \leq e}$. In particular, there are no pairs of composable non-identity relations in $\mathbf{Sc}(T)$.

Given a T -substitution datum $\{U_{\{e^\dagger \leq e\}}\}$ we abuse notation by writing

$$U_{(-)} : \mathbf{Sc}(T) \rightarrow \Omega$$

for the functor $\eta_a \mapsto \eta$, $T_{e^\dagger \leq e} \mapsto U_{e^\dagger \leq e}$ and sending the inclusions $\eta_a \subset T_{e^\dagger \leq e}$ to the composites

$$\eta \xrightarrow{a} T_{e^\dagger \leq e} = \text{lr}(U_{e^\dagger \leq e}) \rightarrow U_{e^\dagger \leq e}.$$

Proposition 3.42. Let $T \in \Omega$ be a tree. There is an isomorphism of categories

$$\begin{aligned} \text{Sub}(T) &\xrightleftharpoons{\quad} \Omega_{T|}^{\text{pt}} \\ \{U_{e^\dagger \leq e}\} &\longmapsto (T \rightarrow \text{colim}_{\mathbf{Sc}(T)} U_{(-)}) \\ \{U_{\varphi(e^\dagger) \leq \varphi(e)}\} &\longleftarrow (T \xrightarrow{\varphi} U) \end{aligned} \tag{3.43}$$

SUBDATAUNDERPLAN EQ

Where $\text{Sub}(T)$ denotes the category of T -substitution data and $\Omega_{T|}^{\text{pt}}$ the category of planar tall maps under T .

Proof. We first claim that (i) the $\text{colim}_{\mathbf{Sc}(T)} U_{(-)}$ indeed exists; (ii) for the canonical datum $\{T_{e^\dagger \leq e}\}$, it is $T = \text{colim}_{\mathbf{Sc}(T)} T_{(-)}$; (iii) the induced map $T \rightarrow \text{colim}_{\mathbf{Sc}(T)} U_{(-)}$ is planar tall.

The argument is by induction on the number of vertices of T , with the base cases of T with 0 or 1 vertices being immediate, since then T is the terminal object of $\mathbf{Sc}(T)$. Otherwise, one can choose a non trivial grafting decomposition so as to write $T = R \sqcup_e S$, resulting in identifications $\mathbf{Sc}(R) \subset \mathbf{Sc}(T)$, $\mathbf{Sc}(S) \subset \mathbf{Sc}(T)$ so that $\mathbf{Sc}(R) \cup \mathbf{Sc}(S) = \mathbf{Sc}(T)$ and $\mathbf{Sc}(R) \cap \mathbf{Sc}(S) = \{\eta_e\}$. The existence of $\text{colim}_{\mathbf{Sc}(T)} U_{(-)}$ is thus equivalent to the existence of the pushout below.

$$\begin{array}{ccc} \eta & \longrightarrow & \text{colim}_{\mathbf{Sc}(R)} U_{(-)} \\ \downarrow & & \downarrow \\ \text{colim}_{\mathbf{Sc}(S)} U_{(-)} & \dashrightarrow & \text{colim}_{\mathbf{Sc}(T)} U_{(-)} \end{array} \tag{3.44}$$

ASSEMBLYGRAFT EQ

By induction, the top right and bottom left colimits exist for any $U_{(-)}$, equal R and S in the case $U_{(-)} = T_{(-)}$, and the maps $R \rightarrow \text{colim}_{\mathbf{Sc}(R)} U_{(-)}$, $S \rightarrow \text{colim}_{\mathbf{Sc}(S)} U_{(-)}$ are planar tall. But it now follows that (3.44) is a grafting pushout diagram, so that the pushout indeed exists. The conditions that $T = \text{colim}_{\mathbf{Sc}(T)} T_{(-)}$ and $T \rightarrow \text{colim}_{\mathbf{Sc}(T)} U_{(-)}$ is planar tall follow.

The fact that the two functors in (3.43) are inverse to each other is clear by the same inductive argument. \square

Remark 3.45. It follows from the previous proof that, writing $U = \text{colim}_{\mathbf{Sc}(T)} U_{(-)}$, one has

$$V(U) = \coprod_{(e^\dagger \leq e) \in V(T)} V(U_{e^\dagger \leq e}). \tag{3.46}$$

VERTEXDECOMP EQ

Alternatively, (3.46) can be regarded as a map $f^* : V(U) \rightarrow V(T)$ induced by the planar tall map $f : T \rightarrow U$. Explicitly, $f^*(U_{u^\dagger \leq u})$ is the unique $T_{t^\dagger \leq t}$ such that $U_{u^\dagger \leq u} \subset U_{t^\dagger \leq t}$. We note that f^* is indeed contravariant in the tall planar map f .

BUILDABLE PROP

The following is a converse of sorts to Proposition 3.42. SUBDATAUNDERPLAN PROP

Proposition 3.47. *Let $U \in \Omega$ be a tree. Then:*

- (i) *given non stick outer subtrees U_i such that $V(U) = \coprod_i V(U_i)$ there is a unique tree T and planar tall map $T \rightarrow U$ such that $\{U_i\} = \{U_{e| \leq e}\}$;*
- (ii) *given multiplicities $m_e \geq 1$ for each edge $e \in U$, there is a unique planar degeneracy $\rho: T \rightarrow U$ such that $\rho^{-1}(e)$ has m_e elements;*
- (iii) *planar tall maps $T \rightarrow U$ are in bijection with collections $\{U_i\}$ of outer subtrees such that $V(U) = \coprod_i V(U_i)$ and U_j is not an inner edge of any U_i whenever $U_j \simeq \eta$ is a stick.*

Proof. We first show (i) by induction on the number of subtrees U_i . The base case $\{U_i\} = \{U\}$ is immediate, setting $T = \text{lr}(U)$. Otherwise, letting e be edge that is both an inner edge of U and a root of some U_i , and one can form a pushout diagram

$$\begin{array}{ccc} \eta & \xrightarrow{e} & V \\ e \downarrow & & \downarrow \\ W & \longrightarrow & U \end{array} \quad (3.48) \quad \text{DECOMPPROOF EQ}$$

inducing a nontrivial partition $\{U_i\} = \{U_i | U_i \hookrightarrow V\} \sqcup \{U_i | U_i \hookrightarrow W\}$. Existence of $T \rightarrow U$ now follows from the induction hypothesis. For uniqueness, the condition that no U_i is a stick guarantees that T possesses a single inner edge mapping to e , and thus admits a compatible decomposition as in (3.48), and thus uniqueness too follows by the induction hypothesis. DECOMPPROOF EQ

For (ii), we argue existence by nested induction on the number of vertices $|V(U)|$ and the sum of the multiplicities m_e . The base case $|V(U)| = 0$, i.e., $U = \eta$ is immediate. Otherwise, writing $m_e = m'_e + 1$, one can form a decomposition (3.48) where either $|V(V)|, |V(W)| < |V(U)|$ or one of V, W is η , so that $T \rightarrow U$ can be built via the induction hypothesis. For uniqueness, note first that by [3, Lemma 5.33] each pre-image $\rho^{-1}(e)$ is linearly ordered and by the “further” claim in [3, Cor. 5.39] the remaining broad relations are precisely the pre-image of the non-identity relations in U , showing that the tree T is uniquely determined. DECOMPPROOF EQ

(iii) follows by combining (i) and (ii). Indeed, any planar tall map $T \rightarrow U$ has a unique decomposition $T \twoheadrightarrow \bar{T} \hookrightarrow U$ as a planar degeneracy followed by a planar inner face, and each of these maps is classified by the data in (b) and (a). □

Lemma 3.49. *Suppose $T_1, T_2 \hookrightarrow T$ are two outer faces with at least one common edge e . Then there exists a unique outer face $T_1 \cup T_2$ such that $V(T_1 \cup T_2) = V(T_1) \cup V(T_2)$.*

Proof. If either of T_1, T_2 is the root or a leaf the result is obvious. Otherwise, one can necessarily choose e to be an inner edge of T , in which case all of T_1, T_2, T admit compatible decompositions (3.48) and the result follows by induction on $|V(T)|$. □ DECOMPPROOF EQ

4 The genuine equivariant operad monad

We now turn to the task of building the monad encoding genuine equivariant operads.

4.1 Wreath product over finite sets

In what follows we will let \mathbf{F} denote the usual skeleton of the category of finite sets and all set maps. Explicitly, its objects are the finite sets $\{1, 2, \dots, n\}$ for $n \geq 0$. However, much as in the discussion in Convention 3.18 we will often find it more convenient to regard the elements of \mathbf{F} as equivalence classes of finite sets equipped with total orders. PLANARCONV CON

Definition 4.1. For a category \mathcal{C} , we let $F \wr \mathcal{C}$ denote the opposite of the Grothendieck construction for the functor

$$\begin{aligned} F^{op} &\longrightarrow \mathbf{Cat} \\ I &\longmapsto \mathcal{C}^I \end{aligned}$$

Explicitly, the objects of $F \wr \mathcal{C}$ are tuples $(c_i)_{i \in I}$ and a map $(c_i)_{i \in I} \rightarrow (d_j)_{j \in J}$ consists of a pair

$$(\phi: I \rightarrow J, (f_i: c_i \rightarrow d_{\phi(i)})_{i \in I}),$$

henceforth abbreviated as $(\phi, (f_i))$.

The following is immediate.

Proposition 4.2. Suppose \mathcal{C} has all finite coproducts. One then has a functor as on the left below. Dually, if \mathcal{C} has all finite products, one has a functor as on the right below.

$$\begin{aligned} F \wr \mathcal{C} &\xrightarrow{\coprod} \mathcal{C} & (F \wr \mathcal{C}^{op})^{op} &\xrightarrow{\prod} \mathcal{C} \\ (c_i)_{i \in I} &\longmapsto \coprod_{i \in I} c_i & (c_i)_{i \in I} &\longmapsto \prod_{i \in I} c_i \end{aligned}$$

Lemma 4.3. Suppose that \mathcal{E} is a bicomplete category such that coproducts commute with limits in each variable. If the leftmost diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{E} \\ k \downarrow & \nearrow \eta & \uparrow G \\ \mathcal{D} & & \end{array} \quad \begin{array}{ccccc} F \wr \mathcal{C} & \xrightarrow{F \wr F} & F \wr \mathcal{E} & \xrightarrow{\prod} & \mathcal{E} \\ F \wr k \downarrow & \nearrow F \wr \eta & \nearrow F \wr G & & \uparrow \prod \circ F \wr G \\ F \wr \mathcal{D} & & & & \end{array} \quad (4.4) \quad \boxed{\text{WRRAN EQ}}$$

is a right Kan extension diagram then so is the composite of the rightmost diagram.

Dually, if in \mathcal{E} products commute with colimits in each variable, and the leftmost diagram

$$\begin{array}{ccc} \mathcal{C}^{op} & \xrightarrow{F} & \mathcal{E} \\ k \downarrow & \nearrow \epsilon & \uparrow G \\ \mathcal{D}^{op} & & \end{array} \quad \begin{array}{ccccc} (F \wr \mathcal{C})^{op} & \xrightarrow{(F \wr F)^{op}} & (F \wr \mathcal{E})^{op} & \xrightarrow{\prod} & \mathcal{E} \\ (F \wr k)^{op} \downarrow & \nearrow & \nearrow (F \wr G)^{op} & & \uparrow \prod \circ (F \wr G)^{op} \\ (F \wr \mathcal{D})^{op} & & & & \end{array} \quad (4.5) \quad \boxed{\text{WRLAN EQ}}$$

is a left Kan extension diagram then so is the composite of the rightmost diagram.

Proof. Unpacking definitions using the pointwise formula for Kan extensions ([2, X.3.1]), the claim concerning (4.4) amounts to showing that for each $(d_i) \in F \wr \mathcal{D}$ one has natural isomorphisms

$$\lim_{((d_i) \rightarrow (kc_j)) \in ((d_i) \downarrow F \wr \mathcal{C})} \left(\coprod_j F(c_j) \right) \simeq \coprod_i \lim_{(d_i \rightarrow kc_i) \in d_i \downarrow \mathcal{C}} (F(c_i)). \quad (4.6) \quad \boxed{\text{POINTKAN EQ}}$$

Noting that the canonical factorizations of each $(\varphi, (f_i)): (d_i)_{i \in I} \rightarrow (kc_j)_{j \in J}$ as

$$(d_i)_{i \in I} \rightarrow (c_{\phi(i)})_{i \in I} \rightarrow (kc_j)_{j \in J}$$

exhibit $\prod_i (d_i \downarrow \mathcal{C})$ as a coreflexive subcategory of $(d_i) \downarrow F \wr \mathcal{C}$, we see that it is an initial subcategory. Therefore

$$\lim_{((d_i) \rightarrow (kc_j)) \in ((d_i) \downarrow F \wr \mathcal{C})} \left(\coprod_j F(c_j) \right) \simeq \lim_{((d_i) \rightarrow (kc_i)) \in \prod_i (d_i \downarrow \mathcal{C})} \left(\coprod_i F(c_i) \right)$$

and hence (4.6) now follows from the assumption that coproducts commute with limits in each variable. \square

Notation 4.7. Using the coproduct functor $F^{i2} = F^{i\{0,1\}} = F \wr F \xrightarrow{u} F$ (where $\coprod_{i \in I} J_i$ is ordered lexicographically) and the simpleton $\{1\} \in F$ one can regard the collection of categories $F^{i\{0,\dots,n\}} \wr \mathcal{C} = F^{in} \wr \mathcal{C}$ as a coaugmented cosimplicial object in \mathbf{Cat} . As such, we will denote by

$$\delta^i: F^{in-1} \wr \mathcal{C} \rightarrow F^{in} \wr \mathcal{C}, \quad 0 \leq i \leq n$$

the cofaces obtained by inserting simpletons $\{1\} \in F$ and by

$$\sigma^i: F^{in+1} \wr \mathcal{C} \rightarrow F^{in} \wr \mathcal{C}, \quad 0 \leq i \leq n$$

the codegeneracies obtained by applying the coproduct $F^{i2} \xrightarrow{u} F$ to adjacent F coordinates.

4.2 Equivariant leaf-root and vertex functors

Definition 4.8. A morphism $T \xrightarrow{\varphi} S$ in Ω_G is called a *quotient* if the underlying morphism of forests

$$\coprod_{[g] \in G/H} T_{[g]} \rightarrow \coprod_{[h] \in G/K} S_{[h]}$$

maps each tree component (or, equivalently, some tree component) isomorphically onto its image component.

We denote the subcategory of G -trees and quotients by Ω_G^q .


Definition 4.9. The *G-symmetric category*, which we will also call the *category of G-corollas*, is the full subcategory $\Sigma_G \subset \Omega_G^q$ of those G -trees that are corollas, i.e. G -trees such that each edge is either a root or a leaf (but not both).

Definition 4.10. The *leaf-root functor* is the functor $\Omega_G^q \xrightarrow{\text{lr}} \Sigma_G$ defined by

$$\text{lr}(T) = \{\text{leaves of } T\} \sqcup \{\text{roots of } T\}$$

with a broad relation $l_1 \cdots l_n \leq r$ holding in $\text{lr}(T)$ iff its image holds in T and similarly for the planar structure \leq_p .

Remark 4.11. Generalizing Remark 3.25, $\text{lr}(T)$ can alternatively be characterized as being the *unique* G -corolla which admits an also unique (tree-wise) tall planar map $\text{lr}(T) \rightarrow T$. Moreover, $\text{lr}(T)$ can usually be regarded as the “smallest inner face” of T , obtained by removing all the inner edges, although this characterization fails when $T = G \cdot_H \eta$ is a stick G -tree. Some examples with $G = \mathbb{Z}/4$ follow.



Remark 4.12. One consequence of the fact that planarizations can not be pushed forward along tree maps (cf. Remark 3.24) is that $\text{lr}: \Omega_G^q \rightarrow \Sigma_G$ is not a categorical fibration. **maybe add to this.**

VG DEF


Definition 4.13. Given $T \in \Omega_G$ we define the set $V_G(T)$ of G -vertices of T to be the orbit set $V(T)/G$, i.e. the quotient of the vertex set $V(T)$ by its G -action.

Furthermore, we will regard $V_G(T)$ as an object in \mathbf{F} by equipping it with its lexicographic order: i.e. vertex equivalence classes $[e^\dagger \leq e]$ are ordered according to the planar order \leq_p of the smallest representative ge , $g \in G$.

VERTEXDECOMP REM

Remark 4.14. Following Remark 3.45, a planar tall map $f: T \rightarrow U$ of G -trees induces a G -equivariant map $f^*: V(U) \rightarrow V(T)$ and thus also a map of orbits $f^*: V_G(U) \rightarrow V_G(T)$. We note, however, that f^* is not in general compatible with the order on V_G , as is indeed the case even in the non-equivariant case.

A minimal example follows.



In $V(T)$ the vertices are ordered as $a < c < d$ while in $V(U)$ they are ordered as $a < e < c < d$ but the map $f^*: V(U) \rightarrow V(T)$ is given by $a \mapsto a, c \mapsto c, d \mapsto d, e \mapsto d$.

Note that each element of $V_G(T)$ corresponds to an unique edge orbit Ge for e not a leaf. As such, we will represent the corresponding G -vertex by $v_{Ge} = (Ge)^\dagger \leq Ge$ (which we interpret as the concatenation of the relations $f^\dagger \leq f$ for $f \in Ge$) and write


$$T_{v_{Ge}} = T_{(Ge)^\dagger \leq Ge} = \coprod_{f \in Ge} T_{f^\dagger \leq f}.$$

We note that $T_{v_{Ge}}$ is always a G -corolla. Indeed, noting that a quotient map $\varphi: T \rightarrow S$ induces quotient maps $T_{v_{ge}} \rightarrow S_{v_{G\varphi(e)}}$ one obtains a functor

$$\begin{aligned} \Omega_G^q &\xrightarrow{V_G} \mathbf{F} \wr \Sigma_G \\ T &\longmapsto (T_{v_{Ge}})_{v_{Ge} \in V_G(T)}. \end{aligned} \tag{4.15}$$

VFUNCTOR EQ

Remark 4.16. The need to introduce the $\mathbf{F} \wr \mathcal{C}$ categories comes from the fact that general quotient maps do not preserve the number of G -vertices. For a simple example, let $G = \mathbb{Z}/4$ and consider the quotient map



sending edges labeled a, b, c to the edges with the same name and the edges a^{+1}, b^{+1} to the edges $a + 1, b + 1$. We note that T has three G -vertices $v_{Gc}, v_{Gb}, v_{Gb^{+1}}$ while S has only two G -vertices v_{Gc} and v_{Gb} . $V(\phi)$ then maps the two corollas $T_{v_{Gb}}$ and $T_{v_{Gb^{+1}}}$ isomorphically onto $T_{S_{Gb}}$ and the corolla $T_{v_{Gc}}$ non-isomorphically onto $S_{v_{Gc}}$.

Definition 3.39 now immediately generalizes.

TUTIONDATUMG DEF

Definition 4.17. Let $T \in \Omega_G$ be a G -tree.

A T -substitution datum is a tuple $\{U_{v_{Ge}}\}_{v_{Ge} \in V_G(T)}$ such that $\text{lr}(U_{v_{Ge}}) = T_{v_{Ge}}$.

Further, a map of T -substitution data $\{U_{v_{Ge}}\} \rightarrow \{V_{v_{Ge}}\}$ is a tuple of planar tall maps $\{U_{v_{Ge}} \rightarrow V_{v_{Ge}}\}$.

UBSDATUMCONV REM

Remark 4.18. To establish the equivariant analogue of Proposition 3.42 we will prefer to repackage equivariant substitution data in terms of non-equivariant terms.

Noting that there are decompositions $U_{v_{Ge}} = \coprod_{ge \in G_e} U_{ge \uparrow \leq ge}$ and letting $G \ltimes V(T)$ denote the Grothendieck construction for the action of G on the non-equivariant vertices $V(T)$ (often called the action groupoid), it is immediate that an equivariant T -substitution datum is the same as a functor $G \ltimes V(T) \rightarrow \Omega$ whose restriction to $V(T) \subset G \ltimes V(T)$ is a (non-equivariant) substitution datum.

AUNDERPLANG PROP

Proposition 4.19. Let $T \in \Omega_G$ be a G -tree. There is an isomorphism of categories

$$\begin{aligned} \text{Sub}(T) &\xrightleftharpoons{\quad} \Omega_{G,T}^{\text{pt}} \\ \{U_{v_{Ge}}\} &\longmapsto (T \rightarrow \text{colim}_{\text{Sc}(T)} U_{(-)}) \end{aligned} \tag{4.20}$$

SUBDATAUNDERPLANG EQ

Proof. This is a minor adaptation of the non-equivariant analogue Proposition 4.19. Since $\text{Sc}(T)$ inherits a G -action, one can form the Grothendieck construction $G \ltimes \text{Sc}(T)$ and by Remark 4.18 equivariant substitution data $\{U_{v_{Ge}}\}$ therefore induce functors $U_{(-)}: G \ltimes \text{Sc}(T) \rightarrow \Omega$. It is then immediate that $\text{colim}_{\text{Sc}(T)} U_{(-)}$ inherits a G -action, provided it exists. The key observation is then that, since $\text{Sc}(T)$ is now a disconnected poset, this colimit is to be interpreted as taken in the category Φ of forests rather than in Ω . \square

Remark 4.21. We will need to know that each of the maps

$$U_{v_{Ge}} \rightarrow U = \text{colim}_{\text{Sc}(T)} U_{(-)}$$

induced by the previous proof is a planar map of G -trees. This requires two observations: (i) the restrictions to each of the constituent non-equivariant trees $U_{ge \uparrow \leq ge}$ is planar by Proposition 4.19; (ii) the restriction to the roots of $U_{v_{Ge}}$ is injective and order preserving since it matches the inclusion of the roots of $T_{v_{Ge}}$, and the map $T \rightarrow U$ is a planar map of G -trees.

PLANARSTRING SEC

4.3 Planar strings

The leaf-root and vertex functors will allow us to reinterpret our results concerning substitution.

Definition 4.22. The category $\Omega_{G,n}$ of *substitution n -strings* is the category whose objects are strings

$$T_0 \xrightarrow{f_1} T_1 \xrightarrow{f_2} \cdots \xrightarrow{f_n} T_n$$

where $T_i \in \Omega_G$ and the f_i are tall planar maps, and arrows are commutative diagrams

$$\begin{array}{ccccccc} T_0 & \xrightarrow{f_1} & T_1 & \xrightarrow{f_2} & \cdots & \xrightarrow{f_n} & T_n \\ q_0 \downarrow & & q_1 \downarrow & & & & q_n \downarrow \\ T'_0 & \xrightarrow{f'_1} & T'_1 & \xrightarrow{f'_2} & \cdots & \xrightarrow{f'_n} & T'_n \end{array} \tag{4.23}$$

PTNARROW EQ

where each q_i is a quotient map.

Notation 4.24. Since compositions of planar tall arrows are planar tall and identity arrows are planar tall it follows that $\Omega_{G,\bullet}$ forms a simplicial object in \mathbf{Cat} , with faces given by composing and degeneracies by inserting identities.

Noting that $\Omega_{G,0} = \Omega_G^q$ and setting $\Omega_{G,-1} = \Sigma_G$, the leaf-root functor $\Omega_G^q \xrightarrow{\text{lr}} \Sigma_G$ makes $\Omega_{G,\bullet}^q$ into an augmented simplicial object and, furthermore, the maps $s_{-1}: \Omega_{G,n}^q \rightarrow \Omega_{G,n+1}^q$ sending $T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_n$ to $\text{lr}(T_0) \rightarrow T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_n$ equip it with extra degeneracies.

IMPOPERATORS NOT

Notation 4.25. We extend the vertex functor to a functor $V_G: \Omega_{G,n+1} \rightarrow \mathbf{F} \wr \Omega_{G,n}$ by

$$V_G(T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_n) = (T_{1,v_{Ge}} \rightarrow \cdots \rightarrow T_{n,v_{Ge}})_{v_{Ge} \in V_G(T_0)} \quad (4.26) \quad \text{VGDEF EQ}$$

where we abuse notation by writing $T_{i,v_{Ge}}$ for $T_{i,(f_i \circ \cdots \circ f_1)(v_{Ge})}$.

The following is a reinterpretation of Proposition 4.19.

Proposition 4.27. *The diagram*

$$\begin{array}{ccc} \Omega_{G,n+1} & \xrightarrow{V_G} & \mathbf{F} \wr \Omega_{G,n} \\ d_{1,\dots,n+1} \downarrow & & \downarrow \text{Fid}_{0,\dots,n} \\ \Omega_{G,0} & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G \end{array} \quad (4.28) \quad \text{PTPULL EQ}$$

is a pullback diagram in \mathbf{Cat} .

Proof. An object in the pullback (4.28) over $T \in \Omega_{G,n} = \Omega_n^q$ is precisely the same as a n -string in $\mathbf{Sub}(T)$, and thus by Proposition 4.19 equivalent to a $n+1$ planar tall string starting at T .

The case of arrows is slightly more subtle. A quotient map $\pi: T \rightarrow T'$ induces a G -equivariant poset map $\pi_*: \mathbf{Sc}(T) \rightarrow \mathbf{Sc}(T')$ (or equivalently, a map of Grothendieck constructions $G \ltimes \mathbf{Sc}(T) \rightarrow G \ltimes \mathbf{Sc}(T')$) and diagrams as on the left below (where v_{Ge} ranges over $V_G(T)$ and $e' = \varphi(e)$) induce diagrams (of functors $\mathbf{Sc}(T) \rightarrow \Omega$) as on the right below.

$$\begin{array}{ccc} T_{v_{Ge}} \rightarrow T_{1,v_{Ge}} \rightarrow \cdots \rightarrow T_{n,v_{Ge}} & T_{(-)} \longrightarrow T_{1,(-)} \longrightarrow \cdots \longrightarrow T_{n,(-)} \\ \downarrow & \downarrow & \downarrow \\ T'_{v_{Ge'}} \rightarrow T'_{1,v_{Ge'}} \rightarrow \cdots \rightarrow T'_{n,v_{Ge'}} & T'_{(-)} \circ \pi_* \rightarrow T'_{1,(-)} \circ \pi_* \rightarrow \cdots \rightarrow T'_{n,(-)} \circ \pi_* \end{array} \quad (4.29) \quad \text{PTNARROWLOC EQ}$$

Passing to colimits then gives the desired commutative diagram (4.23). Moreover, diagrams of the form (4.23) clearly induce diagrams as in (4.29) and it is straightforward to check that these are inverse processes. \square

Remark 4.30. The diagrams (with back and lower slanted faces instances of (4.28))

$$\begin{array}{ccc} \Omega_{G,n+2} & \xrightarrow{\quad} & \mathbf{F} \wr \Omega_{G,n+1} \\ \downarrow & \searrow d_{i+1} & \downarrow \text{Fid}_i \\ & \Omega_{G,n+1} & \xrightarrow{\quad} \mathbf{F} \wr \Omega_{G,n} \\ \downarrow & \swarrow & \downarrow \\ \Omega_{G,0} & \xrightarrow{\quad} & \mathbf{F} \wr \Sigma_G \end{array} \quad \begin{array}{ccc} \Omega_{G,n+1} & \xrightarrow{\quad} & \mathbf{F} \wr \Omega_{G,n} \\ \downarrow & \searrow s_i & \downarrow \text{Fid}_{s_{i-1}} \\ & \Omega_{G,n+2} & \xrightarrow{\quad} \mathbf{F} \wr \Omega_{G,n+1} \\ \downarrow & \swarrow & \downarrow \\ \Omega_{G,0} & \xrightarrow{\quad} & \mathbf{F} \wr \Sigma_G \end{array}$$

commute whenever defined (i.e. $0 \leq i \leq n+1$).

Notation 4.31. We will let

$$V_{G,n}: \Omega_{G,n} \rightarrow \mathbf{F} \wr \Sigma_G$$

be inductively defined by $V_{G,n} = \sigma_0 \circ V_{G,n-1} \circ V_G$.

Remark 4.32. When $n = 2$, $V_{G,2}$ is thus the composite

$$\Omega_{G,2} \xrightarrow{V_G} \mathbf{F} \wr \Omega_{G,1} \xrightarrow{V_G} \mathbf{F} \wr \mathbf{F} \wr \Omega_{G,0} \xrightarrow{V_G} \mathbf{F} \wr \mathbf{F} \wr \mathbf{F} \wr \Sigma_G \xrightarrow{\sigma^0} \mathbf{F} \wr \mathbf{F} \wr \Sigma_G \xrightarrow{\sigma^0} \mathbf{F} \wr \Sigma_G$$

In light of Remarks 3.45 and 4.14, $V_{G,n}(T_0 \rightarrow \cdots \rightarrow T_n)$ is identified with the tuple

$$(T_{n,v_{Ge}})_{v_{Ge} \in V_G(T_n)}, \quad (4.33) \quad \text{VGNISO EQ}$$

though this requires changing the total order in $V_G(T_n)$. Rather than using the order induced by T_n , one instead equips $V_G(T_n)$ with the order induced lexicographically from the maps $V_G(T_n) \rightarrow V_G(T_{n-1}) \rightarrow \cdots \rightarrow V_G(T_0)$, i.e., for $v, w \in V_G(T_n)$ the condition $v < w$ is determined by the lowest i such that the images of $v, w \in V_G(T_i)$ are distinct.

4.4 A monad on spans

WSPAN DEF

Definition 4.34. We will write $\mathbf{WSpan}^l(\mathcal{C}, \mathcal{D})$ (resp. $\mathbf{WSpan}^r(\mathcal{C}, \mathcal{D})$), which we call the category of *left weak spans* (resp. *right weak spans*), to denote the category with objects the spans

$$\mathcal{C} \xleftarrow{k} A \xrightarrow{F} \mathcal{D},$$

arrows the diagrams as on the left (resp. right) below

$$\begin{array}{ccc} & A_1 & \\ k_1 \swarrow & & \searrow F_1 \\ \mathcal{C} & i \downarrow & \mathcal{D} \\ k_2 \swarrow & \Downarrow \varphi & \nearrow F_2 \\ & A_2 & \end{array} \quad \begin{array}{ccc} & A_1 & \\ k_1 \swarrow & & \searrow F_1 \\ \mathcal{C} & i \downarrow & \mathcal{D} \\ k_2 \swarrow & \Downarrow \varphi & \nearrow F_2 \\ & A_2 & \end{array} \quad (4.35)$$

TWISTEDARROWRIGHT EQ

which we write as $(i, \varphi): (k_1, F_1) \rightarrow (k_2, F_2)$, and composition given in the obvious way.

Remark 4.36. There are natural isomorphisms

$$\mathbf{WSpan}^r(\mathcal{C}, \mathcal{D}) \simeq \mathbf{WSpan}^l(\mathcal{C}^{op}, \mathcal{D}^{op}). \quad (4.37)$$

LRSPANISO EQ

Remark 4.38. The terms *left/right* are motivated by the existence of adjunctions (which are seen to be equivalent by using (4.37))

$$\mathbf{Lan}: \mathbf{WSpan}^l(\mathcal{C}, \mathcal{D}) \rightleftarrows \mathbf{Fun}(\mathcal{C}, \mathcal{D}): \iota$$

$$\iota: \mathbf{Fun}(\mathcal{C}, \mathcal{D}) \rightleftarrows \mathbf{WSpan}^r(\mathcal{C}, \mathcal{D}): \mathbf{Ran}$$

where the functors ι denote the obvious inclusions (note the need for the $(-)^{op}$ in the second adjunction) and $\mathbf{Lan}/\mathbf{Ran}$ denote the left/right Kan extension functors.

We will mainly be interested in the span categories $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V}) \simeq \mathbf{WSpan}^r(\Sigma_G, \mathcal{V}^{op})$.

Notation 4.39. Given a functor $\pi: A \rightarrow \Sigma_G$, we let $\Omega_{G,n}^{(A)}$ denote the pullback (in \mathbf{Cat})

$$\begin{array}{ccc} \Omega_{G,n}^{(A)} & \xrightarrow{V_{G,n}^{(A)}} & \mathbf{F} \wr A \\ \downarrow & & \downarrow \\ \Omega_{G,n} & \xrightarrow{V_{G,n}} & \mathbf{F} \wr \Sigma_G \end{array}$$

Explicitly, the objects of $\Omega_{G,n}^{(A)}$ are pairs

$$(T_0 \rightarrow \cdots \rightarrow T_n, (a_{e^\dagger \leq e})_{(e^\dagger \leq e) \in V_G(T_n)}) \quad (4.40)$$

OMEGAGNA EQ

such that $\pi(a_{e^\dagger \leq e}) = T_{n, e^\dagger \leq e}$.

Remark 4.41. Our primary interest here will be in the $\Omega_{G,0}^{(A)}$ construction. Importantly, the composite maps $\Omega_{G,0}^{(A)} \rightarrow \Omega_{G,0} \rightarrow \Sigma_G$ allow us to iterate the $\Omega_{G,0}^{(-)}$ construction. In practice, the role of higher strings $\Omega_{G,n}^{(A)}$ will then be to provide more convenient models for iterated $\Omega_{G,0}^{(-)}$ constructions.

Indeed, the content of Proposition 4.27 is then that there are compatible identifications $\Omega_{G,0}^{(\Omega_{G,n})} \simeq \Omega_{G,n+1}$ which identify $V_G^{(\Omega_{G,n})}$ with V_G .

Moreover, since all squares in the diagram

$$\begin{array}{ccccccc} \Omega_{G,n+1}^{(A)} & \xrightarrow{V_G^{(A)}} & \mathbf{F} \wr \Omega_{G,n}^{(A)} & \xrightarrow{\mathbf{F} \wr V_{G,n}^{(A)}} & \mathbf{F} \wr \mathbf{F} \wr A & \xrightarrow{\sigma^0} & \mathbf{F} \wr A \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_{G,n+1} & \xrightarrow{V_G} & \mathbf{F} \wr \Omega_{G,n} & \xrightarrow{\mathbf{F} \wr V_{G,n}} & \mathbf{F} \wr \mathbf{F} \wr \Sigma_G & \xrightarrow{\sigma^0} & \mathbf{F} \wr \Sigma_G \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_{G,0} & \longrightarrow & \mathbf{F} \wr \Sigma_G & & & & \end{array} \quad (4.42)$$

ALLSQUARES EQ

are pullback squares (the top center square is so by induction, the top right square by direct verification, the total top square by definition of $\Omega_{G,n+1}^{(A)}$ and the bottom left square by Proposition 4.27), we likewise obtain identifications $\Omega_G^{(\Omega_{G,n}^{(A)})} \simeq \Omega_{G,n+1}^{(A)}$.

Proposition 4.43. *For any $A \rightarrow \Sigma_G$ there are functors $d_0^{(A)}: \Omega_{G,1}^{(A)} \rightarrow \Omega_G^{(A)}$ and natural isomorphisms*

$$\begin{array}{ccccc} \Omega_{G,1}^{(A)} & \xrightarrow{V_G} & \mathbf{F} \wr \Omega_G^{(A)} & \xrightarrow{\mathbf{F} \wr V_G} & \mathbf{F} \wr \mathbf{F} \wr A \\ d_0^{(A)} \downarrow & & \nearrow \pi^{(A)} & & \downarrow \sigma^0 \\ \Omega_G^{(A)} & \xrightarrow{V_G} & \mathbf{F} \wr A, & & \end{array} \quad (4.44) \quad \text{SHUFFLEPERMA EQ}$$

both natural in $A \rightarrow \Sigma$. Here naturality of $\pi^{(-)}$ means that for a functor $H: A \rightarrow B$ with corresponding diagram

$$\begin{array}{ccccccc} \Omega_{G,1}^{(A)} & \xrightarrow{V_G^{(A)}} & \mathbf{F} \wr \Omega_{G,0}^{(A)} & \xrightarrow{\mathbf{F} \wr V_G^{(A)}} & \mathbf{F} \wr \mathbf{F} \wr A & \xrightarrow{\sigma^0} & \mathbf{F} \wr A \\ \Omega_{G,1}^{(H)} \downarrow d_0^{(A)} \searrow & & \downarrow & \nearrow \pi^{(A)} & \downarrow & & \downarrow \mathbf{F} \wr H \\ \Omega_{G,1}^{(B)} & \xrightarrow{V_G^{(B)}} & \mathbf{F} \wr \Omega_{G,0}^{(B)} & \xrightarrow{\mathbf{F} \wr V_G^{(B)}} & \mathbf{F} \wr \mathbf{F} \wr B & \xrightarrow{\sigma^0} & \mathbf{F} \wr B \\ d_0^{(B)} \searrow & & \downarrow & \nearrow \pi^{(B)} & \downarrow & & \downarrow \mathbf{F} \wr H \\ \Omega_{G,1}^{(B)} & \xrightarrow{V_G^{(B)}} & \mathbf{F} \wr \Omega_{G,0}^{(B)} & \xrightarrow{\mathbf{F} \wr V_G^{(B)}} & \mathbf{F} \wr \mathbf{F} \wr B & \xrightarrow{\sigma^0} & \mathbf{F} \wr B \end{array} \quad (4.45) \quad \text{PICUBOIDAB EQ}$$

one has an equality

$$(\mathbf{F} \wr H) \pi^{(A)} = \pi^{(B)} \Omega_{G,1}^{(H)}$$

(i.e. the two natural isomorphisms between the two distinct functors $\Omega_{G,1}^{(A)} \Rightarrow \mathbf{F} \wr B$ coincide).

Proof. Informally, using the object description in (4.40), $d_0^{(A)}$ is simply given by the formula

$$d_0^{(A)}(T_0 \rightarrow T_1, (a_{e \dagger \leq e})(e \dagger \leq e) \in V_G(T_1)) = (T_1, (a_{e \dagger \leq e})(e \dagger \leq e) \in V_G(T_1)), \quad (4.46) \quad \text{GENDO EQ}$$

though one must note that since in (4.40) the order in $V_G(T_1)$ is induced lexicographically from the string, the two orders for $V_G(T_1)$ in each side of (4.46) do not coincide.

It now follows that the composites $\sigma^0 \circ (\mathbf{F} \wr V_G^{(A)}) \circ V_G^{(A)}$ and $V_G^{(A)} \circ d_0^{(A)}$ differ by the natural automorphism $\pi^{(A)}$ given by the tuple permutations interchanging the two orders in $V_G(T_1)$ for each $T_0 \rightarrow T_1$.

The commutativity of (4.45) is clear. \square

Definition 4.47. Suppose \mathcal{V} has finite products.

We define an endofunctor N of $\mathbf{Wspan}^r(\Sigma_G, \mathcal{V}^{op})$ by letting $N(\Sigma_G \leftarrow A \rightarrow \mathcal{V}^{op})$ be the span $\Sigma_G \leftarrow \Omega_G^{(A)} \rightarrow \mathcal{V}^{op}$ given composition along the diagram

$$\begin{array}{ccccc} \Omega_{G,0}^{(A)} & \longrightarrow & \mathbf{F} \wr A & \longrightarrow & \mathbf{F} \wr \mathcal{V}^{op} \xrightarrow{\Pi^{op}} \mathcal{V}^{op} \\ \downarrow & & \downarrow & & \\ \Omega_{G,0} & \longrightarrow & \mathbf{F} \wr \Sigma_G & & \\ \downarrow & & & & \\ \Sigma_G & & & & \end{array}$$

and defined on maps of spans in the obvious way.

One has a multiplication $\mu: N \circ N \Rightarrow N$ given by the natural isomorphisms

$$\begin{array}{ccccccc}
 \Sigma & \longleftarrow & \Omega_{G,1}^{(A)} & \xrightarrow{V_G} & F \wr \Omega_G^{(A)} & \xrightarrow{F \wr V_G} & F \wr F \wr A \longrightarrow F \wr F \wr \mathcal{V}^{op} \longrightarrow F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 \parallel & & d_0^{(A)} \downarrow & \nearrow \pi^{(A)} & \downarrow \sigma^0 & \downarrow \sigma^0 & \nearrow \alpha \\
 \Sigma & \longleftarrow & \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op}
 \end{array}
 \quad (4.48)$$

MULTDEFSPAN EQ

where α is an associativity isomorphism for the product Π . We note that naturality of μ follows from the commutativity of (H.45).

Lastly, there is a unit $\eta: id \Rightarrow N$ given by the strictly commutative diagrams

$$\begin{array}{ccccccc}
 \Sigma & \longleftarrow & A & \xlongequal{\quad} & A & \longrightarrow & \mathcal{V}^{op} \xlongequal{\quad} \mathcal{V}^{op} \\
 \parallel & & s_{-1}^{(A)} \downarrow & & \downarrow & & \downarrow \\
 \Sigma & \longleftarrow & \Omega_G^{(A)} & \xrightarrow{V_G} & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op}.
 \end{array}
 \quad (4.49)$$

UNITSPAN EQ

MONSPAN PROP

Proposition 4.50. (N, μ, η) form a monad on $\mathcal{Wspan}^r(\Sigma_G, \mathcal{V}^{op})$.

Proof. The natural transformation component of $\mu \circ (N\mu)$ is given by the composite diagram

$$\begin{array}{ccccccccccc}
 \Omega_{G,2}^{(A)} & \rightarrow & F \wr \Omega_{G,1}^{(A)} & \rightarrow & F^{i2} \wr \Omega_G^{(A)} & \rightarrow & F^{i3} \wr A & \rightarrow & F^{i3} \wr \mathcal{V}^{op} & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_1^{(A)} \downarrow & & \downarrow & & \nearrow F \wr \pi^{(A)} & & \downarrow \sigma^1 & & \downarrow \sigma^1 & & \nearrow F \wr \alpha & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \rightarrow & F \wr \Omega_G^{(A)} & \rightarrow & F^{i2} \wr A & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & & \nearrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel \\
 \Omega_G^{(A)} & \rightarrow & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op}
 \end{array}
 \quad (4.51)$$

ASSOCSPAN1 EQ

whereas the natural transformation component of $\mu \circ (\mu N)$ is given by

$$\begin{array}{ccccccccccc}
 \Omega_{G,2}^{(A)} & \rightarrow & F \wr \Omega_{G,1}^{(A)} & \rightarrow & F^{i2} \wr \Omega_G^{(A)} & \rightarrow & F^{i3} \wr A & \rightarrow & F^{i3} \wr \mathcal{V}^{op} & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & & \nearrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \rightarrow & F \wr \Omega_G^{(A)} & \rightarrow & F^{i2} \wr A & \rightarrow & F^{i2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & & \nearrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel \\
 \Omega_G^{(A)} & \rightarrow & F \wr A & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op}
 \end{array}
 \quad (4.52)$$

ASSOCSPAN2 EQ

That the rightmost sections of (4.51) and (4.52) coincide follows from compatibility of the associativity isomorphisms for Π^{op} .

For the leftmost sections, note first that, in either diagram, the top right and bottom left paths $\Omega_{G,2}^{(A)} \rightarrow F \wr A$ differ only by the induced order on $V_G(T_2)$ for each string $T_0 \rightarrow T_1 \rightarrow T_2$. More explicitly, the top right paths use the order induced lexicographically from the string $T_0 \rightarrow T_1 \rightarrow T_2$ while the bottom left paths use the order induced exclusively by T_2 . The two left sections then coincide since are both given by the permutation interchanging these orders, the only difference being that the intermediate stage of (4.51) uses the order induced lexicographically from $T_0 \rightarrow T_2$ while (4.52) uses the order induced lexicographically from $T_1 \rightarrow T_2$.

As for unit conditions, $\mu \circ (N\eta)$ is represented by

$$\begin{array}{ccccccc}
 \Omega_G^{(A)} & \longrightarrow & F \wr A & = & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} = F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} \\
 s_0^{(A)} \downarrow & & \downarrow & & \downarrow \delta^1 & & \downarrow \delta^1 & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \longrightarrow & F \wr \Omega_G^{(A)} & \longrightarrow & F^{\wr 2} \wr A & \longrightarrow & F^{\wr 2} \wr \mathcal{V}^{op} & \longrightarrow & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & \nearrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel & \\
 \Omega_G^{(A)} & \longrightarrow & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op}
 \end{array} \quad (4.53) \quad \text{UNITSPAN1 EQ}$$

while $\mu \circ (\eta N)$ is represented by

$$\begin{array}{ccccccc}
 \Omega_G^{(A)} & = & \Omega_G^{(A)} & \longrightarrow & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} \longrightarrow \mathcal{V}^{op} = \mathcal{V}^{op} \\
 s_{-1}^{(A)} \downarrow & & \downarrow & & \downarrow \delta^0 & & \downarrow \delta^0 & & \parallel & & \parallel \\
 \Omega_{G,1}^{(A)} & \longrightarrow & F \wr \Omega_G^{(A)} & \longrightarrow & F \wr F \wr A & \longrightarrow & F \wr F \wr \mathcal{V}^{op} & \longrightarrow & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\
 d_0^{(A)} \downarrow & \nearrow \pi^{(A)} & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \nearrow \alpha & & \parallel & \\
 \Omega_G^{(A)} & \longrightarrow & F \wr A & \longrightarrow & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op}
 \end{array} \quad (4.54) \quad \text{UNITSPAN2 EQ}$$

It is straightforward to check that the composites of the left and right sections of both (4.53) and (4.54) are strictly commutative diagrams, and thus that (4.53) and (4.54) coincide. \square UNITSPAN1 EQ
UNITSPAN2 EQ

4.5 The free genuine operad monad

Recalling that $\mathbf{Wspan}^r(\Sigma_G, \mathcal{V}^{op}) \simeq \mathbf{Wspan}^l(\Sigma_G^{op}, \mathcal{V})$, Proposition 4.50 and Remark 4.38 give an adjunction MONSPAN PROP
RANLANADJ REM

$$\mathbf{Lan}: \mathbf{Wspan}^l(\Sigma_G^{op}, \mathcal{V}) \rightleftarrows \mathbf{Fun}(\Sigma_G^{op}, \mathcal{V}): \iota \quad (4.55) \quad \text{LANIOTAADJ EQ}$$

together with a monad N in the leftmost category $\mathbf{Wspan}^l(\Sigma_G^{op}, \mathcal{V})$. We now turn to showing that, under reasonable hypothesis on \mathcal{V} , the composite $\mathbf{Lan} \circ N \circ \iota$ inherits a monad structure from N . The key will be to show that under such conditions the map $\mathbf{Lan} \circ N \Rightarrow \mathbf{Lan} \circ N \circ \iota \circ \mathbf{Lan}$ is a natural isomorphism.

Recall that following Convention 3.18 our model for \mathbf{O}_G consists of totally ordered sets. One therefore has *root functors* PLANARCONV CON

$$\Omega_G^q \xrightarrow{r} \mathbf{O}_G, \quad \Sigma_G \xrightarrow{r} \mathbf{O}_G$$

sending each planar G -tree to its ordered orbital G -set of roots.

Root functors are compatible with the leaf-root functor and the inclusion, i.e. the following commute.

$$\begin{array}{ccc}
 \Omega_G^q & \xrightarrow{lr} & \Sigma_G \\
 \searrow r & & \downarrow r \\
 & & \mathbf{O}_G
 \end{array}
 \quad
 \begin{array}{ccc}
 \Sigma_G & \hookrightarrow & \Omega_G^q \\
 \searrow r & & \downarrow r \\
 & & \mathbf{O}_G
 \end{array} \quad (4.56) \quad \text{ROOTLEAFTROOTCOM EQ}$$

Moreover, the diagrams (4.56) possess some extra structure we will need to make use of. Indeed, both functors are split Grothendieck fibrations: given a map $\varphi: A \rightarrow B$ in \mathbf{O}_G and G -tree T such that $r(T) = B$ we can build a cartesian arrow $\varphi^*(T) \rightarrow T$ by letting $\varphi^*(T)$ to be the pullback G -tree together with the planar structure on roots given by A and on non-equivariant nodes given by their image via $\varphi^*(T) \rightarrow T$. ROOTLEAFTROOTCOM EQ

It now follows that (4.56) are diagrams of split Grothendieck fibrations.

Definition 4.57. A split Grothendieck fibration $A \xrightarrow{r} \mathbf{O}_G$ is called a *root fibration* and a split Grothendieck fibration diagram

$$\begin{array}{ccc} A & \longrightarrow & B \\ & \searrow r & \downarrow r \\ & & \mathbf{O}_G \end{array}$$

is called a *root fibration functor*.

The relevance of root fibrations is given by the following couple of lemmas.

Lemma 4.58. *If $A \rightarrow \Sigma_G$ is a root fibration functor then so is $\Omega_G^{(A)} \rightarrow \Omega_G$, naturally in A .*

Proof. We consider the pullback diagram below.

$$\begin{array}{ccc} \Omega_{G,0}^{(A)} & \xrightarrow{V_G^{(A)}} & \mathbf{F} \wr A \\ \downarrow & & \downarrow \\ \Omega_{G,0} & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G \end{array} \quad (4.59) \quad \text{ROOTIMPLIESROOT EQ}$$

The hypothesis that $A \rightarrow \Sigma_G$ is root fibration implies that the rightmost map in (4.59) is a map of split Grothendieck fibrations over $\mathbf{F} \wr \mathbf{O}_G$.

Since the map V_G sends the chosen cartesian arrows in $\Omega_{G,0}$ (over \mathbf{O}_G) to chosen cartesian arrows of $\mathbf{F} \wr \Sigma_G$ (over $\mathbf{F} \wr \mathbf{O}_G$), the result follows. \square

Example 4.60. Let $G = \mathbb{Z}/8$. The following exemplifies a pull back along the twist map $\varphi: G/2G \rightarrow G/2G$ (i.e., accounting for order, φ is the permutation (12)), with the topmost representation of φ^*T maintaining the chosen generators for each edge orbit from T and the bottom representation choosing instead the generators to be minimal with regard to the planar structure.

$$\begin{array}{ccc} \begin{array}{c} \begin{array}{ccccc} a & a+4 & a+2 & a+6 & \\ \diagdown & & \diagup & & \diagdown \\ & b & & b+2 & \\ & \diagup & & \diagdown & \\ & c & & c+1 & \end{array} & \begin{array}{ccccc} a+1 & a+5 & a+3 & a+7 & \\ \diagdown & & \diagup & & \diagdown \\ & b+1 & & b+3 & \\ & \diagup & & \diagdown & \\ & c+1 & & c+1 & \end{array} \end{array} & \xrightarrow{\varphi^*T} & \begin{array}{ccccc} a & a+4 & a+2 & a+6 & \\ \diagdown & & \diagup & & \diagdown \\ & b & & b+2 & \\ & \diagup & & \diagdown & \\ & c & & c & \end{array} \\ \begin{array}{c} \begin{array}{ccccc} a & a+4 & a+2 & a+6 & \\ \diagdown & & \diagup & & \diagdown \\ & b & & b+2 & \\ & \diagup & & \diagdown & \\ & c & & c+1 & \end{array} & \begin{array}{ccccc} a+1 & a+5 & a+3 & a+7 & \\ \diagdown & & \diagup & & \diagdown \\ & b+1 & & b+3 & \\ & \diagup & & \diagdown & \\ & c+1 & & c+1 & \end{array} \end{array} & \xrightarrow{\varphi^*T} & \begin{array}{ccccc} d & d+4 & d+2 & d+6 & \\ \diagdown & & \diagup & & \diagdown \\ & e & & e+2 & \\ & \diagup & & \diagdown & \\ & f & & f+1 & \end{array} \end{array} \quad (4.61)$$

We note that $(\varphi^*(T))_{v_{Ge}} = \psi^*(T_{v_{Gb}})$ for ψ the permutation (13)(24) encoded by the composite identifications $\{1, 2, 3, 4\} \simeq \{e, e+2, e+3, e+1\} \simeq \{b+1, b+3, b, b+2\} \simeq \{3, 4, 1, 2\}$.

Lemma 4.62. *Suppose that \mathcal{V} is complete and that $A \rightarrow \Sigma_G$ is a root fibration. If the rightmost triangle in*

$$\begin{array}{ccccc} \Omega_{G,0}^{(A)} & \xrightarrow{V_G^{(A)}} & \mathbf{F} \wr A & \longrightarrow & \mathcal{V} \\ \downarrow & & \downarrow & \nearrow & \nearrow \\ \Omega_{G,0} & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G & & \end{array} \quad (4.63)$$

is a right Kan extension diagram then so is the composite diagram.

Proof. Unpacking definitions using the pointwise formula for right Kan extensions ($\frac{\text{McL}}{2}$, X.3.1]), it suffices to check that for each $T \in \Omega_{G,0}$ the functor

$$T \downarrow \Omega_{G,0}^{(A)} \rightarrow V_G(T) \downarrow F \wr A \quad (4.64)$$

LANPULLCOMA EQ

is initial. In the course of the proof of Lemma $\frac{\text{FINWREATPRODLIM LEM}}{4.3}$ it was shown that the subcategory

$$\prod_{v_{Ge} \in V_G(T)} T_{v_{Ge}} \downarrow A$$

is initial in the $V_G(T) \downarrow F \wr A$.

On the other hand, since $\Omega_G^{(A)} \rightarrow \Omega_G$ is a root fibration functor, $T \downarrow \Omega_G^{(A)}$ has an initial subcategory $T \downarrow_{r,\simeq} \Omega_G^{(A)}$ with objects $(S \in \Omega_G^{(A)}, T \rightarrow u(S))$ such that $T \rightarrow u(S)$ is a quotient map that induces an ordered isomorphism on roots. Note that this can be restated as saying that $T \rightarrow u(S)$ is an isomorphism preserving the order of the roots.

The result now follows from the natural isomorphism

$$T \downarrow_{r,\simeq} \Omega_G^{(A)} \simeq \prod_{v_{Ge} \in V_G(T)} T_{v_{Ge}} \downarrow_{r,\simeq} A. \quad (4.65)$$

TDOWNISOA EQ

To see this, we focus first on the case $A = \Sigma_G$. In that case, the left hand side of ($\frac{\text{TDOWNISOA EQ}}{4.65}$) encodes replanarizations of T that preserve the root order. On the other hand, the right hand side encodes replanarizations of all the G -vertices that preserve the order of their roots, or, equivalently, replanarizations of the non-equivariant vertices of T . That these are equivalent is the content of Proposition $\frac{\text{PLANARIZATIONCHAR PROP}}{3.13}$.

Note that $(T \rightarrow S) \in (T \downarrow_{r,\simeq} \Omega_G)$ is then encoded by a tuple $(T_{v_{Ge}} \rightarrow \varphi_{v_{Ge}}^* S_{v_{Ge}})_{v_{Ge} \in V_G(T)}$ where the pullbacks $\varphi_{v_{Ge}}^*$ are needed to correct the root order.

The case of general A follows likewise, using the corresponding pullbacks $\varphi_{v_{Ge}}^*$.

Note: an addendum is needed to show that ($\frac{\text{TDOWNISOA EQ}}{4.65}$) suffices, since $T \downarrow_{r,\simeq} \Omega_G^{(A)}$ is not sent directly to $\prod_{v_{Ge} \in V_G(T)} T_{v_{Ge}} \downarrow_{r,\simeq} A$ \square

Lemma $\frac{\text{ROOTFIBPULL LEM}}{4.58}$ can be interpreted as saying that, if one defines a category $\text{Wspan}_r^l(\Sigma_G^{op}, \mathcal{V})$ of *rooted spans*

$$\Sigma_G^{op} \leftarrow A^{op} \rightarrow \mathcal{V}$$

where $A \rightarrow \Sigma_G$ is a root fibration functor, the monad N built in Proposition $\frac{\text{MONSPAN PROP}}{4.50}$ lifts to a monad N_r in $\text{Wspan}_r^l(\Sigma_G^{op}, \mathcal{V})$, and likewise for the adjunction ($\frac{\text{LANIOTADJ EQ}}{4.55}$).

Corollary 4.66. *Suppose that finite products in \mathcal{V} commute with colimits in each variable. The functors*

$$\text{Lan} \circ N_r \Rightarrow \text{Lan} \circ N_r \circ \iota \circ \text{Lan}, \quad \text{Lan} \circ \iota \Rightarrow id$$

are natural isomorphisms.

Proof. This follows by combining Lemma $\frac{\text{LANPULLCOMA LEM}}{4.62}$ with Lemma $\frac{\text{FINWREATPRODLIM LEM}}{4.3}$. \square

Definition 4.67. The *genuine equivariant operad monad* is the monad \mathbb{F}_G on $\text{Fun}(\Sigma_G^{op}, \mathcal{V})$ given by

$$\mathbb{F}_G = \text{Lan} \circ N_r \circ \iota$$

and with multiplication and unit given by the composites

$$\text{Lan} \circ N_r \circ \iota \circ \text{Lan} \circ N_r \circ \iota \xleftarrow{\simeq} \text{Lan} \circ N_r \circ N_r \circ \iota \Rightarrow \text{Lan} \circ N_r \circ \iota$$

$$id \xleftarrow{\simeq} \text{Lan} \circ \iota \Rightarrow \text{Lan} \circ N_r \circ \iota.$$

Remark 4.68. The functor $\text{Lan} \circ N_r \circ \iota$ is isomorphic to $\text{Lan} \circ N \circ \iota$, and this isomorphism is compatible with the multiplication and unit in Definition 4.67, and we will henceforth simply write N rather than N_r . $\frac{\text{THEMONAD DEF}}{4.67}$

From this point of view, the role of root fibrations is to guarantee that $\text{Lan} \circ N \circ \iota$ is indeed a monad, but unnecessary to describe the monad structure itself.

THEMONAD DEF

Remark 4.69. Since a map

$$\mathbb{F}_G X = \text{Lan} \circ N_r \circ \iota X \rightarrow X$$

is adjoint to a map

$$N_r \circ \iota X \rightarrow \iota X$$

one easily verifies that X is a genuine equivariant operad, i.e. a \mathbb{F}_G -algebra, iff ιX is a N -algebra. Moreover, the bar resolution

$$\mathbb{F}_G^{\bullet+1} X$$

is isomorphic to

$$\text{Lan}(N^{\bullet+1} \iota X).$$

5 Free extensions

Our overall goal in this section will be to produce a description of free genuine operad pushouts, i.e. pushouts of the form

$$\begin{array}{ccc} \mathbb{F}_G A & \longrightarrow & X \\ \downarrow & & \downarrow \\ \mathbb{F}_G B & \longrightarrow & Y \end{array}$$

in the category Op_G of genuine equivariant operads.

5.1 Extensions over general monads

Any monad T on \mathcal{C} one obtains induced monads $T^{\times l}$ on $\mathcal{C}^{\times l}$, and we will make use of several standard relations between these. In particular, any map $\alpha: \underline{l} \rightarrow \underline{m}$ induces a forgetful functor such that for the forgetful functor $\alpha^*: \mathcal{C}^{\times l} \rightarrow \mathcal{C}^{\times n}$ one has $T^{\times l} \alpha^* \simeq \alpha^* T^{\times m}$.

Indeed, we will need to make use of a slightly more general setup. Letting I denote the identity monad on \mathcal{C} , and $K \subset \underline{m}$ be a subset, there is a monad $T^{\times K} \times I^{\times(\underline{m}-K)}$ on $\mathcal{C}^{\times m}$, which we abusively denote simply as $T^{\times K}$. Identities then determine maps of monads $T^J \rightarrow T^{\times K}$ whenever $J \subset K$ and, moreover, there are identifications $T^{\times \alpha^{-1}(K)} \alpha^* \simeq \alpha^* T^{\times K}$. One then has the following.

Proposition 5.1. *The functor*

$$T^{\times \alpha^{-1}(K)} \Rightarrow \alpha^* T^{\times K} \alpha_! \tag{5.2}$$

adjoint to the identification $T^{\times \alpha^{-1}(K)} \alpha^ \simeq \alpha^* T^{\times K}$ is a map of monads on $\mathcal{C}^{\times n}$.*

Proof. We first note that there are identifications of functors $(FG)^{\times K} \simeq F^{\times K} G^{\times K}$ which are compatible with the identifications $F^{\times \alpha^{-1}(K)} \alpha^* \simeq \alpha^* F^{\times K}$ in the sense that the identification $(FG)^{\times \alpha^{-1}(K)} \circ \alpha^* \simeq \alpha^* (FG)^{\times K}$ matches the composite identification $F^{\times \alpha^{-1}(K)} G^{\times \alpha^{-1}(K)} \alpha^* \simeq F^{\times \alpha^{-1}(K)} \alpha^* G^{\times K} \simeq \alpha^* F^{\times K} G^{\times K}$.

Letting η, ϵ denote the unit and counit for the $(\alpha_!, \alpha^*)$ adjunction, (5.2) is then the composite

$$T^{\times \alpha^{-1}(K)} \xrightarrow{\eta} T^{\times \alpha^{-1}(K)} \alpha^* \alpha_! \simeq \alpha^* T^{\times K} \alpha_!.$$

That this is a monad map is the condition that the following multiplication and unit diagrams commute.

$$\begin{array}{ccc} T^{\times \alpha^{-1}(K)} \circ T^{\times \alpha^{-1}(K)} & \longrightarrow & \alpha^* T^{\times K} \alpha_! \circ \alpha^* T^{\times K} \alpha_! \\ \downarrow & & \downarrow \\ T^{\times \alpha^{-1}(K)} & \longrightarrow & \alpha^* T^{\times K} \alpha_! \end{array} \quad \begin{array}{ccc} I^{\times n} & & \\ \downarrow & \searrow & \\ T^{\times \alpha^{-1}(K)} & \longrightarrow & \alpha^* T^{\times K} \alpha_! \end{array}$$

We argue only the case of the leftmost multiplication diagram, with commutativity of the unit diagram following by a similar but simpler argument. Since the precomposition $(-) \circ \alpha^*$ is the left adjoint to the precomposition $(-) \circ \alpha_!$ this follows from the following diagram.

$$\begin{array}{ccccccc}
T^{\times \alpha^{-1}(K)} T^{\times \alpha^{-1}(K)} \alpha^* & \xrightarrow{\simeq} & T^{\times \alpha^{-1}(K)} \alpha^* T^{\times K} & \xrightarrow{\eta} & T^{\times \alpha^{-1}(K)} \alpha^* \alpha_! \alpha^* T^{\times K} & \xrightarrow{\simeq} & \alpha^* T^{\times K} \alpha_! \alpha^* T^{\times K} \\
\downarrow & & \searrow & & \downarrow \epsilon & & \downarrow \epsilon \\
& & & & T^{\times \alpha^{-1}(K)} \alpha^* T^{\times K} & \xrightarrow{\simeq} & \alpha^* T^{\times K} T^{\times K} \\
& & & & & & \downarrow \\
T^{\times \alpha^{-1}(K)} \alpha^* & \xrightarrow{\simeq} & & & & & \alpha^* T^{\times K}
\end{array}$$

□

Remark 5.3. Since $T^{\times K} \alpha_!$ is a right $\alpha^* T^{\times K} \alpha_!$ -module, Proposition 5.1 implies that it is also a right $T^{\times \alpha^{-1}(K)}$ -module or, moreover, a right $T^{\times J}$ -module whenever $\alpha(J) \subset K$.

Remark 5.4. Combining the precomposition and postcomposition adjunctions, the identification $T^{\times \alpha^{-1}(K)} \alpha^* \simeq \alpha^* T^{\times K}$ is then adjoint to a functor $\alpha_! T^{\times \alpha^{-1}(K)} \rightarrow T^{\times K} \alpha_!$ which is readily checked to be a map of right $T^{\times \alpha^{-1}(K)}$ -modules.

More generally, for $\alpha(J) \subset K$, the composite $T^{\times J} \alpha^* \rightarrow T^{\times \alpha^{-1}(K)} \alpha^* \simeq \alpha^* T^{\times K}$ is thus adjoint to a map of right $T^{\times J}$ -modules

$$\alpha_! T^{\times J} \rightarrow T^{\times K} \alpha_!. \quad (5.5)$$

RIGHTMODULETMAP EQ

We now unpack the content of (5.5) when $\alpha: \underline{l} \rightarrow *$ is the unique map to the singleton $* = \underline{1}$. In this case we can instead write $\alpha_! = \coprod$, $\alpha^* = \Delta$, and we thus have commutative diagrams

$$\begin{array}{ccc}
\coprod_J TTA_j \sqcup \coprod_{\underline{n}-J} A_j & \longrightarrow & T(\coprod_J TA_j \sqcup \coprod_{\underline{n}-J} A_j) \\
\downarrow & & \downarrow \\
\coprod_J TA_j \sqcup \coprod_{\underline{n}-J} A_j & \longrightarrow & T(\coprod_J A_j \sqcup \coprod_{\underline{n}-J} A_j)
\end{array} \quad (5.6)$$

RIGHTMODULETMAPAUX EQ

where the vertical maps come from the right $T^{\times J}$ -module structure. Writing \sqcup^a for the coproduct of T -algebras and recalling the canonical identifications $\coprod_K^a(TA_k) \simeq T(\coprod_K A_k)$, (5.6) in fact shows that the right $T^{\times J}$ -module structure on $T \circ \coprod$ in fact codifies the multiplication maps

$$\coprod_J^a TTA_j \sqcup^a \coprod_{\underline{l}-J}^a TA_j \rightarrow \coprod_J^a TA_j \sqcup^a \coprod_{\underline{l}-J}^a TA_j.$$

5.2 Labeled planar strings


We now translate the results in the previous section to the context of the monad N on $\mathbf{WSpan}^l(\Sigma^{op}, \mathcal{V})$. In analogy to the planar string models $\Omega_{G,n}^{(A)}$ for iterations $N^{\circ n+1}$ of the monad N , we will find it convenient to build similar string models $\Omega_{G,n}^{(\underline{A}_J)}$ for $N \circ \coprod \circ (N^{\times J})^{\circ n}$.

Definition 5.7. A l -node labeled G -tree (or just l -labeled G -tree) G -tree is a pair $(T, V_G(T) \rightarrow \{1, \dots, l\})$ with $T \in \Omega_G$, which we think of as a G -tree together with G -vertices labels in $1, \dots, l$.

Further, a tall map $\varphi: T \rightarrow S$ between l -labeled trees is called a *label map* if for each G -vertex v_{Ge} of T with label j , the vertices of the subtree $S_{v_{Ge}}$ are all labeled by j .

Lastly, given a subset $J \subset \underline{l}$, a planar label map $\varphi: T \rightarrow S$ is said to be J -inert if for every G -vertex v_{Ge} of T with label $j \in J$ it is $S_{v_{Ge}} = T_{v_{Ge}}$.

Example 5.8. Consider the 2-labeled trees below (for $G = *$ the trivial group), with black nodes (\bullet) denoting labels by the number 1 and white nodes (\circ) labels by the number 2. The planar map φ (sending $a_i \mapsto a$, $b \mapsto b$, $c \mapsto c$, $d \mapsto d$, $e \mapsto e$) is a label map which is $\{1\}$ -inert.



SUBSDATUMTREESLAB EQ

Definition 5.10. Let $0 \leq s \leq n$ and $J \subset \underline{l}$ be a subset.

We define $\Omega_{G,n,s}^J$ to have as objects n -planar strings

$$T_0 \xrightarrow{f_1} T_1 \xrightarrow{f_2} \dots \xrightarrow{f_s} T_s \xrightarrow{f_{s+1}} T_{s+1} \xrightarrow{f_{s+2}} \dots \xrightarrow{f_n} T_n \quad (5.11)$$

NSTRINGLAB EQ

together with l -labelings of T_s, T_{s+1}, \dots, T_n such that the $f_r, r > s$ are $(\underline{l} - J)$ -inert label maps.

Arrows in $\Omega_{G,n,s}^J$ are quotients of strings $(q_r: T_r \rightarrow T_r')$ such that $q_r, r \leq s$ are label maps.

Informally, $\Omega_{G,n,s}^J$ consists of n -strings such that trees and maps after T_s are l -labeled.

Remark 5.12. Our main case of interest will that of $s = 0$, in which case we abbreviate $\Omega_{G,n}^J = \Omega_{G,n,0}^J$. Indeed, such strings will suffice to build models for $N \circ \coprod (N^{\times J})^{\text{on}}$

However, to unpack the right $N^{\times J}$ -module structure as in Remark 5.3 one further needs to encode composites $NN \circ \coprod (N^{\times J})^{\text{on}-1}$, a role played by strings $\Omega_{G,n,1}^J$.

Notation 5.13. We will further write

$$\Omega_{G,n,-1}^J = \coprod_J \Omega_{G,n} \sqcup \coprod_{\underline{l}-J} \Sigma_G, \quad \Omega_{G,n,n+1}^J = \Omega_{G,n} \quad (5.14)$$

OMEGANMINUSONE EQ

To justify this convention, we note that a string as in (5.11) can be extended by prepending to it the map $\text{lr}(T_0) = T_{-1} \xrightarrow{f_0} T_0$. If one then attempts to define $\Omega_{G,n,-1}^J$ by insisting that T_{-1} also be labeled, it follows that all node labels in each string must coincide, resulting in the coproduct decomposition in (5.14).

There are a number of obvious functors relating the $\Omega_{G,n,s}^J$ categories, which we now make explicit. Given $s \leq s'$ or $J \subset J'$ there are forgetful functors

$$\Omega_{G,n,s}^J \rightarrow \Omega_{G,n,s'}^J \quad \Omega_{G,n,s}^J \rightarrow \Omega_{G,n,s}^{J'} \quad (5.15)$$

NKNFGT EQ

The simplicial operators in Notation 4.24 generalize to operators (where $0 \leq i \leq n$, $-1 \leq j \leq n$)

$$\begin{aligned} d_i: \Omega_{G,n,s}^J &\rightarrow \Omega_{G,n-1,s-1}^J & i < s & & s_j: \Omega_{G,n,s}^J &\rightarrow \Omega_{G,n+1,s+1}^J & j < s \\ d_i: \Omega_{G,n,s}^J &\rightarrow \Omega_{G,n-1,s}^J & s \leq i & & s_j: \Omega_{G,n,s}^J &\rightarrow \Omega_{G,n+1,s}^J & s \leq j \end{aligned}$$

which are compatible with the forgetful functors in the obvious way.

Remark 5.16. For $J \subset J'$ the forgetful functor in (5.15) is a fully faithful inclusion. However, and somewhat subtly, this is not the case for the $s \leq s'$ forgetful functors. Indeed, regarding $T \rightarrow U$ in Examples 5.8 as an object in $\Omega_{*,n,0}^2$, changing the label of the $a_1 \leq a_2$ vertex of T from a \circ -label to a \bullet -label yields an alternate object $\bar{T} \rightarrow U$ of $\Omega_{*,n,0}^2$ forgetting to the same object of $\Omega_{*,n,1}^2$, yet $T \rightarrow U$ and $\bar{T} \rightarrow U$ are not isomorphic.

We note that this is a consequence of the fact that substitution data can replace unary nodes by stumps, which have no nodes.

Generalizing Notation ~~INDVNG NOT~~ 4.31 there is a commutative diagram

$$\begin{array}{ccc} \Omega_{G,n,s}^J & \xrightarrow{V_{G,n}} & \mathbf{F} \wr \Sigma_G^{ul} \\ \downarrow & & \downarrow \\ \Omega_{G,n} & \xrightarrow{V_{G,n}} & \mathbf{F} \wr \Sigma_G \end{array}$$

where for a labeled string it is $V_G(T_0 \rightarrow \dots \rightarrow T_n) = (T_{n,v_{Ge}})_{V_G(T_n)}$, where we regard $T_{n,v_{Ge}} \in \Sigma_G^{ul} \simeq \Omega_{G,-1,-1}^l$ by using the label in 1.1.1. ~~OMEGAGNA NOT~~

We now expand Notation 4.39.

Notation 5.17. Let \underline{A} denote a l -tuple $(\pi_j: A_j \rightarrow \Sigma_G)_l$ of categories over Σ_G . We define $\Omega_{G,n,s}^{(\underline{A}),J}$ by the pullback diagram

$$\begin{array}{ccc} \Omega_{G,n,s}^{(\underline{A}),J} & \xrightarrow{V_{G,n}^{(\underline{A})}} & \mathbf{F} \wr \coprod A_j \\ \downarrow & & \downarrow \\ \Omega_{G,n,s}^J & \xrightarrow{V_{G,n}} & \mathbf{F} \wr \Sigma_G^{ul} \end{array} \quad (5.18) \quad \boxed{\text{LTUPLEAPULL EQ}}$$

Explicitly, an object of $\Omega_{G,n,s}^{(\underline{A}),J}$ consists of a labeled string $T_0 \rightarrow \dots \rightarrow T_n$ as in ~~NSTRINGLAB EQ~~ (5.11) together with a tuple $(a_{v_{Ge}})_{V_G(T_n)}$ such that $a_{v_{Ge}} \in A_j$ if v_{Ge} has label j and $\pi_j(a_{v_{Ge}}) = T_{n,v_{Ge}}$.

The reader may have noticed a certain asymmetry between our definition of the $V_{G,n}$ functors here versus their analogues in §4.3, where they were defined iteratively in terms of simpler functors V_G . This is because of the possibility that $s = -1$, in which case ~~OMEGANMINUSONE EQ~~ (5.14) applies and some caution is needed in that the following result fails.

Proposition 5.19. Suppose $0 \leq s \leq n$. One has a diagram of pullback squares (generalizing ~~ALLSQUARES EQ~~ (4.42))

$$\begin{array}{ccccccc} \Omega_{G,n,s}^{(\underline{A}),J} & \xrightarrow{V_G^{(\underline{A})}} & \mathbf{F} \wr \Omega_{G,n-1,s-1}^{(\underline{A}),J} & \xrightarrow{\mathbf{F} \wr V_{G,n}^{(\underline{A})}} & \mathbf{F} \wr \mathbf{F} \wr \coprod A_j & \xrightarrow{\sigma^0} & \mathbf{F} \wr \coprod A_j \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_{G,n,s}^J & \xrightarrow{V_G} & \mathbf{F} \wr \Omega_{G,n-1,s-1}^J & \xrightarrow{\mathbf{F} \wr V_{G,n}} & \mathbf{F} \wr \mathbf{F} \wr \Sigma_G^{ul} & \xrightarrow{\sigma^0} & \mathbf{F} \wr \Sigma_G^{ul} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega_{G,0} & \xrightarrow{V_G} & \mathbf{F} \wr \Sigma_G & & & & \end{array} \quad (5.20) \quad \boxed{\text{ALLSQUARESJ EQ}}$$

such that the composite of the top squares is ~~LTUPLEAPULL EQ~~ (5.18).

Proof. The V_G functors are defined just as in ~~VGDEF EQ~~ (4.26) via the formula

$$V_G(T_0 \rightarrow T_1 \rightarrow \dots \rightarrow T_n) = (T_{1,v_{Ge}} \rightarrow \dots \rightarrow T_{n,v_{Ge}})_{v_{Ge} \in V_G(T_0)}$$

with the strings $T_{1,v_{Ge}} \rightarrow \dots \rightarrow T_{n,v_{Ge}}$ inheriting the extra structure in the obvious way.

Since the top composite square, top center square and top right square are all pullback squares, it remains only to show that the bottom left square is a pullback. This last claim is simply a variation of Proposition 4.27, and follows from the same proof, since both labels and inertness conditions are inherited when assembling substitution data into trees via ~~SUBDATAUNDERPLAN PROP~~ Proposition 3.42. \square

5.3 Bar constructions on spans

We use the results in the previous sections to obtain a string description of the bar constructions

$$\coprod_J^a N^{\bullet+1} A_j \sqcup^a \coprod_{l=J}^a N A_l.$$

For simplicity, we discuss first the particular case $\coprod^a N^{\bullet+1} A$. Writing the span as $\Sigma_G \leftarrow A \xrightarrow{F} \mathcal{V}$ the identifications $\Omega_{G,0}^{(\Omega_{G,n}^{(A)})} \simeq \Omega_{G,n+1}^{(A)}$ iteratively identify the operator in the bar construction $N^{\bullet+1} A$ as follows.

The top boundaries d_n have natural transformation given by

$$\begin{array}{ccccccc} \Omega_{G,n}^{(A)} & \xrightarrow{V_G^{on}} & F^{in} \wr \Omega_{G,0}^{(A)} & \xrightarrow{F^{in} \wr F_1} & F^{in} \wr \mathcal{V}^{op} & \xrightarrow{\Pi^{on}} & \mathcal{V}^{op} \\ \downarrow & & \downarrow & \swarrow F^{in} \wr m & \parallel & & \parallel \\ \Omega_{G,n-1}^{(A)} & \xrightarrow{V_G^{on}} & F^{in} \wr A & \xrightarrow{F^{in} \wr F} & F^{in} \wr \mathcal{V}^{op} & \xrightarrow{\Pi^{on}} & \mathcal{V}^{op} \end{array} \quad (5.21)$$

where m is the natural transformation component of the multiplication $NA \rightarrow A$, and the remaining differentials d_i for $0 \leq i < n$ are given by

$$\begin{array}{ccccccc} \Omega_{G,n}^{(A)} & \xrightarrow{V_G^{on+1}} & F^{in+1} \wr A & \xrightarrow{F} & F^{in+1} \wr \mathcal{V}^{op} & \xrightarrow{\Pi^{on+1}} & \mathcal{V}^{op} \\ d_i^{(A)} \downarrow & \swarrow \pi_i^{(A)} & \downarrow \sigma^i & & \downarrow \sigma^i & \swarrow \alpha_i & \parallel \\ \Omega_{G,n-1}^{(A)} & \xrightarrow{V_G^{on}} & F^{in} \wr A & \xrightarrow{F} & F^{in} \wr \mathcal{V}^{op} & \xrightarrow{\Pi^{on}} & \mathcal{V}^{op} \end{array} \quad (5.22) \quad \boxed{\text{REMAINDIFF EQ}}$$

where $\pi_i^{(A)}$ interchanges lexicographic orders on the i -th F coordinate of F^{in} and α_i is the natural associativity isomorphism.

Maybe add degeneracies

Similarly, Proposition 5.19 shows that $\Omega_{G,n}^{(A)} \simeq \Omega_{G,0}^{(\Pi \Omega_{G,n-1}^{(A)})}$ so that the top boundaries d_n in the bar construction $N \circ \sqcup \circ (N^{xl})^{on} \underline{A}$ are given by

$$\begin{array}{ccccccc} \Omega_{G,n}^{(\underline{A})} & \xrightarrow{V_G} & F \wr \coprod \Omega_{G,n-1}^{(A_j)} & \xrightarrow{V_G^{on-1}} & F \wr \coprod F^{in-1} \wr \Omega_{G,0}^{(A_j)} & \xrightarrow{F_1} & F \wr \coprod F^{in-1} \wr \mathcal{V}^{op} \xrightarrow{\Pi^{on-1}} F \wr \mathcal{V}^{op} \xrightarrow{\Pi} \mathcal{V}^{op} \\ \downarrow & & \downarrow & \swarrow \underline{m} & \parallel & & \parallel \\ \Omega_{G,n-1}^{(\underline{A})} & \xrightarrow{V_G} & F \wr \coprod \Omega_{G,n-2}^{(A_j)} & \xrightarrow{V_G^{on-1}} & F \wr \coprod F^{in-1} \wr A_j & \xrightarrow{F} & F \wr \coprod F^{in-1} \wr \mathcal{V}^{op} \xrightarrow{\Pi^{on-1}} F \wr \mathcal{V}^{op} \xrightarrow{\Pi} \mathcal{V}^{op} \end{array} \quad (5.23)$$

where \underline{m} stands for the functor induced by the tuple of multiplication maps $m_j: NA_j \rightarrow A_j$, and the other boundaries d_i for $0 \leq i < n$ are given by

$$\begin{array}{ccccccc} \Omega_{G,n}^{(\underline{A})} & \xrightarrow{V_G} & F \wr \coprod \Omega_{G,n-1}^{(A_j)} & \xrightarrow{V_G^{on}} & F \wr \coprod F^{in} \wr A & \xrightarrow{F} & F \wr \coprod F^{in} \wr \mathcal{V}^{op} \xrightarrow{\Pi^{on}} F \wr \mathcal{V}^{op} \xrightarrow{\Pi} \mathcal{V}^{op} \\ d_i^{(\underline{A})} \downarrow & \swarrow \pi_i^{(\underline{A})} & \downarrow \sigma^i & & \downarrow \sigma^i & \swarrow \alpha_i & \parallel \\ \Omega_{G,n-1}^{(\underline{A})} & \xrightarrow{V_G} & F \wr \coprod \Omega_{G,n-2}^{(A_j)} & \xrightarrow{V_G^{on-1}} & F \wr \coprod F^{in-1} \wr A & \xrightarrow{F} & F \wr \coprod F^{in-1} \wr \mathcal{V}^{op} \xrightarrow{\Pi^{on-1}} F \wr \mathcal{V}^{op} \xrightarrow{\Pi} \mathcal{V}^{op} \end{array} \quad (5.24) \quad \boxed{\text{DISJBARDN EQ}} \quad \boxed{\text{REMAINDIFF EQ}}$$

where again $\pi_i^{(\underline{A})}$ interchanges lexicographic orders on the i -th F coordinate and α_i is again an associativity isomorphism. We note that (5.24) follows directly from (5.22) for $0 < i < n$,

but that the case $i = 0$, which uses the $N^{\times l}$ right action on $N \circ \sqcup$ (cf. Remark [TALPHAKMOD REM 5.3](#)), which after unpacked leads to the composite diagram below.

$$\begin{array}{ccccccccccccccc}
\Omega_{G,n}^{(\underline{A})} & \rightarrow & F \wr \coprod \Omega_{G,n-1}^{(A_j)} & \longrightarrow & F \wr \coprod F^{i_{n-1}} A_j & \rightarrow & F \wr \coprod F^{i_{n-1}} \mathcal{V}^{op} & \longrightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \parallel & & \parallel \\
\Omega_{G,n-1}^{(\underline{A})} & \rightarrow & F \wr \Omega_{G,n-1}^{(\underline{A})} & \rightarrow & F^{i_2} \wr \coprod \Omega_{G,n-2}^{(A_j)} & \rightarrow & F^{i_2} \wr \coprod F^{i_{n-2}} A_j & \rightarrow & F^{i_2} \wr \coprod F^{i_{n-2}} \mathcal{V}^{op} & \rightarrow & F^{i_2} \wr \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \rightarrow & \mathcal{V}^{op} \\
d_0^{(\underline{A})} \downarrow & \nearrow \pi_0 & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & & \downarrow \sigma^0 & \nearrow \alpha_0 & \parallel & & \\
\Omega_{G,n-1}^{(\underline{A})} & \longrightarrow & F \wr \coprod \Omega_{G,n-2}^{(A_j)} & \rightarrow & F \wr \coprod F^{i_{n-2}} A_j & \rightarrow & F \wr \coprod F^{i_{n-2}} \mathcal{V}^{op} & \rightarrow & F \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} & & & &
\end{array} \tag{5.25}$$

ASSOCSPANJ1 EQ

Finally, using the inclusions $\Omega_{G,n}^{(\underline{A}),J} \hookrightarrow \Omega_{G,n}^{(\underline{A})}$, one obtains analogous descriptions of the bar constructions $N \circ \sqcup \circ (N^{\times J})^{on} \underline{A}$, depicted below.

$$\begin{array}{ccccccc}
\Omega_{G,n}^{(\underline{A}),J} & \longrightarrow & F \wr \left(\coprod_J F^{i_{n-1}} \wr \Omega_{G,0}^{(A_j)} \sqcup \coprod_{l-J} A_j \right) & \xrightarrow{F_1} & F \wr \coprod F^{i_{n-1}} \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\
\downarrow & & \downarrow & \nearrow \underline{m} & \parallel & & \parallel \\
\Omega_{G,n-1}^{(\underline{A}),J} & \longrightarrow & F \wr \left(\coprod_J F^{i_{n-1}} \wr A_j \sqcup \coprod_{l-J} A_j \right) & \xrightarrow{F} & F \wr \coprod F^{i_{n-1}} \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op}
\end{array} \tag{5.26}$$

$$\begin{array}{ccccccc}
\Omega_{G,n}^{(\underline{A}),J} & \longrightarrow & F \wr \left(\coprod_J F^{i_n} \wr A_j \sqcup \coprod_{l-J} A_j \right) & \xrightarrow{F} & F \wr \coprod F^{i_n} \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op} \\
d_i^{(\underline{A})} \downarrow & \nearrow \pi_i^{(\underline{A})} & \downarrow \sigma^i & & \downarrow \sigma^i & \nearrow \alpha_i & \parallel \\
\Omega_{G,n-1}^{(\underline{A}),J} & \longrightarrow & F \wr \left(\coprod_J F^{i_{n-1}} \wr A_j \sqcup \coprod_{l-J} A_j \right) & \xrightarrow{F} & F \wr \coprod F^{i_{n-1}} \wr \mathcal{V}^{op} & \longrightarrow & \mathcal{V}^{op}
\end{array} \tag{5.27}$$

DISJBARDNJ EQ

5.4 Transferring simplicial colimits of left Kan extensions

Given genuine equivariant operads $X, Y \in \text{Op}_G$ one has an isomorphism

$$X \sqcup^a Y \simeq \text{colim}_{\Delta^{op}} (\mathbb{F}_G^{\bullet+1} X \sqcup^a \mathbb{F}_G^{\bullet+1} Y)$$

so that combining Remarks [REPACKAGERES REM 4.69](#) and Remark [PRECOMPOSTCOMP REM 5.4](#) with the results in the previous section one obtains isomorphisms

$$X \sqcup^a Y \simeq \text{colim}_{\Delta^{op}} (\text{Lan} (N^{\bullet+1} \wr X \sqcup^a N^{\bullet+1} \wr Y)) \tag{5.28}$$

$$\simeq \text{colim}_{\Delta^{op}} (\text{Lan} (N \circ \sqcup \circ (N^{\times 2})^\bullet (\wr X, \wr Y))) \tag{5.29}$$

$$\simeq \text{colim}_{\Delta^{op}} \left(\text{Lan}_{\Omega_{G,\bullet}^{2,op} \rightarrow \Sigma_G^{op}} N_\bullet^{(X,Y)} \right) \tag{5.30}$$

COLIMLAN EQ

where we write $N_\bullet^{(X,Y)}: \Omega_{G,\bullet}^{2,op} \rightarrow \mathcal{V}$ for the induced functor.

The purpose of this section will be show that one can repackage formulas such as [\(5.30\)](#) with a single left Kan extension over a category $\Omega_G^2 = |\Omega_{G,\bullet}^2|$ obtained from $\Omega_{G,\bullet}^2$ via realization in Cat .

We note that $\Omega_{G,\bullet}^2$, together with the corresponding functors to Σ_G , \mathcal{V}^{op} can be viewed as a simplicial object $\Delta^{op} \rightarrow \text{WSpan}^l(\Sigma, G^{op}, \mathcal{V})$, and our first task will be to repackage such functors in terms of Grothendieck constructions.

Lemma 5.31. *Functors $F: \mathcal{D} \rtimes \mathcal{I}_\bullet \rightarrow \mathcal{C}$ are in bijection with lifts*

$$\begin{array}{ccc}
& & \text{WSpan}^l(*, \mathcal{C}) \\
& \nearrow \mathcal{I}_\bullet^F & \downarrow \text{fgt} \\
\mathcal{D} & \xrightarrow{\mathcal{I}_\bullet} & \text{Cat.}
\end{array}$$

where fgt is the functor forgetting the maps to $*$ and \mathcal{C} .

Proof. This is a matter of unpacking notation. The restrictions $F|_{\mathcal{I}_d}$ to the fibers $\mathcal{I}_d \subset D \times \mathcal{I}_\bullet$ are precisely the functors $\mathcal{I}_d^F: \mathcal{I}_d \rightarrow \mathcal{C}$ describing $\mathcal{I}_\bullet^F(d)$.

Furthermore, the images $F((d, i) \rightarrow (d', f_*(i)))$ of the pushout arrows over a fixed arrow $f: d \rightarrow d'$ of \mathcal{D} assemble to a natural transformation

$$\begin{array}{ccc} \mathcal{I}_d & \xrightarrow{\mathcal{I}_d^F} & \mathcal{C} \\ f_* \downarrow & \searrow & \uparrow \\ \mathcal{I}_{d'} & \xrightarrow{\mathcal{I}_{d'}^F} & \mathcal{C} \end{array} \quad (5.32)$$

which describes $\mathcal{I}_\bullet^F(f)$. It is straightforward to check that the associativity and unitality conditions coincide. \square

In the cases of interest we will have $\mathcal{D} = \Delta^{op}$, so that \mathcal{I}_\bullet can be interpreted as an object $\mathcal{I}_\bullet \in \text{Cat}^{\Delta^{op}}$. By recalling the standard cosimplicial object $[\bullet] \in \text{Cat}^\Delta$ given by $[n] = (0 \rightarrow 1 \rightarrow \dots \rightarrow n)$ one obtains the following definition.

Definition 5.33. The left adjoint

$$|-|: \text{Cat}^{\Delta^{op}} \rightleftarrows \text{Cat}: (-)^{[\bullet]}$$

will be called the *realization* functor.

Remark 5.34. More explicitly, one has

$$|\mathcal{I}_\bullet| = \text{coeq} \left(\coprod_{[n] \rightarrow [m]} [n] \times \mathcal{I}_m \rightrightarrows \coprod_{[n]} [n] \times \mathcal{I}_n \right). \quad (5.35) \quad \boxed{\text{REALDEF EQ}}$$

Example 5.36. Any $\mathcal{I} \in \text{Cat}$ induces objects $\mathcal{I}, \mathcal{I}_\bullet, \mathcal{I}^{[\bullet]} \in \text{Cat}^{\Delta^{op}}$ where \mathcal{I} is the constant simplicial object and \mathcal{I}_\bullet is the nerve $N\mathcal{I}$ with each level regarded as a discrete category. It is straightforward to check that $|\mathcal{I}| = |\mathcal{I}_\bullet| = |\mathcal{I}^{[\bullet]}| = \mathcal{I}$.

Lemma 5.37. Given $\mathcal{I}_\bullet \in \text{Cat}^{\Delta^{op}}$ one has an identification $ob(|\mathcal{I}_\bullet|) \simeq ob(\mathcal{I}_0)$. Furthermore, the arrows of $|\mathcal{I}_\bullet|$ are generated by the image of the arrows in $\mathcal{I}_0 \simeq \mathcal{I}_0 \times [0]$ and the image of the arrows in $[1] \times ob(\mathcal{I}_1)$.

For each $i_1 \in \mathcal{I}_1$, we will denote the arrow of $|\mathcal{I}_\bullet|$ induced by the arrow in $[1] \times \{i_1\}$ by

$$d_1(i_1) \xrightarrow{i_1} d_0(i_1).$$

Proof. We write $d_{\hat{k}}, d_{\hat{k}, \hat{l}}$ for the simplicial operators induced by the maps $[0] \xrightarrow{0 \mapsto k} [n]$, $[1] \xrightarrow{0 \mapsto k, 1 \mapsto l} [n]$ which can informally be thought of as the “composite of all faces other than d_k, d_l ”. Using (5.35) one has equivalence relations of objects

$$[n] \times \mathcal{I}_n \ni (k, i_n) \sim (0, d_{\hat{k}}(i_n)) \in [0] \times \mathcal{I}_0$$

and since for any generating relation $(k, i_n) \sim (l, i'_n)$ it is $d_{\hat{k}}(i_n) = d_{\hat{l}}(i'_n)$ the identification $ob(|\mathcal{I}_\bullet|) \simeq ob(\mathcal{I}_0)$ follows.

To verify the claim about generating arrows, note that any arrow of $[n] \times \mathcal{I}_n$ factors as

$$(k, i_n) \rightarrow (l, i_n) \xrightarrow{I_n} (l, i'_n) \quad (5.38) \quad \boxed{\text{FACTORIZATIONREAL EQ}}$$

for $I_n: i_n \rightarrow i'_n$ an arrow of \mathcal{I}_n . The $d_{\hat{l}}$ relation identifies the right arrow in (5.38) with $(0, d_{\hat{l}}(i_n)) \xrightarrow{d_{\hat{l}}(I_n)} (0, d_{\hat{l}}(i'_n))$ in $[0] \times \mathcal{I}_0$ while (if $k < l$) the $d_{\hat{k}, \hat{l}}$ relation identifies the left arrow with $(0, d_{\hat{k}, \hat{l}}(i_n)) \rightarrow (1, d_{\hat{k}, \hat{l}}(i_n))$ in $[1] \times \mathcal{I}_1$. The result follows. \square

Remark 5.39. Given $\mathcal{I}_\bullet \in \mathbf{Cat}^{\Delta^{op}}$, $\mathcal{C} \in \mathbf{Cat}$, the isomorphisms

$$\mathrm{Hom}_{\mathbf{Cat}}(|\mathcal{I}_\bullet|, \mathcal{C}) \simeq \mathrm{Hom}_{\mathbf{Cat}^{\Delta^{op}}}(\mathcal{I}_\bullet, \mathcal{C}^{[\bullet]})$$

together with the fact that $\mathcal{C}^{[\bullet]}$ is always 2-coskeletal show that $|\mathcal{I}_\bullet|$ is determined by the categories $\mathcal{I}_0, \mathcal{I}_1, \mathcal{I}_2$ and maps between them, i.e. by the truncated version of formula (5.35) with $n, m \leq 2$. REALDEF EQ

Indeed, it can be shown that a sufficient set of generating relations in $|\mathcal{I}_\bullet|$ is given by
 (i) the relations in \mathcal{I}_0 (including relations stating that identities of \mathcal{I}_0 are identities of $|\mathcal{I}_\bullet|$);
 (ii) relations stating that for each $i_0 \in \mathcal{I}_0$ the arrow $i_0 = d_1(s_0(i_0)) \xrightarrow{s_0(i_0)} d_1(s_0(i_0)) = i_0$ is an identity; (iii) for each arrow $I_1: i_1 \rightarrow i'_1$ in \mathcal{I}_1 the relation that the square below commutes

$$\begin{array}{ccc} d_1(i_1) & \xrightarrow{i_1} & d_0(i_1) \\ d_1(I_1) \downarrow & & \downarrow d_0(I_1) \\ d_1(i'_1) & \xrightarrow{i'_1} & d_0(i'_1) \end{array}$$

and (iv) for each object $i_2 \in \mathcal{I}_2$ the relation that the following triangle commutes.

$$\begin{array}{ccc} d_{1,2}(i_2) & \xrightarrow{d_1(i_2)} & d_{0,1}(i_2) \\ & \searrow d_2(i_2) & \nearrow d_0(i_2) \\ & d_{0,2}(i_2) & \end{array}$$

Example 5.40. For $\Omega_{G,\bullet}$ the simplicial object of planar strings one has $|\Omega_{G,\bullet}| = \Omega_G^t$, the category of G -trees and tall maps. Indeed, arrows of $\Omega_{G,0}$ and objects of $\Omega_{G,1}$ are naturally identified with the quotient arrows and planar tall maps of Ω_G^t , which are a generating set of arrows. And likewise, relations in $\Omega_{G,0}$, arrows in $\Omega_{G,1}$ and objects in $\Omega_{G,2}$ are identified with the relations of Ω_G^t .

Analogously, for $\Omega_{G,\bullet}^J$ the simplicial object of planar l -labeled strings that are $(\{l\} - J)$ -inert, one has $|\Omega_{G,\bullet}^J| = \Omega_G^{J,t}$, the category of l -labeled G -trees and $(\{l\} - J)$ -inert tall maps.

The following is the key result in this section.

Proposition 5.41. Let $\mathcal{I}_\bullet \in \mathbf{Cat}^{\Delta^{op}}$. Then there is a natural functor

$$\Delta^{op} \ltimes \mathcal{I}_\bullet \xrightarrow{s} |\mathcal{I}_\bullet|. \quad (5.42)$$

Further, s is final.

Remark 5.43. The s in the result above stands for *source*. This is because, for any $\mathcal{I} \in \mathbf{Cat}$, the map $\Delta^{op} \ltimes \mathcal{I}^{[\bullet]} \rightarrow |\mathcal{I}^{[\bullet]}| \simeq \mathcal{I}$ is given by $s(i_0 \rightarrow \dots \rightarrow i_n) = i_0$.

Proof. Recall that $|\mathcal{I}_\bullet|$ is the coequalizer (5.35). Given $(k, g_m) \in [n] \times \mathcal{I}_m$, we will write $[k, g_m]$ for the corresponding object in $|\mathcal{I}_\bullet|$. To simplify notation, we will write objects of \mathcal{I}_n as i_n and implicitly assume that $[k, i_n]$ refers to the class of the object $(k, i_n) \in [n] \times \mathcal{I}_n$. REALDEF EQ

We define s on objects by $s([n], i_n) = [0, i_n]$ and on an arrow $(\phi, I_m): (n, i_n) \rightarrow (m, i'_m)$ as the composite (note that $\phi: [m] \rightarrow [n]$ and $I_m: \phi^*(i_n) \rightarrow i_m$)

$$[0, i_n] \rightarrow [\phi(0), i_n] = [0, \phi^*(i_n)] \xrightarrow{I_m} [0, i'_m]. \quad (5.44)$$

To check associativity, the cases of a pair of either two fiber arrows (i.e. arrows where ϕ is the identity) or two pushforward arrows (i.e. arrows where I_m is the identity) are immediate from (5.44), hence we are left with the case $([n], i_n) \xrightarrow{I_n} ([n], i'_n) \rightarrow ([m], \phi^*(i'_n))$ of a fiber TARGETDEFINITION EQ

arrow followed by a pushforward arrow. Noting that in $\Delta^{op} \ltimes \mathcal{I}_\bullet$ this composite can be rewritten as $([n], i_n) \rightarrow ([m], \phi^*(i_n)) \xrightarrow{\phi^*(I_n)} ([m], \phi^*(i'_n))$ this amounts to checking that

$$\begin{array}{ccccc} [0, i_n] & \longrightarrow & [\phi(0), i_n] & \equiv & [0, \phi^*(i_n)] \\ I_n \downarrow & & I_n \downarrow & & \downarrow \phi^*(I_n) \\ [0, i'_n] & \longrightarrow & [\phi(0), i'_n] & \equiv & [0, \phi^*(i_n)] \end{array} \quad (5.45)$$

commutes in $|\mathcal{I}_\bullet|$, which is the case since the left square is encoded by a square in $[n] \times \mathcal{I}_n$ and the right square is encoded by an arrow in $[m] \times \mathcal{I}_n$.

We now turn to showing that s is final.

Fix $j \in \mathcal{I}_0$. We will show that $[0, j] \downarrow \Delta^{op} \ltimes \mathcal{I}_\bullet$ is indeed connected. By Lemma 5.37 any object in this undercategory has a description (not necessarily unique) as a pair

$$([n], i_n), [0, j] \xrightarrow{f_1} \dots \xrightarrow{f_r} s([n], i_n)$$

where each f_i is a generating arrow of $|\mathcal{I}_\bullet|$ induced by either an arrow I_0 of \mathcal{I}_0 or object $i_1 \in \mathcal{I}_1$.

We will connect this object to the canonical object $(([0], h), [0, h] = [0, h])$, arguing by induction on r . If $n \neq 0$, the map $d_0: ([n], i_n) \rightarrow ([0], d_0^*(i_n))$ and the fact that $s(d_0^*) = id_{[0, d_0^*(i_n)]}$ provides an arrow to an object with $n = 0$ without changing r . If $n = 0$, one can apply the induction hypothesis by lifting f_r to $\Delta^{op} \ltimes \mathcal{I}_\bullet$ according to one of two cases: (i) if f_r is induced by an arrow I_0 of \mathcal{I}_0 , the lift of f_r is simply $([0], i'_0) \xrightarrow{I_0} ([0], i_0)$; (ii) if f_r is induced by $i_1 \in \mathcal{I}_1$ the lift is provided by the map $([1], i_1) \rightarrow ([0], d_0(i_1))$. \square

In practice, we will need to know that s satisfies the following stronger finality condition with respect to left Kan extensions.

Corollary 5.46. *Consider a map $\mathcal{I}_\bullet \rightarrow \mathcal{J}$ between $\mathcal{I}_\bullet \in \text{Cat}^{\Delta^{op}}$ and a constant object $\mathcal{J} = \mathcal{J}_\bullet \in \text{Cat}^{\Delta^{op}}$. Then the source map s*

$$\begin{array}{ccc} \Delta^{op} \ltimes \mathcal{I}_\bullet & \xrightarrow{s} & |\mathcal{I}_\bullet| \\ & \searrow & \swarrow \\ & \mathcal{J} & \end{array}$$

is Lan-final over \mathcal{J} , i.e. the functors $s \downarrow j: (\Delta^{op} \ltimes \mathcal{I}_\bullet) \downarrow j \rightarrow |\mathcal{I}_\bullet| \downarrow j$ are final for all $j \in \mathcal{J}$.

Proof. It is clear that $(\Delta^{op} \ltimes \mathcal{I}_\bullet) \downarrow j \simeq \Delta^{op} \ltimes (\mathcal{I}_\bullet \downarrow j)$ while Lemma 2.2 guarantees that, since $(-) \downarrow j$ is a left adjoint, $|\mathcal{I}_\bullet| \downarrow j \simeq |\mathcal{I}_\bullet \downarrow j|$. One thus reduces to Proposition 5.41. \square

We end this section with two basic lemmas that will allow us to apply Corollary 5.46 to the tree categories we will be interested in.

Lemma 5.47. *Let $\mathcal{I}_\bullet^F \in \text{Span}(*, \mathcal{C})^{\Delta^{op}}$ be such that the diagrams*

$$\begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ d_i \downarrow & \nearrow \delta_i & \uparrow F_{n-1} \\ \mathcal{I}_{n-1} & & \end{array} \quad \begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ s_j \downarrow & \nearrow \sigma_j & \uparrow F_{n+1} \\ \mathcal{I}_{n+1} & & \end{array} \quad (5.48) \quad \text{IDENTSIMPRELSISO EQ}$$

commute up to isomorphism for $0 < i \leq n$, $0 \leq j \leq n$.

Then the functors $\tilde{F}_n: \mathcal{I}_n \rightarrow \mathcal{C}$ given by the composites

$$\mathcal{I}_n \xrightarrow{d_1, \dots, d_n} \mathcal{I}_0 \xrightarrow{F_0} \mathcal{C}$$

assemble to an object $\mathcal{I}_\bullet^{\tilde{F}} \in \text{Span}(*, \mathcal{C})^{\Delta^{op}}$ which is isomorphic to \mathcal{I}_\bullet^F and such that the corresponding diagrams (5.48) for $0 < i \leq n$, $0 \leq j \leq n$ are strictly commutative.

Proof. This follows by a straightforward verification. \square

Lemma 5.49. *A (necessarily unique) factorization*

$$\Delta^{op} \ltimes \mathcal{I}_\bullet \begin{array}{c} \xrightarrow{\quad} \mathcal{C} \\ \searrow s \quad \nearrow \text{dashed} \\ |\mathcal{I}_\bullet| \end{array} \quad (5.50) \quad \text{SOURCEFACT EQ}$$

exists iff for the associated object $\mathcal{I}_\bullet \in \text{Span}(*, \mathcal{C})^{\Delta^{op}}$ (cf. Lemma 5.31) all faces d_i for $0 < i \leq n$ and degeneracies s_j for $0 \leq j \leq n$ are strictly commutative, i.e. they are given by diagrams

$$\begin{array}{ccc} \begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ d_0 \downarrow & \nearrow \varphi_n & \\ \mathcal{I}_{n-1} & \xrightarrow{F_{n-1}} & \end{array} & \begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ d_i \downarrow & \nearrow & \\ \mathcal{I}_{n-1} & \xrightarrow{F_{n-1}} & \end{array} & \begin{array}{ccc} \mathcal{I}_n & \xrightarrow{F_n} & \mathcal{C} \\ s_j \downarrow & \nearrow & \\ \mathcal{I}_{n+1} & \xrightarrow{F_{n+1}} & \end{array} \end{array} \quad (5.51) \quad \text{IDENTSIMPRELS EQ}$$

Proof. For the “if” direction, it suffices to note that s sends all pushout arrows of $\Delta^{op} \ltimes \mathcal{I}_\bullet$ for faces d_i , $0 < i \leq n$ and degeneracies s_j , $0 \leq j \leq n$ to identities and this yields the commutative diagrams (5.51).

For the “only if” direction, this will follow by building a functor $\mathcal{I}_\bullet \xrightarrow{\bar{F}} \mathcal{C}^{[\bullet]}$ together with the naturality of the source map s (recall that $|\mathcal{C}^{[\bullet]}| \simeq \mathcal{C}$). We define $\bar{F}_{n|k \rightarrow k+1}$ as the map

$$F_{n-k} d_{0, \dots, k-1} \xrightarrow{\varphi_{n-k} d_{0, \dots, k-1}} F_{n-k-1} d_{0, \dots, k}. \quad (5.52) \quad \text{EQUIVALENCEDEF EQ}$$

The claim that $s \circ (\Delta^{op} \ltimes \bar{F})$ recovers the horizontal map in (5.50) is straightforward, hence the real task is to prove that (5.52) indeed defines a map of simplicial objects.

$$\varphi_{n-1} d_i = \varphi_n, \quad 1 < i \quad \varphi_{n-1} d_1 = (\varphi_{n-1} d_0) \circ \varphi_n, \quad \varphi_{n+1} s_i = \varphi_n, \quad 0 < i, \quad \varphi_{n+1} s_0 = id_{F_n} \quad (5.53)$$

Next, note that there is no ambiguity in writing simply $\varphi_{n-k} d_{0, \dots, k-1}$ to denote the map (5.52). We now check that $\bar{F}_{n-1} d_i = d_i \bar{F}_n$, $0 \leq i \leq n$, which must be verified after restricting to each $k \rightarrow k+1$, $0 \leq k \leq n-2$. There are three cases, depending on i and k :

- ($i < k+1$) $\varphi_{n-k-1} d_{0, \dots, k-1} d_i = \varphi_{n-k-1} d_{0, \dots, k}$;
- ($i = k+1$) $\varphi_{n-k-1} d_{0, \dots, k-1} d_i = \varphi_{n-k-1} d_{0, \dots, k-1} = (\varphi_{n-k-1} d_0 \circ \varphi_{n-k}) d_{0, \dots, k-1} = (\varphi_{n-k-1} d_{0, \dots, k}) \circ (\varphi_{n-k} d_{0, \dots, k-1})$;
- ($i > k+1$) $\varphi_{n-k-1} d_{0, \dots, k-1} d_i = \varphi_{n-k-1} d_{i-k} d_{0, \dots, k-1} = \varphi_{n-k} d_{0, \dots, k-1}$.

The case of degeneracies is similar. \square

Remark 5.54. One can twist all results by the opposite functor

$$\Delta \xrightarrow{(-)^{op}} \Delta$$

which sends $[n]$ to itself and d_i, s_i to d_{n-i}, s_{n-i} . In doing so, one obtains vertical isomorphisms

$$\begin{array}{ccc} \Delta^{op} \ltimes (\mathcal{J}_\bullet \circ (-)^{op}) & \xrightarrow{s} & |\mathcal{J}_\bullet \circ (-)^{op}| \\ \simeq \downarrow & & \downarrow \simeq \\ \Delta^{op} \ltimes \mathcal{J}_\bullet & \xrightarrow{t} & |\mathcal{J}_\bullet| \end{array}$$

which reinterpret the “source” functor as what one might call the “target” functor, with $t([n], i_n) = [n, i_n]$ rather than $s([n], i_n) = [0, i_n]$.

Corollary 5.46 now says that t is Lan-final and Lemmas 5.47, 5.49 generalize in the obvious way by replacing s with t and d_0 with d_n .

5.5 The category of extension trees

In this section we combine the previous sections to obtain a compact description of free extension pushouts

$$\begin{array}{ccc} \mathbb{F}_G A & \longrightarrow & X \\ \downarrow & & \downarrow \\ \mathbb{F}_G B & \longrightarrow & Y \end{array} \quad (5.55) \quad \boxed{\text{FREEEXT EQ}}$$

as a left Kan extension over a convenient category of trees.

For simplicity, we first explain how to obtain a similar description for the simpler case of a coproduct $X \sqcup^a Y$. By (5.30), one has a description

$$\begin{aligned} X \sqcup^a Y &\simeq \text{colim}_{\Delta^{op}} \left(\text{Lan}_{\Omega_{G,\bullet}^{2,op} \rightarrow \Sigma_G^{op}} N_{\bullet}^{(X,Y)} \right) \\ &\simeq \text{Lan}_{\Delta^{op} \ltimes \Omega_{G,\bullet}^{2,op} \rightarrow \Sigma_G^{op}} N_{\bullet}^{(X,Y)} \end{aligned}$$

where the second identification follows from formal properties of Grothendieck constructions.

Combining the fact that (5.27) consists of natural isomorphisms with (the Remark 5.54 dual of) Lemma 5.47, yields an isomorphic twisted functor $\tilde{N}_{\bullet}^{(X,Y)}$ with strictly commutative s_i and d_i for $i \neq n$. The dual of Lemma 5.49 now says that $\tilde{N}_{\bullet}^{(X,Y)}$ factors via the target map t through $\Omega_{G,\bullet}^{2,op}$ (writing $\tilde{N}^{(X,Y)}$ for the factorization) and thus the dual of Corollary 5.46 finally yields

$$X \sqcup^a Y \simeq \text{Lan}_{\Omega_G^{2,op} \rightarrow \Sigma_G^{op}} \tilde{N}^{(X,Y)}. \quad (5.56)$$

We recall that by Example 5.40, Ω_G^2 is simply the category of $\underline{2}$ -labeled trees and tall label maps.

More generally, one has

$$\coprod_J^a X_j \sqcup^a \coprod_{l=J}^a \mathbb{F}_G X_j \simeq \text{Lan}_{\Omega_G^{J,op} \rightarrow \Sigma_G^{op}} \tilde{N}^{(\underline{X})}. \quad (5.57) \quad \boxed{\text{LANCOPRODDDESC}}$$

where Ω_G^J is the category of \underline{l} -labeled trees and tall $(\underline{l} - J)$ -inert label maps.

Remark 5.58. We note that the twisting $\tilde{N}_{\bullet}^{(X,Y)}$ is fairly harmless. For explicitness, we focus on the simplest case of a “unary coproduct”, in which case (5.57) is simply recovering the genuine equivariant operad X from its bar resolution. In that case $N_{\bullet}^X: \Omega_{G,2}^{op} \rightarrow \mathcal{V}$ is given by the top map in (4.51) or, equivalently, by the top map in (4.52) (we note that, in the notation therein, it is $A = \Sigma_G$). On the other hand, the twisted map $\tilde{N}_2^X: \Omega_{G,2}^{op} \rightarrow \mathcal{V}$ is given by the left bottom composite in either of (4.51), (4.52). Informally, the role of this twisting is therefore simply that of replacing the order on $V_G(T_n)$ induced lexicographically by planar strings $T_0 \rightarrow \dots \rightarrow T_n$ with the simpler order induced directly from T_n .

In what follows we will largely be able to ignore this technicality. Indeed, the role of lexicographic orders in building (5.57) is that of guaranteeing that \tilde{N}_{\bullet} satisfies the necessary simplicial identities, which are ensured by appealing to the bar construction for the monad N .

We now turn to the task of building (5.55) as a left Kan extension. One has a colimit description

$$\mathbb{F}B \coprod_{\mathbb{F}A} X \simeq \text{colim}_{\Delta^{op}} \left(\mathbb{F}B \sqcup \mathbb{F}A \sqcup X \xleftarrow{\quad} \mathbb{F}B \sqcup \mathbb{F}A \sqcup \mathbb{F}A \sqcup \mathbb{F}A \sqcup X \xleftarrow{\quad} \mathbb{F}B \sqcup \mathbb{F}A \sqcup \mathbb{F}A \sqcup \mathbb{F}A \sqcup \mathbb{F}A \sqcup \mathbb{F}A \sqcup X \quad \dots \right) \quad (5.59) \quad \boxed{\text{FREEEXTUSEFCOL EQ}}$$

where all differentials are fold maps of $\mathbb{F}A$ except to the n -th differential d_n , which is induced by the two maps $\mathbb{F}A \rightarrow X$, $\mathbb{F}A \rightarrow \mathbb{F}B$.

By the previous discussion each individual object $X \sqcup (\mathbb{F}A)^{\sqcup 2n+1} \sqcup \mathbb{F}B$ in (5.61) can be described as a left Kan extension over the tree category $\Omega_G^{\{X\}}$ where $\{X\} \subset \{B, A, \dots, A, X\}$ is a singleton. The maps in (5.61) can themselves be encoded as span maps between the $\Omega_G^{\{X\}}$. To see this, we make (5.61) more precise. Firstly, we write $\langle n \rangle$ for the poset

$$-\infty \leq -n \leq -n+1 \leq \dots \leq -1 \leq 0 \leq 1 \leq \dots \leq n-1 \leq n \leq +\infty.$$

The posets $\langle n \rangle$ together with antisymmetric (i.e. such that $f(-x) = -f(x)$) poset maps preserving all three of $-\infty, 0, +\infty$ then form a simplicial object¹ $\langle - \rangle: \Delta^{op} \rightarrow \mathbf{F}$.

(5.55) thus induces a simplicial object $(B, A, X)_{\langle n \rangle} \in \mathbf{F} \wr \mathbf{Fun}(\Sigma_G^{op}, \mathcal{V})$.

Each level of $(\iota B, \iota A, \iota X)_{\langle n \rangle}$ is then a $N^{\times\{+\infty\}}$ -algebra on $(\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V}))^{\times\langle n \rangle}$, compatibly with the simplicial maps. One thus obtains a *bisimplicial* object

$$\Sigma_G^{op} \leftarrow \Omega_{G, \bullet}^{\{+\infty\}(\bullet), op} \xrightarrow{N_{\bullet}^{(B, A, X)}(\bullet)} \mathcal{V}$$

on $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V})$ whose realization along the string direction yields the spans

$$\Sigma_G^{op} \leftarrow \Omega_G^{\{+\infty\}(\bullet), op} \xrightarrow{N^{(B, A, X)}(\bullet)} \mathcal{V} \quad (5.60) \quad \text{PARTREALSPAN EQ}$$

discussed above, except now assembled into a simplicial object in $\mathbf{WSpan}^l(\Sigma_G^{op}, \mathcal{V})$.

All degeneracies s_i and differentials d_i of (5.60) other than the top differential d_n are induced by maps α^* described in §5.1 and thus given by strictly commutative diagrams, so that Lemma 5.49 and Corollary 5.46 can be applied (this time with no need to appeal to Lemma 5.47) so as to allow (5.61) to be repackaged as


$$\mathbb{F}B \coprod_{\mathbb{F}A} X \simeq \mathbf{Lan}_{\Omega_G^{e, op} \rightarrow \Sigma_G^{op}} N^{(B, A, X)} \quad (5.61) \quad \text{FREEEXTUSEFCOL EQ}$$

where we write Ω_G^e for $|\Omega_G^{\{+\infty\}(\bullet)}|$. We now turn to the task of describing Ω_G^e , starting with by defining it directly.

Definition 5.62. The *extension tree category* Ω_G^e is the category whose objects are $\{B, A, X\}$ -labeled trees and whose maps $\varphi: T \rightarrow S$ are tall maps of trees such that

- (i) if $T_{v_{Ge}}$ has an A -label, then $S_{v_{Ge}} = T_{v_{Ge}}$ and $S_{v_{Ge}}$ has an A -label;
- (ii) if $T_{v_{Ge}}$ has a B -label, then $S_{v_{Ge}} = T_{v_{Ge}}$ and $S_{v_{Ge}}$ has either an A -label or a B -label;
- (iii) if $T_{v_{Ge}}$ has a X -label, then $S_{v_{Ge}}$ has only A and X -labels.

Example 5.63. The following is an example of a planar map in Ω_G^e , where black nodes represent X -labeled nodes, grey nodes represent B -labeled nodes and white nodes represent A -labeled nodes.



(5.64) REGALTERNMAP EQ

¹Indeed, we recall that the opposite simplex category Δ^{op} can be equivalently described as the category of *intervals*, i.e. finite ordered posets with distinct top and bottom, along with order maps preserving both top and bottom. $\langle n \rangle$ can then be regarded as obtained by gluing the interval $0 \leq 1 \leq \dots \leq n \leq +\infty$ with its opposite.

Proposition 5.65. *One has an identification*

$$\Omega_G^\varepsilon \simeq |\Omega_G^{\{+\infty\}(\bullet)}|.$$

Proof. We note first that Ω_G^ε contains all label maps that are $\{A, B\}$ -inert. In fact, any map of Ω_G^ε clearly has a unique factorization as such a label map followed by an underlying planar isomorphism of trees that replaces some of the X and B labels with A labels. We will refer to the former as label maps and to the latter as relabel maps.

We recall that $\Omega_G^{\{+\infty\}(n)}$ consists of trees with $2n + 3$ types of labels: X -labels, B -labels and $2n + 1$ distinct types of A -labels. One can equivalently encode such a tree as a string $T_0 \rightarrow \dots \rightarrow T_n$ of relabel maps. Indeed, the A -label nodes of T_n in such a string are partitioned into $2n + 1$ types according to that node's labels on the T_i (which are either all A 's, some X 's and then A 's or some B 's and then A 's). Moreover, a diagram

$$\begin{array}{ccccccc} T_0 & \longrightarrow & T_1 & \longrightarrow & \dots & \longrightarrow & T_n \\ f_0 \downarrow & & f_1 \downarrow & & & & f_n \downarrow \\ T'_0 & \longrightarrow & T'_1 & \longrightarrow & \dots & \longrightarrow & T'_n \end{array}$$

with f_i label maps of Ω_G^ε is then equivalent to a label map $f_n: T_n \rightarrow T'_n$ respecting all $2n + 3$ labels in $\Omega_G^{\{+\infty\}(n)}$. Since the string description above is also compatible with the simplicial structure maps in the obvious way, the result is now clear. \square

HERE

Our next task will be that of identifying a convenient Lan-final subcategory $\bar{\Omega}_G^\varepsilon \hookrightarrow \Omega_G^\varepsilon$.

6 Filtration of Cellular Extensions

As we saw above, we have the the free extension $\mathcal{P}[u]$ given by the pushout

$$\begin{array}{ccc} \mathbb{F}_G X & \longrightarrow & \mathcal{P} \\ u \downarrow & & \downarrow \\ \mathbb{F}_G Y & \longrightarrow & \mathcal{P}[u] \end{array}$$

can be built via Kan extensions over $\Omega_{G,e}$. Thus, in order to filter the map $\mathcal{P} \rightarrow \mathcal{P}[u]$, it will suffice, via naturality of Kan extensions, to filter the category $\Omega_{G,e}$.

We begin by analyzing the objects of this category. These are $(\mathcal{P}; X, Y)$ -alternating G -trees T ; that is, an odd tree T with each odd vertex labeled by \mathcal{P} , and the even vertices labeled by either X or Y .

Adjusting the general notation of the previous section to this setting, given $T \in \Omega_{G,e}$ we let $V_{\mathcal{P}}(T)$, $V_X(T)$, and $V_Y(T)$ denote the G -sets of vertices labeled by \mathcal{P} , X , or Y , respectively, and $V_{G,\mathcal{P}}(T)$ (and similarly) the set $V_{\mathcal{P}}(T)/G$ of orbits. Further, let $V_{G,in}(T) = V_{G,X}(T) \sqcup V_{G,Y}(T)$ denote the set of *inert* or *passive* nodes. Moreover, we let $\mathbb{V}_{G,in}(T) \in \mathbf{F}\mathbf{i}\mathbf{V}$ denote the map $V_{G,in}(T) \rightarrow \mathcal{V}$ which sends T_v to $X(T_v)$ or $Y(T_v)$, depending on the labeling of the vertex v .

Further, we define the *degree* of T , denote $|T|$, to be the sum $|T|_X + |T|_Y$, where $|T|_X$ is define by

$$|T|_X = \frac{|V_X(T)|}{|G.r|} = \sum_{G.v \in V_{G,X}(T)} \frac{|G.v|}{|G.r|}$$

where $G.r$ is the root G -set of T , and similarly for $|T|_Y$. Intuitively, $|T|_X$ is the number of X -labeled vertices in any (every) single tree component of T .

6.1 Filtration Pieces

We begin our filtration of $\Omega_{G,e}$.

Definition 6.1. We define subcategories of Ω_G^e .

1. Let $\Omega_G^e[\leq k]$ (respectively $\Omega_G^e[k]$) be the full subcategory of Ω_G^e spanned by trees T with $|T| \leq k$ (respectively, $|T| = k$).
2. Let $\Omega_G^e[\leq k, -]$ (respectively $\Omega_G^e[k, -]$) be the full subcategory of $\Omega_G^e[\leq k]$ (respectively $\Omega_G^e[k]$) spanned by trees T with $|T|_Y \neq k$.
3. Let $\Omega_G^e[\leq k, 0]$ (respectively $\Omega_G^e[k, 0]$) be the full subcategory of $\Omega_G^e[\leq k]$ (respectively $\Omega_G^e[k]$) spanned by trees T with $|T|_X = 0$ (equivalently, $|T|_Y = k$).
4. If Ξ is any of the above categories, and $C \in \Sigma_G$, let $\Xi(C)$ denote the full subcategory of Ξ spanned by those trees T with $\text{val}(T) \simeq C$.

Remark 6.2. The categories $\Omega_G^e[k]$ and $\Omega_G^e[k, -]$ have only very limited morphisms, as there cannot be any “active substitutions”. Thus, any map $S \rightarrow T$ in these categories just changes some Y -labelings into X -labelings, while the underlying $(\mathcal{P}; Z)$ -alternating tree remains fixed (where here the one passive colour Z encompasses both Y and X).

Lemma 6.3. $\Omega_G^e[\leq k-1]^{op}$ is Lan-final in $\Omega_G^e[\leq k, -]^{op}$ over Σ_G^{op} .


In order to prove this, we will first need a particular construction $T \mapsto T_{\mathcal{P}}^{\lambda}$ on $\Omega_{G,e}$.

Definition 6.4. Given a $(\mathcal{P}; X, Y)$ -alternating G -tree T , let $T_{\mathcal{P}}^{\lambda}$ denote the $(\mathcal{P}; Y)$ -alternating tree created from T by

1. relabelling at X -nodes \mathcal{P} (yielding a (\mathcal{P}, Y) -labeled tree); then
2. collapsing all connected components of \mathcal{P} -labeled nodes.

There is a unique planar-tall map $\partial_{\mathcal{P}} : T_{\mathcal{P}}^{\wedge} \rightarrow T$, and in fact this map factors through all maps of the form $\partial_{\mathcal{P}} : S \rightarrow T$.

Example 6.5. We observe that this construction is symmetric across all tree components, and hence, to give an example, it suffices to show what happens on a single component (i.e. when $G = \{e\}$). Consider the $(\mathcal{P}; X, Y)$ -alternating tree T below, where black nodes are \mathcal{P} -labeled, white nodes filled with \times are X -labeled, and empty white nodes are Y -labeled. After Step (1), we produce the tree $T_{\mathcal{P}}^{\sim}$ in the middle, and collapsing connected \mathcal{P} -components yields the tree $T_{\mathcal{P}}^{\wedge}$ on the right.



Proof of Lemma 6.3. ~~MINUS_LAN_FINAL_LEMMA~~ Fix an arbitrary $C \in \Sigma_G$, and consider an element $q_S : \text{val}(S) \leftarrow C$ in $\Omega_G^e[\leq k, -]^{op} \downarrow C$ (so in particular $S \in \Omega_G^e[\leq k, -]$). We must show that the overcategory

$$(\Omega_G^e[\leq k-1]^{op} \downarrow C) \downarrow (\text{val}(S) \leftarrow C)$$

is non-empty and connected. If in fact $S \in \Omega_G^e[\leq k-1]$, the result is immediate. Otherwise, consider the map

$$S_{\mathcal{P}}^{\wedge} \xrightarrow{\partial_{\mathcal{P}}} S.$$

Since $|S|_Y \neq k$, $|S_{\mathcal{P}}^{\wedge}| \leq k-1$, and hence we have a diagram


$$\begin{array}{ccc} \text{val}(S) & \xleftarrow{\partial_{\mathcal{P}}} & \text{val}(S_{\mathcal{P}}^{\wedge}) \\ & \nwarrow q_S \quad \nearrow q_S & \\ & C & \end{array} \quad (6.6) \quad \boxed{\text{K-1_LAN_FINAL_EQ1}}$$

showing that the desired overcategory is inhabited. Further, given any other element

$$\begin{array}{ccc} \text{val}(S) & \xleftarrow{f} & \text{val}(T) \\ & \nwarrow q_T \quad \nearrow q_S & \\ & C & \end{array} \quad (6.7) \quad \boxed{\text{K-1_LAN_FINAL_EQ2}}$$

in the overcategory, consider the following zig-zag of maps connecting the objects ~~(6.6)~~ ^{~~MINUS_LAN_FINAL_LEMMA~~} and ~~(6.6)~~ ^{~~MINUS_LAN_FINAL_LEMMA~~}

K-1_LAN_FINAL_EQ2
(6.7):



(6.8)

K-1_LAN_FINAL_DIAGRAM

Here, we have omitted the notation “ val ” from the top three rows. To understand this diagram, we first record that we have a factorization:

$$q_S = \tilde{q}_T q_T,$$

Then, if we let $C_S = val(S) = val(S_P^\wedge)$ and $C_T = val(T)$, we have

$$C \xrightarrow{q_T} C_T \xrightarrow{\tilde{q}_T} C_S$$

and hence, by the unique factorization of maps in $\Omega_{G,e}$, a factorization

$$\begin{array}{ccc} C_T & \xrightarrow{\quad} & \tilde{q}_T^*(S_P^\wedge) \\ \tilde{q}_T \downarrow & & \downarrow \tilde{q}_T \\ C_s & \longrightarrow & S_P^\wedge \end{array}$$

(where we are recording $C \rightarrow val(S)$ as a planar-tall map $C \rightarrow S$). A similar analysis shows that the top left trapezoid commutes.

The other regions also commute by a straightforward analysis. Indeed, the top right trapezoid commutes by unique factorization, and finally the middle triangle of ∂_P maps commutes since $(\tilde{q}_T^* S)_P^\wedge = \tilde{q}_T^*(S_P^\wedge)$.

Lastly, we must check that the middle two maps are in fact elements of the appropriate overcategory. This follows from the fact that S_P^\wedge and T have $|-|_Y < k$. Thus, the overcategory in question is connected, as desired. \square

come back: define S_Y^\wedge .

N_FINALITY_LEMMA

Lemma 6.9. $\Omega_G^e[k, 0]^{op}$ is Lan-final in $\Omega_G^e[k]$ over Σ_G^{op} .

Similarly, we need a construction $T \mapsto T_Y^\wedge$ in order to prove this lemma. However, in this case, the analogous notion is much simpler, as T_Y^\wedge has the same underlying $(\mathcal{P}; Z)$ -alternating G -tree, but we just relabel all X -vertices as being Y -labeled.

Proof of Lemma 6.9. ZERO_LAN_FINALITY_LEMMA This follows analogously to Lemma 6.3, by replacing Diagram MINUS_LAN_FINAL_LEMMA K-1_LAN_FINAL_DIAGRAM 6.8 with

the diagram below:

$$\begin{array}{ccccc}
 & & S & & \\
 & \nearrow \partial_Y & \uparrow \tilde{q}_T & \nwarrow f & \\
 & & \tilde{q}_T^*(S) & & \\
 & \nwarrow \partial_Y & \uparrow \partial_Y & \nearrow \partial_Y & \\
 S_Y^\wedge & \xleftarrow{\tilde{q}_T} & \tilde{q}_T^*(S_Y^\wedge) & \xrightarrow{\partial_Y} & T \\
 & \nwarrow q_S & \uparrow q_T & \nearrow q_T & \\
 & & C & &
 \end{array}$$

□

Finally, we show that each layer $\Omega_G^e[\leq k]$ can be built from $\Omega_{G,e}[\leq k-1]$ via a pushout which attaches trees with precise degree k . While dealing with general pushouts of categories requires solving a “word problem” on morphisms, we will only work in cases where the problem collapses. We recall that, given a square of categories

$$\begin{array}{ccc}
 \mathcal{A} & \longrightarrow & \mathcal{C} \\
 \downarrow & & \downarrow \\
 \mathcal{C} & \longrightarrow & \mathcal{D}
 \end{array}$$

if the nerve of this square is a pushout in \mathbf{sSet} , then this is a pushout of categories (since the nerve is the inclusion of a reflective subcategory).

Definition 6.10. We call such squares *nervous pushouts* of categories.

If we further assume that the span of functors is built out of fully-faithful inclusions, these pushouts behave as nicely as possible with left Kan extensions.

Lemma 6.11. *Given any diagram in categories of the form*

$$\begin{array}{ccccc}
 \mathcal{A} & \xrightarrow{f} & \mathcal{C} & & \\
 \downarrow & & \downarrow i & & \\
 \mathcal{B} & \xrightarrow{g} & \mathcal{D} & \xrightarrow{Y} & \mathcal{V} \\
 & & \downarrow j & & \\
 & & \mathcal{D} & &
 \end{array}$$

such that the square is a nervous pushout of fully-faithful functors, then $\text{Lan}_j Y$ is the pushout of the induced span

$$\begin{array}{ccc}
 \text{Lan}_{jif}(Yif) & \longrightarrow & \text{Lan}_{ji}(Yi) \\
 \downarrow & & \\
 \text{Lan}_{jg}(Yg) & &
 \end{array}$$

Proof. By the universal property of left Kan extensions, it suffices to show that, for any functor $Z : \mathcal{V} \rightarrow \mathcal{D}$, the natural map

$$\mathcal{V}^{\mathcal{D}}(Y, Zj) \longrightarrow \mathcal{V}^{\mathcal{B}}(Yg, Zjg) \prod_{\mathcal{V}^{\mathcal{A}}(Yif, Zjif)} \mathcal{V}^{\mathcal{C}}(Yi, Zji)$$

is a bijection. These two sets give the same data: a collection of maps $\Phi_b : Y(b) \rightarrow Z(b)$ and $\Phi_c : Y(c) \rightarrow Z(c)$ for all $b \in \mathcal{B}$ and $c \in \mathcal{C}$, such that $\Phi_b = \Phi_c$ whenever $b = c \in \mathcal{A}$. In general, the compatibilites required on the right are less demanding. However, with the above assumptions, a map $d \rightarrow d'$ in \mathcal{D} is *uniquely* a map in \mathcal{A} , $\mathcal{B} \setminus \mathcal{A}$, or $\mathcal{C} \setminus \mathcal{A}$, and thus all the necessary compatibilities are covered by (at least) one of the $\{\Phi_b\}$ or $\{\Phi_c\}$. □

We can now build our category $\Omega_{G,e}[\leq k]$ inductively.

Lemma 6.12. $\Omega_G^e[\leq k]$ is isomorphic to the pushout below.

$$\begin{array}{ccc} \Omega_G^e[k, -] & \longrightarrow & \Omega_G^e[\leq k, -] \\ \downarrow & & \downarrow \\ \Omega_G^e[k] & \longrightarrow & \Omega_G^e[\leq k] \end{array}$$

In fact, it is a nervous pushout of fully-faithful functors.

Proof. Since maps in Ω_G^e can only increase $|-|$ by adding $|-|_X$, if T is a tree with $|T| = |T|_Y = k$, then any other tree $S \in \Omega_G^e$ connected to T via a zig-zag of maps must have $|S| = k$; that is, if $T \in \Omega_G^e[\leq k] \setminus \Omega_G^e[\leq k, -]$, then the connected component of T is entirely contained in $\Omega_G^e[k]$. Conversely, if $T \in \Omega_G^e[\leq k] \setminus \Omega_G^e[k]$, the connected component of T is entirely contained in $\Omega_G^e[\leq k, -]$. Since the natural induced map

$$\Omega_G^e[k] \sqcup \Omega_G^e[\leq k, -] \rightarrow \Omega_G^e[\leq k]$$

is clearly full and surjective on objects, the result follows from the above discussion and the obvious fully-faithfulness of the span. \square

Abusing notation, we will denote by N^e the restriction of that functor to any of the subcategories of Ω_G^e in the above pushout square.

We can now define the sequencers which will make up our filtration of $\mathcal{P}[u]$:

Definition 6.13. Let \mathcal{P}_k denote the left Kan extension

$$\begin{array}{ccc} \Omega_G^e[\leq k]^{op} & \xrightarrow{N^e} & \mathcal{V} \\ \text{val} \downarrow & \searrow \mathcal{P}_k & \\ \Sigma_G^{op} & & \end{array}$$

Note that by naturality of Lan, we have maps $\mathcal{P}_{k-1} \rightarrow \mathcal{P}_k$.

6.2 Notation

Luis: should this be stated earlier when defining the categorical wreath products?

In order to state our filtration result, we will need to identify another categorical construction. This filtration will be built out of “pushout products over trees of maps of sequences”. This subsection is dedicated to making that statement precise.

Definition 6.14. Given a map $Y_0 \rightarrow Y_1$ of G -symmetric sequences $\mathcal{V}^{\Sigma_G^{op}}$, and $(A, D) \in \mathbf{F} \wr \Sigma_G$, we borrow notation from [1] and define the functor

$$[u]^D : (0 \rightarrow 1)^A \rightarrow \mathcal{V}$$

as the composite

$$(0 \rightarrow 1)^A \rightarrow \mathbf{F} \wr \mathcal{V} \xrightarrow{x} \mathcal{V}$$

where the first map is defined on $\xi : A \rightarrow \{0, 1\}$ by

$$(a \mapsto \xi(a)) \mapsto (A, (a \mapsto Y_{\xi(a)}(D(a))))$$

We recall that, in a general category \mathcal{C} , a subcategory $\mathcal{C}' \subseteq \mathcal{C}$ is called *convex* if whenever $c' \in \mathcal{C}'$ and $c \rightarrow c'$ is an arrow in \mathcal{C} then both c and the map are in \mathcal{C}' .

Q_DEFINITION

Definition 6.15. Let \mathcal{C} be a convex subcategory of $(0 \rightarrow 1)^A$. We define

$$Q_{\mathcal{C}}[u]^D := \text{colim}_{\mathcal{C}} [u]^D.$$

Moreover, given nested convex subcategories $\mathcal{C}' \subseteq \mathcal{C}$, we let

$$[u]^D \square_{\mathcal{C}'}^{\mathcal{C}} : Q_{\mathcal{C}'}[u]^D \rightarrow Q_{\mathcal{C}}[u]^D$$

denote the unique natural map.

In particular, if \mathcal{C} is the full “punctured cube” subcategory $(0 \rightarrow 1)^A \setminus \{(1)_a\}$, we simplify the notation as follows:

$$Q[u]^D := Q_{\mathcal{C}}[u]^D$$

$$[u]^{\square D} := [u]^D \square_{\mathcal{C}}^{(0 \rightarrow 1)^A} : Q[u]^D \rightarrow \bigotimes_{a \in A} Y_1(D(a)).$$

6.3 Filtration Result

We can now state our filtration of the cellular extension $\mathcal{P} \rightarrow \mathcal{P}[u]$.

Theorem 6.16. *Let \mathcal{P} be a genuine G -operad, and suppose we are given a map of G -symmetric sequences $u : Y_0 \rightarrow Y_1$. Then we have a filtration in G -sequences of the cellular extension*

$$\mathcal{P} = \mathcal{P}_0 \rightarrow \mathcal{P}_1 \rightarrow \dots \rightarrow \text{colim}(\mathcal{P}_i) = \mathcal{P}[u],$$

where $\mathcal{P}_{k-1} \rightarrow \mathcal{P}_k$ is given by the pushout

$$\begin{array}{ccc} \text{Lan}_{\Omega_{G,e}[k,-]^{op}} N^e & \longrightarrow & \mathcal{P}_{k-1} \\ \downarrow & & \downarrow \\ \text{Lan}_{\Omega_{G,e}[k]^{op}} N^e & \longrightarrow & \mathcal{P}_k \end{array}$$

Levelwise, for each $C \in \Sigma_G$, in the underlying category $\mathcal{V}^{G \times \Sigma^n}$, we have a filtration on the evaluations at C , where $\mathcal{P}_{k-1}(C) \rightarrow \mathcal{P}_k(C)$ is given by the pushout

$$\begin{array}{ccc} \coprod_{[T] \in \Omega_{G,e}[k,0](C)/\simeq} \left(\bigotimes_{v \in V_{G,\mathcal{P}}(T)} \mathcal{P}(T_v) \otimes Q[u]^{\vee_{G,in}(T)} \right) \otimes_{\text{Aut}(T)} \text{Aut}(C) & \longrightarrow & \mathcal{P}_{k-1}(C) \\ \downarrow & & \downarrow \\ \coprod_{[T] \in \Omega_{G,e}[k,0](C)/\simeq} \left(\bigotimes_{v \in V_{G,\mathcal{P}}(T)} \mathcal{P}(T_v) \otimes \bigotimes_{v \in V_{G,in}(T)} Y_1(T_v) \right) \otimes_{\text{Aut}(T)} \text{Aut}(C) & \longrightarrow & \mathcal{P}_k(C) \end{array}$$

where the left vertical map is the iterated box product

$$\coprod_{V_{G,\mathcal{P}}(T)} \square_{\mathcal{P}(T_v)} \iota_{\mathcal{P}(T_v)} \square[u]^{\square_{\vee_{G,in}(T)}},$$

$\iota_{\mathcal{P}(T_v)}$ is the canonical map $\mathcal{O} \rightarrow \mathcal{P}(T_v)$, out of the initial object, and $\Omega_{G,e}[k,0](C)$ is as in Definition 6.1.

LAN_PUSHOUT_TREE_FORMS_DECOMP_LEMMA

Proof. Combining Lemmas 6.11 and 6.12, we have that \mathcal{P}_k can be computed as the pushout

$$\begin{array}{ccc} \text{Lan}_{\Omega_{G,e}[k,-]^{op}} N^e & \longrightarrow & \text{Lan}_{\Omega_{G,e}[\leq k,-]^{op}} N^e \\ \downarrow & & \downarrow \\ \text{Lan}_{\Omega_{G,e}[k]^{op}} N^e & \longrightarrow & \text{Lan}_{\Omega_{G,e}[\leq k]^{op}} N^e := \mathcal{P}_k \end{array}$$

(6.17)

FILTRATION_LAN_SQUARE

MINUS LAN FINAL LEMMA

By Lemma 6.3, the top right corner can be identified with \mathcal{P}_{k-1} . Thus, it remains to identify the left hand side.

ZERO LAN FINALITY LEMMA

By Lemma 6.9, we may replace the bottom left corner with $\text{Lan}_{\Omega_{G,e}[k,0]^{op}} N^e$. Now, given $T \in \Omega_{G,e}[k,0]$, let $[T]$ denote the isomorphism class of T in $\Omega_{G,e}[k,0]$. With this notation, the bottom left corner can further be identified with

$$\coprod_{[T] \in \Omega_{G,e}[k,0](C)/\sim} N^e(T) \otimes_{\text{Aut}(T)} \text{Aut}(C) = \coprod_{[T]} \left(\bigotimes_{v \in V_{G,\mathcal{P}}(T)} \mathcal{P}(T_v) \otimes \bigotimes_{v \in V_{G,Y}(T)} Y_1(T_v) \right) \otimes_{\text{Aut}(T)} \text{Aut}(C).$$

Next, we observe that the non-invertible morphisms of $\Omega_{G,e}[k,-]^{op} \downarrow C$ are just those which change the labeling of some nodes from X to Y . Given S and T in $\Omega_{G,e}[k,-]$, write $S \sim T$ if they are in the same path component, and again note that this implies $|S| = |T|$, and moreover that S and T forget to the same $(\mathcal{P}; Z)$ -alternating tree. Denote the path component of T by (T) .

We note that the set of path components of those trees with $\text{val}(T) = C$ is equal to the set of isomorphism classes in $\Omega_{G,e}[k,0](C)$, as both are just determined precisely by their underlying $(\mathcal{P}; Z)$ -alternating tree.

To account for the $\text{Aut}(C)$ -action on the indexing category, we note that each connected component of $\Omega_{G,e}[k,-]^{op} \downarrow C$ has an action of $\text{Aut}([T])$. Thus, the top left corner of Diagram (6.17) can be identified with the image of the colimit map below:

$$\begin{array}{c} \coprod_{[T] \in \Omega_{G,e}[k](C)/\sim} \left(\coprod_{S \in (T) \setminus \{T\}} N^e(S) \right) \otimes_{\text{Aut}(T)} \text{Aut}(C) \\ \downarrow \text{colim} \\ \coprod_{[T]} \left(\bigotimes_{v \in V_{G,\mathcal{P}}(T)} \mathcal{P}(T_v) \otimes Q[u]^{\vee_{G,in}(T)} \right) \otimes_{\text{Aut}(T)} \text{Aut}(C) \end{array}$$

where $Q[u]^{\vee_{G,in}(T)}$ is the source of the pushout product map defined in Definition 6.15.

Lastly, this left-side map is induced, via the naturality of Kan extensions, by an inclusion of categories, in particular the product of multiple inclusions of categories, each corresponding to the inclusion of a punctured cube into the full cube. Thus, after taking colimits, we have that the left-side map in Diagram (6.17) is in fact (multiple copies of) the pushout-product maps

$$[u]^{\square_{\vee_{G,in}(T)}} : Q[u]^{\vee_{G,in}(T)} \rightarrow \bigotimes_{v \in V_{G,in}(T)} Y_1(T_v),$$

as desired. \square

References

- BM03 [1] C. Berger and I. Moerdijk. Axiomatic homotopy theory for operads. Commentarii Mathematici Helvetici, 78:805–831, 2003.
- McL [2] S. Mac Lane. Categories for the working mathematician, volume 5 of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1998.
- Pe16b [3] L. A. Pereira. Equivariant dendroidal sets. Available at: <http://www.faculty.virginia.edu/luisalex/>, 2016.
- We12 [4] I. Weiss. Broad posets, trees, and the dendroidal category. Available at: <https://arxiv.org/abs/1201.3987>, 2012.