

# Genuine equivariant operads

## An Elmendorf-Piacenza theorem for operads

Luís Pereira

University of Notre Dame

9/28/2017

# Table of contents

Operads and equivariant operads

Elmendorf-Piacenza theorem

# Table of contents

Operads and equivariant operads

Elmendorf-Piacenza theorem

Trees and equivariant trees

# Table of contents

Operads and equivariant operads

Elmendorf-Piacenza theorem

Trees and equivariant trees

Genuine equivariant operads

# Table of contents

Operads and equivariant operads

Elmendorf-Piacenza theorem

Trees and equivariant trees

Genuine equivariant operads

## Definition

An *operad*  $\mathcal{O}$  consists of:

- ▶  $\Sigma_n$ -spaces  $\mathcal{O}(n)$  for  $n \geq 0$ ;



## Definition

An *operad*  $\mathcal{O}$  consists of:

- ▶  $\Sigma_n$ -spaces  $\mathcal{O}(n)$  for  $n \geq 0$ ;
- ▶ composition maps

$$\begin{aligned} \mathcal{O}(n) \times \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_n) &\rightarrow \mathcal{O}(k_1 + \dots + k_n) \\ f, g_1, \dots, g_n &\mapsto f(g_1, \dots, g_n) \end{aligned}$$



## Definition

An *operad*  $\mathcal{O}$  consists of:

- ▶  $\Sigma_n$ -spaces  $\mathcal{O}(n)$  for  $n \geq 0$ ;
- ▶ composition maps

$$\begin{aligned} \mathcal{O}(n) \times \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_n) &\rightarrow \mathcal{O}(k_1 + \dots + k_n) \\ f, g_1, \dots, g_n &\mapsto f(g_1, \dots, g_n) \end{aligned}$$

- ▶ satisfying associativity, unit and symmetry conditions.



## Definition

An *operad*  $\mathcal{O}$  consists of:

- ▶  $\Sigma_n$ -spaces  $\mathcal{O}(n)$  for  $n \geq 0$ ;
- ▶ composition maps

$$\mathcal{O}(n) \times \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_n) \rightarrow \mathcal{O}(k_1 + \dots + k_n)$$

$$f, g_1, \dots, g_n \mapsto f(g_1, \dots, g_n)$$

- ▶ satisfying associativity, unit and symmetry conditions.

## Example (Depicting composition)

$$\mathcal{O}(3) \times \mathcal{O}(1) \times \mathcal{O}(0) \times \mathcal{O}(3) \rightarrow \mathcal{O}(4)$$

$$f, g_1, g_2, g_3 \mapsto f(g_1, g_2, g_3)$$



## Definition

An *operad*  $\mathcal{O}$  consists of:


- ▶  $\Sigma_n$ -spaces  $\mathcal{O}(n)$  for  $n \geq 0$ ;
- ▶ composition maps

$$\begin{aligned} \mathcal{O}(n) \times \mathcal{O}(k_1) \times \dots \times \mathcal{O}(k_n) &\rightarrow \mathcal{O}(k_1 + \dots + k_n) \\ f, g_1, \dots, g_n &\mapsto f(g_1, \dots, g_n) \end{aligned}$$

- ▶ satisfying associativity, unit and symmetry conditions.

## Example (Depicting composition)

$$\begin{aligned} \mathcal{O}(3) \times \mathcal{O}(1) \times \mathcal{O}(0) \times \mathcal{O}(3) &\rightarrow \mathcal{O}(4) \\ f, g_1, g_2, g_3 &\mapsto f(g_1, g_2, g_3) \end{aligned}$$



## Definition

An *algebra* over an operad  $\mathcal{O}$  is a space  $X$  with compatible maps

$$\mathcal{O}(n) \times X^{\times n} \rightarrow X$$

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Ass</b>	$\text{Ass}(n) = \Sigma_n$	associative monoids
<b>Com</b>	$\text{Com}(n) = *$	commutative monoids

## Definition

An *algebra* over an operad  $\mathcal{O}$  is a space  $X$  with compatible maps

$$\mathcal{O}(n) \times X^{\times n} \rightarrow X$$

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Ass</b>	$\text{Ass}(n) = \Sigma_n$	associative monoids
<b>Com</b>	$\text{Com}(n) = *$	commutative monoids
$E_\infty$ -operads	$\mathcal{O}(n) \simeq E\Sigma_n$ i.e. $\Sigma_n$ -free and contractible	homotopy commutative monoids

## Definition

An *algebra* over an operad  $\mathcal{O}$  is a space  $X$  with compatible maps

$$\mathcal{O}(n) \times X^{\times n} \rightarrow X$$

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Ass</b>	$\text{Ass}(n) = \Sigma_n$	associative monoids
<b>Com</b>	$\text{Com}(n) = *$	commutative monoids
$E_\infty$ -operads	$\mathcal{O}(n) \simeq E\Sigma_n$ i.e. $\Sigma_n$ -free and contractible	homotopy commutative monoids

# Equivariant Operads

## Definition

A  $G$ -operad  $\mathcal{O}$  is an operad in  $\mathbf{Top}^G$  (also, a  $G$ -object in operads).

- ▶  $\mathcal{O}(n)$  is a  $G \times \Sigma_n$ -space;
- ▶ composition maps are  $G$ -equivariant.

# Equivariant Operads

## Definition

A  $G$ -operad  $\mathcal{O}$  is an operad in  $\text{Top}^G$  (also, a  $G$ -object in operads).

- ▶  $\mathcal{O}(n)$  is a  $G \times \Sigma_n$ -space;
- ▶ composition maps are  $G$ -equivariant.

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Com</b>	$\text{Com}(n) = *$	commutative $G$ -monoids

# Equivariant Operads

## Definition

A  $G$ -operad  $\mathcal{O}$  is an operad in  $\text{Top}^G$  (also, a  $G$ -object in operads).

- ▶  $\mathcal{O}(n)$  is a  $G \times \Sigma_n$ -space;
- ▶ composition maps are  $G$ -equivariant.

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Com</b>	$\text{Com}(n) = *$	commutative $G$ -monoids
$G$ -trivial $E_\infty$	$\mathcal{O}(n) \simeq E\Sigma_n$ with $G$ -trivial action	<b>not</b> the homotopy commutative $G$ -monoids



# Equivariant Operads

## Definition

A  $G$ -operad  $\mathcal{O}$  is an operad in  $\text{Top}^G$  (also, a  $G$ -object in operads).

- ▶  $\mathcal{O}(n)$  is a  $G \times \Sigma_n$ -space;
- ▶ composition maps are  $G$ -equivariant.

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Com</b>	$\text{Com}(n) = *$	commutative $G$ -monoids
$G$ -trivial $E_\infty$	$\mathcal{O}(n) \simeq E\Sigma_n$ with $G$ -trivial action	<b>not</b> the homotopy commutative $G$ -monoids
$G$ - $E_\infty$	$\mathcal{O}(n) \simeq E\mathcal{F}_n^\Gamma$	homotopy commutative $G$ -monoids

# Equivariant Operads

## Definition

A  $G$ -operad  $\mathcal{O}$  is an operad in  $\text{Top}^G$  (also, a  $G$ -object in operads).

- ▶  $\mathcal{O}(n)$  is a  $G \times \Sigma_n$ -space;
- ▶ composition maps are  $G$ -equivariant.

## Example

$\mathcal{O}$	$\mathcal{O}(n)$	$\mathcal{O}$ -algebras
<b>Com</b>	$\text{Com}(n) = *$	commutative $G$ -monoids
$G$ -trivial $E_\infty$	$\mathcal{O}(n) \simeq E\Sigma_n$ with $G$ -trivial action	<b>not</b> the homotopy commutative $G$ -monoids
$G$ - $E_\infty$	$\mathcal{O}(n) \simeq E\mathcal{F}_n^\Gamma$	homotopy commutative $G$ -monoids

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

For  $\Gamma \leq G \times \Sigma_n$

- ▶ A  $G$ -trivial  $E_\infty$  operad has fixed points

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \leq G \\ \emptyset & \text{otherwise} \end{cases}$$

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

For  $\Gamma \leq G \times \Sigma_n$

- ▶ A  $G$ -trivial  $E_\infty$  operad has fixed points

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \leq G \\ \emptyset & \text{otherwise} \end{cases}$$

- ▶ A  $G$ - $E_\infty$  operad has fixed points

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \cap \Sigma_n = * \\ \emptyset & \text{otherwise} \end{cases}$$

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

For  $\Gamma \leq G \times \Sigma_n$

- ▶ A  $G$ -trivial  $E_\infty$  operad has fixed points


$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \leq G \\ \emptyset & \text{otherwise} \end{cases}$$

- ▶ A  $G$ - $E_\infty$  operad has fixed points

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \cap \Sigma_n = * \\ \emptyset & \text{otherwise} \end{cases}$$




$$\Gamma \leq G$$



$$\Gamma \cap \Sigma_n = *$$




$$\Gamma \leq G$$



$$\Gamma \cap \Sigma_n = *$$



$$\Gamma \leq G$$




$$\Gamma \cap \Sigma_n = *$$






$$\Gamma \leq G$$




$$\Gamma \cap \Sigma_n = *$$

### Definition

$\Gamma \leq G \times \Sigma_n$  such that  $\Gamma \cap \Sigma_n = *$  is called a *graph subgroup*.



$$\Gamma \leq G$$



$$\Gamma \cap \Sigma_n = *$$


## Definition

$\Gamma \leq G \times \Sigma_n$  such that  $\Gamma \cap \Sigma_n = *$  is called a *graph subgroup*.

- ▶ These are graphs of partial homomorphisms  $G \geq H \rightarrow \Sigma_n$ ;



$$\Gamma \leq G$$



$$\Gamma \cap \Sigma_n = *$$


## Definition

$\Gamma \leq G \times \Sigma_n$  such that  $\Gamma \cap \Sigma_n = *$  is called a *graph subgroup*.

- ▶ These are graphs of partial homomorphisms  $G \geq H \rightarrow \Sigma_n$ ;
- ▶ Correspond to  $H$ -sets  $X$  with  $n$  elements;



$$\Gamma \leq G$$




$$\Gamma \cap \Sigma_n = *$$


## Definition

$\Gamma \leq G \times \Sigma_n$  such that  $\Gamma \cap \Sigma_n = *$  is called a *graph subgroup*.

- ▶ These are graphs of partial homomorphisms  $G \geq H \rightarrow \Sigma_n$ ;
- ▶ Correspond to  $H$ -sets  $X$  with  $n$  elements;
- ▶ Given  $H$ -set  $X$ , write  $\Gamma_X \leq G \times \Sigma_n$  for the graph subgroup.



$$\Gamma \leq G$$



$$\Gamma \cap \Sigma_n = *$$

## Definition

$\Gamma \leq G \times \Sigma_n$  such that  $\Gamma \cap \Sigma_n = *$  is called a *graph subgroup*.

- ▶ These are graphs of partial homomorphisms  $G \supseteq H \rightarrow \Sigma_n$ ;
- ▶ Correspond to  $H$ -sets  $X$  with  $n$  elements;
- ▶ Given  $H$ -set  $X$ , write  $\Gamma_X \leq G \times \Sigma_n$  for the graph subgroup.

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

$G$ -trivial $E_\infty$		$G$ - $E_\infty$	
$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \leq G$ otherwise	$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \cap \Sigma_n = *$ otherwise

- $G$ - $E_\infty$  is “as contractible as possible while being  $\Sigma_n$ -free”;

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

$G$ -trivial $E_\infty$		$G$ - $E_\infty$	
$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \leq G$ otherwise	$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \cap \Sigma_n = *$ otherwise

- ▶  $G$ - $E_\infty$  is “as contractible as possible while being  $\Sigma_n$ -free”;
- ▶  $G$ -trivial  $E_\infty$  is “contractible only for graphs of trivial  $H$ -sets”;

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

$G$ -trivial $E_\infty$		$G$ - $E_\infty$	
$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \leq G$ otherwise	$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \cap \Sigma_n = *$ otherwise

- ▶  $G$ - $E_\infty$  is “as contractible as possible while being  $\Sigma_n$ -free”;
- ▶  $G$ -trivial  $E_\infty$  is “contractible only for graphs of trivial  $H$ -sets”;
- ▶ both  $\Sigma_n$ -free;



# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

$G$ -trivial $E_\infty$		$G$ - $E_\infty$	
$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \leq G$ otherwise	$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \cap \Sigma_n = *$ otherwise

- ▶  $G$ - $E_\infty$  is “as contractible as possible while being  $\Sigma_n$ -free”;
- ▶  $G$ -trivial  $E_\infty$  is “contractible only for graphs of trivial  $H$ -sets”;
- ▶ both  $\Sigma_n$ -free;
- ▶ both  $G$ -contractible.

# $G$ -trivial $E_\infty$ vs. $G$ - $E_\infty$

$G$ -trivial $E_\infty$		$G$ - $E_\infty$	
$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \leq G$ otherwise	$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * \\ \emptyset \end{cases}$	$\Gamma \cap \Sigma_n = *$ otherwise

- ▶  $G$ - $E_\infty$  is “as contractible as possible while being  $\Sigma_n$ -free”;
- ▶  $G$ -trivial  $E_\infty$  is “contractible only for graphs of trivial  $H$ -sets”;
- ▶ both  $\Sigma_n$ -free;
- ▶ both  $G$ -contractible.

# Why do graph subgroups matter?

Given:

- ▶ (strict) commutative  $G$ -ring spectrum  $R$ , i.e. Com-algebra;
- ▶ finite  $G$ -set  $X$  with  $n$  elements;

# Why do graph subgroups matter?

Given:

- ▶ (strict) commutative  $G$ -ring spectrum  $R$ , i.e. Com-algebra;
- ▶ finite  $G$ -set  $X$  with  $n$  elements;

there are  $G$ -equivariant *norm maps*

$$N^X R \rightarrow R \quad \text{or} \quad R^{\wedge n} \rightarrow R$$



## Why do graph subgroups matter?

Given:

- ▶ (strict) commutative  $G$ -ring spectrum  $R$ , i.e. Com-algebra;
- ▶ finite  $G$ -set  $X$  with  $n$  elements;

there are  $G$ -equivariant *norm maps*

$$N^X R \rightarrow R \quad \text{or} \quad R^{\wedge n} \rightarrow R$$

**Note:** The  $G$ -action on  $N^X R$  is given by the  $\Gamma_X$ -action on  $R^{\wedge n}$ .



## Why do graph subgroups matter?

Given:

- ▶ (strict) commutative  $G$ -ring spectrum  $R$ , i.e. Com-algebra;
- ▶ finite  $G$ -set  $X$  with  $n$  elements;

there are  $G$ -equivariant *norm maps*

$$N^X R \rightarrow R \quad \text{or} \quad R^{\wedge n} \rightarrow R$$

**Note:** The  $G$ -action on  $N^X R$  is given by the  $\Gamma_X$ -action on  $R^{\wedge n}$ .

Replacing Com with a general operad  $\mathcal{O}$  get

$$\mathcal{O}(X)^G \wedge N^X R \rightarrow R \quad \text{or} \quad \mathcal{O}(n)^{\Gamma_X} \wedge R^{\wedge n} \rightarrow R$$

# Why do graph subgroups matter?

Given:

- ▶ (strict) commutative  $G$ -ring spectrum  $R$ , i.e. Com-algebra;
- ▶ finite  $G$ -set  $X$  with  $n$  elements;

there are  $G$ -equivariant *norm maps*

$$N^X R \rightarrow R \quad \text{or} \quad R^{\wedge n} \rightarrow R$$

**Note:** The  $G$ -action on  $N^X R$  is given by the  $\Gamma_X$ -action on  $R^{\wedge n}$ .

Replacing Com with a general operad  $\mathcal{O}$  get

$$\mathcal{O}(X)^G \wedge N^X R \rightarrow R \quad \text{or} \quad \mathcal{O}(n)^{\Gamma_X} \wedge R^{\wedge n} \rightarrow R$$

i.e.  $\mathcal{O}(X)^G = \mathcal{O}(n)^{\Gamma_X}$  is “the space of norm maps for the  $G$ -set  $X$ ”.

## Why do graph subgroups matter?

Given:

- ▶ (strict) commutative  $G$ -ring spectrum  $R$ , i.e. Com-algebra;
- ▶ finite  $G$ -set  $X$  with  $n$  elements;

there are  $G$ -equivariant *norm maps*

$$N^X R \rightarrow R \quad \text{or} \quad R^{\wedge n} \rightarrow R$$

**Note:** The  $G$ -action on  $N^X R$  is given by the  $\Gamma_X$ -action on  $R^{\wedge n}$ .

Replacing Com with a general operad  $\mathcal{O}$  get

$$\mathcal{O}(X)^G \wedge N^X R \rightarrow R \quad \text{or} \quad \mathcal{O}(n)^{\Gamma_X} \wedge R^{\wedge n} \rightarrow R$$

i.e.  $\mathcal{O}(X)^G = \mathcal{O}(n)^{\Gamma_X}$  is “the space of norm maps for the  $G$ -set  $X$ ”.



# Why do graph subgroups matter?

## Hence:

- Com has *unique* norm maps for each  $H$ -set  $X$ ;
- $G\text{-}E_\infty$ -operads have *homotopy unique* norm maps for each  $H$ -set  $X$ .

## Theorem (Costenoble-Waner)

$G\text{-}E_\infty$ -operads encode  $G$ -ring spectra with all norm maps and equivariant infinite loop spaces.

# Why do graph subgroups matter?

## Hence:

- ▶ Com has *unique* norm maps for each  $H$ -set  $X$ ;
- ▶  $G$ - $E_\infty$ -operads have *homotopy unique* norm maps for each  $H$ -set  $X$ .

## Theorem (Costenoble-Waner)

*$G$ - $E_\infty$ -operads encode  $G$ -ring spectra with all norm maps and equivariant infinite loop spaces.*

On the other hand,  $G$ -trivial  $E_\infty$ -operads encode  $G$ -ring spectra without non-trivial norm maps.

# Why do graph subgroups matter?

## Hence:

- ▶ Com has *unique* norm maps for each  $H$ -set  $X$ ;
- ▶  $G$ - $E_\infty$ -operads have *homotopy unique* norm maps for each  $H$ -set  $X$ .

## Theorem (Costenoble-Waner)

$G$ - $E_\infty$ -operads encode  $G$ -ring spectra with all norm maps and equivariant infinite loop spaces.

On the other hand,  $G$ -trivial  $E_\infty$ -operads encode  $G$ -ring spectra without non-trivial norm maps.

## Weak equivalences of equivariant operads

**Upshot:** weak equivalences of  $G$ -operads should account for norm maps.

### Definition

A map of  $G$ -operads  $\mathcal{O} \rightarrow \mathcal{O}'$  is called a *graph equivalence* if the maps

$$\mathcal{O}(n)^{\Gamma} \xrightarrow{\sim} \mathcal{O}'(n)^{\Gamma}, \quad \Gamma \cap \Sigma_n = *$$

are weak equivalences.

## Weak equivalences of equivariant operads

**Upshot:** weak equivalences of  $G$ -operads should account for norm maps.

### Definition

A map of  $G$ -operads  $\mathcal{O} \rightarrow \mathcal{O}'$  is called a *graph equivalence* if the maps

$$\mathcal{O}(n)^{\Gamma} \xrightarrow{\sim} \mathcal{O}'(n)^{\Gamma}, \quad \Gamma \cap \Sigma_n = *$$

are weak equivalences.

- These are more restrictive than “genuine equivalences”

$$\mathcal{O}(n)^{\Gamma} \xrightarrow{\sim} \mathcal{O}'(n)^{\Gamma}, \quad \Gamma \leq G$$

## Weak equivalences of equivariant operads

**Upshot:** weak equivalences of  $G$ -operads should account for norm maps.

### Definition

A map of  $G$ -operads  $\mathcal{O} \rightarrow \mathcal{O}'$  is called a *graph equivalence* if the maps

$$\mathcal{O}(n)^\Gamma \xrightarrow{\sim} \mathcal{O}'(n)^\Gamma, \quad \Gamma \cap \Sigma_n = *$$

are weak equivalences.

- ▶ These are more restrictive than “genuine equivalences”

$$\mathcal{O}(n)^\Gamma \xrightarrow{\sim} \mathcal{O}'(n)^\Gamma, \quad \Gamma \leq G$$

- ▶ Graph equivalences distinguish between  $G$ -trivial  $E_\infty$  and  $G$ - $E_\infty$ , while genuine equivalences do not.

## Weak equivalences of equivariant operads

**Upshot:** weak equivalences of  $G$ -operads should account for norm maps.

### Definition

A map of  $G$ -operads  $\mathcal{O} \rightarrow \mathcal{O}'$  is called a *graph equivalence* if the maps

$$\mathcal{O}(n)^\Gamma \xrightarrow{\sim} \mathcal{O}'(n)^\Gamma, \quad \Gamma \cap \Sigma_n = *$$

are weak equivalences.

- ▶ These are more restrictive than “genuine equivalences”

$$\mathcal{O}(n)^\Gamma \xrightarrow{\sim} \mathcal{O}'(n)^\Gamma, \quad \Gamma \leq G$$

- ▶ Graph equivalences distinguish between  $G$ -trivial  $E_\infty$  and  $G$ - $E_\infty$ , while genuine equivalences do not.

## Definition (Blumberg-Hill)

A  $G$ -operad  $\mathcal{O}$  is called a  $N_\infty$ -operad if

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \in \mathcal{F}_n \\ \emptyset & \text{otherwise} \end{cases}$$

where each  $\mathcal{F}_n$  is a family of graph subgroups of  $G \times \Sigma_n$ .

- $N_\infty$ -operads are  $G$ -operads “with only some norm maps”;



## Definition (Blumberg-Hill)

A  $G$ -operad  $\mathcal{O}$  is called a  $N_\infty$ -operad if

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \in \mathcal{F}_n \\ \emptyset & \text{otherwise} \end{cases}$$

where each  $\mathcal{F}_n$  is a family of graph subgroups of  $G \times \Sigma_n$ .

- ▶  $N_\infty$ -operads are  $G$ -operads “with only some norm maps”;




## Definition (Blumberg-Hill)

A  $G$ -operad  $\mathcal{O}$  is called a  $N_\infty$ -operad if

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \in \mathcal{F}_n \\ \emptyset & \text{otherwise} \end{cases}$$

where each  $\mathcal{F}_n$  is a family of graph subgroups of  $G \times \Sigma_n$ .

- ▶  $N_\infty$ -operads are  $G$ -operads “with only some norm maps”;



- ▶  $\mathcal{F} = \{\mathcal{F}_n\}$  is not arbitrary.  
Must satisfy *indexing system* closure conditions (BH);


## Definition (Blumberg-Hill)

A  $G$ -operad  $\mathcal{O}$  is called a  $N_\infty$ -operad if

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \in \mathcal{F}_n \\ \emptyset & \text{otherwise} \end{cases}$$

where each  $\mathcal{F}_n$  is a family of graph subgroups of  $G \times \Sigma_n$ .

- ▶  $N_\infty$ -operads are  $G$ -operads “with only some norm maps”;



- ▶  $\mathcal{F} = \{\mathcal{F}_n\}$  is not arbitrary.  
Must satisfy *indexing system* closure conditions (BH);
- ▶ Building  $N_\infty$ -operads for all indexing systems  $\mathcal{F}$  is not trivial.


## Definition (Blumberg-Hill)

A  $G$ -operad  $\mathcal{O}$  is called a  $N_\infty$ -operad if

$$\mathcal{O}(n)^\Gamma \simeq \begin{cases} * & \Gamma \in \mathcal{F}_n \\ \emptyset & \text{otherwise} \end{cases}$$

where each  $\mathcal{F}_n$  is a family of graph subgroups of  $G \times \Sigma_n$ .

- ▶  $N_\infty$ -operads are  $G$ -operads “with only some norm maps”;



- ▶  $\mathcal{F} = \{\mathcal{F}_n\}$  is not arbitrary.  
Must satisfy *indexing system* closure conditions (BH);
- ▶ Building  $N_\infty$ -operads for all indexing systems  $\mathcal{F}$  is not trivial.

# Elmendorf-Piacenza theorem

## Definition

The *orbital category*  $O_G$  is the category of orbital  $G$ -sets  $G/H$ .

## Theorem (Elmendorf-Piacenza)

*There is a Quillen equivalence*

$$\begin{array}{ccc} \mathrm{Top}^{O_G^{op}} & \xrightleftharpoons{\quad} & \mathrm{Top}^G \\ (G/H \mapsto Y(G/H)) & \longmapsto & Y(G) \\ (G/H \mapsto X^H) & \xleftarrow{\quad} & X \end{array}$$

# Elmendorf-Piacenza theorem

## Definition

The *orbital category*  $O_G$  is the category of orbital  $G$ -sets  $G/H$ .

## Theorem (Elmendorf-Piacenza)

There is a Quillen equivalence

$$\begin{array}{ccc} \mathrm{Top}^{O_G^{op}} & \xrightleftharpoons{\quad} & \mathrm{Top}^G \\ (G/H \mapsto Y(G/H)) & \longmapsto & Y(G) \\ (G/H \mapsto X^H) & \longleftarrow & X \end{array}$$

► w.e.s on  $\mathrm{Top}^{O_G^{op}}$  are levelwise;



# Elmendorf-Piacenza theorem

## Definition

The *orbital category*  $O_G$  is the category of orbital  $G$ -sets  $G/H$ .

## Theorem (Elmendorf-Piacenza)

There is a Quillen equivalence

$$\begin{array}{ccc} \mathrm{Top}^{op}_G & \xrightleftharpoons{\quad} & \mathrm{Top}^G \\ (G/H \mapsto Y(G/H)) & \longmapsto & Y(G) \\ (G/H \mapsto X^H) & \longleftarrow & X \end{array}$$

- ▶ w.e.s on  $\mathrm{Top}^{op}_G$  are levelwise;
- ▶ w.e.s on  $\mathrm{Top}^G$  are genuine, i.e. detected on all fixed points.

# Elmendorf-Piacenza theorem

## Definition

The *orbital category*  $O_G$  is the category of orbital  $G$ -sets  $G/H$ .

## Theorem (Elmendorf-Piacenza)

There is a Quillen equivalence

$$\begin{array}{ccc} \mathrm{Top}^{O_G^{op}} & \xrightleftharpoons{\quad} & \mathrm{Top}^G \\ (G/H \mapsto Y(G/H)) & \longmapsto & Y(G) \\ (G/H \mapsto X^H) & \longleftarrow & X \end{array}$$

- ▶ w.e.s on  $\mathrm{Top}^{O_G^{op}}$  are levelwise;
- ▶ w.e.s on  $\mathrm{Top}^G$  are genuine, i.e. detected on all fixed points.



# Generalized Elmendorf-Piacenza theorems

## Theorem (Bergner, Bergner-Gutierrez)

*More generally, there are Quillen equivalences*

$$\blacktriangleright \text{sCat}^{O_G^{op}} \xrightleftharpoons{\quad} \text{sCat}^G$$

*for sCat the category of simplicial categories;*

# Generalized Elmendorf-Piacenza theorems

## Theorem (Bergner, Bergner-Gutierrez)

*More generally, there are Quillen equivalences*

- ▶  $\text{sCat}^{O_G^{op}} \rightleftarrows \text{sCat}^G$   
for  $\text{sCat}$  the category of simplicial categories;
- ▶  $\text{sOp}^{O_G^{op}} \rightleftarrows \text{sOp}^G$   
for  $\text{sOp}$  the category of simplicial operads.

# Generalized Elmendorf-Piacenza theorems

## Theorem (Bergner, Bergner-Gutierrez)

*More generally, there are Quillen equivalences*

- ▶  $\text{sCat}^{O_G^{op}} \rightleftarrows \text{sCat}^G$   
for  $\text{sCat}$  the category of simplicial categories;
- ▶  $\text{sOp}^{O_G^{op}} \rightleftarrows \text{sOp}^G$   
for  $\text{sOp}$  the category of simplicial operads.

## Remark

These results use genuine equivalences, which in the  $\text{sOp}$  case **ignore all non trivial norm maps.**



## Generalized Elmendorf-Piacenza theorems

### Theorem (Bergner, Bergner-Gutierrez)

*More generally, there are Quillen equivalences*

- ▶  $\text{sCat}^{O_G^{op}} \rightleftarrows \text{sCat}^G$   
for  $\text{sCat}$  the category of simplicial categories;
- ▶  $\text{sOp}^{O_G^{op}} \rightleftarrows \text{sOp}^G$   
for  $\text{sOp}$  the category of simplicial operads.

### Remark

These results use genuine equivalences, which in the  $\text{sOp}$  case **ignore all non trivial norm maps**.

# The plan

**Main Goal:** Formulate an Elmendorf-Piacenza result for operads with graph equivalences.

**Issue:** The levels of  $\underline{\mathcal{O}} \in \mathbf{sOp}_G^{op}$  only encode trivial norm maps.

# The plan

**Main Goal:** Formulate an Elmendorf-Piacenza result for operads with graph equivalences.

**Issue:** The levels of  $\underline{\mathcal{O}} \in \mathbf{sOp}^{Op}_G$  only encode trivial norm maps.

**Idea:** Replace  $\mathbf{sOp}^{Op}_G$  with a (larger) category where the levels of objects encode all norm maps.

# The plan

**Main Goal:** Formulate an Elmendorf-Piacenza result for operads with graph equivalences.

**Issue:** The levels of  $\underline{\mathcal{O}} \in \mathbf{sOp}_G^{op}$  only encode trivial norm maps.

**Idea:** Replace  $\mathbf{sOp}_G^{op}$  with a (larger) category where the levels of objects encode all norm maps.

$$\mathbf{sOp}_G \rightleftarrows \mathbf{sOp}^G$$

$\mathbf{sOp}_G$  is the category of *genuine equivariant operads*.

# The plan

**Main Goal:** Formulate an Elmendorf-Piacenza result for operads with graph equivalences.

**Issue:** The levels of  $\underline{\mathcal{O}} \in \mathbf{sOp}_G^{op}$  only encode trivial norm maps.

**Idea:** Replace  $\mathbf{sOp}_G^{op}$  with a (larger) category where the levels of objects encode all norm maps.

$$\mathbf{sOp}_G \rightleftarrows \mathbf{sOp}^G$$

$\mathbf{sOp}_G$  is the category of *genuine equivariant operads*.

**The key:** Each genuine equivariant operad  $\mathcal{P} \in \mathbf{sOp}_G$  will come with composition maps.

To define these, need to understand how composition interacts with norm maps for each  $\mathcal{O} \in \mathbf{sOp}^G$ .



# The plan

**Main Goal:** Formulate an Elmendorf-Piacenza result for operads with graph equivalences.

**Issue:** The levels of  $\underline{\mathcal{O}} \in \mathbf{sOp}_G^{op}$  only encode trivial norm maps.

**Idea:** Replace  $\mathbf{sOp}_G^{op}$  with a (larger) category where the levels of objects encode all norm maps.

$$\mathbf{sOp}_G \xrightleftharpoons{\quad} \mathbf{sOp}^G$$

$\mathbf{sOp}_G$  is the category of *genuine equivariant operads*.

**The key:** Each genuine equivariant operad  $\mathcal{P} \in \mathbf{sOp}_G$  will come with composition maps.

To define these, need to understand how composition interacts with norm maps for each  $\mathcal{O} \in \mathbf{sOp}^G$ .

**Solution:** Those interactions are encoded by *equivariant trees*.

# The plan

**Main Goal:** Formulate an Elmendorf-Piacenza result for operads with graph equivalences.

**Issue:** The levels of  $\mathcal{O} \in \mathbf{sOp}_G^{op}$  only encode trivial norm maps.

**Idea:** Replace  $\mathbf{sOp}_G^{op}$  with a (larger) category where the levels of objects encode all norm maps.

$$\mathbf{sOp}_G \rightleftarrows \mathbf{sOp}^G$$

$\mathbf{sOp}_G$  is the category of *genuine equivariant operads*.

**The key:** Each genuine equivariant operad  $\mathcal{P} \in \mathbf{sOp}_G$  will come with composition maps.

To define these, need to understand how composition interacts with norm maps for each  $\mathcal{O} \in \mathbf{sOp}^G$ .

**Solution:** Those interactions are encoded by *equivariant trees*.

# The dendroidal category $\Omega$


## Definition (Moerdijk-Weiss)

A tree diagram  $T$  defines a colored operad such that:

- ▶ objects are the edges of  $T$ ;
- ▶ operations are generated by the nodes of  $T$ .

## Example

▶ objects:  $a, b, c, d, e, f, g, h, r$




# The dendroidal category $\Omega$

## Definition (Moerdijk-Weiss)

A tree diagram  $T$  defines a colored operad such that:

- ▶ objects are the edges of  $T$ ;
- ▶ operations are generated by the nodes of  $T$ .

## Example



- ▶ objects:  $a, b, c, d, e, f, g, h, r$

- ▶ generating operations:

$$ab \rightarrow f, \emptyset \rightarrow g, cde \rightarrow h, fgh \rightarrow r$$


# The dendroidal category $\Omega$

## Definition (Moerdijk-Weiss)

A tree diagram  $T$  defines a colored operad such that:

- ▶ objects are the edges of  $T$ ;
- ▶ operations are generated by the nodes of  $T$ .

## Example



- ▶ objects:  $a, b, c, d, e, f, g, h, r$
- ▶ generating operations:  
 $ab \rightarrow f, \emptyset \rightarrow g, cde \rightarrow h, fgh \rightarrow r$
- ▶ composite operations:  
 $abgh \rightarrow r, fh \rightarrow r, fgcd \rightarrow r, abh \rightarrow r$   
 $abgcde \rightarrow r, fcde \rightarrow r, abcde \rightarrow r$


# The dendroidal category $\Omega$

## Definition (Moerdijk-Weiss)

A tree diagram  $T$  defines a colored operad such that:

- ▶ objects are the edges of  $T$ ;
- ▶ operations are generated by the nodes of  $T$ .

## Example





- ▶ objects:  $a, b, c, d, e, f, g, h, r$
- ▶ generating operations:  
 $ab \rightarrow f, \emptyset \rightarrow g, cde \rightarrow h, fgh \rightarrow r$
- ▶ composite operations:  
 $abgh \rightarrow r, fh \rightarrow r, fgcd \rightarrow r, abh \rightarrow r$   
 $abgcde \rightarrow r, fcde \rightarrow r, abcde \rightarrow r$

## Definition (Moerdijk-Weiss)

The *dendroidal category*  $\Omega$  consists of trees and operad maps.


## Example



## Definition (Moerdijk-Weiss)

The *dendroidal category*  $\Omega$  consists of trees and operad maps.

## Example






## Definition (Moerdijk-Weiss)

The *dendroidal category*  $\Omega$  consists of trees and operad maps.


## Example



## Definition (Moerdijk-Weiss)

The *dendroidal category*  $\Omega$  consists of trees and operad maps.

## Example




$$\begin{array}{ccc}
 \mathrm{Hom}(T, \mathcal{O}) & \longrightarrow & \mathrm{Hom}(C, \mathcal{O}) \\
 \mathcal{O}(3) \times \mathcal{O}(2) \times \mathcal{O}(0) \times \mathcal{O}(3) & \longrightarrow & \mathcal{O}(5)
 \end{array}$$

## Definition (Moerdijk-Weiss)

The *dendroidal category*  $\Omega$  consists of trees and operad maps.

## Example




$$\begin{array}{ccc} \text{Hom}(T, \mathcal{O}) & \longrightarrow & \text{Hom}(C, \mathcal{O}) \\ \mathcal{O}(3) \times \mathcal{O}(2) \times \mathcal{O}(0) \times \mathcal{O}(3) & \longrightarrow & \mathcal{O}(5) \end{array}$$

Maps in  $\Omega$  encode operadic composition, associativity, etc.

## Definition (Moerdijk-Weiss)

The *dendroidal category*  $\Omega$  consists of trees and operad maps.

## Example



$$\begin{array}{ccc} \text{Hom}(T, \mathcal{O}) & \longrightarrow & \text{Hom}(C, \mathcal{O}) \\ \mathcal{O}(3) \times \mathcal{O}(2) \times \mathcal{O}(0) \times \mathcal{O}(3) & \longrightarrow & \mathcal{O}(5) \end{array}$$

Maps in  $\Omega$  encode operadic composition, associativity, etc.

# The category $\Omega_G$ of $G$ -trees

**Want:** A generalization  $\Omega_G$  of  $\Omega$  that encodes compositions of norm maps;

First guess: Consider  $\Omega^G$ , the category of  $G$ -objects in  $\Omega$ .

# The category $\Omega_G$ of $G$ -trees

**Want:** A generalization  $\Omega_G$  of  $\Omega$  that encodes compositions of norm maps;

**First guess:** Consider  $\Omega^G$ , the category of  $G$ -objects in  $\Omega$ .

## Remark

As it turns out  $\Omega^G \hookrightarrow \Omega_G$ , but  $\Omega^G \neq \Omega_G$ .

# The category $\Omega_G$ of $G$ -trees


**Want:** A generalization  $\Omega_G$  of  $\Omega$  that encodes compositions of norm maps;

**First guess:** Consider  $\Omega^G$ , the category of  $G$ -objects in  $\Omega$ .

## Remark

As it turns out  $\Omega^G \hookrightarrow \Omega_G$ , but  $\Omega^G \neq \Omega_G$ .


## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )



$$\mathrm{Hom}(C, \mathcal{O}) = \mathcal{O}(6)^{\Gamma_{\{a, b, a+2, a+1, b+1, a+3\}}}$$




## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )





$$\mathrm{Hom}(C, \mathcal{O}) = \mathcal{O}(6)^{\Gamma_{\{a, b, a+2, a+1, b+1, a+3\}}} = \mathcal{O}(G \amalg G/2)^G$$

## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )





$$\mathrm{Hom}(C, \mathcal{O}) = \mathcal{O}(6)^{\Gamma_{\{a,b,a+2,a+1,b+1,a+3\}}} = \mathcal{O}(G \amalg G/2)^G$$

## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )





$$\mathrm{Hom}(C, \mathcal{O}) = \mathcal{O}(6)^{\Gamma_{\{a, b, a+2, a+1, b+1, a+3\}}} = \mathcal{O}(G \amalg G/2)^G$$

## Example ( $G = \mathbb{Z}_4$ . Example in $\Omega^G$ )





## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )





$$\text{Hom}(T, \mathcal{O}) = \mathcal{O}(2)^{\Gamma_{\{c, c+1\}}} \times \mathcal{O}(3)^{\Gamma_{\{a, b, a+2\}}}$$

## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )







$$\text{Hom}(T, \mathcal{O}) = \mathcal{O}(2)^{\Gamma_{\{c, c+1\}}} \times \mathcal{O}(3)^{\Gamma_{\{a, b, a+2\}}} = \mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$$

## Example ( $G = \mathbb{Z}/4$ . Example in $\Omega^G$ )



$$\text{Hom}(T, \mathcal{O}) = \mathcal{O}(2)^{\Gamma_{\{c, c+1\}}} \times \mathcal{O}(3)^{\Gamma_{\{a, b, a+2\}}} = \mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$$




## Example ( $G = \mathbb{Z}/4$ . Map in $\Omega^G$ )


 $\text{Hom}(T, \mathcal{O})$ 

 $\text{Hom}(C, \mathcal{O})$







## Example ( $G = \mathbb{Z}/4$ . Map in $\Omega^G$ )


 $\text{Hom}(T, \mathcal{O})$ 

 $\text{Hom}(C, \mathcal{O})$ 
 $\mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$ 

 $\mathcal{O}(G \amalg G/2G)^G$

## Example ( $G = \mathbb{Z}/4$ . Map in $\Omega^G$ )


 $\text{Hom}(T, \mathcal{O})$ 
 $\text{Hom}(C, \mathcal{O})$ 
 $\mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$ 
 $\mathcal{O}(G \amalg G/2G)^G$

## Example ( $G = \mathbb{Z}_4$ . Map in $\Omega^G$ )

$$\mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G} \longrightarrow \mathcal{O}(G \amalg G/2G)^G$$

## Example (general $G$ , $K_1 \leq H \leq G$ , $K_2 \leq H \leq G$ )

$$\mathcal{O}(G/H)^G \times \mathcal{O}(H/K_1 \amalg H/K_2)^H \longrightarrow \mathcal{O}(G/K_1 \amalg G/K_2)^G$$



## Example ( $G = \mathbb{Z}/4$ . Map in $\Omega^G$ )

$$\mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G} \longrightarrow \mathcal{O}(G \amalg G/2G)^G$$

## Example (general $G$ , $K_1 \leq H \leq G$ , $K_2 \leq H \leq G$ )


$$\mathcal{O}(G/H)^G \times \mathcal{O}(H/K_1 \amalg H/K_2)^H \longrightarrow \mathcal{O}(G/K_1 \amalg G/K_2)^G$$

$$\Omega_G \neq \Omega^G$$



$$\text{Hom}(T, \mathcal{O}) = \mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$$



Example ( $G = \mathbb{Z}_4$ . Example not in  $\Omega^G$ )



$$\text{Hom}(S, \mathcal{O}) = \mathcal{O}(2G \amalg 2G/2G)^{2G}$$




$$\Omega_G \neq \Omega^G$$





$$\text{Hom}(T, \mathcal{O}) = \mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$$

Example ( $G = \mathbb{Z}_4$ . Example not in  $\Omega^G$ )





$$\text{Hom}(S, \mathcal{O}) = \mathcal{O}(2G \amalg 2G/2G)^{2G}$$

$$\Omega_G \neq \Omega^G$$



$$\text{Hom}(T, \mathcal{O}) = \mathcal{O}(G/2G)^G \times \mathcal{O}(2G \amalg 2G/2G)^{2G}$$

Example ( $G = \mathbb{Z}_4$ . Example not in  $\Omega^G$ )



$$\text{Hom}(S, \mathcal{O}) = \mathcal{O}(2G \amalg 2G/2G)^{2G}$$

## Definition (Bonventre-P.)

A  $G$ -tree is a  $G$ -object in the category of forests that is  $G$ -indecomposable.

$\Omega_G$  if the category of  $G$ -trees and  $G$ -operad maps.



## Definition (Bonventre-P.)

A  $G$ -tree is a  $G$ -object in the category of forests that is  $G$ -indecomposable.

$\Omega_G$  if the category of  $G$ -trees and  $G$ -operad maps.

- ▶ The orbital representation of a  $G$ -tree “always looks like a tree”.
- ▶ Maps in  $\Omega_G$  encode compositions of norm maps, as well as associativity, etc.

## Definition (Bonventre-P.)

A  $G$ -tree is a  $G$ -object in the category of forests that is  $G$ -indecomposable.

$\Omega_G$  if the category of  $G$ -trees and  $G$ -operad maps.

- ▶ The orbital representation of a  $G$ -tree “always looks like a tree”.
- ▶ Maps in  $\Omega_G$  encode compositions of norm maps, as well as associativity, etc.

# Genuine equivariant operads

## Definition (Bonventre-P.)

A *genuine equivariant operad*  $\mathcal{P} \in \mathbf{sOp}_G$  has levels  $\mathcal{P}(X)$  for each  $H$ -set  $X$ .

Further, the  $\mathcal{P}(X)$  mimic the combinatorics of the norm map spaces  $\mathcal{O}(n)^{\Gamma_X} = \mathcal{O}(X)^G$  of  $\mathcal{O} \in \mathbf{sOp}^G$ . Explicitly:

- ▶ for  $K \leq H$ , there are restriction maps  $\mathcal{P}(X) \rightarrow \mathcal{P}(X|_K)$ ;

# Genuine equivariant operads

## Definition (Bonventre-P.)

A *genuine equivariant operad*  $\mathcal{P} \in \mathbf{sOp}_G$  has levels  $\mathcal{P}(X)$  for each  $H$ -set  $X$ .

Further, the  $\mathcal{P}(X)$  mimic the combinatorics of the norm map spaces  $\mathcal{O}(n)^{\Gamma_X} = \mathcal{O}(X)^G$  of  $\mathcal{O} \in \mathbf{sOp}^G$ . Explicitly:

- ▶ for  $K \leq H$ , there are restriction maps  $\mathcal{P}(X) \rightarrow \mathcal{P}(X|_K)$ ;
- ▶ for  $g \in G$  there are isomorphisms  $\mathcal{P}(X) \simeq \mathcal{P}(gX)$ ;



# Genuine equivariant operads

## Definition (Bonventre-P.)

A *genuine equivariant operad*  $\mathcal{P} \in \mathbf{sOp}_G$  has levels  $\mathcal{P}(X)$  for each  $H$ -set  $X$ .

Further, the  $\mathcal{P}(X)$  mimic the combinatorics of the norm map spaces  $\mathcal{O}(n)^{\Gamma_X} = \mathcal{O}(X)^G$  of  $\mathcal{O} \in \mathbf{sOp}^G$ . Explicitly:

- ▶ for  $K \leq H$ , there are restriction maps  $\mathcal{P}(X) \rightarrow \mathcal{P}(X|_K)$ ;
- ▶ for  $g \in G$  there are isomorphisms  $\mathcal{P}(X) \simeq \mathcal{P}(gX)$ ;
- ▶ there are composition maps as encoded by  $G$ -trees.



# Genuine equivariant operads

## Definition (Bonventre-P.)

A *genuine equivariant operad*  $\mathcal{P} \in \mathbf{sOp}_G$  has levels  $\mathcal{P}(X)$  for each  $H$ -set  $X$ .

Further, the  $\mathcal{P}(X)$  mimic the combinatorics of the norm map spaces  $\mathcal{O}(n)^{\Gamma_X} = \mathcal{O}(X)^G$  of  $\mathcal{O} \in \mathbf{sOp}^G$ . Explicitly:

- ▶ for  $K \leq H$ , there are restriction maps  $\mathcal{P}(X) \rightarrow \mathcal{P}(X|_K)$ ;
- ▶ for  $g \in G$  there are isomorphisms  $\mathcal{P}(X) \simeq \mathcal{P}(gX)$ ;
- ▶ there are composition maps as encoded by  $G$ -trees.

## Composition in genuine equivariant operads

Example (general  $G$ ,  $K_1 \leq H \leq G$ ,  $K_2 \leq H \leq G$ )

$$\mathcal{O}(G/H)^G \times \mathcal{O}(H/K_1 \amalg H/K_2)^H \longrightarrow \mathcal{O}(G/K_1 \amalg G/K_2)^G$$

Hence, a genuine equivariant operad  $\mathcal{P} \in \text{Op}_G$  has maps

$$\mathcal{P}(G/H) \times \mathcal{P}(H/K_1 \amalg H/K_2) \rightarrow \mathcal{P}(G/K_1 \amalg G/K_2)$$

# Composition in genuine equivariant operads

Example (general  $G$ ,  $K_1 \leq H \leq G$ ,  $K_2 \leq H \leq G$ )

$$\mathcal{O}(G/H)^G \times \mathcal{O}(H/K_1 \amalg H/K_2)^H \longrightarrow \mathcal{O}(G/K_1 \amalg G/K_2)^G$$

Hence, a genuine equivariant operad  $\mathcal{P} \in \text{Op}_G$  has maps

$$\mathcal{P}(G/H) \times \mathcal{P}(H/K_1 \amalg H/K_2) \rightarrow \mathcal{P}(G/K_1 \amalg G/K_2)$$

Similarly, for  $L \leq K \leq H$  there are also maps

$$\mathcal{P}(H/K) \times \mathcal{P}(K/L) \rightarrow \mathcal{P}(H/L)$$





## Composition in genuine equivariant operads

Example (general  $G$ ,  $K_1 \leq H \leq G$ ,  $K_2 \leq H \leq G$ )

$$\mathcal{O}(G/H)^G \times \mathcal{O}(H/K_1 \amalg H/K_2)^H \longrightarrow \mathcal{O}(G/K_1 \amalg G/K_2)^G$$

Hence, a genuine equivariant operad  $\mathcal{P} \in \text{Op}_G$  has maps

$$\mathcal{P}(G/H) \times \mathcal{P}(H/K_1 \amalg H/K_2) \rightarrow \mathcal{P}(G/K_1 \amalg G/K_2)$$

Similarly, for  $L \leq K \leq H$  there are also maps

$$\mathcal{P}(H/K) \times \mathcal{P}(K/L) \rightarrow \mathcal{P}(H/L)$$

More generally, compositions in  $\mathcal{P} \in \text{Op}_G$  have the form

$$\begin{aligned}
 & \mathcal{P}(H/K_1 \amalg \cdots \amalg H/K_n) \times \\
 & \times \mathcal{P}(K_1/L_{11} \amalg \cdots \amalg K_1/L_{1m_1}) \times \cdots \times \mathcal{P}(K_n/L_{n1} \amalg \cdots \amalg K_n/L_{nm_n}) \\
 & \quad \downarrow \\
 & \mathcal{P}(H/L_{11} \amalg \cdots \amalg H/L_{1m_1} \amalg \cdots \amalg H/L_{n1} \amalg \cdots \amalg H/L_{nm_n})
 \end{aligned}$$

# Main result

## Theorem (Bonventre-P.)

*There is a Quillen equivalence*

$$\mathrm{sOp}_G \rightleftarrows \mathrm{sOp}^G$$

*where w.e.s on  $\mathrm{sOp}_G$  are levelwise and w.e.s on  $\mathrm{sOp}^G$  are graph equivalences.*

## Remarks:

- It follows that for  $\mathcal{O} \in \mathrm{sOp}^G$ ,  $\mathrm{ho}(\mathcal{O}) \in \mathrm{Op}_G$ ;

# Main result

## Theorem (Bonventre-P.)

*There is a Quillen equivalence*

$$\mathrm{sOp}_G \rightleftarrows \mathrm{sOp}^G$$

*where w.e.s on  $\mathrm{sOp}_G$  are levelwise and w.e.s on  $\mathrm{sOp}^G$  are graph equivalences.*

## Remarks:

- ▶ It follows that for  $\mathcal{O} \in \mathrm{sOp}^G$ ,  $\mathrm{ho}(\mathcal{O}) \in \mathrm{Op}_G$ ;
- ▶ (the proof of) this result can be used to build explicit models for the  $N\mathcal{F}$ -operads of Blumberg-Hill (this generalizes the construction of universal  $E\mathcal{F}$ -spaces).

# Main result

## Theorem (Bonventre-P.)

*There is a Quillen equivalence*

$$\mathrm{sOp}_G \rightleftarrows \mathrm{sOp}^G$$

*where w.e.s on  $\mathrm{sOp}_G$  are levelwise and w.e.s on  $\mathrm{sOp}^G$  are graph equivalences.*

### Remarks:

- ▶ It follows that for  $\mathcal{O} \in \mathrm{sOp}^G$ ,  $\mathrm{ho}(\mathcal{O}) \in \mathrm{Op}_G$ ;
- ▶ (the proof of) this result can be used to build explicit models for the  $N\mathcal{F}$ -operads of Blumberg-Hill (this generalizes the construction of universal  $E\mathcal{F}$ -spaces).

Thanks for listening.

- [BH15] Andrew J. Blumberg and Michael A. Hill. Operadic multiplications in equivariant spectra, norms, and transfers. *Adv. Math.*, 285:658–708, 2015.
- [BP17] Peter Bonventre and Luís A. Pereira. Genuine equivariant operads. arXiv preprint: 1707.02226, 2017.
- [CM11] Denis-Charles Cisinski and Ieke Moerdijk. Dendroidal sets as models for homotopy operads. *J. Topol.*, 4(2):257–299, 2011.
- [CM13a] Denis-Charles Cisinski and Ieke Moerdijk. Dendroidal Segal spaces and  $\infty$ -operads. *J. Topol.*, 6(3):675–704, 2013.
- [CM13b] Denis-Charles Cisinski and Ieke Moerdijk. Dendroidal sets and simplicial operads. *J. Topol.*, 6(3):705–756, 2013.
- [CW91] Steven R. Costenoble and Stefan Waner. Fixed set systems of equivariant infinite loop spaces. *Trans. Amer. Math. Soc.*, 326(2):485–505, 1991.
- [MW09] I. Moerdijk and I. Weiss. On inner Kan complexes in the category of dendroidal sets. *Adv. Math.*, 221(2):343–389, 2009.
- [Per17] Luís Alexandre Pereira. Equivariant dendroidal sets. arXiv preprint: 1702.08119, 2017.