

A short proof of Fermat's Last Theorem
Estimation of numerical errors

Rouben Rostamian

Department of Mathematics and Statistics
University of Maryland, Baltimore County
Baltimore, Maryland 21250

`rostamian@umbc.edu`

November 26, 2004

Overview

Overview of the material.

Graphics

Here we include three images, one each of PDF, PNG, and JPG types.



Outline of the talk

- Introduction

Outline of the talk

- Introduction
- Statement of the main theorem

Outline of the talk

- Introduction
- Statement of the main theorem
- Technical lemmata

Outline of the talk

- Introduction
- Statement of the main theorem
- Technical lemmata
- Proof of the main theorem

Outline of the talk

- Introduction
- Statement of the main theorem
- Technical lemmata
- Proof of the main theorem
- Conclusions

Some other slide

If you click here, you will jump to the slide labeled “intro”.

Clicking [here](#) will also take you to the “intro” slide.

Theorems and such

Definition

A triangle that has a right angle is called a *right triangle*.

Theorem

In a right triangle, the square of the hypotenuse equals the sum of the squares of the two other sides.

Proof.

We leave the proof as an exercise to our astute reader. We also suggest that the reader generalize the proof to non-Euclidean geometries. □

Title

This is a test.

V. Jikov, S. Kozlov and O. Olenik, Homogenization of differential operators and integral functionals, Springer, 1994.

Splitting a slide into columns

The line you are reading goes all the way across the slide. From the left margin to the right margin. Now we are going to split the slide into two columns.

Here is the first column. We put an itemized list in it.

- This is an item
- This is another item
- Yet another item

Here is the second column. We will put a picture in it.



The line you are reading goes all the way across the slide. From the left