The rank-four, eleven-element excluded minors for representability over $\mathbb{F}_5$

A computational classification
Revised exposition, 5 November 2026

Abstract

There are exactly 35 isomorphism classes of rank-four, eleven-element excluded minors for representability over $\mathbb{F}_5$. The proof is an exhaustive construction. Every such matroid has a representable ten-element deletion. For each possible deletion, we recover all nonrepresentable extensions whose other deletions are representable, then test their contractions. The essential observation is that, when the ground set has more than twice the rank many elements, compatible basis data for the single-element deletions already determine a matroid. This turns the extension search into an exact intersection of finite tables. We prove that each stage preserves every possible excluded minor, explain how the tables are constructed and exhausted, and give the computational results that complete the classification. Separate certificates verify nonrepresentability and all immediate minors of each of the 35 resulting matroids.

Contents

1 The theorem and the structure of its proof **2**

2 Step 1: finding every possible parent **3**

3 Step 2: recovering an extension from its deletions **4**

4 Constructing and exhausting the extension tables **6**

5 Step 3: testing contractions and certifying the list **9**

6 The computation and the proof of the theorem **11**

A Details of the representation and SAT encodings **12**

B Exact catalogue **14**

C Computational records and reproduction **18**

1 The theorem and the structure of its proof

A matroid is *quinary* if it is representable over $\mathbb{F}_5 = \mathbb{Z}/5\mathbb{Z}$. An *excluded minor* for quinary matroids is a nonquinary matroid every proper minor of which is quinary. We classify these obstructions at one specified rank and size.

Theorem 1.1 (Computational classification)**.** *There are exactly 35 isomorphism classes of rank-four, eleven-element excluded minors for representability over $\mathbb{F}_5$. Representatives $M_1, \ldots, M_{35}$ are specified in Appendix B.*

The defining condition has a useful finite form. A matroid M is an excluded minor precisely when

$$M \text{ is nonquinary}, \qquad M \setminus a \text{ and } M/a \text{ are quinary for every } a \in E(M).$$

Indeed, every proper minor is a minor of one of these immediate deletions or contractions, and field representability is closed under taking minors. For an eleven-element matroid there are therefore 22 immediate minors to consider. This criterion is easy to apply to a given matroid. The main difficulty is to find a finite list that is guaranteed to contain *every* matroid satisfying it.

We construct such a list by reversing deletion. Choose an element e of an unknown excluded minor M and put $N = M \setminus e$. The matroid N is quinary. We call it a *parent*, since the task is to recover M by adding e to it. The proof proceeds in three steps.

1. *Find all possible parents.* We prove that N must be a connected simple rank-four matroid on ten elements, and generate a complete quinary catalogue of these matroids (Section 2).

2. *Recover all possible extensions of each parent.* For a fixed N, we construct exactly the nonquinary extensions M for which every deletion $M \setminus a$ is quinary. The recovery theorem is proved in Section 3; its finite tables and their exhaustive solution are developed in Section 4.

3. *Impose the remaining minor conditions.* We test every contraction of every recovered matroid and identify isomorphic outputs. This leaves precisely the desired excluded minors (Section 5).

The second step is the heart of the argument. A quinary extension of N can be obtained by appending a column to a representation of N; the desired M, however, is nonquinary. Instead we construct its *different deletions* from representations, and require their basis data to agree wherever they overlap. A local-to-global lemma then shows that these compatible data form a matroid. We can thus enumerate nonrepresentable matroids using only finite lists of representable ones.

Two inclusions will finish the proof. The construction must contain every excluded minor, giving an upper bound on their number. Each listed object must also be an excluded minor, and the objects must be pairwise nonisomorphic, giving the matching lower bound. The computation in Section 6 supplies both: the exhaustive construction leaves 35 classes, and separate certificates verify all 35. Its numerical results are organized there according to the three steps above; implementation records are collected in Appendix C.

For comparison, Brettell's classification through ten elements gives 5, 92, 219, and 130 rank-four excluded minors at sizes seven through ten [1]. We reproduce those smaller catalogues as checks on the method. The use of compatible extensions and of inequivalent representations has precedents in the computations of Brettell and Pendavingh [2]. Here the proof of coverage rests on generating all possible parents and imposing compatibility with every deletion.

2 Step 1: finding every possible parent

This step establishes the starting point of the search: a complete list of matroids N that could occur as $M \setminus e$. We first determine which parents are needed, then prove that the represented generation procedure reaches all of them.

2.1 Why connected simple parents suffice

We use the standard notation $r_M(X)$ for the rank of a subset and $\mathcal{B}(M)$ for the bases of M. All matroids are finite. A simple matroid has no loops or parallel pairs.

Lemma 2.1 (Possible deletions)**.** *If M is a rank-four, eleven-element excluded minor for quinary matroids, then for every $e \in E(M)$ the deletion $M \setminus e$ is a connected simple rank-four quinary matroid on ten elements.*

Proof. An excluded minor for representability over a field is 3-connected when it has at least four elements. To recall the reason, representability over a fixed field is closed under direct sums and 2-sums. A disconnected matroid decomposes as a direct sum; a connected matroid with a 2-separation decomposes as a 2-sum. The summands are isomorphic to proper minors [3]. If the matroid were an excluded minor, all summands would be representable, forcing the matroid itself to be representable. For the 2-sum construction, choose representations in which the common element is a unit vector, identify its one-dimensional span in the two vector spaces, and delete its column.

Thus M is 3-connected. It has no loops, parallel pairs, or coloops, so $N = M \setminus e$ is simple and has rank four. It is quinary by minor-minimality. It remains to prove that N is connected. If $E(N) = A \dot\cup B$ were a nontrivial direct-sum partition, choose the names so that $|A| \geq 2$. Since $r(N) = r(M)$, we would have

$$r_M(A) + r_M(B) = r(M), \qquad r_M(A) + r_M(B \cup \{e\}) - r(M) \leq 1.$$

Both A and $B \cup \{e\}$ have at least two elements. This contradicts 3-connectivity of M. □

Let $P = \{0, \ldots, 9\}$ and take a new label $e = 10$. Let $\mathcal{P}$ contain one labelled representative on P of each connected simple rank-four quinary matroid on ten elements. For each $N \in \mathcal{P}$, define $\mathcal{D}(N)$ to be the set of simple rank-four matroids M on $P \cup \{e\}$ satisfying

$$M \setminus e = N, \qquad M \text{ is nonquinary},$$
$$M \setminus a \text{ is quinary for every } a \in P \cup \{e\}.$$

We call these *deletion-minimal nonquinary extensions* of N. The word "deletion" is essential: contractions have not yet been tested.

Proposition 2.2 (Reduction of the classification)**.** *The rank-four, eleven-element quinary excluded minors are exactly, up to isomorphism, the members of*

$$\bigcup_{N \in \mathcal{P}} \{M \in \mathcal{D}(N) : M/a \text{ is quinary for every } a \in E(M)\}.$$

Proof. Every matroid in the displayed set is nonquinary and has quinary immediate deletions and contractions, so it is an excluded minor. Conversely, let M be any excluded minor of the specified rank and size. By Lemma 2.1, a deletion $M \setminus e$ is isomorphic to a member N of $\mathcal{P}$. Relabel this deletion as N and its missing element as e. The resulting matroid belongs to $\mathcal{D}(N)$, and all its contractions are quinary. Its isomorphism class therefore occurs in the displayed set. □

This proposition specifies exactly what the rest of the computation must produce. It suffices to construct $\mathcal{P}$, compute each $\mathcal{D}(N)$ completely, and apply the contraction test. In particular, a parent may be discarded for being disconnected, but not merely for failing to be 3-connected.

2.2 Generating a complete represented catalogue

A full-rank $r \times k$ matrix over $\mathbb{F}_5$ represents a matroid whose bases are its nonsingular r-column submatrices. In a simple representation, its columns specify distinct points of $\mathrm{PG}(r-1,5)$ and span the ambient space. We represent a projective point by the vector whose first nonzero entry is 1. There are $(5^r-1)/4$ such points, in particular 156 in rank four.

Starting with I_r, append every projective point not already used, and repeat until the required size is reached. Isomorphic repetitions can be removed during growth, provided the equivalence used respects the *representation*. Two matrices are *field-isomorphic* when one can be obtained from the other by invertible row operations, nonzero column scalings, and a permutation of columns. Abstractly isomorphic matroids can have representations that are not field-isomorphic. All such represented classes must be retained.

Proposition 2.3 (Completeness of represented generation)**.** *The procedure just described, retaining one matrix from each field-isomorphism class at each size, generates every simple rank-r quinary representation up to field isomorphism.*

Proof. Take any simple rank-r representation. Choose r columns forming a basis, transform them to I_r, and order the remaining columns. Appending those columns in that order gives a chain of simple rank-preserving represented extensions from I_r to the chosen matrix. Inductively, each intermediate representation is field-isomorphic to a retained one. Such an isomorphism carries all possible appended projective points bijectively to possible appended points of the retained representation, and extends the permutation of old labels by fixing the new label. Thus replacing an intermediate matrix by its retained representative cannot lose any later represented extension class. Induction on the number of columns proves the assertion. □

After a represented level is complete, forgetting the matrices and deduplicating by abstract matroid isomorphism gives the required parent classes. We generate all simple rank-four represented levels through size ten in this way, then retain the connected ten-element parents for $\mathcal{P}$. The same procedure in rank three provides the catalogue that will be needed to test contractions in Section 5.

Step 1 is now justified: every possible deletion of an excluded minor occurs in $\mathcal{P}$. We next explain how to recover its possible nonquinary extensions. This requires more than appending columns to the parent itself; it requires matching the representable deletions of the prospective extension.

3 Step 2: recovering an extension from its deletions

Fix one parent $N \in \mathcal{P}$. Our goal in this section is an exact description of $\mathcal{D}(N)$ that can be computed from finite lists of quinary matroids. We first describe the unknown data of an extension, then express the conditions on its deletions in those data. The main proposition will show that these conditions already enforce the matroid axioms.

3.1 The unknown basis data

The deletion equation $M \setminus e = N$ fixes the status of every four-set avoiding e: it is a basis of M exactly when it is a basis of N. The only unknown four-sets are therefore

$$T \cup \{e\}, \qquad T \in \binom{P}{3}.$$

There are $\binom{10}{3} = 120$ of these, compared with the $\binom{10}{4} = 210$ already fixed four-sets. Introduce a Boolean variable x_T for each triple, with $x_T = 1$ meaning that $T \cup \{e\}$ is a basis. An assignment x specifies a nonempty four-subset family

$$\mathcal{B}_N(x) = \mathcal{B}(N) \ \cup \{T \cup \{e\} : T \in \binom{P}{3},\ x_T = 1\}.$$

For the moment this is only a proposed basis family; an arbitrary assignment need not define a matroid.

Consider how another deletion reads these variables. For $f \in P$, the four-sets of $M \setminus f$ that contain e are indexed by the triples avoiding f. Write

$$\pi_f(x) = x\big|_{\binom{P\setminus\{f\}}{3}}$$

for the restriction to those coordinates. The four-sets avoiding both e and f are already determined by $N \setminus f$. Thus $\pi_f(x)$ is exactly the extra basis data needed to specify $M \setminus f$ as an extension of $N \setminus f$.

For example, the bit $x_{\{0,1,2\}}$ records whether $\{0, 1, 2, 10\}$ is a basis. This bit is visible in the deletion of 3 and in the deletion of 4, but disappears in the deletion of 0. If independently chosen extensions for the deletions of 3 and 4 disagree on that bit, they cannot be deletions of one matroid. A single assignment x enforces agreement on every such shared four-set at once.

To express which deletions are permitted, let Q be any fixed labelled simple rank-four quinary matroid on a set U. Define its *represented extension table* $\mathcal{R}(Q)$ to be the set of Boolean patterns z on $\binom{U}{3}$ for which

$$\mathcal{B}(Q) \ \cup \{T \cup \{e\} : z_T = 1\}$$

is the basis family of a simple rank-four quinary extension of Q. All labels in this definition are fixed. In Section 4 we will construct this table completely from representations of Q.

The meaning of table membership is now precise:

- $\pi_f(x) \in \mathcal{R}(N \setminus f)$ says that the proposed deletion avoiding f is simple, rank four, and quinary;
- $x \in \mathcal{R}(N)$ says that the proposed full extension of N is itself simple, rank four, and quinary.

The first statement is meaningful because a connected parent has no coloops, so every $N \setminus f$ still has rank four.

Consequently the assignments we want should form the finite set

$$\mathcal{S}(N) = \big\{x \in \{0,1\}^{\binom{P}{3}} : \pi_f(x) \in \mathcal{R}(N \setminus f) \text{ for every } f \in P\big\} \setminus \mathcal{R}(N).$$

The intersection requires all the other deletions to be quinary; the subtraction removes the extensions that are quinary themselves. What still needs proof is that an assignment satisfying these conditions actually defines a matroid. Without that implication, the displayed expression would merely describe consistent tables of four-subsets.

3.2 Why compatible deletions force the matroid axioms

The basis-exchange axiom involves two bases and hence at most twice the rank many elements. If the ground set is larger, the whole exchange question can be seen inside one deletion. This gives the needed local-to-global principle.

Lemma 3.1 (Deletion compatibility)**.** *Let $|E| = n > 2r$ and let $\varnothing \neq \mathcal{B} \subseteq \binom{E}{r}$. Suppose that for each $a \in E$ the family of members of $\mathcal{B}$ avoiding a is the basis family of a rank-r matroid on $E \setminus \{a\}$. Then $\mathcal{B}$ is the basis family of a rank-r matroid on E.*

Proof. Take $A, B \in \mathcal{B}$ and $b \in A \setminus B$. We must find $c \in B \setminus A$ such that $(A \setminus \{b\}) \cup \{c\} \in \mathcal{B}$. Since $|A \cup B| \leq 2r < n$, there is an element $a \notin A \cup B$. Both A and B are bases of the specified deletion avoiding a. Basis exchange in that matroid supplies the required c, and its exchanged basis also belongs to $\mathcal{B}$. This proves every ordered instance of basis exchange. Together with nonemptiness and the common size of the members, it proves that $\mathcal{B}$ is a matroid basis family. □

Here $n = 11$ and $r = 4$, so the lemma applies. In particular, the representations witnessing different deletions do not have to fit together as matrices. They need only prescribe compatible bases. The lemma then supplies the abstract matroid, even when it has no representation over $\mathbb{F}_5$.

Proposition 3.2 (Exact recovery of deletion-minimal extensions)**.** *For a fixed parent $N \in \mathcal{P}$, the map $x \mapsto M_x$, where $\mathcal{B}(M_x) = \mathcal{B}_N(x)$, is a bijection from $\mathcal{S}(N)$ to $\mathcal{D}(N)$.*

Proof. First take $x \in \mathcal{S}(N)$. The proposed basis family $\mathcal{B}_N(x)$ is nonempty because it contains $\mathcal{B}(N)$. Its deletion avoiding e is the rank-four quinary matroid N. For each $f \in P$, the condition $\pi_f(x) \in \mathcal{R}(N \setminus f)$ says that its deletion avoiding f is also the basis family of a simple rank-four quinary matroid. Lemma 3.1 therefore makes $\mathcal{B}_N(x)$ a matroid basis family.

Call this matroid M_x. Each pair of elements lies in one of its simple deletions, so M_x is simple. All its deletions are quinary by construction. If M_x were quinary, its pattern would belong to $\mathcal{R}(N)$ by the definition of that table, contrary to the subtraction in $\mathcal{S}(N)$. Hence $M_x \in \mathcal{D}(N)$.

Conversely, take $M \in \mathcal{D}(N)$. Its bases containing e give a unique pattern x. For every $f \in P$, the matroid $M \setminus f$ is a simple quinary extension of $N \setminus f$; it has rank four since it contains the rank-four restriction $N \setminus f$. Thus $\pi_f(x) \in \mathcal{R}(N \setminus f)$. Since M is nonquinary, $x \notin \mathcal{R}(N)$. Therefore $x \in \mathcal{S}(N)$, and $M = M_x$. Finally, different patterns give different basis families, proving injectivity. □

This proposition is the bridge between the mathematical problem and the computation. It accounts for all four requirements on an output: validity as a matroid, simplicity, representability of every deletion, and nonrepresentability of the full matroid. To finish Step 2, it remains to build the tables $\mathcal{R}(Q)$ without omissions and exhaust the finite conditions defining $\mathcal{S}(N)$.

4 Constructing and exhausting the extension tables

Proposition 3.2 reduces each parent search to a finite problem about quinary extensions of nine- and ten-element matroids. There are two obligations: every represented extension must occur in its table, and every simultaneous solution of the table conditions must be considered. We address them in that order.

4.1 Putting all representations on the required labels

The represented catalogues of Section 2 retain all field-isomorphism classes. A table $\mathcal{R}(Q)$, however, describes extensions of one *labelled* matroid Q. To use the catalogue for this purpose, we need all representations on those labels, up to row operations and column scaling alone. We call the latter equivalence *labelled projective equivalence*.

First choose one matroid isomorphism aligning each unlabelled represented class of Q with the chosen labels. There may be many alignments, and they can give different labelled extension patterns. The missing alignments are recovered by applying all automorphisms of Q to the aligned matrices.

Lemma 4.1 (Complete labelled representations)**.** *Align one matrix from every field-isomorphism class representing Q with the fixed labels of Q, and close the resulting set under* $\mathrm{Aut}(Q)$*. Up to labelled projective equivalence, this gives every representation of the labelled matroid Q.*

Proof. Take any representation on those labels and forget its labels. It belongs to one of the represented classes in the complete catalogue. Choose a field isomorphism from the catalogue matrix to this representation. It induces an alignment of the underlying matroid with Q. The chosen alignment of that same catalogue matrix differs from it by an automorphism of Q. Applying that automorphism to the aligned matrix therefore reaches the desired representation, up to row operations and column scaling. □

The computation closes under generators of the full automorphism group, which reaches the same orbit as applying the entire group. Repeated labelled projective classes are recognized by a canonical matrix normalization. The normalization and its proof are given in Appendix A.1; its purpose is to remove repetitions without discarding a representation class. Denote the resulting complete labelled set of matrices by $\mathcal{L}(Q)$.

4.2 Building a table by appending projective points

For each $V \in \mathcal{L}(Q)$ append every point v of $\mathrm{PG}(3,5)$ whose direction is not already a column of V. The resulting matrix has rank four, and its column matroid is simple. Record the pattern

$$\rho(V,v)_T = 1 \quad \Longleftrightarrow \quad \det[\, V_T \mid v \,] \neq 0, \qquad T \in \binom{E(Q)}{3},$$

where V_T consists of the three old columns indexed by T. Collect all these rows, merging duplicates.

Proposition 4.2 (Completeness of the extension table)**.** *The rows obtained in this way are exactly $\mathcal{R}(Q)$.*

Proof. Every appended point gives a simple rank-four quinary extension, so every constructed row belongs to $\mathcal{R}(Q)$. Conversely, take any extension represented by a row of $\mathcal{R}(Q)$, and choose a matrix representing it. Its restriction to Q is labelled projectively equivalent to a member V of $\mathcal{L}(Q)$, by Lemma 4.1. Apply that equivalence to the entire matrix. The new column is a nonzero projective point not already used, since the extension is simple. That point is among those appended to V, so the extension's row occurs in the construction. □

This proves the first obligation of the section. In particular, a deletion $N \setminus f$ is matched to its catalogue representative and all its matrices are aligned back to the actual labels $P \setminus \{f\}$ before its table is formed. Equality of bits in different tables consequently refers to the same elements, as required by Proposition 3.2.

All determinant calculations take place in $\mathbb{F}_5$. For a fixed old triple the determinant is a linear form in v, so its cofactors allow all 156 projective points to be tested in one finite calculation. This speeds up table construction while computing exactly the determinant bits in the displayed definition.

4.3 Compressing the coordinates without changing the solutions

The 120 triple coordinates give the simplest proof of the recovery theorem, but many must agree. We now identify those repetitions to obtain the smaller problems actually solved by the program.

Let M be any rank-preserving matroid extension of N. If a triple T is dependent in N, then $r_M(T \cup \{e\}) \leq 3$, so its bit is zero. If T is independent and $H = \mathrm{cl}_N(T)$, then H is a rank-three hyperplane and

$$\mathrm{cl}_M(T) = \mathrm{cl}_M(H).$$

Indeed, $T \subseteq H \subseteq \mathrm{cl}_M(T)$, which proves both closure inclusions. Hence

$$T \cup \{e\} \in \mathcal{B}(M) \quad \Longleftrightarrow \quad e \notin \mathrm{cl}_M(H).$$

Thus independent triples spanning the same hyperplane have the same bit. Since every hyperplane has a spanning triple, one bit y_H for each rank-three hyperplane H determines all the triple bits of a genuine extension.

Write $\mathcal{H}(N)$ for these hyperplanes. In a table for the deletion of f, the visible hyperplanes are

$$C_f = \{\mathrm{cl}_N(T) : T \in \binom{P \setminus \{f\}}{3},\ r_N(T) = 3\}.$$

The closure of such a triple in $N \setminus f$ is $\mathrm{cl}_N(T) \setminus \{f\}$. Consequently every row of $\mathcal{R}(N \setminus f)$ is constant on triples assigned the same coordinate in C_f, and is zero on dependent triples. Projecting onto one bit per visible hyperplane therefore loses no row information. Let $A_f \subseteq \{0,1\}^{C_f}$ be this projected table, and let $A \subseteq \{0,1\}^{\mathcal{H}(N)}$ be the analogous projection of $\mathcal{R}(N)$.

The problem actually solved is

$$\widehat{\mathcal{S}}(N) = \{y \in \{0,1\}^{\mathcal{H}(N)} : y|_{C_f} \in A_f \text{ for every } f \in P\} \setminus A.$$

Expanding y_H back to every independent triple spanning H, and inserting zeros on dependent triples, takes each solution to one in $\mathcal{S}(N)$. Conversely, Proposition 3.2 makes every solution of $\mathcal{S}(N)$ a genuine matroid extension, so the closure argument above shows that it has a unique such compression. Thus $\widehat{\mathcal{S}}(N)$ and $\mathcal{S}(N)$ have exactly the same solutions in different coordinates.

Every hyperplane coordinate is seen by at least one deletion: choose $f \notin H$ and a spanning triple of H. That triple avoids f, so $H \in C_f$. The union of the coordinate sets C_f is therefore all of $\mathcal{H}(N)$.

4.4 Why the finite join is exhaustive

It remains to impose all the tables simultaneously. For relations U and V on coordinate sets S and T, their *natural join* is the finite relation

$$U \bowtie V = \{u \cup v : u \in U,\ v \in V,\ u|_{S\cap T} = v|_{S\cap T}\}.$$

In words, take every pair of rows that agree on their shared coordinates and combine them into a row on $S \cup T$. Grouping rows by their overlap values implements exactly this operation.

Start with the single empty assignment and join the ten relations A_f, one at a time. After processing any collection of deletions, the current rows are exactly the assignments on the coordinates seen so far that satisfy all the processed relations. This is true initially, and the definition of join proves the induction step. After all ten joins, every coordinate has been seen, so the rows are exactly

$$\{y : y|_{C_f} \in A_f \text{ for every } f \in P\}.$$

Removing the rows in A now gives $\widehat{\mathcal{S}}(N)$ and hence, by the recovery theorem, every member of $\mathcal{D}(N)$.

The implementation may change the join order and remove rows of A as soon as all coordinates have appeared. Neither change alters the result: a completed assignment cannot be changed by later joins, only accepted or rejected. Resource limits are handled separately. An interrupted or bounded calculation is recorded as `UNKNOWN`, and cannot count as an exhaustive parent search. Only a finished calculation establishes that every solution has been considered. A second method solves the same finite relations using SAT; its encoding is described in Appendix A.3.

Step 2 is therefore justified: complete representation lists give complete tables, and completed joins give exactly $\mathcal{D}(N)$. At this stage every output is nonquinary and all its deletions are quinary. The sole remaining minor conditions are the contractions.

5 Step 3: testing contractions and certifying the list

Proposition 2.2 tells us how to finish the classification once the sets $\mathcal{D}(N)$ have been recovered. We must test every contraction and then pass from labelled matroids to isomorphism classes. We also verify each final object by certificates that can be checked without rerunning the search.

5.1 The remaining minor conditions

For a simple rank-four matroid M, every contraction M/a has rank three. It may have parallel elements. Its *simplification* $\mathrm{si}(M/a)$ deletes loops and keeps one element of each parallel class.

Lemma 5.1 (The contraction test)**.** *For $M \in \mathcal{D}(N)$, the matroid M is an excluded minor if and only if every $\mathrm{si}(M/a)$ is a quinary rank-three matroid.*

Proof. A matroid is quinary if and only if its simplification is. One direction follows because a simplification is a restriction; for the other, extend a representation of the simplification by zero columns for loops and scalar copies of columns for parallel elements. Thus the stated condition is equivalent to all contractions of M being quinary. Membership in $\mathcal{D}(N)$ already supplies nonrepresentability of M and representability of all its deletions. The immediate-minor criterion from Section 1 now gives the result. □

This is where the rank-three represented catalogue from Step 1 is used. Each simplification has at most ten elements. A simple rank-three matroid on three elements is free and is represented by I_3. At sizes four through ten, exact isomorphism comparison with the corresponding complete quinary catalogue decides the test. In particular, this step needs no assumption about a classification of rank-three excluded minors of arbitrary size.

After applying this test, Proposition 2.2 shows that the retained matroids are exactly the required excluded minors, with possible repetitions from their different parents or labellings. Exact matroid isomorphism removes these repetitions. The final distinctness check also uses

colored incidence graphs: one color consists of element vertices, the other of vertices for all bases or, when fewer, all nonbases, with edges given by membership. A color-preserving graph isomorphism induces exactly a matroid isomorphism, since either subset family determines the bases. Canonical forms of these graphs therefore give an exact second check of pairwise nonisomorphism.

5.2 Certifying that every listed object is an excluded minor

The construction proves coverage of the target class. For the opposite inclusion, each final basis family is validated and given two kinds of evidence: representations of all its immediate minors, and a proof that the matroid itself has no representation.

The positive evidence is direct. For each $a \in E(M)$ we save matrices for $M \setminus a$ and M/a, in the order of their remaining labels. An independent modular Gaussian-elimination routine computes the basis family of each matrix and compares it with that of the required minor. It checks every subset of the appropriate size, so agreement establishes equality of the represented matroid and the specified minor. Sage performs a second matrix check. These witnesses certify all the proper-minor conditions using only finite-field arithmetic.

Nonrepresentability requires a different form of evidence. We turn existence of a representation into a finite propositional formula Φ_M and check a certificate that this formula is unsatisfiable. Here is the mathematical condition encoded by the formula. Choose a basis $B = (b_1, \ldots, b_r)$ of M and reduce its columns to the identity. The remaining entries form a block Z, and fundamental exchanges prescribe its support:

$$Z_{ij} \neq 0 \quad \Longleftrightarrow \quad (B \setminus \{b_i\}) \cup \{j\} \in \mathcal{B}(M), \qquad j \notin B.$$

The unknown nonzero entries range over $\{1, 2, 3, 4\}$. For every r-subset S, the determinant of the columns on S must be nonzero exactly when $S \in \mathcal{B}(M)$. These conditions impose the entire basis family.

Proposition 5.2 (Meaning of the nonrepresentability certificate)**.** *The exact finite-field encoding Φ_M of these determinant conditions is satisfiable if and only if M is quinary.*

Proof. Any representation of M can be reduced on B to identity form. Its remaining entries have the prescribed support and satisfy every determinant condition, so they give a solution. Conversely, a solution supplies a full-rank matrix whose nonsingular r-column submatrices are exactly $\mathcal{B}(M)$; this matrix represents M. The encoding uses exact addition and multiplication tables modulo five, so its satisfying assignments are precisely the solutions of these field conditions. The harmless scaling normalization and the arithmetic clauses are justified in Appendix A. □

For each final matroid, the SAT solver produces a DRAT refutation of Φ_M. DRAT-trim checks this refutation against the saved formula [5]. By Proposition 5.2, a successful check proves that M is nonquinary. Together with the 22 minor matrices, it certifies that M is an excluded minor. These certificates use the final basis records directly; they do not need the deletion tables or their enumeration results.

At this point the logical roles are distinct. Complete parent generation and exhaustive recovery show that every excluded minor reaches the final list. Certificates show that every entry of that list is an excluded minor. Exact isomorphism tests determine how many different classes the list represents. We now give the computational facts that establish these statements for the present rank and size.

6 The computation and the proof of the theorem

The completed run of 27 September 2026 records the outcomes in Table 1. Each row corresponds to a specified mathematical set or operation in the preceding sections.

Stage	Objects counted	Number
Generation	Simple rank-four quinary ten-element parent classes	58,756
Step 1	Connected parent classes in $\mathcal{P}$	58,672
Step 2	Labelled extensions, counted as pairs (N, M) with $M \in \mathcal{D}(N)$	659
Step 3	Classes after all contraction tests and isomorphism deduplication	35

Table 1: The successive sets in the exhaustive classification.

6.1 Evidence that the construction is complete

The represented rank-four generation retains 64,508 field-isomorphism classes at size ten. Abstract deduplication gives the 58,756 simple parent classes in the first row. Exactly 84 are disconnected and are discarded by Lemma 2.1; the remaining 58,672 form $\mathcal{P}$. Aligning the represented classes and closing their automorphism orbits yields 71,547 labelled projective classes over the full ten-element parent catalogue. The complete nine-element labelled sets supply the other extension tables. All generated levels are listed in Appendix C.

Every one of the 58,672 connected-parent problems finishes by exhaustive natural join, with no unresolved case. The merged logs account for every parent index exactly once, with each parent either excluded for being disconnected or having its finite problem exhausted. The shards have matching input and critical-source fingerprints. By Propositions 4.2 and 3.2, this yields exactly the pairs (N, M) with $M \in \mathcal{D}(N)$: there are 659. This is a count of parent-labelled extensions; it is not yet a count of isomorphism classes of excluded minors.

Testing all contractions using the complete rank-three catalogues, then deduplicating by isomorphism, leaves 35 classes. The independent incidence-graph check confirms that these 35 are pairwise nonisomorphic. For each of them the separate determinant formula has a checked DRAT refutation, and all 22 saved minor matrices pass both modular Gaussian and Sage verification. Thus there are 35 checked refutations and $35 \cdot 22 = 770$ checked matrices.

Proof of Theorem 1.1. To show that there are at most 35 classes, take any rank-four, eleven-element quinary excluded minor M. By Lemma 2.1, a deletion of M is isomorphic to a parent $N \in \mathcal{P}$. The represented generation is complete by Proposition 2.3. After aligning the labels, the recovery theorem, Proposition 3.2, places M among the solutions of the search for N. Its tables are complete by Proposition 4.2, and the recorded search exhausts them. Since every contraction of M is quinary, the complete contraction test of Lemma 5.1 retains it. Hence its class is among the 35 final classes.

To show that there are at least 35 classes, take the 35 final representatives. The checked refutation for each proves its nonrepresentability by Proposition 5.2. Its 22 checked matrices prove representability of every immediate minor, so it is an excluded minor. The independent isomorphism check proves that the representatives belong to 35 different classes. Combining the two bounds proves the classification. □

6.2 Checks on the computation and consequences

The same procedure reproduces the published rank-four catalogues at sizes seven through ten: 5, 92, 219, and 130 classes, respectively [1]. Each comparison is by exact matroid isomorphism with the published archive. At sizes seven and eight the deletion-compatibility inequality $n > 2r$ does not hold, so those calibration runs impose basis exchange explicitly. The rank-four nine-element target does satisfy the inequality.

A separate SAT enumeration of the finite table conditions agrees with the join on every connected-parent problem for the nine-element target. At the ten-element target, the complete sets of 5,991 labelled deletion-minimal candidates agree. These checks test the implementation; the proof of coverage is the parent reduction and recovery argument above. The two enumeration methods share their table construction, and the eleven-element target is exhausted by the join method.

Duality gives a further classification immediately.

Corollary 6.1. *There are exactly 35 isomorphism classes of rank-seven, eleven-element excluded minors for representability over* $\mathbb{F}_5$.

Proof. Field representability is closed under duality, and duality interchanges deletion and contraction. It therefore maps excluded minors to excluded minors and gives a bijection between the rank-four and rank-seven classes on eleven elements. □

Every matroid in the rank-four catalogue has a triangle, so none is paving. This is a property of the resulting classification; no paving restriction is used in the search. Basis and triangle counts accompany the exact descriptions in Appendix B.

A Details of the representation and SAT encodings

The main argument uses matrix normalization to enumerate complete labelled representation sets, and an exact field encoding to certify nonrepresentability. This appendix supplies the details of those two operations. The last subsection describes the alternative SAT enumeration used to cross-check the finite joins.

A.1 A normal form for labelled representations

Fix an ordered basis $B = (b_1, \ldots, b_r)$ of a labelled matroid Q. Reduce a representation on B so that its basis columns form I_r, and call the remaining block Z. The support of Z is determined by fundamental exchanges:

$$Z_{ij} \neq 0 \quad \Longleftrightarrow \quad (B \setminus \{b_i\}) \cup \{j\} \in \mathcal{B}(Q).$$

Its support graph has one vertex for each row, one for each nonbasis column, and an edge for each nonzero entry. Fix a deterministic spanning forest of this graph. By row and column scaling we make every forest entry equal to 1, compensating on basis columns to preserve I_r.

Lemma A.1 (Normal form)**.** *With the basis and support forest fixed, this normalization gives one matrix for each labelled projective equivalence class.*

Proof. To establish existence, root each nontrivial tree at a row vertex and choose its scale. Traverse the tree: whenever the scale at one endpoint of an edge is known, the nonzero entry on that edge uniquely determines the scale at the other endpoint making the entry 1. There is no cycle in the tree, so no conflicting choice is imposed. Isolated vertices require no condition. Inverse scaling of the basis columns restores the identity.

For uniqueness, two projectively equivalent representations reduced on the same labelled basis differ by row and column scalings: the row map must be diagonal because it fixes each basis-column direction. Suppose both have forest entries 1. On every tree edge, the row multiplier times the column multiplier must be 1. Connectivity then forces a common row multiplier throughout a component and its inverse on all column vertices in that component. Their product is also 1 on every nonforest edge, so all entries agree. This proves uniqueness, including when the support graph is disconnected. □

In the labelled-orbit calculation of Lemma 4.1, every new matrix is normalized before comparison with those already seen. The lemma proves that this comparison removes exactly projectively equivalent repetitions. The full automorphism group of Q is obtained from the colored element–basis/nonbasis incidence graph described in Section 5. Its action on element vertices is exactly $\mathrm{Aut}(Q)$, which supplies the generators used in the orbit calculation.

A.2 Compiling the determinant conditions to a formula

For the formula Φ_M in Proposition 5.2, choose a basis and form $[I_r \mid Z]$ as above. The support is fixed by the input matroid. Forest entries may be set to 1 by Lemma A.1, so only the other nonzero entries need variables, each with domain $\{1, 2, 3, 4\}$.

Let S be an r-subset of the ground set and put

$$I(S) = \{i : b_i \notin S\}, \qquad J(S) = S \setminus B.$$

These sets have the same size. Expansion on the identity columns shows that the full determinant on S is, up to sign, $\det Z[I(S), J(S)]$, with the empty determinant equal to 1. It is therefore enough to impose

$$\det Z[I(S), J(S)] \neq 0 \quad \Longleftrightarrow \quad S \in \mathcal{B}(M) \qquad \text{for every } S \in \binom{E(M)}{r}.$$

Every maximal minor is constrained, so these conditions determine the full labelled matroid.

Here is an explicit finite propositional encoding of the field arithmetic. For an expression u with allowed values $D \subseteq \mathbb{F}_5$, introduce indicators $p_{u,a}$ for $a \in D$, requiring exactly one to be true. If $w = u + v$, add the clauses

$$\neg p_{u,a} \ \vee \ \neg p_{v,b} \ \vee \ p_{w,a+b}$$

for every allowed pair a, b, where the sum is taken modulo five. Use the corresponding clauses with ab for multiplication. The output indicators also satisfy the exactly-one condition. Thus selected input values force precisely the correct output value. Constants are substituted directly. A cached Laplace expansion evaluates the determinants, and zero or nonzero requirements constrain their indicators. Induction over the arithmetic expressions proves that the formula represents exactly the field conditions used in Proposition 5.2.

For each listed matroid the saved evidence consists of its basis record, the chosen basis and normalization, the DIMACS formula, a DRAT refutation, and the proof-checker's output. The successful refutation certifies the supplied formula's unsatisfiability; the equivalence with representability is supplied by the mathematical encoding above.

A.3 The alternative SAT enumeration of tables

The natural-join search in Section 4 has an alternative implementation as a propositional satisfiability problem. It uses the same hyperplane coordinates and finite relations A_f. A

reduced binary decision diagram expresses membership in each relation. If a node asks about y and has low and high children l, h, its variable v is constrained by

$$v \iff ((\neg y \wedge l) \vee (y \wedge h)).$$

An exact four-clause encoding is

$$\neg v \vee y \vee l, \qquad \neg v \vee \neg y \vee h,$$
$$v \vee y \vee \neg l, \qquad v \vee \neg y \vee \neg h.$$

Requiring each root enforces all table memberships. A blocking clause excludes each represented full row in A. After an assignment to the hyperplane coordinates is found, another blocking clause excludes that assignment from subsequent enumeration. A final UNSAT answer then establishes exhaustion of this parent's solution set. A timeout is recorded as `UNKNOWN`. The SAT/table-join comparisons reported in Section 6 compare the complete assignments obtained by these two enumeration procedures.

B Exact catalogue

All representatives have ground set $E = \{0, 1, \ldots, 10\}$ and rank four. Each M_i is specified by the listed family $\mathcal{H}_i$ of hyperplanes of size at least four, according to the rule

$$\mathcal{B}(M_i) = \{S \in \binom{E}{4} : S \not\subseteq H \text{ for every } H \in \mathcal{H}_i\}.$$

Thus a four-set is a nonbasis exactly when it is contained in a listed hyperplane. This specifies the whole basis family: every dependent four-set of a rank-four matroid lies in a hyperplane, while hyperplanes of size three contain no four-set and may be omitted. The descriptions have been checked by reconstructing the basis records from them.

The notation M_i abbreviates the stable identifier `GF5-r4-n11-` followed by i in four digits. For example, M_1 is `GF5-r4-n11-0001`. Proof directories are zero-indexed, so its certificates are in `matroid-0000`. Basis and triangle counts identify the corresponding records in the machine-readable catalogue. Across the catalogue there are between 210 and 262 bases, and the triangle distribution is

triangles	4	5	6	7	8	10	12	13
classes	3	5	5	10	9	1	1	1

M_1: 250 bases; 6 triangles. $\{0,1,2,5,7,9\}$, $\{0,1,4,10\}$, $\{0,2,3,4,8\}$, $\{0,2,6,10\}$, $\{0,3,5,10\}$, $\{0,3,6,9\}$, $\{0,4,5,6\}$, $\{0,6,7,8\}$, $\{0,8,9,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8\}$, $\{1,2,5,10\}$, $\{1,3,7,9\}$, $\{1,4,7,9\}$, $\{1,6,7,8,9,10\}$, $\{2,3,4,7,10\}$, $\{2,3,4,9\}$, $\{2,6,7,8\}$, $\{2,8,9,10\}$, $\{3,6,7,8\}$, $\{3,8,9,10\}$, $\{4,5,6,7,8\}$, $\{4,5,6,9\}$, $\{4,5,6,10\}$, $\{4,8,9,10\}$, $\{5,8,9,10\}$.

M_2: 233 bases; 7 triangles. $\{0,1,2,6\}$, $\{0,1,4,7\}$, $\{0,1,8,9,10\}$, $\{0,2,3,4,5\}$, $\{0,2,7,9\}$, $\{0,3,6,9\}$, $\{0,3,7,10\}$, $\{0,4,6,8\}$, $\{0,5,6,10\}$, $\{0,5,7,8\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,8\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,4,7,8\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,8,9,10\}$, $\{6,7,8,9,10\}$.

M_3: 252 bases; 7 triangles. $\{0,1,2,5,7\}$, $\{0,1,3,6\}$, $\{0,1,4,9,10\}$, $\{0,2,3,4,8\}$, $\{0,2,6,9\}$, $\{0,3,5,7\}$, $\{0,4,5,6,7\}$, $\{0,5,7,8,9\}$, $\{0,5,7,10\}$, $\{0,6,8,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8\}$, $\{1,2,5,9\}$, $\{1,2,5,10\}$, $\{1,3,6,7\}$, $\{1,3,6,8,10\}$, $\{1,3,6,9\}$, $\{1,7,8,9\}$, $\{2,3,4,7,10\}$, $\{2,3,4,9\}$, $\{2,6,8,10\}$, $\{2,7,8,9\}$, $\{3,5,9,10\}$, $\{3,7,8,9\}$, $\{4,5,6,8,10\}$, $\{4,5,6,9\}$, $\{4,7,8,9\}$, $\{6,7,8,9,10\}$.

M_4: **250 bases; 5 triangles.** $\{0,1,2,5,7\}$, $\{0,1,3,10\}$, $\{0,1,4,8\}$, $\{0,2,3,4,8,9\}$, $\{0,2,6,10\}$, $\{0,3,5,7\}$, $\{0,4,5,7,8\}$, $\{0,4,6,8\}$, $\{0,4,8,10\}$, $\{0,5,6,7,9\}$, $\{0,5,7,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8\}$, $\{1,2,5,9\}$, $\{1,2,5,10\}$, $\{1,6,7,8,9,10\}$, $\{2,3,4,7,10\}$, $\{2,6,7,9\}$, $\{3,5,8,10\}$, $\{3,6,7,9\}$, $\{4,5,9,10\}$, $\{4,6,7,9\}$.

M_5: **247 bases; 7 triangles.** $\{0,1,2,6,8\}$, $\{0,1,7,9\}$, $\{0,2,3,4,5,7\}$, $\{0,3,6,8,9,10\}$, $\{0,4,6,8\}$, $\{0,5,6,8\}$, $\{0,6,7,8\}$, $\{1,2,3,4,5,6\}$, $\{1,2,6,7\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,3,7,8\}$, $\{1,4,7,10\}$, $\{1,5,8,9,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9\}$, $\{2,3,4,5,10\}$, $\{2,7,8,9,10\}$, $\{4,6,7,9\}$, $\{4,8,9,10\}$, $\{5,6,7,10\}$.

M_6: **241 bases; 8 triangles.** $\{0,1,2,7,9,10\}$, $\{0,1,6,8\}$, $\{0,2,3,4,5,8\}$, $\{0,3,7,9,10\}$, $\{0,4,7,9,10\}$, $\{0,5,6,7,9,10\}$, $\{0,7,8,9,10\}$, $\{1,2,3,4,5,6\}$, $\{1,3,8,9\}$, $\{1,4,8,10\}$, $\{1,5,7,8\}$, $\{2,3,4,5,7\}$, $\{2,3,4,5,9\}$, $\{2,3,4,5,10\}$, $\{2,6,8,10\}$, $\{3,6,7,8\}$, $\{4,6,8,9\}$.

M_7: **249 bases; 7 triangles.** $\{0,1,2,6,8\}$, $\{0,1,7,9,10\}$, $\{0,2,3,4,5,7\}$, $\{0,3,6,8,9\}$, $\{0,4,6,8\}$, $\{0,5,6,8,10\}$, $\{0,6,7,8\}$, $\{1,2,3,4,5,6\}$, $\{1,2,6,7\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,3,7,8\}$, $\{1,4,8,10\}$, $\{1,5,8,9\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9\}$, $\{2,3,4,5,10\}$, $\{2,7,8,9,10\}$, $\{3,7,9,10\}$, $\{4,6,7,9,10\}$, $\{5,7,9,10\}$.

M_8: **252 bases; 7 triangles.** $\{0,1,2,5,7\}$, $\{0,1,3,6\}$, $\{0,1,8,9\}$, $\{0,2,3,4,8\}$, $\{0,3,5,7\}$, $\{0,4,5,6,7\}$, $\{0,5,7,8\}$, $\{0,5,7,9,10\}$, $\{0,6,8,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8,9\}$, $\{1,2,5,10\}$, $\{1,3,6,7\}$, $\{1,3,6,8,9\}$, $\{1,3,6,10\}$, $\{1,4,8,9\}$, $\{1,7,8,9,10\}$, $\{2,3,4,7,9,10\}$, $\{4,5,6,8\}$, $\{4,5,6,9\}$, $\{4,5,6,10\}$, $\{6,7,9,10\}$.

M_9: **255 bases; 6 triangles.** $\{0,1,2,5,7\}$, $\{0,1,3,8\}$, $\{0,1,6,9,10\}$, $\{0,2,3,4,8\}$, $\{0,2,6,9\}$, $\{0,3,5,8\}$, $\{0,3,6,8,9\}$, $\{0,3,7,8,10\}$, $\{0,4,5,6,9\}$, $\{0,6,7,9\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8,9\}$, $\{1,2,5,10\}$, $\{1,3,7,9\}$, $\{1,7,8,10\}$, $\{2,3,4,7\}$, $\{2,3,4,9\}$, $\{2,3,4,10\}$, $\{2,6,7,8,10\}$, $\{3,5,9,10\}$, $\{4,5,6,7\}$, $\{4,5,6,8\}$, $\{4,5,6,10\}$, $\{4,7,8,9,10\}$, $\{5,7,8,10\}$.

M_{10}: **239 bases; 8 triangles.** $\{0,1,2,6,8\}$, $\{0,1,4,9,10\}$, $\{0,1,5,7\}$, $\{0,2,3,4,5\}$, $\{0,2,7,9\}$, $\{0,3,6,8,9\}$, $\{0,3,7,10\}$, $\{0,4,6,7,8\}$, $\{0,5,6,8\}$, $\{0,6,8,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,5,7,8\}$, $\{1,5,7,9\}$, $\{1,5,7,10\}$, $\{2,3,4,5,8,10\}$, $\{2,3,4,5,9\}$, $\{4,6,7,9\}$, $\{4,6,7,10\}$, $\{5,6,9,10\}$, $\{7,8,9,10\}$.

M_{11}: **212 bases; 10 triangles.** $\{0,1,2,6,8,9\}$, $\{0,1,4,10\}$, $\{0,1,5,7\}$, $\{0,2,3,4,5\}$, $\{0,2,7,10\}$, $\{0,3,6,10\}$, $\{0,3,7,9\}$, $\{0,4,6,7\}$, $\{0,5,8,10\}$, $\{1,2,3,4,5,6,7,8\}$, $\{1,2,6,8,10\}$, $\{1,5,7,9\}$, $\{1,5,7,10\}$, $\{2,3,4,5,9,10\}$, $\{4,6,7,9\}$, $\{4,6,7,10\}$, $\{7,8,9,10\}$.

M_{12}: **242 bases; 8 triangles.** $\{0,1,2,6,8\}$, $\{0,1,3,10\}$, $\{0,1,4,9\}$, $\{0,1,5,7\}$, $\{0,2,3,4,5,9\}$, $\{0,4,6,7,9\}$, $\{0,4,8,9\}$, $\{0,4,9,10\}$, $\{0,5,8,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,9,10\}$, $\{1,5,7,8\}$, $\{1,5,7,9\}$, $\{1,5,7,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,10\}$, $\{3,7,8,9,10\}$, $\{4,6,7,8\}$, $\{4,6,7,10\}$.

M_{13}: **254 bases; 5 triangles.** $\{0,1,2,5,7\}$, $\{0,1,4,8\}$, $\{0,2,3,4,8,9\}$, $\{0,3,5,7\}$, $\{0,4,5,7,8\}$, $\{0,4,6,8\}$, $\{0,4,8,10\}$, $\{0,5,6,7,9,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8\}$, $\{1,2,5,9\}$, $\{1,2,5,10\}$, $\{1,3,7,8\}$, $\{1,6,8,9,10\}$, $\{2,3,4,7,10\}$, $\{2,6,9,10\}$, $\{3,6,9,10\}$, $\{4,6,9,10\}$.

M_{14}: **257 bases; 4 triangles.** $\{0,1,2,5,7,9\}$, $\{0,1,4,10\}$, $\{0,2,3,4,8\}$, $\{0,3,6,7,9,10\}$, $\{0,4,7,9\}$, $\{0,5,6,8\}$, $\{0,7,8,9\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8\}$, $\{1,2,5,10\}$, $\{1,3,7,8\}$, $\{1,6,8,9,10\}$, $\{2,3,4,7\}$, $\{2,3,4,9\}$, $\{2,3,4,10\}$, $\{2,6,9,10\}$, $\{4,5,7,8,10\}$, $\{4,6,9,10\}$, $\{5,6,9,10\}$.

M_{15}: **262 bases; 4 triangles.** $\{0,1,2,5,7\}$, $\{0,1,4,10\}$, $\{0,2,3,4,8\}$, $\{0,3,5,7,10\}$, $\{0,4,5,7\}$, $\{0,5,6,7,8,9\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8\}$, $\{1,2,5,9\}$, $\{1,2,5,10\}$, $\{1,3,7,9\}$, $\{1,4,7,8\}$, $\{1,6,8,9,10\}$, $\{2,3,4,7\}$, $\{2,3,4,9,10\}$, $\{2,6,8,9\}$, $\{2,7,8,10\}$, $\{3,6,8,9\}$, $\{4,6,7,10\}$, $\{4,6,8,9\}$.

M_{16}: **210 bases; 13 triangles.** $\{0,1,2,7\}$, $\{0,1,4,9,10\}$, $\{0,1,5,8\}$, $\{0,2,3,4,5,6\}$, $\{0,2,8,9\}$, $\{0,3,7,9\}$, $\{0,3,8,10\}$, $\{0,4,7,8\}$, $\{0,6,7,10\}$, $\{1,2,3,4,5,6,7,8\}$, $\{1,2,7,9\}$, $\{1,2,7,10\}$, $\{1,5,8,9\}$, $\{1,5,8,10\}$, $\{2,3,4,5,6,9\}$, $\{2,3,4,5,6,10\}$, $\{4,7,8,9\}$, $\{4,7,8,10\}$, $\{5,7,9,10\}$, $\{6,8,9,10\}$.

M_{17}: **245 bases; 7 triangles.** $\{0,1,2,5,7,9\}$, $\{0,1,4,8\}$, $\{0,2,3,4,8,9\}$, $\{0,2,6,9\}$, $\{0,2,9,10\}$, $\{0,3,5,7\}$, $\{0,4,5,7,8\}$, $\{0,4,6,8\}$, $\{0,4,8,10\}$, $\{0,5,6,7,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8,10\}$, $\{1,3,7,8,9\}$, $\{1,4,7,9\}$, $\{1,6,7,9\}$, $\{1,7,9,10\}$, $\{2,3,4,7,10\}$, $\{3,5,8,9\}$, $\{3,6,8,9,10\}$.

M_{18}: **242 bases; 6 triangles.** $\{0,1,2,5,8,10\}$, $\{0,1,3,10\}$, $\{0,1,4,9,10\}$, $\{0,1,6,10\}$, $\{0,1,7,10\}$, $\{0,2,3,4,9\}$, $\{0,3,5,7\}$, $\{0,4,5,6,9\}$, $\{0,4,7,9\}$, $\{0,4,8,9\}$, $\{0,6,7,8\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,5,9\}$, $\{1,7,8,9\}$, $\{2,3,4,8\}$, $\{2,3,4,10\}$, $\{2,7,9,10\}$, $\{3,5,7,8\}$, $\{3,5,7,9\}$, $\{3,5,7,10\}$, $\{3,6,8,9,10\}$, $\{4,5,6,8\}$, $\{4,5,6,10\}$, $\{4,7,8,10\}$.

M_{19}: **244 bases; 8 triangles.** $\{0,1,2,6,7,9\}$, $\{0,1,3,9\}$, $\{0,1,4,8,9\}$, $\{0,1,5,9,10\}$, $\{0,2,3,4,5,7\}$, $\{0,2,7,8\}$, $\{0,2,7,10\}$, $\{0,6,8,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,6,8\}$, $\{1,2,6,10\}$, $\{1,3,8,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9\}$, $\{2,3,4,5,10\}$, $\{2,8,9,10\}$, $\{3,6,7,8,9\}$, $\{4,6,7,9,10\}$, $\{5,6,7,9\}$, $\{5,7,8,10\}$.

M_{20}: **256 bases; 6 triangles.** $\{0,1,2,5,7\}$, $\{0,1,3,9,10\}$, $\{0,1,4,6\}$, $\{0,2,3,4,6\}$, $\{0,3,5,7\}$, $\{0,4,5,6,7\}$, $\{0,4,6,8,10\}$, $\{0,4,6,9\}$, $\{0,5,7,8,9\}$, $\{0,5,7,10\}$, $\{1,2,3,4,5\}$, $\{1,2,5,6\}$, $\{1,2,5,8,9\}$, $\{1,2,5,10\}$, $\{1,3,6,7,8\}$, $\{1,4,7,9\}$, $\{1,4,8,10\}$, $\{2,3,4,7\}$, $\{2,3,4,8,10\}$, $\{2,3,4,9\}$, $\{2,6,7,9,10\}$, $\{3,5,6,10\}$, $\{3,5,8,9\}$, $\{4,5,8,9,10\}$, $\{4,7,8,10\}$, $\{5,6,8,9\}$.

M_{21}: **226 bases; 8 triangles.** $\{0,1,2,6,8,9\}$, $\{0,1,4,7\}$, $\{0,1,5,10\}$, $\{0,2,3,4,5\}$, $\{0,2,7,10\}$, $\{0,3,7,9\}$, $\{0,3,8,10\}$, $\{0,4,6,10\}$, $\{0,5,7,8\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,10\}$, $\{1,3,8,10\}$, $\{1,4,7,8\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,8,9,10\}$, $\{2,7,8,9\}$, $\{3,6,8,10\}$, $\{3,7,8,10\}$, $\{6,7,9,10\}$.

M_{22}: **261 bases; 4 triangles.** $\{0,1,2,5,7\}$, $\{0,1,6,8,9\}$, $\{0,2,3,4,8\}$, $\{0,2,6,9\}$, $\{0,3,5,7\}$, $\{0,3,6,9\}$, $\{0,4,5,7\}$, $\{0,4,6,9\}$, $\{0,5,6,7,9,10\}$, $\{0,5,7,8\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8,10\}$, $\{1,2,5,9\}$, $\{1,3,7,10\}$, $\{1,4,7,9\}$, $\{2,3,4,7\}$, $\{2,3,4,9,10\}$, $\{2,7,8,9\}$, $\{3,5,8,9\}$, $\{3,6,7,8\}$, $\{4,7,8,10\}$.

M_{23}: **220 bases; 8 triangles.** $\{0,1,2,6\}$, $\{0,1,4,7\}$, $\{0,1,5,9\}$, $\{0,1,8,10\}$, $\{0,2,3,4,5,10\}$, $\{0,2,8,9\}$, $\{0,3,6,8\}$, $\{0,5,7,8\}$, $\{0,6,7,9,10\}$, $\{1,2,3,4,5,6,7,8\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,3,9,10\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,9\}$, $\{3,6,8,9\}$, $\{3,6,8,10\}$, $\{4,8,9,10\}$, $\{5,7,8,9\}$, $\{5,7,8,10\}$.

M_{24}: **220 bases; 8 triangles.** $\{0,1,2,6\}$, $\{0,1,4,7\}$, $\{0,1,8,9,10\}$, $\{0,2,3,4,5\}$, $\{0,2,7,9\}$, $\{0,3,6,9\}$, $\{0,3,7,10\}$, $\{0,4,6,8\}$, $\{0,5,6,10\}$, $\{0,5,7,8\}$, $\{1,2,3,4,5,6,7,8\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,9,10\}$, $\{4,6,8,9\}$, $\{4,6,8,10\}$, $\{5,7,8,9\}$, $\{5,7,8,10\}$, $\{6,7,9,10\}$.

M_{25}: **243 bases; 7 triangles.** $\{0,1,2,6\}$, $\{0,1,4,7\}$, $\{0,1,8,9,10\}$, $\{0,2,3,4,5\}$, $\{0,2,7,10\}$, $\{0,2,8,9\}$, $\{0,3,6,10\}$, $\{0,3,8,9\}$, $\{0,4,6,8,9\}$, $\{0,5,7,8,9\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,8\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,4,7,8\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9,10\}$, $\{3,7,8,10\}$, $\{5,6,8,10\}$, $\{6,7,9,10\}$.

M_{26}: **243 bases; 7 triangles.** $\{0,1,2,6\}$, $\{0,1,4,7\}$, $\{0,1,8,9,10\}$, $\{0,2,3,4,5\}$, $\{0,2,8,9\}$, $\{0,3,6,8,9\}$, $\{0,3,7,10\}$, $\{0,4,6,10\}$, $\{0,4,8,9\}$, $\{0,5,7,8,9\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,8\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,4,7,8\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9,10\}$, $\{2,7,8,10\}$, $\{5,6,8,10\}$, $\{6,7,9,10\}$.

M_{27}: **228 bases; 12 triangles.** $\{0,1,2,7\}$, $\{0,1,4,9\}$, $\{0,1,5,8,10\}$, $\{0,2,3,4,5,6\}$, $\{0,2,8,9,10\}$, $\{0,3,7,9\}$, $\{0,3,8,10\}$, $\{0,4,7,8,10\}$, $\{0,6,8,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,7,8\}$, $\{1,2,7,9\}$, $\{1,2,7,10\}$, $\{1,3,9,10\}$, $\{1,6,8,9\}$, $\{2,3,4,5,6,8\}$, $\{2,3,4,5,6,9\}$, $\{2,3,4,5,6,10\}$, $\{5,7,8,9\}$, $\{6,7,9,10\}$.

M_{28}: **248 bases; 5 triangles.** $\{0,1,2,5,8\}$, $\{0,1,7,9\}$, $\{0,2,3,4\}$, $\{0,2,7,9\}$, $\{0,3,6,7,9,10\}$, $\{0,4,5,6\}$, $\{0,4,7,8,9\}$, $\{0,5,7,9\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,5,9,10\}$, $\{1,3,8,10\}$, $\{2,3,4,8\}$, $\{2,3,4,9\}$, $\{2,3,4,10\}$, $\{2,6,8,9\}$, $\{3,6,7,8\}$, $\{4,5,6,8,10\}$, $\{4,5,6,9\}$.

M_{29}: **216 bases; 8 triangles.** $\{0,1,2,9\}$, $\{0,1,6,7,8,10\}$, $\{0,2,3,4,5,10\}$, $\{0,3,8,9\}$, $\{0,4,7,9\}$, $\{0,5,6,9\}$, $\{1,2,3,4,5,6,7,8\}$, $\{1,3,9,10\}$, $\{1,6,7,8,9\}$, $\{2,3,4,5,9\}$, $\{2,7,9,10\}$, $\{4,6,9,10\}$, $\{5,8,9,10\}$.

M_{30}: **256 bases; 5 triangles.** $\{0,1,2,5,7\}$, $\{0,1,4,9,10\}$, $\{0,2,3,4,8\}$, $\{0,2,6,9,10\}$, $\{0,3,5,9,10\}$, $\{0,4,5,6\}$, $\{0,6,7,8\}$, $\{0,7,9,10\}$, $\{0,8,9,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8,9\}$, $\{1,2,5,10\}$, $\{1,6,7,8,10\}$, $\{2,3,4,7,10\}$, $\{2,3,4,9\}$, $\{2,6,7,8\}$, $\{3,6,7,8,9\}$, $\{4,5,6,7,8\}$, $\{4,5,6,9\}$, $\{4,5,6,10\}$.

M_{31}: **239 bases; 7 triangles.** $\{0,1,2,6,8,9\}$, $\{0,1,4,7\}$, $\{0,2,3,4,5,10\}$, $\{0,3,6,8\}$, $\{0,4,6,8\}$, $\{0,5,6,8\}$, $\{0,5,7,9\}$, $\{0,6,7,8,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,10\}$, $\{1,3,9,10\}$, $\{1,4,7,8\}$, $\{1,4,7,9\}$, $\{1,4,7,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9\}$, $\{2,7,9,10\}$, $\{3,7,8,9\}$, $\{4,6,9,10\}$, $\{5,8,9,10\}$.

M_{32}: **241 bases; 7 triangles.** $\{0,1,2,5,8\}$, $\{0,1,3,7\}$, $\{0,1,4,9,10\}$, $\{0,2,3,4\}$, $\{0,2,6,7,9\}$, $\{0,3,5,8,9\}$, $\{0,3,6,10\}$, $\{0,4,5,6,8\}$, $\{0,5,7,8\}$, $\{0,5,8,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,5,9\}$, $\{1,2,5,10\}$, $\{1,3,7,8,9\}$, $\{1,3,7,10\}$, $\{2,3,4,8,9\}$, $\{2,3,4,10\}$, $\{2,6,7,8,10\}$, $\{3,6,8,9\}$, $\{3,8,9,10\}$, $\{4,5,6,9\}$, $\{4,5,6,10\}$, $\{5,7,9,10\}$.

M_{33}: **241 bases; 6 triangles.** $\{0,1,2,5,8,9\}$, $\{0,1,3,10\}$, $\{0,2,3,4\}$, $\{0,2,6,10\}$, $\{0,3,6,7,9\}$, $\{0,4,5,6\}$, $\{0,4,7,8\}$, $\{0,4,9,10\}$, $\{0,5,7,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,5,10\}$, $\{1,3,8,9\}$, $\{1,4,8,9,10\}$, $\{1,6,8,9\}$, $\{1,7,8,9\}$, $\{2,3,4,8\}$, $\{2,3,4,9,10\}$, $\{3,6,7,8,10\}$, $\{4,5,6,8\}$, $\{4,5,6,9,10\}$, $\{4,7,9,10\}$.

M_{34}**: 258 bases; 5 triangles.** $\{0,1,2,5,7,9\}$, $\{0,1,4,10\}$, $\{0,2,3,4,8\}$, $\{0,3,5,7\}$, $\{0,3,6,9,10\}$, $\{0,4,5,6,7\}$, $\{0,5,7,8\}$, $\{0,5,7,10\}$, $\{1,2,3,4,5,6\}$, $\{1,2,5,8,10\}$, $\{1,3,7,9\}$, $\{1,4,7,9\}$, $\{1,6,7,8,9\}$, $\{1,7,9,10\}$, $\{2,3,4,7\}$, $\{2,3,4,9\}$, $\{2,3,4,10\}$, $\{2,6,7,10\}$, $\{3,5,8,9\}$, $\{3,7,8,10\}$, $\{4,5,6,8\}$, $\{4,5,6,9\}$, $\{4,5,6,10\}$, $\{4,8,9,10\}$.

M_{35}**: 238 bases; 8 triangles.** $\{0,1,2,6,8\}$, $\{0,1,5,7,9\}$, $\{0,2,3,4,5,10\}$, $\{0,3,6,8,9\}$, $\{0,4,6,7,8\}$, $\{0,5,6,8\}$, $\{0,6,8,10\}$, $\{1,2,3,4,5,6,7\}$, $\{1,2,6,9\}$, $\{1,2,6,10\}$, $\{1,3,9,10\}$, $\{1,5,7,8,10\}$, $\{2,3,4,5,8\}$, $\{2,3,4,5,9\}$, $\{2,7,9,10\}$, $\{4,6,7,9\}$, $\{4,6,7,10\}$, $\{4,8,9,10\}$, $\{5,6,9,10\}$.

C Computational records and reproduction

This appendix identifies the saved evidence for each step of the proof. All paths are relative to the repository root. The exact catalogue is `results/r4-n11/excluded.jsonl`, with SHA-256 digest

`7940989ae051b32496dbfd7e4717d405431bcfd89da69d0dd522517552c76441.`

Its equivalent hyperplane descriptions are in `results/r4-n11/hyperplanes.json`.

C.1 Parent coverage and exhaustive search

Table 2 gives the completed represented levels. The rank-four size-four entry is the seed I_4; rank-three generation starts from I_3. The counts distinguish represented classes, which must be retained during generation, from abstract matroid classes used to index the parent searches.

Size	Rank 3 field	Rank 3 abstract	Rank 4 field	Rank 4 abstract
4	2	2	1	1
5	4	4	3	3
6	11	9	12	11
7	22	18	58	44
8	42	34	460	303
9	92	82	5,001	3,763
10	174	168	64,508	58,756

Table 2: Complete simple represented catalogues. "Field" counts field-isomorphism classes; "abstract" counts matroid classes.

The main scripts correspond to the proof as follows.

Parent generation. `scripts/generate_parents.py` generates all represented rank-four and rank-three levels through size ten. The completed abstract lists index the parents and supply the contraction comparisons.

Labelled representations. `scripts/prepare_representations.py` aligns the represented classes and closes under automorphisms. The normalization and point-extension tables are implemented in `gf5minors/projective.py`.

Recovery and contraction testing. `scripts/search_compatible_deletions.py` builds the tables for each parent and tests contractions of the recovered matroids. The exact join is in `gf5minors/table_join.py`; the alternative SAT solver is in `gf5minors/table_sat.py`.

Merging and counting. `scripts/merge_search.py` checks parent coverage and removes isomorphic repetitions. `scripts/final_audit.py` checks certificate coverage and distinctness by canonical incidence graphs.

The four ranges in Table 3 partition $[0, 58756)$. Their logs record either exhaustion or the disconnected-parent exclusion for every index. The merge checks the ranges for gaps and overlaps, terminal statuses, and agreement of input and critical-source fingerprints.

Start	Stop	Parents	Candidates	Seconds
0	14,689	14,689	514	1794.54
14,689	29,378	14,689	106	1629.19
29,378	44,067	14,689	29	1430.24
44,067	58,756	14,689	10	1299.00

Table 3: Exhaustive target-search ranges. Candidate counts precede contraction filtering and global isomorphism deduplication.

The global coverage record is `results/r4-n11/summary.json`. Detailed parent logs and input manifests are in `runs/r4n11/shard-00/` through `runs/r4n11/shard-03/`. The final audit and statistics are `results/r4-n11/verification.json` and `results/r4-n11/statistics.json`. The target search used four Slurm tasks, each requesting one CPU and 4 GB of memory. The longest range took about 30 minutes.

C.2 Certificates and their replay

`scripts/certify_nonrepresentation.py` constructs the separate determinant formulas and checks their DRAT proofs. The 35 proof directories under `results/r4-n11/nonrepresentation/` are named `matroid-0000` through `matroid-0034`. For M_i, the directory index is $i-1$. Its files `representation.cnf` and `proof.drat` can be replayed with

```
vendor/drat-trim/drat-trim FORMULA PROOF
```

A successful check prints `s VERIFIED`.

`scripts/certify_minors.py` constructs and checks the immediate-minor matrices. All 770 are stored in `results/r4-n11/minors/minor-witnesses.jsonl`. The remaining column labels are in increasing parent order, renumbered to $0, \ldots, 9$. The pure modular checker `gf5minors.matroid.verify_representation` can check these matrices directly from the basis records, independently of the search and the represented catalogues.

The completed computation records Sage 10.9 and, in a separate environment, Python 3.11.16, python-sat 1.9.dev15, and NumPy 2.4.6. Sage supplies represented extensions, isomorphism, and graph operations [4]. The proof checker is pinned to DRAT-trim commit `2e3b2dc0ecf938addbd779d42877b6ed69d9a985`. The computational proof relies on the correctness of parent generation, representation orbits, table construction, exhaustive enumeration, and isomorphism handling. The DRAT checks certify the final determinant formulas; coverage of the classification is supplied by the exhaustive construction.

On the original cluster, run computation through Slurm with the existing `my-sage` and `matroid1` environments. `slurm/research.slurm` selects the environment and caps numerical-library threads at one. Full reproduction commands, in dependency order, are given in `docs/REPRODUCING.md`. Use fresh output directories for new runs. The reproducibility archive `results/r4-n11-reproducibility.tar.gz` and its adjacent JSON manifest preserve the original result and its inputs; that archive predates this exposition.

References

[1] N. Brettell, *The excluded minors for* GF(5)*-representable matroids on ten elements*, Advances in Applied Mathematics **166** (2025), 102864. doi:10.1016/j.aam.2025.102864. Preprint: arXiv:2307.14614, version 2, 2024.

[2] N. Brettell and R. Pendavingh, *Computing excluded minors for classes of matroids representable over partial fields*, Electronic Journal of Combinatorics **31**(3) (2024), P3.20. arXiv:2302.13175.

[3] J. Oxley, *Matroid Theory*, second edition, Oxford Graduate Texts in Mathematics 21, Oxford University Press, 2011.

[4] The Sage Developers, *SageMath, the Sage Mathematics Software System*, version 10.9, used in the computation. Linear matroids reference manual.

[5] N. Wetzler, M. J. H. Heule, and W. A. Hunt, Jr., *DRAT-trim: Efficient checking and trimming using expressive clausal proofs*, Theory and Applications of Satisfiability Testing (SAT 2014), Lecture Notes in Computer Science **8561**, 422–429. Software and documentation.