

TYLENOL COLD PLUS HEAD CONGESTION SEVERE- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, film coated
Kenvue Brands LLC

Tylenol® Cold + Head Congestion Severe

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 4 hours ▪ swallow whole–do not crush, chew or dissolve ▪ do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

- each caplet contains: **sodium 3 mg**
- store between 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, croscarmellose sodium, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sucralose, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-566-25
 TYLENOL[®]
 FOR ADULTS

COLD + HEAD
 CONGESTION SEVERE

Acetaminophen,
 Phenylephrine HCl, Guaifenesin
 Pain Reliever–Fever Reducer,
 Nasal Decongestant, Expectorant

- HEADACHE + BODY ACHES
- FEVER + SORE THROAT
- NASAL CONGESTION
- MUCUS + CHEST
 CONGESTION

Actual Size

24 CAPLETS



NDC 50580-566-25

TYLENOL[®]

FOR ADULTS

COLD + HEAD CONGESTION **SEVERE**

Acetaminophen,

Phenylephrine HCl, Guaifenesin

Pain Reliever-Fever Reducer,


Nasal Decongestant, Expectorant

- HEADACHE + BODY ACHES
- FEVER + SORE THROAT
- NASAL CONGESTION
- **MUCUS + CHEST
CONGESTION**



Actual Size

24 CAPLETS



Important: Read all product information before using. Keep this box for important information.

Made in Italy

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

©J&JCI 2020

30047053/
123662

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Guafenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Inactive ingredients

camellia wax, croscarmellose sodium, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sucralose, titanium dioxide, triacetin

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have ■ liver disease ■ heart disease

■ high blood pressure ■ thyroid disease ■ diabetes

■ trouble urinating due to an enlarged prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if ■ nervousness, dizziness, or sleepiness occur

■ pain, nasal congestion or cough gets worse or lasts more than 7 days

■ fevers get worse or lasts more than 3 days ■ redness or swelling is present

■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control

Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

adults and children 12 years

and over ■ take 2 caplets every 4 hours

■ swallow whole—do not crush, chew or dissolve

■ do not take more than 10 caplets in 24 hours

children under 12 years ask a doctor

Other information

■ each caplet contains: sodium 3 mg ■ store between 20°-25°C (68°-77°F)

■ do not use if blister unit is torn or broken

How can we help?
1-877-895-3665



TYLENOL COLD PLUS HEAD CONGESTION SEVERE

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-566
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients	
Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics			
Color	white	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	TYLENOL;1072
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-566-25	2 in 1 CARTON	07/15/2013	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/15/2013	

Labeler - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC