

AZATHIOPRINE- azathioprine tablet

Oceanside Pharmaceuticals


Azathioprine Tablets, USP 75mg and 100mg

WARNING - MALIGNANCY Chronic immunosuppression with this purine antimetabolite increases *risk of malignancy* in humans. Reports of malignancy include post-transplant lymphoma and hepatosplenic T-cell lymphoma (HSTCL) in patients with inflammatory bowel disease. Physicians using this drug should be very familiar with this risk as well as with the mutagenic potential to both men and women and with possible hematologic toxicities. Physicians should inform patients of the risk of malignancy with Azathioprine Tablets. See **WARNINGS**.

DESCRIPTION

Azathioprine, an immunosuppressive antimetabolite, is available in tablet form for oral administration. Each scored tablet contains 75 mg or 100 mg azathioprine and the inactive ingredients lactose monohydrate, pregelatinized starch, povidone, corn starch, magnesium stearate, and stearic acid.

Azathioprine is chemically 1 *H*-purine, 6-[(1-methyl-4-nitro-1 *H*-imidazol-5-yl)thio]-. The structural formula of azathioprine is:



It is an imidazolyl derivative of 6-mercaptopurine and many of its biological effects are similar to those of the parent compound.

Azathioprine is insoluble in water, but may be dissolved with addition of one molar equivalent of alkali. Azathioprine is stable in solution at neutral or acid pH but hydrolysis to mercaptopurine occurs in excess sodium hydroxide (0.1N), especially on warming. Conversion to mercaptopurine also occurs in the presence of sulfhydryl compounds such as cysteine, glutathione, and hydrogen sulfide.

CLINICAL PHARMACOLOGY

Azathioprine is well absorbed following oral administration. Maximum serum radioactivity occurs at 1 to 2 hours after oral ³⁵S-azathioprine and decays with a half-life of 5 hours.

This is not an estimate of the half-life of azathioprine itself, but is the decay rate for all ^{35}S -containing metabolites of the drug. Because of extensive metabolism, only a fraction of the radioactivity is present as azathioprine. Usual doses produce blood levels of azathioprine, and of mercaptopurine derived from it, which are low (<1 mcg/mL). Blood levels are of little predictive value for therapy since the magnitude and duration of clinical effects correlate with thiopurine nucleotide levels in tissues rather than with plasma drug levels. Azathioprine and mercaptopurine are moderately bound to serum proteins (30%) and are partially dialyzable. See **OVERDOSAGE**.

Azathioprine is metabolized to 6-mercaptopurine (6-MP). Both compounds are rapidly eliminated from blood and are oxidized or methylated in erythrocytes and liver; no azathioprine or mercaptopurine is detectable in urine after 8 hours. Activation of 6-mercaptopurine occurs via hypoxanthine-guanine phosphoribosyltransferase (HGPRT) and a series of multi-enzymatic processes involving kinases to form 6-thioguanine nucleotides (6-TGNs) as major metabolites. The cytotoxicity of azathioprine is due, in part, to the incorporation of 6-TGN into DNA.

6-MP undergoes two major inactivation routes. One is thiol methylation, which is catalyzed by the enzyme thiopurine S-methyltransferase (TPMT), to form the inactive metabolite methyl-6-MP (6-MeMP). Another inactivation pathway is oxidation, which is catalyzed by xanthine oxidase (XO) to form 6-thiouric acid. The nucleotide diphosphatase (NUDT15) enzyme is involved in conversion of the 6-TGNs to inactive 6-TG monophosphates. TPMT activity correlates inversely with 6-TGN levels in erythrocytes and presumably other hematopoietic tissues, since these cells have negligible xanthine oxidase (involved in the other inactivation pathway) activities.

Genetic polymorphisms influence TPMT and NUDT15 activity. Several published studies indicate that patients with reduced TPMT or NUDT15 activity receiving usual doses of 6-MP or azathioprine, accumulate excessive cellular concentrations of active 6-TGNs, and are at higher risk of severe myelosuppression. Because of the risk of toxicity, patients with TPMT or NUDT15 deficiency require alternative therapy or dose modification (see **DOSAGE AND ADMINISTRATION**).

Approximately 0.3% (1:300) of patients of European or African ancestry have two loss-of-function alleles of the TPMT gene and have little or no TPMT activity (homozygous deficient or poor metabolizers), and approximately 10% of patients have one loss-of-function TPMT allele leading to intermediate TPMT activity (heterozygous deficient or intermediate metabolizers). The TPMT*2, TPMT*3A, and TPMT*3C alleles account for about 95% of individuals with reduced levels of TPMT activity. NUDT15 deficiency is detected in $<1\%$ of patients of European or African ancestry. Among patients of East Asian ancestry (i.e., Chinese, Japanese, Vietnamese), 2% have two loss-of-function alleles of the NUDT15 gene, and approximately 21% have one loss-of-function allele. The p.R139C variant of NUDT15 (present on the *2 and *3 alleles) is the most commonly observed, but other less common loss-of-function NUDT15 alleles have been observed.

Inhibition of xanthine oxidase (XO) may cause increased plasma concentrations of azathioprine or its metabolites leading to toxicity (**see PRECAUTIONS: Drug Interactions**). Proportions of metabolites are different in individual patients, and this presumably accounts for variable magnitude and duration of drug effects. Renal clearance is probably not important in predicting biological effectiveness or toxicities, although dose reduction is practiced in patients with poor renal function.

Homograft Survival: The use of azathioprine for inhibition of renal homograft rejection

is well established, the mechanism(s) for this action are somewhat obscure. The drug suppresses hypersensitivities of the cell-mediated type and causes variable alterations in antibody production. Suppression of T-cell effects, including ablation of T-cell suppression, is dependent on the temporal relationship to antigenic stimulus or engraftment. This agent has little effect on established graft rejections or secondary responses.

Alterations in specific immune responses or immunologic functions in transplant recipients are difficult to relate specifically to immunosuppression by azathioprine. These patients have subnormal responses to vaccines, low numbers of T-cells, and abnormal phagocytosis by peripheral blood cells, but their mitogenic responses, serum immunoglobulins, and secondary antibody responses are usually normal.

Immunoinflammatory Response: Azathioprine suppresses disease manifestations as well as underlying pathology in animal models of autoimmune disease. For example, the severity of adjuvant arthritis is reduced by azathioprine.

The mechanisms whereby azathioprine affects autoimmune diseases are not known. Azathioprine is immunosuppressive, delayed hypersensitivity and cellular cytotoxicity tests being suppressed to a greater degree than are antibody responses. In the rat model of adjuvant arthritis, azathioprine has been shown to inhibit the lymph node hyperplasia, which precedes the onset of the signs of the disease. Both the immunosuppressive and therapeutic effects in animal models are dose-related. Azathioprine is considered a slow-acting drug and effects may persist after the drug has been discontinued.

INDICATIONS AND USAGE

Azathioprine Tablets are indicated as an adjunct for the prevention of rejection in renal homotransplantation. It is also indicated for the management of active rheumatoid arthritis to reduce signs and symptoms.

Renal Homotransplantation: Azathioprine Tablets are indicated as an adjunct for the prevention of rejection in renal homotransplantation. Experience with over 16,000 transplants shows a 5-year patient survival of 35% to 55%, but this is dependent on donor, match for HLA antigens, anti-donor or anti-B-cell alloantigen antibody, and other variables. The effect of Azathioprine Tablets on these variables has not been tested in controlled trials.

Rheumatoid Arthritis: Azathioprine Tablets are indicated for the treatment of active rheumatoid arthritis (RA) to reduce signs and symptoms. Aspirin, non-steroidal anti-inflammatory drugs and/or low dose glucocorticoids may be continued during treatment with Azathioprine Tablets. The combined use of Azathioprine Tablets with disease modifying anti-rheumatic drugs (DMARDs) has not been studied for either added benefit or unexpected adverse effects. The use of Azathioprine Tablets with these agents cannot be recommended.

CONTRAINDICATIONS

Azathioprine Tablets should not be given to patients who have shown hypersensitivity to the drug.

Azathioprine Tablets should not be used for treating rheumatoid arthritis in pregnant women. Patients with rheumatoid arthritis previously treated with alkylating agents (cyclophosphamide, chlorambucil, melphalan, or others) may have a prohibitive risk of malignancy if treated with Azathioprine Tablets.

WARNINGS

Malignancy: Patients receiving immunosuppressants, including Azathioprine Tablets, are at increased risk of developing lymphoma and other malignancies, particularly of the skin. Physicians should inform patients of the risk of malignancy with Azathioprine Tablets. As usual for patients with increased risk for skin cancer, exposure to sunlight and ultraviolet light should be limited by wearing protective clothing and using a sunscreen with a high protection factor.

Post-transplant: Renal transplant patients are known to have an increased risk of malignancy, predominantly skin cancer and reticulum cell or lymphomatous tumors. The risk of post-transplant lymphomas may be increased in patients who receive aggressive treatment with immunosuppressive drugs, including Azathioprine Tablets. Therefore, immunosuppressive drug therapy should be maintained at the lowest effective levels.

Rheumatoid Arthritis: Information is available on the risk of malignancy with the use of Azathioprine Tablets in rheumatoid arthritis (see **ADVERSE REACTIONS**). It has not been possible to define the precise risk of malignancy due to Azathioprine Tablets. The data suggest the risk may be elevated in patients with rheumatoid arthritis, though lower than for renal transplant patients. However, acute myelogenous leukemia as well as solid tumors have been reported in patients with rheumatoid arthritis who have received Azathioprine Tablets.

Inflammatory Bowel Disease: Postmarketing cases of hepatosplenic T-cell lymphoma (HSTCL), a rare type of T-cell lymphoma, have been reported in patients treated with Azathioprine Tablets. These cases have had a very aggressive disease course and have been fatal. The majority of reported cases have occurred in patients with Crohn's disease or ulcerative colitis and the majority were in adolescent and young adult males. Some of the patients were treated with Azathioprine Tablets as monotherapy and some had received concomitant treatment with a TNF α blocker at or prior to diagnosis. The safety and efficacy of Azathioprine Tablets for the treatment of Crohn's disease and ulcerative colitis have not been established.

Cytopenias: Severe leukopenia, thrombocytopenia, anemias including macrocytic anemia and/or pancytopenia may occur in patients being treated with Azathioprine Tablets. Severe bone marrow suppression may also occur. Hematologic toxicities are dose-related and may be more severe in renal transplant patients whose homograft is undergoing rejection. It is suggested that patients on Azathioprine Tablets have complete blood counts, including platelet counts, weekly during the first month, twice monthly for the second and third months of treatment, then monthly or more frequently if dosage alterations or other therapy changes are necessary. Delayed hematologic suppression may occur. Prompt reduction in dosage or temporary withdrawal of the drug may be necessary if there is a rapid fall in or persistently low leukocyte count, or other evidence of bone marrow depression. Leukopenia does not correlate with therapeutic effect; therefore the dose should not be increased intentionally to lower the white blood cell count.

TPMT or NUDT15 Deficiency: Patients with thiopurine S-methyl transferase (TPMT) or nucleotide diphosphatase (NUDT15) deficiency may be at an increased risk of severe and life-threatening myelotoxicity if receiving conventional doses of Azathioprine Tablets (see **CLINICAL PHARMACOLOGY**). Death associated with pancytopenia has been reported in patients with absent TPMT activity receiving azathioprine. In patients with severe myelosuppression, consider evaluation for TPMT and NUDT15 deficiency (see **PRECAUTIONS: Laboratory Tests**). Consider alternative therapy in patients with homozygous TPMT or NUDT15 deficiency and reduced dosages in patients with heterozygous deficiency (see **DOSAGE AND ADMINISTRATION**).

Serious Infections: Patients receiving immunosuppressants, including Azathioprine Tablets, are at increased risk for bacterial, viral, fungal, protozoal, and opportunistic infections, including reactivation of latent infections. These infections may lead to serious, including fatal outcomes.

Progressive Multifocal Leukoencephalopathy: Cases of JC virus-associated infection resulting in progressive multifocal leukoencephalopathy (PML), sometimes fatal, have been reported in patients treated with immunosuppressants, including Azathioprine Tablets. Risk factors for PML include treatment with immunosuppressant therapies and impairment of immune function. Consider the diagnosis of PML in any patient presenting with new-onset neurological manifestations and consider consultation with a neurologist as clinically indicated. Consider reducing the amount of immunosuppression in patients who develop PML. In transplant patients, consider the risk that the reduced immunosuppression represents to the graft.

Effect on Sperm in Animals: Azathioprine Tablets has been reported to cause temporary depression in spermatogenesis and reduction in sperm viability and sperm count in mice at doses 10 times the human therapeutic dose; ¹ a reduced percentage of fertile matings occurred when animals received 5 mg/kg. ²

Pregnancy: Azathioprine Tablets can cause fetal harm when administered to a pregnant woman. Azathioprine Tablets should not be given during pregnancy without careful weighing of risk versus benefit. Whenever possible, use of Azathioprine Tablets in pregnant patients should be avoided. This drug should not be used for treating rheumatoid arthritis in pregnant women. ³

Azathioprine Tablets are teratogenic in rabbits and mice when given in doses equivalent to the human dose (5 mg/kg daily). Abnormalities included skeletal malformations and visceral anomalies. ²

Postmarketing cases of intrahepatic cholestasis of pregnancy (ICP) have been reported in women treated with azathioprine during pregnancy. ICP symptoms and evaluated bile acid levels improved following azathioprine discontinuation. Discontinue Azathioprine Tablets if ICP develops in a pregnant woman.

Limited immunologic and other abnormalities have occurred in a few infants born of renal allograft recipients on Azathioprine Tablets. In a detailed case report,⁴ documented lymphopenia, diminished IgG and IgM levels, CMV infection, and a decreased thymic shadow were noted in an infant born to a mother receiving 150 mg azathioprine and 30 mg prednisone daily throughout pregnancy. At 10 weeks most features were normalized. DeWitte et al reported pancytopenia and severe immune deficiency in a preterm infant whose mother received 125 mg azathioprine and 12.5 mg prednisone daily. ⁵ There have been two published reports of abnormal physical findings. Williamson

and Karp described an infant born with preaxial polydactyly whose mother received azathioprine 200 mg daily and prednisone 20 mg every other day during pregnancy.⁶ Tallent et al described an infant with a large myelomeningocele in the upper lumbar region, bilateral dislocated hips, and bilateral talipes equinovarus. The father was on long-term azathioprine therapy.⁷

Benefit versus risk must be weighed carefully before use of Azathioprine Tablets in patients of reproductive potential. There are no adequate and well-controlled studies in pregnant women. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing age should be advised to avoid becoming pregnant.

PRECAUTIONS

General: A gastrointestinal hypersensitivity reaction characterized by severe nausea and vomiting has been reported. These symptoms may also be accompanied by diarrhea, rash, fever, malaise, myalgias, elevations in liver enzymes, and occasionally, hypotension. Symptoms of gastrointestinal toxicity most often develop within the first several weeks of therapy with Azathioprine Tablets and are reversible upon discontinuation of the drug. The reaction can recur within hours after rechallenge with a single dose of Azathioprine Tablets.

Information for Patients: Patients being started on Azathioprine Tablets should be informed of the necessity of periodic blood counts while they are receiving the drug and should be encouraged to report any unusual bleeding or bruising to their physician. They should be informed of the danger of infection while receiving Azathioprine Tablets and asked to report signs and symptoms of infection to their physician. Careful dosage instructions should be given to the patient, especially when Azathioprine Tablets are being administered in the presence of impaired renal function or concomitantly with allopurinol (see **Drug Interactions** subsection and **DOSAGE AND ADMINISTRATION**). Patients should be advised of the potential risks of the use of Azathioprine Tablets during pregnancy and during the nursing period. The increased risk of malignancy following therapy with Azathioprine Tablets should be explained to the patient.

Laboratory Tests:

Complete Blood Count (CBC) Monitoring: Patients on Azathioprine Tablets should have complete blood counts, including platelet counts, weekly during the first month, twice monthly for the second and third months of treatment, then monthly or more frequently if dosage alterations or other therapy changes are necessary.

TPMT and NUDT15 Testing: Consider genotyping or phenotyping patients for TPMT-deficiency and genotyping for NUDT15 deficiency in patients with severe myelosuppression. TPMT and NUDT15 testing cannot substitute for complete blood count (CBC) monitoring in patients receiving Azathioprine Tablets. Accurate phenotyping (red blood cell TPMT activity) results are not possible in patients who have received recent blood transfusions (see **CLINICAL PHARMACOLOGY, WARNINGS:**

Cytopenias, ADVERSE REACTIONS and DOSAGE AND ADMINISTRATION sections).

Drug Interactions:

Use with xanthine oxidase (XO) inhibitors: One of the pathways for inactivation of

azathioprine is inhibited by XO inhibitors (allopurinol or febuxostat). Patients receiving Azathioprine Tablets and allopurinol concomitantly should have a dose reduction of Azathioprine Tablets, to approximately 1/3 to 1/4 the usual dose. Concomitant use of Azathioprine Tablets with febuxostat is not recommended. Inhibition of XO may cause increased plasma concentrations of azathioprine or its metabolite, 6-MP, leading to toxicity. It is recommended that a further dose reduction or alternative therapies be considered for patients with low or absent TPMT activity receiving Azathioprine Tablets and xanthine oxidase inhibitors because both TPMT and XO inactivation pathways are affected (see **CLINICAL PHARMACOLOGY, WARNINGS, PRECAUTIONS: Laboratory Tests** and **ADVERSE REACTIONS** sections).

Use with Aminosalicylates: There is in vitro evidence that aminosalicylate derivatives (e.g., sulphasalazine, mesalazine, or olsalazine) inhibit the TPMT enzyme. Concomitant use of these agents with Azathioprine Tablets should be done with caution.

Use with Other Agents Affecting Myelopoiesis: Drugs which may affect leukocyte production, including co-trimoxazole, may lead to exaggerated leukopenia, especially in renal transplant recipients.

Use with Angiotensin-Converting Enzyme Inhibitors: The use of angiotensin-converting enzyme inhibitors to control hypertension in patients on azathioprine has been reported to induce anemia and severe leukopenia.

Use with Warfarin: Azathioprine Tablets may inhibit the anticoagulant effect of warfarin.

Use with ribavirin: The use of ribavirin for hepatitis C in patients receiving azathioprine has been reported to induce severe pancytopenia and may increase the risk of azathioprine-related myelotoxicity. Inosine monophosphate dehydrogenase (IMDH) is required for one of the metabolic pathways of azathioprine. Ribavirin is known to inhibit IMDH, thereby leading to accumulation of an azathioprine metabolite, 6-methylthioinosine monophosphate (6-MTITP), which is associated with myelotoxicity (neutropenia, thrombocytopenia, and anemia). Patients receiving azathioprine with ribavirin should have complete blood counts, including platelet counts, monitored weekly for the first month, twice monthly for the second and third months of treatment, then monthly or more frequently if dosage or other therapy changes are necessary.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

See **WARNINGS** section.

Pregnancy:

Teratogenic Effects: See **WARNINGS** section.

Nursing Mothers: The use of Azathioprine Tablets in nursing mothers is not recommended. Azathioprine or its metabolites are transferred at low levels, both transplacentally and in breast milk.^{8, 9, 10} Because of the potential for tumorigenicity shown for azathioprine, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use: Safety and efficacy of azathioprine in pediatric patients have not been established.

ADVERSE REACTIONS

The principal and potentially serious toxic effects of Azathioprine Tablets are hematologic and gastrointestinal. The risks of secondary infection and malignancy are also significant (see **WARNINGS**). The frequency and severity of adverse reactions depend on the dose and duration of Azathioprine Tablets as well as on the patient's underlying disease or concomitant therapies. The incidence of hematologic toxicities and neoplasia encountered in groups of renal homograft- recipients is significantly higher than that in studies employing Azathioprine Tablets for rheumatoid arthritis. The relative incidences in clinical studies are summarized below:

| Toxicity | Renal Homograft | Rheumatoid Arthritis |
|-----------------------------|--------------------|-------------------------|
| Leukopenia (any degree) | >50% | 28% |
| <2500 cells/mm ³ | 16% | 5.3% |
| Infections | 20% | <1% |
| Neoplasia | | * |
| Lymphoma | 0.5% | |
| Others | 2.8% | |

*Data on the rate and risk of neoplasia among persons with rheumatoid arthritis treated with azathioprine are limited. The incidence of lymphoproliferative disease in patients with RA appears to be significantly higher than that in the general population. In one completed study, the rate of lymphoproliferative disease in RA patients receiving higher than recommended doses of azathioprine (5 mg/kg per day) was 1.8 cases per 1000 patient-years of follow-up, compared with 0.8 cases per 1000 patient-years of follow-up in those not receiving azathioprine. However, the proportion of the increased risk attributable to the azathioprine dosage or to other therapies (i.e., alkylating agents) received by patients treated with azathioprine cannot be determined.

Hematologic: Leukopenia and/or thrombocytopenia are dose-dependent and may occur late in the course of therapy with Azathioprine Tablets. Dose reduction or temporary withdrawal may result in reversal of these toxicities. Infection may occur as a secondary manifestation of bone marrow suppression or leukopenia, but the incidence of infection in renal homotransplantation is 30 to 60 times that in rheumatoid arthritis. Anemias, including macrocytic anemia, and/or bleeding have been reported.

Patients with low or absent TPMT or NUDT15 activity are at increased risk for severe, life-threatening myelosuppression from Azathioprine Tablets (see **CLINICAL PHARMACOLOGY, WARNINGS: Cytopenias** and **PRECAUTIONS: Laboratory Tests, DOSAGE AND ADMINISTRATION**).

Gastrointestinal: Nausea and vomiting may occur within the first few months of therapy with Azathioprine Tablets, and occurred in approximately 12% of 676 rheumatoid arthritis patients. The frequency of gastric disturbance often can be reduced by administration of the drug in divided doses and/or after meals. However, in some patients, nausea and vomiting may be severe and may be accompanied by symptoms such as diarrhea, fever, malaise, and myalgias (see **PRECAUTIONS**). Vomiting with abdominal pain may occur rarely with a hypersensitivity pancreatitis. Hepatotoxicity manifest by elevation of serum alkaline phosphatase, bilirubin, and/or serum transaminases is known to occur following azathioprine use, primarily in allograft

recipients. Hepatotoxicity has been uncommon (less than 1%) in rheumatoid arthritis patients. Hepatotoxicity following transplantation most often occurs within 6 months of transplantation and is generally reversible after interruption of Azathioprine Tablets. A rare, but life-threatening hepatic veno-occlusive disease associated with chronic administration of azathioprine has been described in transplant patients and in one patient receiving Azathioprine Tablets for panuveitis.^{11, 12, 13} Periodic measurement of serum transaminases, alkaline phosphatase, and bilirubin is indicated for early detection of hepatotoxicity. If hepatic veno-occlusive disease is clinically suspected, Azathioprine Tablets should be permanently withdrawn.

Others: Additional side effects of low frequency have been reported. These include skin rashes, alopecia, fever, arthralgias, diarrhea, steatorrhea, negative nitrogen balance, reversible interstitial pneumonitis, hepatosplenic T-cell lymphoma (see **WARNINGS - Malignancy**), and Sweet's Syndrome (acute febrile neutrophilic dermatosis).

Postmarketing Experience

The following adverse reactions have been identified during postapproval use of Azathioprine Tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a casual relationship to drug exposure.

- intrahepatic cholestasis of pregnancy (see **WARNINGS, Pregnancy**).

To report SUSPECTED ADVERSE REACTIONS, contact Oceanside Pharmaceuticals at 1-800-321-4576 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

The oral LD₅₀s for single doses of Azathioprine Tablets in mice and rats are 2500 mg/kg and 400 mg/kg, respectively. Very large doses of this antimetabolite may lead to marrow hypoplasia, bleeding, infection, and death. About 30% of Azathioprine Tablets are bound to serum proteins, but approximately 45% is removed during an 8-hour hemodialysis.¹⁴ A single case has been reported of a renal transplant patient who ingested a single dose of 7500 mg Azathioprine Tablets. The immediate toxic reactions were nausea, vomiting, and diarrhea, followed by mild leukopenia and mild abnormalities in liver function. The white blood cell count, SGOT, and bilirubin returned to normal 6 days after the overdose.

DOSAGE AND ADMINISTRATION

Renal Homotransplantation: The dose of Azathioprine Tablets required to prevent rejection and minimize toxicity will vary with individual patients; this necessitates careful management. The initial dose is usually 3 to 5 mg/kg daily, beginning at the time of transplant. Azathioprine Tablets are usually given as a single daily dose on the day of, and in a minority of cases 1 to 3 days before, transplantation. Dose reduction to maintenance levels of 1 to 3 mg/kg daily is usually possible. The dose of Azathioprine Tablets should not be increased to toxic levels because of threatened rejection. Discontinuation may be necessary for severe hematologic or other toxicity, even if rejection of the homograft may be a consequence of drug withdrawal.

Rheumatoid Arthritis: Azathioprine Tablets are usually given on a daily basis. The initial

dose should be approximately 1.0 mg/kg (50 to 100 mg) given as a single dose or on a twice-daily schedule. The dose may be increased, beginning at 6 to 8 weeks and thereafter by steps at 4-week intervals, if there are no serious toxicities and if initial response is unsatisfactory. Dose increments should be 0.5 mg/kg daily, up to a maximum dose of 2.5 mg/kg per day. Therapeutic response occurs after several weeks of treatment, usually 6 to 8; an adequate trial should be a minimum of 12 weeks. Patients not improved after 12 weeks can be considered refractory. Azathioprine Tablets may be continued long-term in patients with clinical response, but patients should be monitored carefully, and gradual dosage reduction should be attempted to reduce risk of toxicities.

Maintenance therapy should be at the lowest effective dose, and the dose given can be lowered decrementally with changes of 0.5 mg/kg or approximately 25 mg daily every 4 weeks while other therapy is kept constant. The optimum duration of maintenance Azathioprine Tablets has not been determined. Azathioprine Tablets can be discontinued abruptly, but delayed effects are possible.

Patients with TPMT and/or NUDT15 Deficiency: Consider testing for TPMT and NUDT15 deficiency in patients who experience severe bone marrow toxicities. Early drug discontinuation may be considered in patients with abnormal CBC results that do not respond to dose reduction (see **CLINICAL PHARMACOLOGY, WARNINGS: Cytopenias, and PRECAUTIONS: Laboratory Tests**).

Homozygous deficiency in either TPMT or NUDT15:

Because of the risk of increased toxicity, consider alternative therapies for patients who are known to have TPMT or NUDT15 deficiency (see **CLINICAL PHARMACOLOGY, WARNINGS: Cytopenias, and PRECAUTIONS: Laboratory Tests**).

Heterozygous deficiency in TPMT and/or NUDT15:

Because of the risk of increased toxicity, dosage reduction is recommended in patients known to have heterozygous deficiency of TPMT or NUDT15. Patients who are heterozygous for both TPMT and NUDT15 deficiency may require more substantial dosage reductions (see **CLINICAL PHARMACOLOGY, WARNINGS: Cytopenias, and PRECAUTIONS: Laboratory Tests**).

Use in Renal Dysfunction: Relatively oliguric patients, especially those with tubular necrosis in the immediate postcadaveric transplant period, may have delayed clearance of Azathioprine Tablets or its metabolites, may be particularly sensitive to this drug, and are usually given lower doses.

Procedures for proper handling and disposal of this immunosuppressive antimetabolite drug should be considered. Several guidelines on this subject have been published.¹⁵⁻²¹ There is no general agreement that all of the procedures recommended in the guidelines are necessary or appropriate.

HOW SUPPLIED

Azathioprine Tablets, USP are available in:

75 mg, triangle-shaped, yellow, scored tablets,
100 count bottles (NDC 68682-231-01)

100 mg, diamond-shaped, yellow, scored tablets,

100 count bottles (NDC 68682-241-01)

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Store in a dry place and protect from light.

Dispense in a tight, light-resistant container as defined in the USP.

REFERENCES

1. Clark JM. The mutagenicity of azathioprine in mice, *Drosophila melanogaster*, and *Neurospora crassa*. *Mutat Res*. 1975; 28:87-99.
2. Data on file, Sebela Ireland Ltd.
3. Tagatz GE, Simmons RL. Pregnancy after renal transplantation. *Ann Intern Med*. 1975; 82:113-114. Editorial Notes.
4. Cote' CJ, Meuwissen HJ, Pickering RJ. Effects on the neonate of prednisone and azathioprine administered to the mother during pregnancy. *J Pediatr*. 1974; 85:324-328.
5. DeWitte DB, Buick MK, Cyran SE, et al. Neonatal pancytopenia and severe combined immunodeficiency associated with antenatal administration of azathioprine and prednisone. *J Pediatr*. 1984; 105:625-628.
6. Williamson RA, Karp LE. Azathioprine teratogenicity: review of the literature and case report. *Obstet Gynecol*. 1981; 58:247-250.
7. Tallent MB, Simmons RL, Najarian JS. Birth defects in child of male recipient of kidney transplant. *JAMA*. 1970; 211:1854-1855.
8. Data on file, Sebela Ireland Ltd.
9. Saarikoski S, Seppälä M. Immunosuppression during pregnancy: transmission of azathioprine and its metabolites from the mother to the fetus. *Am J Obstet Gynecol*. 1973; 115:1100-1106.
10. Coulam CB, Moyer TP, Jiang NS, et al. Breast-feeding after renal transplantation. *Transplant Proc*. 1982; 14: 605-609.
11. Read AE, Wiesner RH, LaBrecque DR, et al. Hepatic veno-occlusive disease associated with renal transplantation and azathioprine therapy. *Ann Intern Med*. 1986; 104:651-655.
12. Katzka DA, Saul SH, Jorkasky D, et al. Azathioprine and hepatic veno-occlusive disease in renal transplant patients. *Gastroenterology*. 1986; 90:446-454.
13. Weitz H, Gokel JM, Loeshke K, et al. Veno-occlusive disease of the liver in patients receiving immunosuppressive therapy. *Virchows Arch A Pathol Anat Histol*. 1982; 395:245-256.
14. Schusziarra V, Ziekursch V, Schlamp R, et al. Pharmacokinetics of azathioprine under haemodialysis. *Int J Clin Pharmacol Biopharm*. 1976; 14:298-302.
15. Recommendations for the safe handling of parenteral antineoplastic drugs. Washington, DC: Division of Safety; Clinical Center Pharmacy Department and Cancer Nursing Services, National Institute of Health; 1992. US Dept of Health and Human

Services. Public Health Service Publication NIH 92-2621.

16. AMA Council on Scientific Affairs. Guidelines for handling parenteral antineoplastics. *JAMA*. 1985; 253:1590-1592.

17. National Study Commission on Cytotoxic Exposure. Recommendations for handling cytotoxic agents. 1987. Available from Louis P. Jeffrey, Chairman, National Study Commission on Cytotoxic Exposure. Massachusetts College of Pharmacy and Allied Health Sciences, 179 Longwood Avenue, Boston, MA 02115.

18. Clinical Oncological Society of Australia. Guidelines and recommendations for safe handling of antineoplastic agents. *Med J Aust*. 1983; 1:426-428.

19. Jones RB, Frank R, Mass T. Safe handling of chemotherapeutic agents: a report from The Mount Sinai Medical Center. *CA Cancer J for Clinicians*. 1983; 33:258-263.

20. American Society of Hospital Pharmacists. ASHP technical assistance bulletin on handling cytotoxic and hazardous drugs. *Am J Hosp Pharm*. 1990; 47:1033-1049.

21. Yodaiken RE, Bennett D. OSHA Work-Practice guidelines for personnel dealing with cytotoxic (antineoplastic) drugs. *Am J Hosp Pharm*, 1996; 43:1193-1204.

Distributed by:

Oceanside Pharmaceuticals, a division of Bausch Health US, LLC

Bridgewater, NJ 08807 USA

© 2022 Bausch Health Companies Inc. or its affiliates

Manufactured by:

Alcami Corporation

Wilmington, NC 28405 USA

Rev. 08/2024 PC 5423C

AZATHIOPRINE TABLETS, USP 75mg

100 Tablets

NDC 68682-231-01

75 mg

Rx only

Distributed by:

Oceanside Pharmaceuticals, a division of Bausch Health US, LLC

Bridgewater, NJ 08807 USA

Manufactured by:

Alcami Corporation


Wilmington, NC 28405 USA

NDC 68682-**231**-01 Rx only

**AZATHIOPRINE
TABLETS, USP**

75 mg

100 Tablets



OCEANSIDE
PHARMACEUTICALS



Each scored, yellow tablet contains: Azathioprine, USP 75 mg

Usual Dosage: For indications, dosage, precautions, etc., see accompanying package insert.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature] in a dry place and protect from light.
Dispense in a tight, light-resistant container as defined in the USP.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Distributed by:

Oceanside Pharmaceuticals, a division of Bausch Health US, LLC
Bridgewater, NJ 08807 USA

Manufactured by: Alcami Corporation
Wilmington, NC 28405 USA

© 2023 Bausch Health Companies Inc. or its affiliates
PC5421B

AZATHIOPRINE TABLETS, USP 100 mg

100 Tablets

NDC 68682-241-01

100 mg

Rx only

Distributed by:

Oceanside Pharmaceuticals, a division of Bausch Health US, LLC

Bridgewater, NJ 08807 USA

Manufactured by:

Alcami Corporation

Wilmington, NC 28405 USA


© 2023 Bausch Health Companies Inc. or its affiliates

NDC 68682-**241**-01 Rx only

**AZATHIOPRINE
TABLETS, USP**

100 mg

100 Tablets



OCEANSIDE
PHARMACEUTICALS



Each scored, yellow tablet contains: Azathioprine, USP 100 mg

Usual Dosage: For indications, dosage, precautions, etc., see accompanying package insert.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature] in a dry place and protect from light.
Dispense in a tight, light-resistant container as defined in the USP.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Distributed by:

Oceanside Pharmaceuticals, a division of Bausch Health US, LLC
Bridgewater, NJ 08807 USA

Manufactured by: Alcami Corporation
Wilmington, NC 28405 USA

© 2023 Bausch Health Companies Inc. or its affiliates
PC5422B

AZATHIOPRINE

azathioprine tablet

Product Information

| | | | |
|-------------------------|-------------------------|--------------------|---------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:68682-231 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| AZATHIOPRINE (UNII: MRK240IY2L) (AZATHIOPRINE - UNII:MRK240IY2L) | AZATHIOPRINE | 75 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| POVIDONE (UNII: FZ989GH94E) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| MAGNESIUM STEARATE (UNII: 70097M6I3O) | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | |

Product Characteristics

| | | | |
|----------|----------|--------------|----------|
| Color | yellow | Score | 2 pieces |
| Shape | TRIANGLE | Size | 14mm |
| Flavor | | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:68682-231-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 02/04/2023 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA075252 | 02/04/2023 | |

AZATHIOPRINE

azathioprine tablet

| Product Information | | | | |
|--|--|--|----------------------|--------------------|
| Product Type | | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:68682-241 |
| Route of Administration | | ORAL | | |
| | | | | |
| Active Ingredient/Active Moiety | | | | |
| Ingredient Name | | | Basis of Strength | Strength |
| AZATHIOPRINE (UNII: MRK240IY2L) (AZATHIOPRINE - UNII:MRK240IY2L) | | | AZATHIOPRINE | 100 mg |
| | | | | |
| Inactive Ingredients | | | | |
| Ingredient Name | | | | Strength |
| STEARIC ACID (UNII: 4ELV7Z65AP) | | | | |
| POVIDONE (UNII: FZ989GH94E) | | | | |
| STARCH, CORN (UNII: O8232NY3SJ) | | | | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | | | | |
| LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X) | | | | |
| | | | | |
| Product Characteristics | | | | |
| Color | yellow | Score | 2 pieces | |
| Shape | DIAMOND | Size | 14mm | |
| Flavor | | Imprint Code | | |
| Contains | | | | |
| | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:68682-241-01 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 02/04/2023 | |
| | | | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| ANDA | ANDA075252 | | 02/04/2023 | |

Labeler - Oceanside Pharmaceuticals (832011691)

Registrant - Alcami Carolinas Corporation (832395235)

Establishment

| Name | Address | ID/FEI | Business Operations |
|------------------------------|---------|-----------|----------------------------|
| Alcami Carolinas Corporation | | 079646861 | pack(68682-231, 68682-241) |

| Establishment | | | |
|------------------------------|---------|-----------|--------------------------------|
| Name | Address | ID/FEI | Business Operations |
| Alcami Carolinas Corporation | | 831351445 | analysis(68682-231, 68682-241) |

| Establishment | | | |
|------------------------------|---------|-----------|-----------------------------------|
| Name | Address | ID/FEI | Business Operations |
| Alcami Carolinas Corporation | | 832395235 | manufacture(68682-231, 68682-241) |

Revised: 10/2024

Oceanside Pharmaceuticals