


CS 4720/5720 Design and Analysis of Algorithms Homework #1 Student: (Robert Denim Horton)

Answers to homework problems:


1.

- (a) Given the formal definition of a pivotal node, a node can be defined as such when the node of interest exists along every shortest possible path in a given pair of nodes. For example, given a graph with the set of nodes S_0 with defined nodes $\{A, B, C, D, E\}$. The graph could be represented as;




As we can see in this diagram all the nodes are pivotal nodes. Starting with node pivotal node A , its only pivotal node pair is E and B , for node B its only pivotal node pair is C and A , for node C its only pivotal node pair is B and D , its only pivotal node pair is C and E , and lastly for node E its only pivotal node pair is D and A . Here we can see that every node that exists has at least one pair of nodes that also exists in the graph.

- (b) For a pivotal node to have two different pairs of pivotal nodes, we can again define a node with two different pairs of pivotal nodes as a node that is along the fastest path for two nodes but with the added implication that this node is along one other pair of pivotal nodes. With the same set, S_0 , from part a we can construct a graph to be represented as,



We can see that for every node in the graph, it has at least two different pairs of pivotal nodes. Node A has its first pivotal nodes as E and B and second pair of pivotal nodes C and D , node B has its first set of pivotal nodes C and A and second pair of pivotal nodes E and D , node C has its first set of pivotal nodes E and A and its second set of pivotal nodes B and D , node D has its first set of pivotal nodes B and A and second pair of pivotal nodes C and E , node E has its first pair of pivotal nodes D and A and its second set of pivotal nodes C and B and its second set of pivotal nodes A and D .

- (c) For a graph comprised of atleast 4 nodes where a single node is pivotla for any pair of nodes that comprise a pivotal pair. So with a different set, S_1 , of nodes $\{A, B, C, D, X\}$ we can use a graph to represnt a graph where node X is the node of intrest,



Here we that node X has a pivotla pair with every node that is not X in the graph. So node X has pivotal pairs A and C , pivotal pair B and D , pivotal pair A and B , pivotal pair C and B , pivotal pair D and A , and pivotal pair C and D . So six possible pairs in total.

2.

- (a) Give an example (together with an explanation) of a graph in which more than half of all nodes are gatekeepers.
- (b) Give an example (together with an explanation) of a graph in which there are no gatekeepers, but in which every node is a local gatekeeper

3.

- (a) Describe an example of a graph where the diameter is more than three times as large as the average distance.
- (b) Describe how you could extend your construction to produce graphs in which the diameter exceeds the average distance by as large a factor as you'd like. (That is, for every number c , can you produce a graph in which the diameter is more than c times as large as the average distance?)