

## 1 Overview

This library implements a number of functions for mathematical operations, it has dependencies upon the scala-nlp-breeze matrix library, and its purpose is to explore the implementation and usage of a variety of tools and techniques starting with some statistical tools. This document is structured with the main headings corresponding to the subject areas implemented in the library, and minor headings corresponding to the major package names in the project. Each subsection below the minor headings is also decomposed into individual subject areas, and the lowest level heading represents a class.

## 2 Methods of Counting

### 2.1 package au.id.cxd.math.count

The count package contains a series of modules dedicated to methods of counting.

#### 2.1.1 Factorial

The factorial operation is provided as  $n!$  implementing:

$$\prod_{i=1}^{n-1} (n-i)$$

The Factorial implementation will memoize results, allowing for efficient reuse during runtime.

#### 2.1.2 Choose

The choose module implements  $\binom{n}{m}$ , how many ways can m items be selected with replacement from a set of n items.

Determined as:

$$\frac{n!}{m!n!}$$

#### 2.1.3 Permutation

The method of selecting m ordered items from a set of n ordered items  $P\binom{n}{m}$ .

$$\frac{n!}{(n-m)!}$$

## 3 Probability

### 3.1 package au.id.cxd.math.probability

This package provides a series of modules that support operations for inference via probability, and for estimation of distributions.

### 3.1.1 Inequalities

The class TchebysheffInequality implements a simple estimation of a pdf using the inequality rule:

$$P(\mu - k\sigma < Y < \mu + k\sigma) \geq 1 - \frac{1}{k^2}$$

Which can be restated as:

$$P(lower < Y < upper) \geq 1 - \frac{1}{k^2}$$

The value of  $k$  is derived from either upper and lower bounds since:

$$k = \frac{upper - \mu}{\sigma}$$

After determining the value of  $k$  the probability can be estimated by substituting

$$p = 1 - \frac{1}{k^2}$$

### 3.1.2 Discrete Distributions

The discrete distributions packages contains the following.

#### Binomial

The binomial distribution (class name Binomial) has the parameters  $p$  for the prior proportion of successes and  $n$  for the total number of trials and calculates the probability of  $y$  successes

$$P(y; n; p) = \sum_{i=1}^n \binom{n}{y_i} p_i^y (1-p)^{n-y_i}$$

The properties of the binomial are:

Mean:  $\mu = np$

Variance:  $\sigma^2 = np(1-p) = npq$

#### Geometric Distribution

The geometric distribution (class name Geometric) with 1 parameter for probability  $p$  and  $y \geq 1$ . The variable represents the  $n$ th trial where the success occurs (for instance if  $y=2$  then the trial was successful on the 2nd attempt). The parameter  $p$  represents the probability of success. The probability function calculates the probability of success at the  $n$ th trial.

$$P(y; p) = p(1-p)^{y-1}$$

The simple properties of the distribution are:

Mean:  $\mu = \frac{1}{p}$

Variance:  $\sigma^2 = \frac{1-p}{p^2}$

### Hyper Geometric Distribution

The Hypergeometric (class name HyperGeometric) distribution represents the probability of choosing  $y$  number of events of the same kind from a subset of  $r$  like items within a population of all  $N$  possible items (of different kinds) for the sample of size  $n$  containing the mixed items. The constraints are such that  $r \leq n \leq N$  and  $y \leq r \leq n$ . The parameters are  $y, r, n, N$ . The probability distribution is defined as follows.

$$P(y; r, n, N) = \frac{\binom{r}{y} \binom{N-r}{n-y}}{\binom{N}{n}}$$

The simple properties of the distribution are:

Mean:  $\mu = \frac{nr}{N}$

Variance:  $\sigma^2 = n \left( \frac{r}{N} \right) \left( \frac{N-r}{N} \right) \left( \frac{N-n}{N-1} \right)$

### Negative Binomial Distribution

The Negative Binomial Distribution (class name NegativeBinomial) provides the probability of the  $n$ th success or potentially  $n$ th failure of a bernoulli trial. The parameters are  $r$  representing the  $(r-1)$  initial trials that where the successful and  $y$  the total number of trials before the next success  $r$  occurs. The distribution is calculated as follows:

$$P(y; r) = \binom{y-1}{r-1} p^r q^{y-r}$$

where  $y = r, r+1, \dots$

The simple properties of the distribution are:

Mean:  $\mu = \frac{r}{p}$

Variance:  $\sigma^2 = \frac{r(1-p)}{p^2}$

### The Poisson Distribution

The Poisson Distribution (class name Poisson) provides the probability of an event occurring a certain number of times within an interval. It is commonly used to model a number of events occurring in a certain period of time. We can use the two parameters  $\lambda$  and  $y$  to represent the number of events, and period of time respectively. The distribution is defined as:

$$P(y; \lambda) = \frac{\lambda^y e^{-\lambda}}{y!}$$

where  $y \geq 0$  and  $\lambda > 0$

The simple properties of the distribution are:

Mean:  $\mu = \lambda$

Variance:  $\sigma^2 = \lambda$

### 3.1.3 Functions

#### Package au.id.cxd.math.function

The package au.id.cxd.math.function contains a set of supporting functions.

### The Gamma Function

The gamma function (class name `GammaFn`) supports the gamma distribution and is implemented very simply as

$$\Gamma(\alpha) = (\alpha - 1)!$$

### The Beta Function

The beta function (class name `BetaFn`) supports the beta distribution and is built upon the gamma function in the following manner:

$$\beta(\alpha, \beta) = \frac{(\alpha - 1)!(\beta - 1)!}{(\alpha + \beta - 1)!}$$

#### 3.1.4 Continuous Distributions

##### Package `au.id.cxd.math.probability.continuous`

The package `au.id.cxd.math.probability.continuous` contains a set of classes that can be used to model continuous distributions and their simple properties.

##### Uniform Distribution

The uniform distribution (class name `Uniform`) provides a uniform output for  $y$  between a *min* and *max* and is implemented as follows.

$$f(y; min, max) = \begin{cases} \frac{1}{max-min} & \text{if } min \leq y \leq max \\ 0 & \text{otherwise} \end{cases}$$

The distribution has the following simple properties:

Mean:  $\mu = \frac{max+min}{2}$

Variance:  $\sigma^2 = \frac{(max-min)^2}{12}$

##### Gamma Distribution

The Gamma distribution (class name `Gamma`) is suitable for modelling distributions that are left skewed and is useful for things such as time measurements (similar to the Poisson distribution for discrete events). The distribution is defined as:

$$f(y; \alpha, \beta) = \begin{cases} \frac{y^{\alpha-1} e^{-y/\beta}}{\beta^\alpha \Gamma(\alpha)} & \text{for } 0 \leq y \leq \infty \\ 0 & \text{elsewhere} \end{cases}$$

The simple properties of the distribution are:

Mean:  $\mu = \alpha\beta$

Variance:  $\sigma^2 = \alpha\beta^2$

The Gamma distribution also has an important relationship with the chi-square distribution where  $\nu$  is the degrees of freedom, if the gamma distribution has the parameters  $\alpha = \nu/2$  and  $\beta = 2$  then it is also a  $\chi^2$  distribution with  $\nu$  degrees of freedom.

**Exponential Distribution**

The Exponential distribution (class name Exponential) is an instance of the Gamma distribution where the  $\alpha$  parameter equals 1, leaving the parameter  $\beta$  to be provided. The distribution can be used to model processes with "lifetimes" or decay. The distribution is defined as:

$$f(y;\beta) = \begin{cases} \frac{1}{\beta} e^{-y/\beta}, & 0 \leq y < \infty \\ 0 & \text{elsewhere} \end{cases}$$

The exponential distribution has the following simple properties:

Mean:  $\mu = \beta$

Variance:  $\sigma^2 = \beta^2$

**Beta Distribution****Normal Distribution****Chi-square Distribution****F Distribution****3.1.5 Inference****3.1.6 Regression**