Name: Marcus Bell

Q1. Scarlet Bakery bakes bagels and muffins. A dozen bagels uses 2 kg of flour and 1 hour of oven time; a dozen muffins uses 1 kg of flour and 1 hour of oven time. Each day 40 kg of flour and 30 oven hours are available, and at least 5 dozen muffins must be baked. Profit is $9 per dozen bagels and $7 per dozen muffins. Formulate an LP that maximizes daily profit.

(a) Define the decision variables. [8 pts]

x1 = bagels, x2 = muffins

4/8

Give units and a period: dozens per day.

(b) Write the objective function. [8 pts]

max z = 9x1 + 7x2

8/8

(c) Write all constraints. [14 pts]

2x1 + x2 <= 40; x1 + x2 <= 30; x2 >= 5

10.5/14

Forgot the non-negativity constraints x1, x2 >= 0.

Name: Marcus Bell

Q2. Convert to standard form: max z = x1 + 4x2 subject to -x1 + x2 <= 3, 3x1 + x2 >= 6, x1 >= 0, x2 free.

(a) Write the constraints as equalities with slack/surplus variables. [10 pts]

-x1 + x2 + s1 = 3; 3x1 + x2 - s2 = 6; s1, s2 >= 0

10/10

(b) Handle the free variable x2 and rewrite the objective. [10 pts]

x2 = x2+ - x2-, x2+, x2- >= 0; max z = x1 + 4x2+ - 4x2-

10/10

Name: Marcus Bell

Q3. Simplex: max z = 2x1 + 3x2 subject to x1 + 2x2 <= 8, 3x1 + 2x2 <= 12, x1, x2 >= 0.

(a) Choose the entering and leaving variables for the first pivot. [10 pts]

x2 enters (-3); ratios 8/2 = 4, 12/2 = 6, so s1 leaves.

10/10

(b) Carry out pivots to optimality. [10 pts]

Pivot 2: x1 enters, x2 leaves (ratio 8).

5/10

(c) State the optimal solution and value; explain how you know it is optimal. [10 pts]

x1 = 0, x2 = 4, z = 12 - optimal after one pivot.

4/10

Name: Marcus Bell

Q4. Graphical method: max z = 5x1 + 5x2 subject to x1 + x2 <= 8, x1 <= 5, x2 <= 6, x1, x2 >= 0.

(a) List the corner points of the feasible region. [10 pts]

7/10

(0, 0), (5, 0), (5, 4), (2, 6), (0, 6)

x1 = 5 meets x1 + x2 = 8 at x2 = 3, not 4.

(b) Find all optimal solutions and explain. [10 pts]

4/10

Unique optimum (5, 3), z = 40.

The objective 5x1 + 5x2 is parallel to the binding constraint x1 + x2 = 8, so the whole edge from (2, 6) to (5, 3) is optimal.