Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Acetylcysteine	Acetadote	Acetaminophen overdose	300 mg/kg total dose administered as a 21-hour IV infusion regimen consisting of 3 doses. First load over 1 hour, second over 4 hours, third over 16 hours		Hepatoprotective by restoring hepatic glutathione and enhancing the nontoxic sulfate conjugation of acetaminophen	
Acyclovir	Zovirax	Antiviral	200 mg orally every 4 hours for 5 doses/day for 10 days		Inhibits DNA synthesis and viral replication by competing with deoxyguanosine triphosphate for viral DNA polymerase and being incorporated into viral DNA	
Albuterol sulfate	Proventil Ventolin Proair	Asthma COPD Bronchospasm	1-2 inhalations (90mcg HFA) every 4-6 hours		Short acting beta 2 adrenergic agonist that relaxes bronchial smooth muscle	
Alendronate	Fosamax	Osteoporosis	10 mg orally once daily OR 70 mg once weekly		Adheres to sites of bone reabsorption, inhibiting osteoclast activity. Normal bone is formed on top of alendronate, which is then incorporated into the bone matrix	
Allopurinol	Zyloprim	Gout	200-300 mg by mouth once a day		Competitive inhibition of xanthine oxidase which prevents the conversion of hypoxanthine to xanthine to uric acid	
Alprazolam	Xanax Xanax XR	Anxiety	0.25-0.5 mg by mouth three times a day prn	CIV	Binds to the GABA(A) receptor in the central nervous system and facilitates the action of GABA, an inhibitory neurotransmitter	Concomitant use with opioids increases the risk of profound sedation, respiratory depression, coma, and death. Risk of abuse, misuse and addiction which can lead to overdose or death. Continued use may lead to dependence, avoid abrupt discontinuation to prevent withdrawal reactions.
Alteplase	Activase	Acute ischemic stroke	Load with 0.09 mg/kg IV over 1 minute, followed by 0.81 mg/kg continuous infusion over 60 minutes (MAX 90 mg)		Initiates local fibrinolysis by binding to fibrin in the thrombus and converts entrapped plasminogen to plasmin	

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Amiodarone	Cordarone, Pacerone Nexterone	Arrhythmia	200-400 mg by mouth once a day		Class III antiarrhythmic that lengthens cardiac action potential by prolonging intranodal conduction and refractoriness of the AV node	Use only intended for life-threatening arrhythmias due to risk of toxicity. Pulmonary and hepatic toxicity may occur during therapy. May exacerbate arrhythmias and therefore should be initiated where continuous ECG and cardiac resuscitation are available.
Amitriptyline	Elavil	Depression	10-100 mg by mouth in divided doses or at bedtime		Prevents reuptake of norepinephrine and serotonin at the presynaptic nerve terminal which increases activity at post synaptic terminals	Can increase suicidal thoughts and actions in children, adolescents, and young adults
Amlodipine	Norvasc	Hypertension	5 mg by mouth once daily		Dihydropyridine calcium channel blocker. Inhibits calcium influx in vascular smooth muscle (more than cardiac muscle)	
Amoxicillin	Amoxil Moxatag Trimox	Bacterial infection	250-500 mg by mouth every 8 hours OR 500-875 mg every 12 hours		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme	
Amoxicillin/ Clavulanate	Augmentin, Augmentin XR	Bacterial infection	500/125 mg by mouth every 8 hours OR 875/125 mg every 12 hours		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme and inhibition of beta-lactamase	
Amphetamine/ Dextroamphetamine salts	Adderall Adderall XR	ADHD	5-20 mg (XR) by mouth once a day	CII	Releases norepinephrine and dopamine from storage sites causing stimulation on adrenergic receptors	High potential for abuse and dependence. Misuse may cause sudden death or serious cardiovascular adverse events
Ampicillin/ Sulbactam	Unasyn	Bacterial infection	1.5 g IV infusion every 6 hours (dose is combination of ampicillin and sulbactam)		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme and inhibition of beta-lactamase	
Anastrozole	Arimidex	Breast cancer	1 mg by mouth once a day		Selective non-steroidal aromatase inhibitor which prevents the conversion of androstenedione to estrone and testosterone to estradiol. This decreases tumor mass or delays tumor progression	
Apixaban	Eliquis	Anticoagulation	5 mg by mouth two times a day		Selectively inhibits Factor Xa which reduces thrombin formation	Premature discontinuation increases risk of thrombosis Risk of epidural or spinal hematomas when used in patients undergoing epidural or spinal anesthesia resulting in permanent paralysis

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Aripiprazole	Abilify Abilify Maintena	Schizophrenia	10 mg by mouth once a day		Action MAY be combination of partial agonist activity at dopamine type 2 receptor and serotonin (5-HT1A and antagonism effects at 5-HT2 receptors	Increased mortality in elderly patients with dementia-related psychosis. Increased risk of suicidal thoughts and actions in children, adolescents, and young adults.
Atenolol	Tenormin	Hypertension	50 mg by mouth once a day		Cardioselective beta 1 adrenergic blocker	Abrupt discontinuation in patients with coronary artery disease may cause severe exacerbation of angina, myocardial infarction, or ventricular arrhythmias
Atomoxetine	Strattera	ADHD	80 mg by mouth once a day OR 40 mg by mouth two times a day		MAY work by selective inhibition of the pre-synaptic norepinephrine transporter	Can increase suicidal thoughts and actions in children, adolescents, and young adults
Atorvastatin	Lipitor	Dyslipidemia	10 mg by mouth once a day		Selective competitive inhibitor of HMG-CoA reductase	
Azelastine	Astelin Astepro	Rhinitis	2 sprays per nostril twice a day		Histamine H1 receptor antagonist in peripheral tissues	
Azithromycin	Zithromax Zmax	Bacterial infection	500 mg by mouth on day 1, 250 mg by mouth on days 2-5		Suppresses protein synthesis by binding to the 50 S ribosomal subunit	
Baclofen	Lioresal	Musculoskeletal pain/spasm	10 mg by mouth three times a day as needed		MAY work through hyperpolarization of afferent terminals resulting in inhibition of monosynaptic and polysynaptic reflexes	Abrupt discontinuation of intrathecal baclofen can result in rhabdomyolysis, multiple organ-system failure, and death
Benzonatate	Tessalon	Antitussive	100 mg three time daily as needed		Topical anesthetic activity in respiratory stretch receptors	
Bimatoprost (ophthalmic)	Lumigan Latisse	Glaucoma	1 drop into affected eye(s) once daily in the evening		Synthetic prostaglandin analog which reduces intraocular pressure by increasing the outflow of the aqueous humor	
Brimonidine Tartrate (ophthalmic)	Alphagan-P	Glaucoma	1 drop into affected eye(s) three times daily 8 hours apart		Alpha 2 adrenergic agonist which reduced production of aqueous humor and increased outflow	
Budesonide	Pulmicort	Asthma	360 mcg inhaled two times a day		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Budesonide/ Formoterol	Symbicort	Asthma COPD	2 inhalations of 160 mcg/4.5 mcg MDI two times a day		Stimulates synthesis of anti-inflammatory proteins and enzymes and long-acting beta 2 adrenergic receptor agonist	
Bupropion	Wellbutrin Wellbutrin SR Wellbutrin XL Zyban*	Depression Smoking cessation*	150 mg ER once a day		Blocks neuronal uptake of norepinephrine and dopamine in the central nervous system	Can increase suicidal thoughts and actions in children, adolescents, and young adults

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Buspirone	BuSpar	Anxiety	5 mg by mouth three times a day as needed		MAY alter brain chemistry by increasing metabolism of norepinephrine, moderate presynaptic dopamine agonist, serotonin agonist	
Butalbital, Acetaminophen and Caffeine	Fioricet	Tension headache	Capsule – 50 mg/300 mg/40 mg Tablet – 50 mg/325 mg/40 mg Dose - 1-2 capsule/tablet every four hours as needed not to exceed 6 capsules/tablets per day		Butalbital – CNS depressant by activating inhibitory GABA receptors and inhibition of AMPA receptors Acetaminophen – Inhibits CNS prostaglandin synthesis and block pain impulse generation Caffeine – CNS stimulant by inhibition of phosphodiesterase increasing levels of c-AMP	Acetaminophen has been associated with liver failure typically at dosing exceeding 4000 mg/day
Carbamazepine	Tegretol Tegretol XR Carbatrol ER Equetro ER	Seizure	400 mg by mouth three times a day		Na+ channel antagonist; delays recovery of inactive Na+ channels	Serious and fatal dermatologic skin reactions can occur with a strong association in patients with HLA-B*1502 variance. Can cause aplastic anemia and agranulocytosis
Carbidopa/Levodopa	Sinemet Sinemet CR Rytary	Parkinson's disease	25/100 mg by mouth three times a day		Levodopa is a precursor to dopamine Carbidopa is a DOPA-decarboxylase inhibitor and prevents the breakdown of levodopa before it crosses the blood brain barrier Combined these increase the dopamine available to the central nervous system	
Cariprazine	Vraylar	Antipsychotic	1.5 mg by mouth once daily		Partial agonist at D2 and 5-HT1A receptors, high affinity for dopamine, binds to histamine receptors	Increased mortality in elderly patients with dementia related psychosis Antidepressants increase the risk of suicidal thoughts and actions
Carisoprodol	Soma	Pain relief of musculoskeletal conditions	250-350 mg by mouth three days per day and at bedtime	CIV	Central depressant resulting in musculoskeletal relaxation	
Carvedilol	Coreg, Coreg CR	Hypertension	6.25 mg by mouth two times a day		Non-selective beta- and alpha-1 adrenergic blocking activity	
Cefazolin	Ancef	Bacterial infection	1-2 g IV every 8 hours		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme	
Cefdinir	Omnicef	Bacterial infections	600 mg by mouth daily in 1or 2 divided doses		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme	

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Cefepime	Maxipime	Bacterial infection	1-2 g IV every 8 hours		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme	
Ceftriaxone	Rocephin	Bacterial infection	1-2 g IV/IM every 24 hours		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme	
Celecoxib	Celebrex	Pain Osteoarthritis	200 mg by mouth daily		Inhibits prostaglandin synthesis through selective inhibition of cyclooxygenase 2	May cause increased risk of serious and fatal cardiovascular thrombotic events. Have been associated with gastrointestinal bleeding, ulceration, and perforation.
Cephalexin	Keflex	Bacterial infection	500 mg by mouth every 12 hours		Inhibits bacterial cell wall synthesis via acylation transpeptidase enzyme	
Chlorhexidine	Peridex Periogard	Gingivitis	Swish oral rinse for 30 seconds two times a day		Disrupts the plasma membrane of the bacterial cell	
Chlorthalidone	Hygroton	Edema	12.5 mg by mouth once a day		Inhibits reabsorption of sodium and chloride at the cortical diluting segment of the ascending loop of Henle	
Ciprofloxacin	Cipro Cipro XR	Bacterial infection	500 mg by mouth two times a day		Prevents synthesis of bacterial DNA by interfering with the enzyme DNA gyrase	May cause serious adverse reactions including tendonitis, tendon rupture, peripheral neuropathy, central nervous system effects, and exacerbation of myasthenia gravis
Cisatracurium	Nimbex	Paralytic	Initial IV bolus: 0.1-0.2 mg/kg, Continuous infusion: 1-3 mcg/kg/min		Competes with acetylcholine for cholinergic receptors inhibiting neuromuscular transmission	
Citalopram	Celexa	Depression	20 mg by mouth once a day		Blocks neuronal uptake of serotonin in the central nervous system	Antidepressants increase the risk of suicidal thoughts and actions in children, adolescents, and young adults
Clindamycin	Cleocin	Bacterial infection	300 mg by mouth every 6 hours		Suppresses protein synthesis by binding to the 50 S ribosomal subunit	Can cause severe and fatal colitis due to C. difficile
Clonazepam	Klonopin	Panic disorder	0.25 mg by mouth two times a day	CIV	Binds to the GABA(A) receptor in the central nervous system and facilitates the action of GABA, an inhibitory neurotransmitter	Concomitant use with opioids increases the risk of respiratory depression, coma, and death. Risk of abuse, misuse and addiction which can lead to overdose or death. Continued use may lead to dependence, avoid abrupt discontinuation to prevent withdrawal reactions
Clonidine	Catapres Catapres-TTS Nexiclon XR	Hypertension	0.1 mg by mouth two times a day		Alpha 2 adrenergic receptor agonist resulting in reduced sympathetic outflow from the CNS	
Clonidine ER	Kapvay	ADHD	0.1 mg by mouth two times a day		Alpha 2 adrenergic receptor agonist resulting in reduced sympathetic outflow from the CNS	

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Clopidogrel	Plavix	Reduction of atherosclerotic events in patients with atherosclerosis	75 mg by mouth once a day		Inhibits platelet aggregation by irreversibly modifying platelet receptors	Do not use in patients identified with two loss of function alleles of the CYP2C219 gene
Clozapine	Clozaril Versacloz	Schizophrenia	Initial 12.5 mg by mouth 1-2 times a day (MAX 900 mg/day)		Antagonism of dopamine type 2 and 5-HT2A receptors. Antagonist at adrenergic, cholinergic, histaminergic, and serotonergic receptors	Severe neutropenia Orthostatic hypotension, bradycardia and syncope Seizures Myocarditis and cardiomyopathy and mitral valve incompetence Increased mortality in elderly patients with dementia-related psychosis
Codeine phosphate with Guaifenesin	Mytussin AC Robitussin AC	Antitussive	5-10 mL by mouth every 4-6 hours not to exceed 60 mL per 24 hours	CV	Guaifenesin thins bronchial mucus Codeine suppresses the cough reflex in the medulla oblangata	
Colestipol Hydrochloride	Colestid	Dyslipidemia	Tablets – 2 grams by mouth 1-2 times per day Granules – 5 grams by mouth 1-2 times per day		Binds bile acids which are then excreted in the feces	
Cyanocobalamin Vitamin B-12	Nascobal	Vitamin B12 deficiency	1000 mcg intramuscular or subcutaneously once per month		Vitamin that is required for the synthesis of DNA and maintenance of normal erythropoiesis	
Cyclobenzaprine	Amrix Flexeril Fexmid	Musculoskeletal pain/spasm	10 mg by mouth 3 times a day		Acts in the brain stem to reduce tonic somatic motor activity	
Cyclosporin Emulsion (ophthalmic)	Restasis	Keratoconjunctivitis sicca (Dry eye syndrome)	Instill 1 drop in each eye twice daily at 12 hour intervals		Immunomodulator that may block the release of proinflammatory cytokines	
Dapagliflozin	Farxiga	Diabetes	5-10 mg by mouth in the morning		Inhibits sodium-glucose co-transporter 2 (SGLT2) which increases urinary glucose excretion	
Desvenlafaxine	Pristiq	Depression	50 mg by mouth once a day		Blocks neuronal uptake of serotonin and norepinephrine in the central nervous system	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Dexmedetomidine	Precedex	Sedation	0.2-1.5 mcg/kg/hour		Alpha 2 adrenergic receptor agonist	

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Diazepam	Valium	Anxiety	2 mg by mouth every 4 hours as needed	CIV	Binds to the GABA(A) receptor in the central nervous system and facilitates the action of GABA, an inhibitory neurotransmitter	Concomitant use with opioids increases the risk for respiratory depression, coma, and death. Risk of abuse, misuse and addiction which can lead to overdose or death. Continued use may lead to dependence, avoid abrupt discontinuation to prevent withdrawal reactions
Diclofenac	Voltaren Voltaren XR	Pain	50 mg by mouth three times a day		Inhibits prostaglandin synthesis through non-specific inhibition of cyclooxygenase	May cause increased risk of serious and fatal cardiovascular thrombotic events. Have been associated with gastrointestinal bleeding, ulceration, and perforation.
Diclofenac (topical)	Voltaren Gel	Osteoarthritis	Apply 2 g to affected area of upper extremity or 4 g to lower extremity up to four times a day, maximum total body dose 32 g/day		Inhibits prostaglandin synthesis through non-specific inhibition of cyclooxygenase	
Dicyclomine	Bentyl	Irritable bowel syndrome	20 mg by mouth up to four times a day 30 minutes before meals		Inhibition of acetylcholine at postganglionic parasympathetic receptor sites in multiple tissues	
Digoxin	Lanoxin Digitek	Heart failure Atrial fibrillation	0.125-0.25 mg by mouth once a day (dose individualized based on serum levels)		Inhibits membrane bound Na+/K+ ATPase which increases intracellular Na+ concentration and a slightly decreased intracellular K+ concentration leading to increased extracellular Ca++ influx increased contraction force/velocity of heart	
Diltiazem	Cardizem, Cardizem CD Cartia Dilacor Dilt-XR Tiazac	Hypertension	240 mg by mouth divided 1-2 times daily		Calcium channel blocker which inhibits calcium influx in vascular smooth muscle (more than in cardiac muscle)	
Dipyridamole and Aspirin	Aggrenox	Ischemic stroke prevention	200 mg/25 mg capsule by mouth twice daily		Dipyridamole inhibits platelet aggregation Aspirin inhibits prostaglandin and thromboxane A2 production	
Divalproex sodium	Depakote Depakote ER	Seizures Bipolar disorder	250 mg by mouth three times a day		MAY be related to voltage gated sodium channel blockade and increased brain levels of GABA	Serious and fatal hepatotoxicity Life threatening pancreatitis Do not use in pregnancy due to neural tube defects Increased risk of liver failure and death in patients with mitochondrial disease

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Dobutamine	Dobutrex	Heart failure	Initial dose: 2-5 mcg/kg/minute IV titrated to clinical effect		Stimulates beta-2 and alpha-1 receptors to increase myocardial contractility and heart rate	
Dofetilide	Tikosyn	Atrial fibrillation Atrial flutter	500 mg by mouth twice daily		Class III antiarrhythmic Blockade of cardiac ion channel carrying the rapid component of the delayed rectifier potassium current	To minimize risk of inducing an arrythmia, hospitalize patients when initiating therapy for continuous EKG monitoring and monitoring renal function
Donepezil	Aricept	Alzheimer's disease	5-10 mg by mouth at bedtime		Reversible acetylcholinesterase inhibitor working centrally to increase acetylcholine available at cholinergic synapses	
Dopamine	Intropin	Shock	1-20 mcg/kg/minute IV titrated to clinical effect		Agonist at dopaminergic, alpha-, and beta-adrenergic receptors; precursor for norepinephrine synthesis	Extravasation can result in peripheral ischemia. Treat with phentolamine
Doxycycline	Vibramycin Doryx	Bacterial infection	100 mg by mouth every 12 hours		Prevents bacterial protein synthesis by competitively binding to the 30 S ribosomal subunit	
Duloxetine	Cymbalta	Depression	40-60 mg by mouth one time a day		Blocks neuronal uptake of serotonin and norepinephrine in the central nervous system	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Empagliflozin	Jardiance	Diabetes	10-25 mg by mouth daily in the morning		Inhibits sodium-glucose co-transporter 2 (SGLT2) which increases urinary glucose excretion	
Enoxaparin	Lovenox	Anticoagulation	1 mg/kg subcutaneous every 12 hours OR 1.5 mg/kg once daily		Inhibits Factor Xa (and to a lesser extent IIa) which reduces thrombin formation	Risk of epidural or spinal hematomas when used in patients undergoing epidural or spinal anesthesia resulting in permanent paralysis
Ephedrine	Akovaz Emerphed	Hypotension caused by anesthesia	IV bolus: 5-10 mg once, may repeat dose (MAX total 50 mg)		Releases norepinephrine from tissue stores resulting in alpha- and beta-adrenergic stimulation	
Epinephrine	EpiPen Auvi-Q Adrenalin	Anaphylaxis	Inject 0.3 mg IM, may repeat dose if patient not responding		Sympathomimetic catecholamine that acts on both alpha- and beta-adrenergic receptors causing vasoconstriction (alpha), cardiac stimulation (beta-1), and relaxation of bronchial smooth muscles (beta-2)	
Epoprostenol	Veletri Flolan	Pulmonary hypertension	Initial 2 ng/kg/min IV, titrate up in increments of 1-2 ng/kg/min every 15 minutes. Usual dose is 25-40 ng/kg/min.		Directly stimulates vasodilation of all vascular beds. It activates intracellular adenylate cyclase which inhibits platelet aggregation. Also known as prostacyclin (PGI_2).	

Generic **P1 P2 P3**		**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Escitalopram		Lexapro	Depression	10 mg by mouth once a day		Blocks neuronal uptake of serotonin in the central nervous system	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Esomeprazole		Nexium	GERD	20 mg by mouth once a day		Inhibits H+/K+ ATPase enzyme system at the secretory surface of the gastric parietal cell	
Estradiol	Oral	Estrace	Menopausal vasomotor symptoms	0.5 mg by mouth once a day		Replaces estrogen post menopause. Promotes growth and development of vagina, uterus, and fallopian tubes, and enlargement of the breast	Increased risk of endometrial cancer with unopposed systemic estrogen in women with a uterus Please note – some vaginal products are absorbed systemically
	Topical	Vivelle-Dot Alora Climara		Product dependent: Patches are typically applied 1-2 times per week per product labeling			
	Vaginal	Femring Estring Vagifem Divigel Many other products		Product dependent			
Eszopiclone		Lunesta	Insomnia	2 mg by mouth at bedtime	CIV	MOA unknown, thought to bind or interact with GABA receptor complexes	Sleepwalking, sleep driving, and engaging in other activities while not fully awake
Evolocumab		Repatha	Dyslipidemia	140 mg subcutaneously every 2 weeks or 420 mg subcutaneously once per month		Monoclonal antibody that binds PCSK9 to increase the number of LDL receptors available to clear LDL from the blood	
Ezetimibe		Zetia	Dyslipidemia	10 mg by mouth once a day		Inhibits intestinal absorption of cholesterol and related phytosterols	
Famotidine		Pepcid	GERD	20 mg by mouth two times a day		Competitive inhibition of histamine at H2 receptors of the gastric parietal cells, which inhibits gastric acid secretion	
Fenofibrate (nanocrystallized)		Tricor	Dyslipidemia	145 mg by mouth once a day		Activates peroxisome proliferator activated receptor-a resulting in increased lipolysis and elimination of triglycerides	

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Fentanyl Transdermal Patch	Duragesic	Chronic pain	1 patch applied topically every 72 hours, remove and replace at the end of 72 hour period	CII	Opioid receptor agonist	Serious, life-threatening respiratory depression may occur. Can be addictive, abused, and misused which can lead to overdose and death. Accidental exposure can be fatal, especially in children. Concomitant use with CYP3A4 inhibitors, benzodiazepines, alcohol, or other CNS depressants can result in respiratory depression and coma. Use in pregnancy can cause neonatal withdrawal. Required REMS to ensure benefits outweigh risks. Exposure of application site to heat sources may increase absorption and result in overdose.
Finasteride	Proscar	BPH	5 mg by mouth once a day		Competitively inhibits the enzyme 5-alpha-reducatese, resulting in decreased metabolism of testosterone to DHT	
Fluconazole	Diflucan	Candida infection	150 mg by mouth as a single dose or may require daily treatment		Inhibition of fungal ergosterol synthesis via inhibition of fungal C14 demethylase	
Fluoxetine	Prozac Sarafem*	Depression Premenstrual dysphoria disorder*	20 mg by mouth once a day		Blocks neuronal uptake of serotonin in the central nervous system	Antidepressants increase risk of suicidal thoughts and actions in children, adolescents, and young adults
Fluticasone furoate/Vilanterol	Breo Ellipta	COPD	1 inhalation of 100/25 mcg daily		Stimulates synthesis of anti-inflammatory proteins and enzymes and long-acting beta 2 adrenergic receptor agonist	
Fluticasone furoate/ Umeclidinium bromide/ Vilanterol	Trelegy Ellipta	COPD	1 inhalation of 100/62.5 /25 mcg daily		Stimulates synthesis of anti-inflammatory proteins and enzymes, competitively inhibits action of M3 receptors in bronchial smooth muscle, and long-acting beta 2 adrenergic receptor agonist	
Fluticasone propionate (nasal)	Flonase	Allergic rhinitis	2 sprays in each nostril once daily		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Fluticasone/ Salmeterol	Advair	Asthma	1 diskus inhalation or 2 HFA inhalations two times a day		Stimulates synthesis of anti-inflammatory proteins and enzymes and long-acting beta 2 adrenergic receptor agonist	

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Folic acid	Generic/No brand name	Folic acid deficiency	Up to 1 mg by mouth daily		Vitamin that is required for the synthesis of thymidylate, purine nucleotides, and maintenance of normal erythropoiesis	
Fremanezumab-vfrm	Ajovy	Migraine	225 mg subcutaneously monthly or 675 mg subcutaneously every 3 months		Humanized monoclonal antibody that binds to calcitonin gene related peptide (CGRP) ligand and blocks binding to the receptor	
Furosemide	Lasix	Edema	20 mg by mouth two times a day		Loop diuretic which inhibits reabsorption of sodium and chloride in the proximal and distal tubules and in the loop of Henle	
Gabapentin	Neurontin Gralise	Neuropathic pain	300 mg by mouth three times a day	CV in ND, but not controlled federally – verify in your state of practice	Ca2+ channel antagonist; binds with alpha 2 subunit of Ca2+ channel	
Glimepiride	Amaryl	Diabetes	1-4 mg by mouth once a day		Stimulates insulin synthesis and release from beta cells in the pancreatic islets	
Glipizide	Glucotrol, Glucotrol XL	Diabetes	5-15 mg by mouth once a day OR 10-20 mg by mouth two times a day		Stimulates insulin synthesis and release from beta cells in the pancreatic islets	
Guanfacine ER	Intuniv	ADHD	1 mg by mouth once a day		Alpha 2 adrenergic receptor agonist	
Heparin	Generic/No brand name	Anticoagulation	5,000 units subcutaneous every 8 hours		Inactivates thrombin and activated Factor Xa through an antithrombin dependent mechanism	
Hydrochlorothiazide	Microzide	Edema	25 mg by mouth once a day		Inhibits reabsorption of sodium and chloride at the distal renal tubule	
Hydrocodone/ Acetaminophen	Lortab Vicodin Norco	Pain	1-2 tablets (5/300 mg per tablet) by mouth every 4-6 hours as needed	CII	Opioid receptor agonist/inhibition of prostaglandin synthesis in the central nervous system	Opioids can be addictive, abused, and misused which can lead to overdose and death. REMS. Serious life-threatening respiratory depression may occur. Accidental exposure can be fatal, especially in children. Use in pregnancy can cause neonatal withdrawal. 3A4 interactions. Acetaminophen can cause hepatotoxicity. Concomitant use with benzodiazepines, alcohol, or other CNS depressants.

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Hydrocortisone sodium succinate	Solu-Cortef	Inflammatory disorders	100 mg IV, may require additional dosing		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Hydroxychloroquine	Plaquenil	Rheumatoid arthritis	200-400 mg by mouth once a day		MOA not well understood	
Hydroxyzine hydrochloride	Atarax	Pruritis	25 mg by mouth up to 4 times a day		Histamine 1 receptor agonist in peripheral tissue	
Ibuprofen	Advil Motrin	Pain	400-800 mg by mouth 3-4 times a day		Inhibits prostaglandin synthesis through non-specific COX-1 and COX-2 inhibition	May increase risk of serious cardiovascular thrombotic events, myocardial infarction, stroke, gastrointestinal bleeding, ulcerations, and perforation
Icosapent ethyl	Vascepa	Dyslipidemia	2 g by mouth two times a day with food		Eicosapentaenoic acid (EPA), the active metabolite, reduces hepatic VLDL-triglyceride synthesis and/or secretion and increases triglyceride clearance from circulating VLDL particles	
Insulin aspart	Novolog	Diabetes	Short acting Individualized dosing administered within 5-10 minutes of meals		Activates specific enzymes, promotes the transport of monosaccharides across cell membranes and facilitates the transformation of intracellular amino acids into proteins	
Insulin degludec	Tresiba	Diabetes	Long acting Individualized dosing administered once daily		Activates specific enzymes, promotes the transport of monosaccharides across cell membranes and facilitates the transformation of intracellular amino acids into proteins	
Insulin glargine	Lantus Toujeo Basaglar Semglee	Diabetes	Long acting Individualized dosing administered once daily at bedtime		Activates specific enzymes, promotes the transport of monosaccharides across cell membranes and facilitates the transformation of intracellular amino acids into proteins	
Irbesartan	Avapro	Hypertension	150-300 mg by mouth once a day		Angiotensin II antagonist. Blocks the binding of angiotensin II to the ATI receptor	Fetal toxicity: discontinue as soon as pregnancy is detected
Iron sucrose	Venofer	Iron deficiency	100 mg IV per dose; repeat doses may be given until requirements are met		Used by body for normal erythropoiesis	
Isosorbide mononitrate	Imdur	Angina pectoris	30-120 mg by mouth in the morning		Nitric oxide donor; vasodilation	

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Ketorolac	Toradol	Pain	30 mg IV every 6 hours for a maximum of 5 days		Inhibits prostaglandin synthesis through non-specific COX-1 and COX-2 inhibition	Only indicated short-term, up to 5 days. Serious cardiovascular events thrombotic events, serious GI bleeding, ulceration and perforation, renal risk, risk of bleeding Contraindicated for intrathecal or epidural administration. Severe hypersensitivity reactions may occur. Contraindicated in patients with advanced renal impairment. Contraindicated in labor and delivery. Contraindicated in patients currently receiving aspirin or NSAIDs. Adjust dose for patients over 65 years and patients weighing less than 50kg
Lactulose	Kristalose Constulose Enulose	Constipation	10-20 g by mouth daily		Increases osmotic pressure in colon and increases pH resulting in increased water content of stool	
Lamotrigine	Lamictal, Lamictal XR Lamictal ODT Lamictal CD	Bipolar disorder Seizures	25 mg by mouth once a day		Na+ antagonist; AMPA receptor antagonist	Serious life-threatening rashes have been reported
Lansoprazole	Prevacid	GERD	15 mg by mouth once a day		Inhibits H+/K+ ATPase enzyme system at the secretory surface of the gastric parietal cell	
Latanoprost	Xalatan	Glaucoma	One drop into affected eye once a day in the evening		Prostaglandin F2-alpha analog believed to reduce intraocular pressure by increasing the outflow of the aqueous humor	
Levetiracetam	Keppra	Seizure	500 mg by mouth two times a day		Decreases glutamate release by inhibiting synaptic vesicle glycoprotein, SV2	
Levofloxacin	Levaquin	Bacterial infection	500 mg by mouth once a day x 10-14 days OR 750 mg by mouth once a day x 5 days		Inhibition of bacterial DNA gyrase and topoisomerase IV	Increased risk of tendonitis and tendon rupture. Risk is enhanced with patients over 60 years of age, in patients taking corticosteroid drugs and patients with kidney, heart or lung transplants. May exacerbate myasthenia gravis
Levothyroxine	Synthroid Levoxyl Levothyroid	Hypothyroidism	25-300 mcg by mouth once a day		Synthetic form of thyroxine involved in normal metabolism, growth, and development	Do not use for treatment of obesity or for weight loss in euthyroid patients. Large doses may cause serious and fatal toxicity

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Linaclotide	Linzess	Irritable bowel syndrome with constipation	290 mcg by mouth once daily on an empty stomach 30 minutes before the first meal of the day		Linaclotide and its active metabolite bind and agonize guanylate cyclase-C on the luminal surface of intestinal epithelium resulting in increased GI transit	Risk of serious dehydration in pediatric patients under 2 years of age - Contraindicated in patients under 2 years
Linezolid	Zyvox	Bacterial infection	600 mg by mouth every 12 hours		Inhibition of bacterial protein synthesis by binding to the 23 S site on the 50 S ribosomal subunit	
Lisdexamfetamine	Vyvanse	ADHD	30 mg by mouth once a day in the morning	CII	Releases norepinephrine and dopamine from storage sites causing stimulation on adrenergic receptors	CNS stimulants have a high potential for abuse and dependence
Lisinopril	Zestril Prinivil	Hypertension	10 mg by mouth once a day		Angiotensin-converting enzyme inhibitor	Fetal toxicity: discontinue as soon as pregnancy is detected
Lisinopril/Hydrochlorothiazide	Zestoretic Prinzide	Hypertension	10/12.5 mg by mouth once a day		Angiotensin-converting enzyme inhibitor and lowers plasma volume by decreasing sodium and water reabsorption	Fetal toxicity: discontinue as soon as pregnancy is detected
Lithium	Lithonate Lithotabs Lithobid	Bipolar disorder	300 mg by mouth three times a day		MAY increase reuptake of norepinephrine and dopamine or antagonize their release from nerve terminals	Toxicity is closely related to serum levels
Lorazepam	Ativan	Anxiety	0.5 mg 2-3 times a day	CIV	Binds to the GABA(A) receptor in the central nervous system and facilitates the action of GABA, an inhibitory neurotransmitter	Concomitant use with opioids increases the risk of respiratory depression, coma, and death. Abuse, misuse, and addiction. Dependence and withdrawal reactions.
Losartan	Cozaar	Hypertension	50 mg by mouth once a day		Angiotensin II antagonist, blocks the binding of angiotensin II to the AT1 receptor	Fetal toxicity: discontinue drug as soon as pregnancy is detected
Meloxicam	Mobic	Pain	7.5 mg by mouth once a day		Inhibits prostaglandin synthesis through non-specific inhibition of cyclooxygenase	May cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke. NSAIDs have been involved in rare cases of GI bleeding incidents
Memantine	Namenda	Alzheimer's disease	5-10 mg daily		Non-competitive N-methyl-D-aspartate (NMDA) receptor antagonist	
Meropenem	Merrem	Bacterial infection	500 mg IV every 6 hours		Inhibits bacterial cell wall synthesis via inhibition of peptidoglycan transpeptidase enzyme	
Metformin	Glucophage Glucophage XR Glumetza Riomet	Diabetes	500 mg by mouth two times a day with food OR 850 mg by mouth once a day with morning meal		Decreases intestinal absorption of glucose, suppresses hepatic gluconeogenesis, and improves insulin sensitivity	May cause serious or fatal lactic acidosis

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Methadone	Methadose Dolophine	Pain	2.5-10 mg every 8-12 hours as needed	CII	Mu-opioid receptor agonist	Addiction, abuse, and misuse. REMS compliant education encouraged. Life-threatening respiratory depression. Life-threatening QT prolongation. Accidental exposure, Neonatal withdrawal syndrome. Many interactions with cytochrome P450 inhibitors. Risks from concomitant use with benzodiazepines or other CNS depressants.
Methenamine	Hiprex	Urinary tract infection prevention	1 gram by mouth two times a day		Hydrolyzed to formaldehyde which has nonspecific bactericidal action, may also aid in maintaining urine acidity and suppressing bacterial growth	
Methocarbamol	Robaxin	Musculoskeletal pain/spasm	750 mg by mouth four times a day		MAY work in the spinal cord by inhibiting the flexor and crossed extensor reflexes	
Methotrexate	Trexall	Rheumatoid arthritis	7.5-25 mg by mouth once weekly		Anti-inflammatory AND folate antimetabolite that inhibits DNA synthesis, repair, and cellular replication	May cause bone marrow suppression, renal, GI, hepatic, pulmonary toxicities, hypersensitivity, and dermatologic reactions, Contraindicated in pregnancy
Methylphenidate	Ritalin, Methylin Metadate Concerta	ADHD	20 mg by mouth two times a day, before breakfast and lunch	CII	Releases norepinephrine and dopamine from storage sites causing stimulation on adrenergic receptors	CNS stimulants have a high potential for abuse and dependence
Methylprednisolone (oral)	Medrol	Inflammatory disorders	Individualized – normal doses range from 4-48 mg by mouth per day		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Methylprednisolone sodium succinate (parenteral)	Solu-Medrol	Inflammatory disorders	Individualized		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Metoprolol succinate	Toprol XL	Hypertension	25-100 mg by mouth once a day		Cardioselective beta 1 adrenergic blocker	
Metoprolol tartrate	Lopressor	Hypertension	50 mg by mouth twice a day		Cardioselective beta 1 adrenergic blocker	Avoid abrupt discontinuation due to risk of myocardial infarction
Metronidazole	Flagyl	Antibiotic	250-500 mg by mouth three times daily for 7 days		Converted to unstable intermediates in susceptible bacteria which then bind to bacterial DNA causing DNA strand breakage and cell death	
Micafungin	Mycamine	Fungal infection	100 mg once a day		Inhibits synthesis of 1, 3-beta-D-glucan, an essential component of fungal cell walls	
Midazolam	Versed	Sedation	Individualized	CIV	Binds to the GABA(A) receptor in the central nervous system	Associated with respiratory depression and respiratory arrest. Concomitant use

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
						with opioids may result in profound sedation, respiratory depression, coma, and death. Abuse, misuse, and addiction. Dependence and withdrawal reactions
Minocycline	Minocin Dynacin Solodyn ER	Antibiotic	200 mg by mouth initially then 100 mg every 12 hours		Prevents protein synthesis by binding to the 30 S ribosomal subunit	
Mirabegron	Myrbetriq	Overactive bladder	25 mg by mouth once a day		Beta-3 adrenergic agonist	
Mirtazapine	Remeron	Depression	15 mg by mouth once a day		Central pre-synaptic alpha 2 antagonist. Antagonist of 5-HT2 and 5-HT3 receptors. Inhibitor of histamine H1 receptors and alpha 1 and muscarinic receptors	Antidepressants can increase suicidal thoughts and actions in children, adolescents, and young adults
Montelukast	Singulair	Allergic rhinitis	10 mg by mouth once a day		CysLT1 receptor antagonist	May cause serious neuropsychiatric events
Morphine	MS Contin Roxanol Kadian	Pain	ER formulations – 15 mg by mouth once a day Dosing is individualized	CII	Opioid receptor agonist	Monitor for at least 24 hours after first dose if given intrathecal or epidural route. Do not consume alcohol with ER capsules. Addiction, abuse, misuse. REMS. Serious or life-threatening respiratory depression. Neonatal withdrawal syndrome. Accidental exposure can be fatal, especially in children. Use in pregnancy can cause neonatal withdrawal syndrome. Concomitant use with benzodiazepines, alcohol, or other CNS depressants can result in respiratory depression, coma, and death
Mupirocin	Bactroban	Bacterial infection	Apply three times a day		Inhibits bacterial protein synthesis by reversibly and specifically binding to the bacterial isoleucyl transfer-RNA synthesis	
Naloxone	Narcan	Opioid overdose	0.4-2 mg IV/IM/subQ every 2-3 minutes as needed up to 10 mg		Selective mu-opioid receptor antagonist	
Nitrofurantoin	Macrodantin Macrobid Furadantin	Urinary tract infection	100 mg by mouth two times a day		Inhibition of acetyl-coenzyme A which interferes with carbohydrate metabolism	
Norepinephrine	Levophed	Shock	0.05-0.15 mcg/kg/minute, titrated to goal mean arterial pressure		Sympathomimetic amine that acts on both alpha- and beta-1 adrenergic receptors causing peripheral	

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
					vasoconstriction (alpha) and inotropic stimulation of the heart and coronary artery vasodilation (beta)	
Norethindrone/Ethinyl estradiol/Ferrous fumarate	Loestrin FE Loestrin FE 24 Microgestin FE Junel FE Minastrin 24	Contraception	One tablet by mouth once a day		Suppresses LH and FSH resulting in inhibition of ovulation. Thickening of cervical mucous may also contribute	Cigarette smoking increases risk of CV events. Increased risk of endometrial and breast cancers, cardiovascular disease, and dementia. Discuss risks and benefits with patients
Norgestimate/Ethinyl estradiol	Ortho-Cyclen Sprintec Estarylla	Contraception	One tablet by mouth once a day		Suppresses LH and FSH resulting in inhibition of ovulation. Thickening of cervical mucous may also contribute	Cigarette smoking increases risk of CV events
Nortriptyline	Pamelor	Depression	50 mg by mouth daily in single or divided doses		Prevents reuptake of norepinephrine and serotonin at the presynaptic nerve terminal which increases activity at post synaptic terminals	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Nystatin (topical)	Nystop	Fungal infection	Apply to affected area 2-3 times a day		Binds to ergosterol in the fungal cell membrane which causes pores in the membrane to leak cell contents and leads to cell death	
Ofloxacin Ophthalmic	Ocuflox	Antibiotic	1-2 drops into conjunctival sac every 2-4 hours while awake for two days followed by four times daily for the next five days		Inhibits DNA Gyrase	
Olanzapine	Zyprexa Zyprexa Relprevv	Schizophrenia	10-15 mg by mouth once a day		MAY be related to dopamine D2 and serotonin 5-HT2 receptor antagonism in CNS	Increased mortality in elderly patients with dementia-related psychosis. Post injection delirium and sedation syndrome with extended release injectable formulation
Olmesartan	Benicar	Hypertension	20 mg by mouth once a day		Angiotensin II antagonist. Blocks the binding of angiotensin II to the AT1 receptor	Fetal toxicity: discontinue drug as soon as pregnancy is detected
Omega-3-Acid Ethyl Esters	Lovasa	Dyslipidemia	4 grams daily in 1 or 2 doses with meals		MAY be inhibition of Acyl CoA, decreases lipogenesis in the liver	
Omeprazole	Prilosec	GERD	20 mg by mouth once a day		Inhibits H+/K+ ATPase enzyme system at the secretory surface of the gastric parietal cell	
Ondansetron	Zofran	Nausea and vomiting	8 mg by mouth three times a day		Selective 5-HT3 receptor antagonism	
Oxcarbazepine	Trileptal Oxtellar XR	Seizure	300 mg by mouth two times a day		Na+ channel antagonist; delays recovery of inactive Na+ channels	

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Oxybutynin	Ditropan Ditropan XL	Overactive bladder	5 mg XR by mouth once a day		Inhibition of the muscarinic action of acetylcholine on smooth muscle resulting in relaxation of bladder	
Oxycodone	OxyContin Roxicodone	Pain	Immediate release product: 5 mg by mouth every 4-6 hours as needed Extended release product: 10-20 mg by mouth two times a day	CII	Opioid receptor agonist	Opioids can be addictive, abused and misused which can lead to overdose and death. REMS. Serious life-threatening respiratory depression may occur. Accidental exposure can be fatal, especially in children. Use in pregnancy can cause neonatal withdrawal syndrome. Use with benzodiazepines, alcohol, or other CNS depressants can result in respiratory depression, coma, and death
Oxycodone/Acetaminophen	Percocet Roxicet Endocet	Pain	1 tablet by mouth every 6 hours as needed	CII	Opioid receptor agonist and inhibition of prostaglandin synthesis in CNS	Opioids can be addictive, abused and misused which can lead to overdose and death. REMS. Serious life-threatening respiratory depression may occur. Accidental exposure can be fatal, especially in children. Use in pregnancy can cause neonatal withdrawal syndrome. Use with benzodiazepines, alcohol, or other CNS depressants can result in respiratory depression, coma, and death
Oxytocin	Pitocin	Labor induction	Initial: 0.5-1 milliunit/min IV; gradually increase dose in increments of 1-2 milliunits until desired contraction pattern has been established		Increases intracellular concentrations of calcium in the uterine myometrial tissue prompting uterine contractions	Oxytocin is not indicated for elective induction of labor
Paliperidone	Invega	Antipsychotic	6 mg by mouth once daily in the morning		Direct antagonism of dopamine (D2) and serotonin (5-HT2) receptor in the central nervous system	
Pantoprazole	Protonix	GERD	40 mg by mouth once a day		Inhibits H+/K+ ATPase enzyme system at the secretory surface of the gastric parietal cell	
Paroxetine	Paxil Paxil XR Pexeva	Depression	20 mg by mouth once a day		Blocks neuronal uptake of serotonin in the central nervous system	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Phentermine Hydrochloride	Adipex-P Ionamin	Weight management	1 tablet/capsule take before breakfast or 1-2 hours after breakfast	CIV	Sympathomimetic amine	

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Phenylephrine	Generic/No brand name	Shock	0.5-2 mcg/kg/minute continuous IV infusion, titrate to desired MAP		Alpha adrenergic agonist that produces peripheral vasoconstriction and increased arterial pressure	
Phenytoin	Dilantin	Seizure	100 mg by mouth three times a day		Na+ channel antagonist; delays recovery of inactive Na+ channels	Cardiovascular risk associated with rapid injection
Phytonadione Vitamin K	Mephyton	Warfarin overdose	2.5-10 mg by mouth or up to 25 mg orally		Promotes liver synthesis of clotting factors. Exact MOA not well understood	Serious hypersensitivity reactions or anaphylaxis may occur with intramuscular or intravenous administration
Pioglitazone	Actos	Diabetes	15-30 mg by mouth once a day		May decrease insulin resistance by binding to peroxisome proliferator-activated receptor-gamma (PPAR)	Contraindicated in patients with CHF
Piperacillin/Tazobactam	Zosyn	Bacterial infection	3.375 g IV every 6 hours		Inhibition of bacterial cell wall synthesis via inhibiting peptidoglycan transpeptidase + beta-lactamase inhibition	
Potassium chloride	Klor-Con K-Dur Micro-K	Hypokalemia	20-80 mEq by mouth per day in single or divided doses		Main intracellular cation of most body tissues	
Pramipexole	Mirapex	Restless leg syndrome	0.125 mg by mouth once a day		D3 receptor agonist	
Pravastatin	Pravachol	Dyslipidemia	20 mg by mouth once a day		Selective competitive inhibitor of HMG-CoA reductase	
Prazosin	Minipress	Hypertension	1 mg by mouth 2-3 times a day		Selective antagonism of postsynaptic alpha 1 adrenergic receptors	
Prednisolone Acetate (Ophthalmic)	Pred Forte Pred Mild Omnipred	Ophthalmic anti-inflammatory	2 drops in affected eye four times daily		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Prednisone	Deltasone	Inflammatory disorders	2.5-10 mg by mouth daily		Stimulates synthesis of anti-inflammatory proteins and enzymes	
Pregabalin	Lyrica	Neuropathic pain	75 mg by mouth two times a day OR 50 mg by mouth three times a day	CV	Ca2+ channel antagonist; binds with alpha 2 subunit of Ca2+ channel	
Progesterone	Prometrium	Endometrial hyperplasia prophylaxis	200 mg by mouth once a day in the evening for 12 days every 28 day cycle		Increases synthesis of RNA by interacting with DNA causing transformation of proliferative endometrium into secretory endometrium	Increases risk of CV disorders, breast cancer, endometrial cancer, and in women over age of 65, probable dementia
Propofol	Diprivan	Sedation	5 mcg/kg/minute titrated to clinical effect	CIV in ND, but not controlled federally	May work through GABA-a receptor modulation increasing time chloride channels are open and increasing Cl- in cells	

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
Propranolol	Inderal Inderal LA Innopran XL	Hypertension	80 mg by mouth per day in single or divided doses		Non-selective beta-adrenergic blocker	
Quetiapine	Seroquel Seroquel XR	Schizophrenia	25 mg by mouth two times a day		Related to dopamine D2 and serotonin 5-HT2 receptor antagonism in CNS	Increased mortality in elderly patients with dementia-related psychosis. Increased risk of suicidal actions and thoughts in children, adolescents, and young adults
Ramipril	Altace	Hypertension	2.5-20 mg by mouth divided 1-2 times a day		Angiotensin-converting enzyme inhibitor	Fetal toxicity: discontinue drug as soon as pregnancy is detected
Rimegepant	Nurtec	Migraine	Acute treatment – 75 mg by mouth as a single dose Preventative treatment – 75 mg by mouth every other day		Calcitonin gene-related peptide receptor antagonist	
Risperidone	Risperdal Risperdal Consta Perseris	Schizophrenia	1 mg by mouth two times a day		Related to dopamine D2 and serotonin 5-HT2 receptor antagonism in CNS	Increased mortality in elderly patients with dementia related psychosis
Rivaroxaban	Xarelto	Anticoagulation	Extremely variable (reference indication and renal function)		Factor Xa inhibitor	Premature discontinuation of therapy increases the risk of thrombotic events. May cause spinal or epidural hematoma
Rizatriptan Benzoate	Maxalt Maxalt MLT	Migraine	5-10 mg by mouth at the first sign of a migraine		Selective agonist of vascular 5-HT1 receptor causing vasoconstriction	
Rocuronium	Zemuron	Tracheal intubation	0.45-0.6 mg/kg IV (individualized)		Blocks acetylcholine from binding to receptors inhibiting depolarization	
Ropinirole	Requip Requip XL	Parkinson's disease	0.25 mg by mouth three times a day		Dopamine agonist activity (high affinity to D3 receptor)	
Rosuvastatin	Crestor	Dyslipidemia	10-20 mg by mouth once a day (up to 40 mg)		HMG-CoA reductase inhibitor	
Sacubitril/Valsartan	Entresto	Heart failure	97/103 mg by mouth two times a day		Prodrug metabolized to the active metabolite LBQ657, which inhibits neprilysin, increasing levels of peptides and angiotensin receptor blocker to selectively block the AT1 receptor and inhibit angiotensin-II dependent aldosterone release	Fetal toxicity: discontinue drug as soon as pregnancy is detected
Semaglutide	Ozempic	Diabetes	Initial dose 0.25 mg injected subQ once weekly		Human glucagon-like peptide-1(GLP-1) receptor agonist	Contraindicated in patients with personal or family history of medullary thyroid cancer or multiple endocrine neoplasia syndrome type 2
Sertraline	Zoloft	Depression	50 mg by mouth once a day		Blocks neuronal uptake of serotonin in the central nervous system	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Sildenafil	Revatio	Pulmonary hypertension	20 mg by mouth three times a day		Phosphodiesterase type 5 inhibitor which prevents degradation of cGMP resulting in smooth muscle relaxation	
Sildenafil	Viagra	Erectile dysfunction	50 mg by mouth 1 hour prior to sexual activity		Phosphodiesterase type 5 inhibitor which prevents degradation of cGMP resulting in smooth muscle relaxation	
Simvastatin	Zocor	Dyslipidemia	20-40 mg by mouth in the evening		HMG-CoA reductase inhibitor	
Sitagliptin	Januvia	Diabetes	100 mg by mouth once a day		Dipeptidyl peptidase-4 enzyme inhibitor	
Solifenacin	Vesicare	Overactive bladder	5-10 mg by mouth once a day		Inhibits muscarinic receptors	
Sotalol	Betapace	Arrhythmia	80 mg by mouth two times a day		Beta-blocker which contains both Class II and Class III antiarrhythmic properties	Can cause life-threatening ventricular tachycardia associated with QT prolongation
Spironolactone	Aldactone	Edema	100 mg by mouth once a day		K sparing diuretic – aldosterone antagonist (primarily at the aldosterone-dependent Na/K exchange sites in the distal convoluted tubule)	
Succinylcholine	Quelicin Anectine	Tracheal intubation	0.6 mg/kg IV		Depolarization of the end motor plate	Risk of cardiac arrest from hyperkalemic rhabdomyolysis
Sucralfate	Carafate	Duodenal ulcer	1 g by mouth four times a day on an empty stomach		May form ulcer-adherent complex that covers ulcer site and protects it against further attack by acid, pepsin, and bile	
Sulfamethoxazole/Trimethoprim	Bactrim Septra	Bacterial infection	DS – 1 tablet by mouth two times a day		Inhibition of bacterial dihydrofolic acid and tetrahydrofolic acid synthesis	
Sumatriptan	Imitrex	Migraine	50 mg by mouth at first sign of migraine		Selective agonist of vascular 5-HT1 receptor causing vasoconstriction	
Tadalafil	Cialis Adcirca*	Erectile dysfunction	10 mg by mouth one hour prior to sexual activity		Phosphodiesterase type 5 inhibitor which prevents degradation of cGMP resulting in smooth muscle relaxation	
Tamoxifen	Nolvadex	Breast cancer	20-40 mg by mouth in two divided doses		Competes with estrogen for binding sites in target tissues	
Tamsulosin	Flomax	Benign prostatic hypertrophy (BPH)	0.4 mg by mouth once a day		Alpha 1A-adrenoreceptor antagonist	
Telmisartan	Micardis	Hypertension	20-80 mg by mouth daily		Blocks vasoconstrictor and aldosterone secreting effects of angiotensin II by selectively blocking	

Generic **P1 P2 P3**	**Brand** **P1 P2 P3**	**Indication** **P1 P2 P3**	**Dosing** **P2 P3**	**Controlled** **P2 P3**	**Mechanism of Action** **P3**	**Black Box Warning** **P3**
					the binding of angiotensin II to the ATI receptor in tissues	
Terbinafine	Lamisil	Antifungal	250 mg by mouth daily for 6-12 weeks for nail infection Topical cream – apply to affected area 1-2 times daily for 7 days		Inhibits squalene epoxidase, an enzyme found in fungi	
Testosterone	Androgel	Male hypogonadism	50 mg applied topically once a day to upper arms and shoulders	CIII	Stimulation of mRNA resulting in synthesis of specific proteins and enzymes that carry out its actions	Certain formulations may increase blood pressure. Secondary exposure has caused virilization in children. Serious pulmonary oil microembolism reactions can occur after injection
Tirzepatide	Mounjaro Zepbound	Diabetes Obstructive sleep apnea Weight management	Initial dose 2.5 mg subcutaneously once weekly		Glucose-dependent insulinotropic polypeptide (GIP) receptor agonist and human glucagon-like peptide-1(GLP-1) receptor agonist	Contraindicated in patients with personal or family history of medullary thyroid cancer or multiple endocrine neoplasia syndrome type 2
Tizanidine	Zanaflex	Musculoskeletal pain/spasm	4 mg by mouth every 6-8 hours		Alpha 2-adrenergic receptor agonist	
Tolterodine	Detrol Detrol LA	Overactive bladder	4 mg by mouth once a day in an LA capsule		Muscarinic receptor antagonist	
Topiramate	Topamax Trokendi XR	Seizure	50 mg by mouth once a day		Na+ channel antagonist; GABA-a receptor agonist; AMPA receptor antagonist	
Tramadol	Ultram Ultram ER	Pain	50-100 mg by mouth every 4-6 hours as needed	CIV	Opioid receptor agonist AND blocks reuptake of norepinephrine and serotonin	Addiction, abuse, and misuse. REMS. Life-threatening respiratory depression. Accidental exposure. Ultra-rapid metabolism increases risk of respiratory depression in children. Neonatal opioid withdrawal syndrome. Interactions with drugs affecting cytochrome P450 enzymes. Risks from concomitant use with benzodiazepines or other CNS depressants
Trazodone	Desyrel	Depression	150 mg by mouth daily in single or divided doses		Inhibits serotonin reuptake by brain synaptosomes	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Triamcinolone Acetonide (topical)	Kenalog	Inflammatory disorders	Apply 2-4 times a day		Stimulates synthesis of anti-inflammatory properties and enzymes	
Triamterene/ Hydrochlorothiazide	Dyazide Maxzide	Edema	1-2 doses by mouth two times a day		Inhibits reabsorption of Na+ in exchange for K+ at the distal renal tubule/ inhibits the reabsorption of Na+ and Cl- at the distal renal tubule	Hyperkalemia can occur

Generic P1 P2 P3	Brand P1 P2 P3	Indication P1 P2 P3	Dosing P2 P3	Controlled P2 P3	Mechanism of Action P3	Black Box Warning P3
Ursodiol	Actigall	Gall bladder disease	300 mg two times a day by mouth with food		Dissolves cholesterol containing gall stones. Replaces/displaces toxic concentrations of endogenous hydrophobic bile acids that tend to accumulate in cholestatic liver disease	
Valacyclovir	Valtrex	Viral infection	1 g by mouth two times a day		Hepatic metabolism to acyclovir which inhibits viral DNA synthesis	
Vancomycin	Vancocin	Bacterial infection	Individualized doses Monitor trough levels		Inhibits bacterial cell wall synthesis via inhibition of peptidoglycan transglycosylase	Risk of embryo-fetal toxicity due to some excipients
Venlafaxine	Effexor Effexor XR	Depression	75 mg XR by mouth once a day		Blocks neuronal uptake of serotonin and norepinephrine in the central nervous system	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Vilazodone Hydrochloride	Viibryd	Depression	40 mg by mouth one daily with food		Selective inhibition of serotonin reuptake in the CNS and partial agonist of 5-HT (1A) receptors	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Vortioxetine	Trintellix	Depression	20 mg by mouth once a day		Inhibits the reuptake of serotonin; antagonizes 5-HT3, 5-HT1D, and 5-HT7 receptors; agonizes 5-HT1A receptors; and is a 5-HT1B receptor partial agonist	Antidepressants increase suicidal thoughts and actions in children, adolescents, and young adults
Warfarin	Coumadin Jantoven	Anticoagulation	5 mg by mouth once a day. Individualize dose regimen by monitoring INR		Antagonizes vitamin K resulting in a decreased hepatic synthesis of vitamin K dependent clotting factors	Increases the risk of fatal bleeding. Perform regular monitoring of INR
Zolpidem	Ambien Ambien CR Intermezzo Zolpimist	Insomnia	5-12.5 mg by mouth at bedtime	CIV	Selective agonist of GABA-BZ1 receptor	Complex sleep behaviors including sleep-walking, sleep-driving, and engaging in other activities while not fully awake