

Problem 1

The metric is given by

$$ds^2 = d\theta^2 + \sin^2 \theta d\phi^2. \quad (1)$$

From the definition of the Christoffel symbols,

$$\Gamma^\mu_{\rho\sigma} = \frac{1}{2}g^{\mu\nu} \left(\frac{\partial g_{\nu\sigma}}{\partial x^\rho} + \frac{\partial g_{\rho\nu}}{\partial x^\sigma} - \frac{\partial g_{\rho\sigma}}{\partial x^\nu} \right), \quad (2)$$

we can easily calculate that,

$$\begin{aligned} \Gamma^\theta_{\phi\phi} &= -\frac{1}{2}g^{\theta\theta} \frac{\partial g_{\phi\phi}}{\partial \theta} = -\frac{1}{2}\partial_\theta(\sin^2 \theta) = -\sin \theta \cos \theta, \\ \Gamma^\phi_{\theta\phi} &= \frac{1}{2}g^{\phi\phi} \frac{\partial g_{\phi\phi}}{\partial \theta} = \frac{1}{2} \frac{1}{\sin^2 \theta} \partial_\theta(\sin^2 \theta) = \cot \theta. \end{aligned} \quad (3)$$

Problem 2

Given $ds^2 = dx^2 + dy^2 + dz^2$ and for a two-sphere with radius R , we have

$$\begin{cases} x = R \sin \theta \cos \phi \\ y = R \sin \theta \sin \phi \\ z = R \cos \theta \end{cases}. \quad (4)$$

Then,

$$\begin{aligned} dx &= R \cos \theta \cos \phi d\theta - R \sin \theta \sin \phi d\phi, \\ dy &= R \cos \theta \sin \phi d\theta + R \sin \theta \cos \phi d\phi, \\ dz &= -R \sin \theta d\theta. \end{aligned} \quad (5)$$

Therefore,

$$\begin{aligned} ds^2 &= dx^2 + dy^2 + dz^2 \\ &= R^2 d\theta^2 + R^2 \sin^2 \theta d\phi^2. \end{aligned} \quad (6)$$

Problem 3

In the polar coordinates, we have $\vec{x} = r\hat{r}$ and

$$x = r \cos \theta, \quad y = r \sin \theta. \quad (7)$$

Therefore,

$$d\vec{x} = (dx, dy) = (\cos \theta dr - r \sin \theta d\theta, \sin \theta dr + r \cos \theta d\theta). \quad (8)$$

And

$$d\vec{x} \cdot d\vec{x} = dx^2 + dy^2 = dr^2 + r^2 d\theta^2. \quad (9)$$

Also

$$\begin{aligned}\vec{x} \cdot d\vec{x} &= r \cos^2 \theta dr - r^2 \sin \theta \cos \theta d\theta + r \sin^2 \theta dr + r^2 \sin \theta \cos \theta d\theta \\ &= r dr.\end{aligned}\tag{10}$$

Problem 4

Pove

$$(\xi^n)_\alpha^\beta = K^n (xCx)^{n-1} C_{\alpha\rho} x^\rho x^\beta, \tag{11}$$

by induction, where the matrix $\xi_\alpha^\nu = KC_{\alpha\rho} x^\rho x^\nu$ and $xCx \equiv C_{\mu\nu} x^\mu x^\nu$.

First,

$$\begin{aligned}(\xi^2)_\alpha^\beta &= \xi_\alpha^\nu \xi_\nu^\beta = K^2 C_{\alpha\rho} x^\rho x^\nu C_{\nu\sigma} x^\sigma x^\beta \\ &= K^2 (xCx) C_{\alpha\rho} x^\rho x^\beta.\end{aligned}\tag{12}$$

Suppose equation (11) holds for $n-1$,

$$(\xi^{n-1})_\alpha^\beta = K^{n-1} (xCx)^{n-2} C_{\alpha\rho} x^\rho x^\beta, \tag{13}$$

then,

$$\begin{aligned}(\xi^n)_\alpha^\beta &= (\xi^{n-1})_\alpha^\nu \xi_\nu^\beta = (K^{n-1} (xCx)^{n-2} C_{\alpha\rho} x^\rho x^\nu) (KC_{\nu\sigma} x^\sigma x^\beta) \\ &= K^n (xCx)^{n-1} C_{\alpha\rho} x^\rho x^\beta.\end{aligned}\tag{14}$$

Problem 5

Prove that

$$R_{\mu\rho} = \partial_\mu \partial_\rho \log \sqrt{-g} - \frac{1}{\sqrt{-g}} \partial_\sigma (\sqrt{-g} \Gamma_{\mu\rho}^\sigma) + \Gamma_{\mu\alpha}^\sigma \Gamma_{\rho\sigma}^\alpha. \tag{15}$$

From the definition of Ricci tensor, L.H.S. equals

$$R_{\mu\rho} = R_{\mu\rho\alpha}^\alpha = \Gamma_{\mu\alpha,\rho}^\alpha - \Gamma_{\mu\rho,\alpha}^\alpha + \Gamma_{\mu\alpha}^\sigma \Gamma_{\sigma\rho}^\alpha - \Gamma_{\mu\rho}^\sigma \Gamma_{\sigma\alpha}^\alpha. \tag{16}$$

Fisrt expand the Christoffel symbol of the form

$$\Gamma_{\mu\alpha}^\alpha = \frac{1}{2} g^{\alpha\beta} \left(\frac{\partial g_{\alpha\beta}}{\partial x^\mu} + \frac{\partial g_{\mu\beta}}{\partial x^\alpha} - \frac{\partial g_{\mu\alpha}}{\partial x^\beta} \right) = \frac{1}{2} g^{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial x^\mu}. \tag{17}$$

In the last step, the last two terms inside the bracket, when dotted with $g^{\alpha\beta}$, gives zero, because they are antisymmetric in α, β , while the metric $g^{\alpha\beta}$ is symmetric in α, β . Now consider

$$\begin{aligned}\partial_\mu \partial_\rho \log \sqrt{-g} &= \partial_\rho \left(\frac{\partial g_{\alpha\beta}}{\partial x^\mu} \frac{\partial g}{\partial g_{\alpha\beta}} \frac{\partial}{\partial g} (\log \sqrt{-g}) \right) \\ &= \partial_\rho \left(\frac{\partial g_{\alpha\beta}}{\partial x^\mu} (g g^{\alpha\beta}) \left(\frac{1}{2g} \right) \right) \\ &= \partial_\rho \left(\frac{1}{2} g^{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial x^\mu} \right) = \Gamma_{\mu\alpha,\rho}^\alpha.\end{aligned}\tag{18}$$

Next consider

$$\begin{aligned}
-\frac{1}{\sqrt{-g}}\partial_\sigma(\sqrt{-g}\Gamma^\sigma_{\mu\rho}) &= -\frac{1}{\sqrt{-g}}\partial_\sigma(\sqrt{-g})\Gamma^\sigma_{\mu\rho} - \Gamma^\sigma_{\mu\rho,\sigma} \\
&= -\frac{1}{2}g^{\alpha\beta}\frac{\partial g_{\alpha\beta}}{\partial x^\sigma}\Gamma^\sigma_{\mu\rho} - \Gamma^\alpha_{\mu\rho,\alpha} \\
&= -\Gamma^\alpha_{\sigma\alpha}\Gamma^\sigma_{\mu\rho} - \Gamma^\alpha_{\mu\rho,\alpha}.
\end{aligned} \tag{19}$$

Therefore, R.H.S. of (15) equals

$$\Gamma^\alpha_{\mu\alpha,\rho} - \Gamma^\alpha_{\mu\rho,\alpha} + \Gamma^\sigma_{\mu\alpha}\Gamma^\alpha_{\rho\sigma} - \Gamma^\alpha_{\sigma\alpha}\Gamma^\sigma_{\mu\rho}. \tag{20}$$

Compare (16) and (20), we see that equation (15) is proved.

Problem 6

First consider

$$\begin{aligned}
\partial_\mu\partial_\rho\log\sqrt{-g} &= -\frac{1}{2}\partial_\mu\partial_\rho\log(1-K(xCx)) = \partial_\mu\left(\frac{KC_{\rho\sigma}x^\sigma}{1-K(xCx)}\right) \\
&= \frac{KC_{\rho\mu}(1-K(xCx)) + 2K^2C_{\mu\nu}C_{\rho\sigma}x^\nu x^\sigma}{(1-K(xCx))^2} \\
&= \frac{KC_{\rho\mu}}{1-K(xCx)} + \frac{2K^2C_{\mu\nu}C_{\rho\sigma}x^\nu x^\sigma}{(1-K(xCx))^2} \\
&= \frac{K}{1-K(xCx)}g_{\mu\rho} + \frac{K^2}{(1-K(xCx))^2}C_{\mu\alpha}C_{\rho\beta}x^\alpha x^\beta.
\end{aligned} \tag{21}$$

Next consider in N dimensional spacetime,

$$\begin{aligned}
-\frac{1}{\sqrt{-g}}\partial_\alpha(\sqrt{-g}\Gamma^\alpha_{\mu\rho}) &= \partial_\alpha\left(\frac{1}{\sqrt{-g}}\right)\sqrt{-g}\Gamma^\alpha_{\mu\rho} - \partial_\alpha\Gamma^\alpha_{\mu\rho} \\
&= \partial_\alpha(\sqrt{1-K(xCx)})\frac{Kx^\alpha g_{\mu\rho}}{\sqrt{1-K(xCx)}} - \partial_\alpha(Kx^\alpha g_{\mu\rho}) \\
&= -\frac{K^2(xCx)}{1-K(xCx)}g_{\mu\rho} - K\partial_\alpha(x^\alpha)g_{\mu\rho} - Kx^\alpha\partial_\alpha g_{\mu\rho} \\
&= -\frac{K^2(xCx)}{1-K(xCx)}g_{\mu\rho} - K\delta_\alpha^\alpha g_{\mu\rho} \\
&\quad - Kx^\alpha\left(\frac{KC_{\mu\alpha}C_{\rho\beta}x^\beta + KC_{\rho\alpha}C_{\mu\beta}x^\beta}{1-K(xCx)} + \frac{2K^2C_{\mu\nu}C_{\rho\sigma}C_{\alpha\beta}x^\nu x^\sigma x^\beta}{(1-K(xCx))^2}\right) \\
&= -\frac{K^2(xCx)}{1-K(xCx)}g_{\mu\rho} - KNg_{\mu\rho} - \frac{2K^2}{(1-K(xCx))^2}C_{\mu\alpha}C_{\rho\beta}x^\alpha x^\beta
\end{aligned} \tag{22}$$

Then consider

$$\begin{aligned}
\Gamma_{\mu\alpha}^{\sigma}\Gamma_{\rho\sigma}^{\alpha} &= K^2 x^{\sigma} x^{\alpha} g_{\mu\alpha} g_{\rho\sigma} \\
&= K^2 x^{\sigma} x^{\alpha} \left(C_{\mu\alpha} + \frac{K C_{\mu\nu} C_{\alpha\beta} x^{\nu} x^{\beta}}{1 - K(xCx)} \right) \left(C_{\rho\sigma} + \frac{K C_{\rho\nu} C_{\sigma\beta} x^{\nu} x^{\beta}}{1 - K(xCx)} \right) \\
&= K^2 C_{\mu\alpha} C_{\rho\sigma} x^{\alpha} x^{\sigma} + \frac{2K^3(xCx) C_{\rho\alpha} C_{\mu\beta} x^{\alpha} x^{\beta}}{1 - K(xCx)} \\
&\quad + \frac{K^4(xCx)^2 C_{\rho\alpha} C_{\mu\beta} x^{\alpha} x^{\beta}}{(1 - K(xCx))^2} \\
&= \left(K^2 + \frac{2K^3(xCx)}{1 - K(xCx)} + \frac{K^4(xCx)^2}{(1 - K(xCx))^2} \right) C_{\mu\alpha} C_{\rho\beta} x^{\alpha} x^{\beta} \\
&= \frac{K^2}{(1 - K(xCx))^2} C_{\mu\alpha} C_{\rho\beta} x^{\alpha} x^{\beta}. \tag{23}
\end{aligned}$$

Now add them up and we obtain

$$R_{\mu\rho} = \frac{K}{1 - K(xCx)} g_{\mu\rho} - \frac{K^2(xCx)}{1 - K(xCx)} g_{\mu\rho} - KN g_{\mu\rho} = K(1 - N) g_{\mu\rho}. \tag{24}$$