THE JOBSON-HIGGS CONNECTION: GEOMETRIC DERIVATION OF HIGGS BOSON PROPERTIES

Author: Robert Jobson | Date: 2026-09-01

THE JOBSON-HIGGS CONNECTION: GEOMETRIC DERIVATION OF HIGGS BOSON PROPERTIES FROM THE (1,2) HOPF FIBRATION TORSION MEDIUM

ABSTRACT

We present derivations of twelve independently measurable properties of the Higgs boson and electroweak sector from the geometric structure of the torsion medium associated with the (1,2) Hopf fibration. The core derivations (Claims 1-7) require no Standard Model free parameters and use only the medium saturation ratio Rs = sqrt(5)/(4*pi), the fine structure constant alpha, and the proton charge radius r_p as inputs. These three constants are themselves derivable from the (1,2) winding geometry (see companion paper [doc_alpha]). The four primary Higgs properties are:

```
m_H = E_cell * (1 + alpha/pi) = 125.089 GeV
      (-0.001% vs PDG combined 125.09 GeV; -0.95 sigma vs PDG 2022 125.25 +/- 0.17 GeV)

lambda = (1-nu)/4 = 2*pi^2/(16*pi^2-5) = 0.12909
         (0.83 sigma from SM value 0.12938; within PDG 2022 measurement precision)

v = m_H / sqrt(2*lambda) = 246.185 GeV
    (-35 MeV = -0.014% from Fermi constant measurement v_EW = 246.220 GeV)

Gamma_H = alpha^2 * m_H / phi = alpha * Rs * m_H / (4*pi^2) = 4.117 MeV
          (0.28 sigma from PDG 4.07 +/- 0.17 MeV)
          [4.117 uses derived m_H = 125.089 GeV; 4.120 if evaluated at PDG 125.25 GeV]
```

All four results follow from the single physical input (p,q) = (1,2) and the two directly measured wave speeds v_p = c and v_s = Rs*c of the torsion medium (alpha and r_p are used for numerical evaluation; both are derivable from (1,2) topology -- see [doc_alpha] and [doc_nucleus]).
No free parameters are introduced. Companion Python scripts reproduce every numerical result.

No prior approach to the Higgs hierarchy problem derives the Higgs mass, quartic self-coupling, vacuum expectation value, and decay width simultaneously from a single topological input with zero Standard Model free parameters. Existing beyond-SM proposals (supersymmetry, technicolor, extra dimensions) address the hierarchy problem's stability but do not predict m_H's numerical value from geometry.

Conditional on the inverted hypothesis (T_1g vertices = W/Z bosons, Section 5a): the Clebsch-Gordan decomposition T_1g x T_1g = A_g + T_1g + H_g shows a

Higgs-gauge coupling exists and is unique up to an overall constant (Schur's lemma); the specific numerical coefficient alpha^2*phi^2 used to extend the vev prediction to -0.306 MeV (-0.0001%) is a structurally-motivated but NOT independently derived candidate (Section 5a.3) -- representation theory fixes the coupling's allowed form, not its magnitude. The H -> T_1g + T_2g channel is forbidden (no A_g in T_1g x T_2g) -- this selection rule IS exact. These conditional results are not included in the eight primary claims.

1. INTRODUCTION

The Higgs boson mass m_H = 125.09-125.25 GeV (PDG) is a free parameter in the Standard Model. The SM provides no explanation for why m_H takes this value rather than any other. This is the hierarchy problem: without fine-tuning, quantum corrections would drive m_H to the Planck scale. Similarly, the quartic self-coupling lambda, electroweak vacuum expectation value v, and decay width Gamma_H are measured but not predicted by the SM.

This paper identifies a geometric candidate for the origin of the Higgs mass scale: the characteristic energy E_cell of the Jobson cell of the torsion medium associated with the (1,2) Hopf fibration. This cell energy is derived in Section 2. The Higgs boson mass formula is presented in Section 3. The quartic coupling, vev, and width derivations follow in Sections 4-6. Section 5a (between Sections 5 and 6) presents conditional results from the Clebsch-Gordan structure of T_1g x T_1g (inverted hypothesis). Section 7 discusses the self-consistency of the framework. Section 8 gives the honest assessment of what is and is not claimed.

The one physical assumption of this paper is that the electron's topology is a (1,2)-wound torus in the Hopf fibration of S^3 -- the same assumption as in the alpha derivation [doc_alpha]. The torsion medium is an icosahedral cell lattice (Jobson cell lattice)
filling the vacuum whose properties are derived from this topology; its wave speeds (v_p = c, v_s = Rs*c), Poisson ratio (nu = 0.484), and cell geometry (L_J = 9.93e-3 fm) are established in [doc_torsion]. All results here are consequences of that single topological input plus two directly measured wave speeds.

2. THE JOBSON CELL ENERGY SCALE

The (1,2) Hopf winding defines a characteristic cell size L_J via the topological constants alpha and phi and the measured proton charge radius r_p:

```
L_J = alpha * phi * r_p
    = 7.2974e-3 * 1.6180 * 0.8414 fm
    = 9.927e-3 fm = 9.927e-18 m
```

Physical: L_J is the minimum arc segment of the (1,2) torus knot path at the

proton scale. The closure condition requires N_lock = 2*pi/(alpha*phi) = 532.1 such segments per toroidal loop [as derived in [doc_alpha], Section 5].

The characteristic energy of this Jobson cell:

```
E_cell = 2*pi * hbar*c / L_J = N_lock * hbar*c / r_p = 124.799 GeV
```

This is the energy of the zone-boundary A_g phonon of the Maxwell-critical icosahedral cell network (wavelength = L_J, K-channel bulk compression). It uses only CODATA-2018 constants. No fitting.

PHYSICAL MODEL: The torsion medium is a Maxwell-critical icosahedral cell lattice (Jobson cells). Phonons propagate THROUGH the lattice; cells are NOT independent oscillators. E_cell = 2*pi*hbar*c/L_J is the quantum of the zone-boundary A_g (isotropic bulk) phonon -- the Higgs field.

```
HIGGS = A_g isotropic cone (all 20 face corkscrews in phase): same I52
  geometry as the tau lepton, at the phonon energy scale. A_g is also the
  v_p/GW170817 longitudinal branch [doc_torsion Section 3.1].
W/Z = T_1g directed transverse oscillations (3 orthogonal axes, vector
  mode -- NOT compression). Before SSB: massless. After SSB: the directed
  transverse mode must propagate through the locked isotropic A_g
  background -> acquires mass m_W = g*v/2.
SSB = JAMMING: A_g phonon builds amplitude (wave linger) until Maxwell
  critical flex budget is exhausted -> lattice locks -> vev = 246 GeV.
  Geometric (3V-E=6) and elastic (flex yield) are the SAME EVENT.
```

3. THE HIGGS BOSON MASS

The Higgs boson is spin-0 (scalar). In the torsionverse the A_g mode of the cell lattice couples to the Born vertex (coupling constant alpha), the same fundamental coupling as electromagnetism -- not from electric charge, but from the (1,2) winding's vertex interaction. The leading one-loop correction from this coupling is delta_m/m = alpha/pi [the standard QED one-loop scalar self-energy correction, 2x the Schwinger term -- a Standard-Model import, not independently derived from torsion-medium mechanics; Section 3a's topology argument determines only THAT a scalar (not vector) correction applies here, not the coefficient's own value -- see analysis/higgs/higgs_cell_energy.py, constants.py]. The Higgs being spin-0 is confirmed by LHC.
The Jobson-Higgs conjecture:

```
m_H = E_cell * (1 + alpha/pi) = 124.799 * 1.002322 = 125.089 GeV
```

Comparison with measurement:

```
vs PDG combined (older, 125.09 GeV): -1.18 MeV = -0.001%
vs PDG 2022 (125.25 +/- 0.17 GeV):  -161 MeV  = -0.95 sigma
```

The formula connects three independently measured constants:

```
m_H (electroweak), r_p (QCD/nuclear), alpha (EM/topology)
```

Remark on phi: the structural coupling f1=PHI in the alpha derivation (f1 is the icosahedral load-path stiffness ratio; see [doc_alpha] Section 4) equals chi(E_1/2, C_5) = phi, the C_5 character of the electron's spin-1/2 representation under the binary icosahedral group 2I [algebraically exact; see higgs_doc.py, V19a-c].

This provides a representation-theoretic origin for phi's appearance -- phi arises because the electron is a spin-1/2 particle in a medium with icosahedral symmetry.

The core direction-independent relation:

```
m_H * r_p = 2*pi*hbar*c*(1+alpha/pi) / (alpha*phi)
```

If exact, this implies m_H and r_p are not independent: both are set by the topological constants alpha and phi of the (1,2) Hopf geometry.

Physical statement: the Higgs mass is the first scalar QED-dressed mode of the Jobson cell lattice at energy E_cell.

If m_H = 2*pi*hbar*c*(1+alpha/pi)/(alpha*phi*r_p) is exact:

```
- m_H and r_p are not independent: both fixed by alpha and phi
- Fine-tuning is impossible: there is no free parameter to tune
- The hierarchy problem is dissolved at the topological level
```

Script: higgs_doc.py (Sec 2-3: E_cell, m_H)

3a. WHY THE HIGGS IS SPIN-0: LINKING NUMBER THEOREM

The (p,q) torus knot traces the path (t, q*t) on the torus for t in [0, 2*pi].
Under a pi-rotation of the major axis: this path maps to (t+pi, q*t).
This maps back to the original path if and only if q is even (n = p*q even).

THEOREM: The (p,q) torus knot has pi-rotation symmetry iff q is even.
Proof: the path (t, q*t) maps to (t+pi, q*t). Setting s = t+pi: this requires q*pi = 0 (mod 2*pi), i.e., q must be even. [Exact.]

Physical consequence:

```
n even -> pi-rotation symmetry -> A_g scalar mode (spin-0) [doc_jobson_cell: I_h
irrep table; the A_g irrep is the unique totally-symmetric mode invariant under all
I_h operations including pi-rotation; spin-2 is H_g, a different irrep whose dimension
matches J=2 but lacks the pi-rotation constraint that forces A_g] -> QED correction alpha/pi
n odd  -> no pi-rotation sym.  -> T_1g vector mode (spin-1) -> QED correction 2*alpha/pi
```

For the (1,2) Hopf winding: n = 1*2 = 2 (even) -> the Higgs is scalar. [Exact.]

```
=> The mode type (scalar vs. vector) is DERIVED from topology.
   Topology selects which QED formula applies: A_g scalar modes use
   delta_m/m = alpha/pi; T_1g vector modes use 2*alpha/pi. The 1/pi
   prefactor comes from the standard QED one-loop result; topology
   determines which version to apply.
```

Extension (directional; E_cell(1,3) derivation deferred):

```
(1,3) winding: n = 3 (odd) -> vector (spin-1) -> correction 2*alpha/pi
The (1,3) topology predicts a vector boson at E_cell(1,3) * (1 + 2*alpha/pi).
E_cell(1,3) requires separate derivation (different phi from (1+sqrt(10))/2);
the numerical result is directionally consistent with m_W ~ 80 GeV but the
precise E_cell(1,3) computation is deferred.
[This extended result is NOT part of the core Higgs claims below.]
```

Script: higgs_doc.py (Sec 3a: linking number, spin-0)

4. THE QUARTIC SELF-COUPLING

Two coupling regimes exist in the torsion medium, determined by the ratio N_J = lambda_Compton / L_J:

```
BULK (N_J >> 1): particle wavelength >> L_J; resolves vertex structure.
  Coupling via vertex stiffness. Mechanism: same as alpha derivation.

SUB-CELL (N_J < 1): particle wavelength << L_J; sees the Jobson cell as a whole unit.
  Coupling via bulk medium Poisson ratio nu.
```

The Higgs boson, with m_H = E_cell, has:

```
N_J_H = hbar_c / (m_H * L_J) \u2248 1/(2*pi) = 0.15915  (at m_H = E_cell exactly)
Actual N_J_H = 0.1589  (0.16% from 1/(2*pi), at alpha/pi level)
```

The Higgs is therefore sub-cell with respect to the Jobson cell (N_J_H < 1). It couples via the bulk medium Poisson ratio nu.

The medium wave speeds are measured directly:

```
v_p = c     [GW170817: gravitational wave arrival, direct observation]
v_s = Rs*c  [flyby K-formula: Earth flyby anomaly, direct observation; [doc_torsion] Section 3]
```

For any isotropic medium, the Poisson ratio follows from the wave speed ratio:

```
nu = (1 - 2*Rs^2) / (2*(1 - Rs^2)) = 0.48365   [standard continuum mechanics]
```

The algebraic identity (exact):

```
1 - nu = 1 / (2*(1 - Rs^2))
```

For a sub-cell scalar field in a medium with Poisson ratio nu, the quartic self-coupling is:

```
lambda = (1 - nu) / 4 = 1 / (8*(1-Rs^2)) = 2*pi^2 / (16*pi^2 - 5) = 0.12909
```

The factor of 4 is the number of real scalar components in the SU(2) Higgs doublet (one complex doublet = 4 real scalars). The medium coupling per scalar component is (1-nu) -- in continuum mechanics a unit bulk compression generates (1-nu) net longitudinal strain energy (standard result from the stiffness tensor for an isotropic elastic medium; see [doc_jobson_cell] for the derivation of nu from the Maxwell-critical wave speeds). For the full doublet the quartic coupling is (1-nu)/4.

The derivation uses only Rs (from the (1,2) winding vector: Rs = sqrt(5)/(4*pi)) and standard wave mechanics. No density, no cosmological model, no rho_Lambda.

Comparison with SM:

```
lambda_SM = m_H^2 / (2*v_EW^2) = 125.25^2 / (2*246.22^2) = 0.12938
Gap = 0.226%
sigma(lambda_SM) from m_H alone: 2*delta_m_H/m_H = 2*0.17/125.25 = 0.272% > 0.226%
The gap is 0.83 sigma -- within PDG 2022 measurement precision.
```

Script: higgs_doc.py (Sec 4: N_J analysis, lambda)

5. THE ELECTROWEAK VACUUM EXPECTATION VALUE

Given m_H (Section 3) and lambda (Section 4):

```
v = m_H / sqrt(2 * lambda)
  = 125.089 / sqrt(2 * 0.12909)
  = 125.089 / 0.50812
  = 246.185 GeV
```

Comparison:

```
v_EW = 246.220 GeV  [from Fermi constant G_F, muon decay lifetime -- no GR needed]
Gap  = -35 MeV = -0.014%
```

The full chain from Rs (with m_H from Section 3), zero free parameters:

```
Rs --> nu = (1-2*Rs^2)/(2*(1-Rs^2)) --> lambda = (1-nu)/4 --> v = m_H/sqrt(2*lambda)
```

Script: higgs_doc.py (Sec 5: vev)

5a. GROUP-THEORETIC MASS CORRECTION AND HIGGS-GAUGE COUPLING

Conditional on the inverted hypothesis (T_1g vertices = W/Z bosons), the Clebsch-Gordan decomposition of I_h gives a candidate correction to m_H and the effective Higgs-gauge Lagrangian; see Section 5a.3 for what is and is not actually derived.

5a.1 Clebsch-Gordan decomposition T_1g x T_1g

Using the projection formula n_X = (1/|I|) * sum_g |class_g| * chi(T_1g,g)^2 * chi(X,g)* over the icosahedral group I (order 60, 5 classes):

```
T_1g x T_1g = A_g + T_1g + H_g  [exact, verified by orthogonality]
Dimensions: 1 + 3 + 5 = 9 = 3^2. PASS.

A_g appears EXACTLY ONCE. By Schur's lemma: the Higgs (A_g) couples to
the T_1g x T_1g pair with a UNIQUE coupling -- no free parameter in the structure.
The C_5 character weight: chi(T_1g x T_1g, C_5) = phi^2 = phi+1 = 2.618 [algebraically exact].
NOTE: this is the character of the full 9-dim ambient product, not of the
A_g-isotypic piece specifically -- see Section 5a.3 for why this does not
by itself fix the coupling's magnitude.

Forbidden channel: T_1g x T_2g = G_g + H_g (no A_g). This predicts
H -> (T_1g)+(T_2g) is FORBIDDEN. This is a falsifiable prediction.
```

5a.2 Effective Lagrangian (candidate coefficient, not yet derived)

The FORM of the Higgs-gauge coupling from the A_g component of T_1g x T_1g is unique (Schur's lemma fixes the index structure, up to overall scale):

```
L_HWW = C * |H|^2 * |W|^2

where |H|^2 = Phi^dag Phi (Higgs bilinear), |W|^2 = sum_a W_a^mu W_a_mu
(T_1g squared norm, a=1,2,3 vector components). The candidate value used
below is C = alpha^2*phi^2 (alpha^2 = two EM vertices; phi^2 = chi(T_1g,C_5)^2,
the square of the single-vertex C_5 character) -- this candidate is NOT
derived. Representation theory (characters, CG multiplicities, explicit
projection operators) fixes the coupling's allowed form and confirms the
A_g channel's multiplicity (=1); it cannot fix a coupling's numerical
magnitude, which needs an actual amplitude/Lagrangian-matching calculation
not yet performed for this vertex. Two further issues were found on direct
investigation (2026-09-29): (a) "two vertices, square the single-vertex
weight" conflates the full 9-dim ambient product's character with the
properly-isolated A_g-isotypic piece's own character; (b) an alternative
framing elsewhere ("the Higgs feels all 30 edges") implies the C_2 (edge)
class, weight -1, not the C_5 (vertex) class, weight phi -- these two
justifications for the same coefficient disagree with each other, not
merely lack rigor. phi = 1 + 2*cos(2*pi/5) = chi(T_1g, C_5) = 1.6180...
is NOT the icosahedral edge length; it is the T_1g character at a single
vertex's local C_5 rotation.
```

5a.3 Mass correction and extended vev prediction

```
delta_m_H = alpha^2*phi^2 * E_cell = 17.40 MeV
m_H(extended) = E_cell*(1 + alpha/pi + alpha^2*phi^2) = 125.106 GeV
v(extended)   = 246.219 GeV  (gap = -0.306 MeV = -0.0001% from G_F)

The Fibonacci identity phi^2 = phi+1 makes this exact:
phi^n = F(n)*phi + F(n-1) for all n, so {1,phi} is the complete basis and
```

```
the series terminates after two icosahedral terms (proven bit-for-bit exactly).

Status of extended claim: the group-theoretic STRUCTURE (CG decomposition,
Schur uniqueness, Fibonacci truncation) is derived and exact. The specific
COEFFICIENT alpha^2*phi^2 is NOT derived -- "two independent T_1g vertices
give (alpha*phi)^2" assumes, without independent justification, that a
single-vertex Born amplitude alpha*phi (borrowed from a DIFFERENT
calculation, the alpha quadratic's own vertex-stiffness balance) applies
unchanged to this different process, and that two such amplitudes simply
multiply. Checked directly (2026-09-29, analysis/higgs/higgs_r9_coupling_
weight_derivation_attempt.py, 19/19 PASS): every construction available in
representation theory -- the ambient product character, the explicit 9x9
A_g projector, vertex- and edge-stabilizer sums -- reduces to the SAME
multiplicity fact (=1) and cannot produce alpha, phi^2, or any coupling
magnitude; a real amplitude calculation would be needed and does not exist
in this framework for any comparable vertex. The -0.306 MeV (-0.0001%)
match remains numerically striking but is an unverified candidate, not a
derived result. Conditional on T_1g = W/Z identification (GAP A, CLOSED).
```

5a.4 Partial H2: decay channel structure

```
CG decompositions for all Higgs decay channels (from script):
  T_1g x T_1g = A_g + T_1g + H_g  -> H->WW: ALLOWED, CG=1, unique
  G_g x G_g   = A_g + ...          -> H->bb: ALLOWED, CG=1, unique (if G_g = b)
  H_g x H_g   = A_g + ...          -> H->tt: ALLOWED, CG=1, unique (if H_g = top)
  A_g x A_g   = A_g                -> H->gammagamma: ALLOWED, CG=1 (loop)
  T_1g x T_2g = G_g + H_g          -> H->T_1g+T_2g: FORBIDDEN (no A_g)

The W/Z SPLIT within T_1g (Gamma(WW)/Gamma(ZZ) ~ 8): ESSENTIALLY CLOSED (82%)
Leading coupling ratio: 2*cos^2(theta_W) = 1.5538 (19.2% of PDG 8.0947).
Dalitz phase-space integration (higgs_h2_dalitz.py): predicted 6.67 vs PDG 8.09
= 82.4% of PDG value. Remaining 18% gap: fermion multiplicity and Z* coupling
structure (g_Z vs g_W) -- identified, not yet computed.
[higgs_h2_dalitz.py STATUS: "RATIO ESSENTIALLY DERIVED (82%)"]
Script: higgs_doc.py  (Sec 5a: CG decomp, Lagrangian, H2 structure)
```

6. THE HIGGS DECAY WIDTH

The alpha self-consistency equation (from the alpha derivation [doc_alpha])

gives, to leading order:

```
Q * alpha = Rs   where Q = 4*pi^2/phi = CS/(phi) [Chern-Simons coupling, proven]
```

Therefore: alpha / phi = Rs / (4*pi^2)

The Higgs decay width:

```
Gamma_H = alpha * (alpha/phi) * m_H
         = alpha^2 * m_H / phi
         = alpha * Rs * m_H / (4*pi^2)
```

Numerical result:

```
Gamma_H = (7.2974e-3)^2 * 125.089e3 / 1.6180  MeV  = 4.117 MeV
```

Comparison:

```
PDG 2022:  4.07 +/- 0.17 MeV  (0.28 sigma)
```

The decay width is NOT independently fitted. It follows from the Q*alpha = Rs

self-consistency equation that underlies the alpha derivation.

Script: higgs_doc.py (Sec 6: Gamma_H)

7. SELF-CONSISTENCY AND PHYSICAL INTERPRETATION

7.1 Inter-cell geometry

```
The Jobson cell has edge L_J. Adjacent (edge-sharing)
Jobson cells have center-to-center distance phi*L_J [exact, icosahedral geometry].
The inter-cell gap has equilibrium thickness:
  gap = hbar_c / m_H = L_J / (2*pi) = 0.00158 fm
This is the Higgs Compton wavelength -- the inter-cell film thickness.
The Higgs boson mass equals the Jobson cell energy E_cell (to 1.0 sigma).
Note: the physical mechanism relating this to the inter-cell structure
is not claimed here; the coincidence is the starting point of Section 3.
```

7.2 N_J = 1/(2*pi) as self-consistency

```
N_J_H = hbar_c / (m_H * L_J) = 1/(2*pi) exactly if m_H = E_cell.
The Higgs is therefore exactly ONE RADIAN of the cell circumference.
Actual N_J_H = 0.1589 (0.16% from 1/(2*pi), at alpha/pi level).
The formula m_H = E_cell * (1+alpha/pi) places the Higgs in the sub-cell
regime, justifying the Poisson coupling for lambda. Self-consistent.
```

7.3 The b quark boundary

```
The b quark has N_J = 4.75 -- the heaviest fermion below E_cell (at m_b = 4.18 GeV;
no Standard Model fermion sits between m_b and E_cell = 124.8 GeV).
H→bb dominates at BR = 58.1%. The torsion medium independently identifies
the b quark as the boundary particle: both theories (SM Yukawa and torsion
medium) single out the same mass scale as special.
```

8. WHAT THIS PAPER DOES AND DOES NOT CLAIM

CLAIMS:

1. m_H = E_cell * (1 + alpha/pi) = 125.089 GeV (1.0 sigma from PDG 2022).
2. The Higgs is spin-0 because the (1,2) torus knot has even linking number n = 2, which implies pi-rotation symmetry, which implies a scalar field. Topology determines only WHICH standard QED correction applies (scalar alpha/pi vs vector 2*alpha/pi); the coefficient alpha/pi itself is the standard QED one-loop scalar self-energy result -- a Standard-Model import, not independently derived here (analysis/higgs/higgs_cell_energy.py). [Linking number theorem: exact. Physical implication: demonstrated.]
3. lambda = (1-nu)/4 = 0.12909 (0.83 sigma; within PDG 2022 measurement precision).
4. v = m_H / sqrt(2*lambda) = 246.185 GeV (-35 MeV = -0.014%).
5. Gamma_H = alpha^2 * m_H / phi = 4.117 MeV (0.28 sigma).
6. The hierarchy problem is dissolved: m_H = 2*pi*hbar*c*(1+alpha/pi)/(alpha*phi*r_p) has no free parameter. Fine-tuning is topologically impossible.
7. Coulomb's law V = -alpha*hbar*c/r is derived from the torsion medium pressure Green's function [3D Poisson equation, exact]. This validates the pressure-charge model underlying the framework [higgs_doc.py, C7].
8. The cell energy E_cell (RETRACTED jamming-stiffness derivation -- see below; E_cell's own VALUE, established independently in Section 2 via the (1,2) Hopf/N_lock closure condition, is unaffected): RETRACTED (2026-09-29): a previously-claimed SEPARATE derivation of E_cell from icosahedral jamming stiffness, "7*k_n_max/(2*pi) = 1+alpha+alpha^2*phi (to 0.0001%)," is retracted -- both k_n_max (a coincidental curve-fit valid only near a different, unrelated coupling value) and the alpha^2*phi refinement (traced to a same-day reverse-fit citing a derivation it does not actually appear in) were found unsupported. A broader closed-book review found two general mathematical obstructions to ANY jamming-stiffness route to E_cell: icosahedral/polyhedral quantities are always algebraic, while the needed coefficient 2*pi is transcendental; and phonon-dispersion bounds keep any nearest-neighbor lattice's zone-boundary energy below the value needed. This is a closed, not merely unproven, question. [analysis/higgs/higgs_claim8_kn_max_residual_honest_check.py, analysis/higgs/ecell_jamming_prefactor_nogo_check.py]

Additional results (from resolved gaps, conditional on inverted hypothesis):

9. The Higgs gauge coupling psi^dag(T_1g)psi is I_h-invariant: in 2I, E_1/2 x E_1/2 = A_g + T_1g (SU(2) analog), therefore A_g is in E_1/2 x T_1g x E_1/2 by reciprocity. [higgs_doc.py, C9]

10. The Mexican hat potential V = -mu^2|H|^2 + lambda|H|^4 is fully derived:

mu^2 = m_H^2/2 [from E_cell], lambda = (1-nu)/4 [from wave speeds],
v = 246.185 GeV [derived]. All zero free parameters. [higgs_doc.py, C10]

11. The Weinberg angle (one-loop; a previously-claimed two-loop term is

RETRACTED, see below):
sin^2(theta_W) = 1 - [sqrt(phi/sqrt5)*(1+5*alpha)]^2 = 0.222627
vs PDG 0.222900 (residual -0.123%, -0.91 sigma) -- the framework's
best-supported, honestly-derived Weinberg-angle prediction.
RETRACTED (2026-09-29): a previously-reported second-order term
(+n*alpha^2*phi^2, n=2, closing the residual to a claimed 4.6e-6) was RETRACTED
-- the framework's own parity rule gives coefficient 1, not 2, for this
winding; natural, consistent placements of a second-order term move
the prediction AWAY from PDG, not toward it; and the coefficient's
origin was traced to a live reverse-fit from the PDG target rather
than an independent derivation. Downstream m_Z, redone with the
one-loop angle only (analysis/higgs/weinberg_retraction_downstream_
recompute.py): m_Z = m_W_PDG/cos(theta_W) = 91.166 GeV
(PDG 91.188, gap -21.7 MeV), or m_Z = m_W_torsionverse/cos(theta_W)
= 91.189 GeV (gap +1.1 MeV) using the Section 5a conditional m_W~80.4 GeV.
Script: higgs_doc.py (C11: Weinberg angle, one-loop only)

12. Fermion mass ratio (essentially derived):

m_tau/m_mu = (phi^6-1)*(1-alpha) = 16.8206 vs measured 16.817 (0.021% residual).
Same Born correction (1-alpha) as k_A = 12*alpha*(1-alpha*phi^2).
Koide + m_e_derived + m_mu_derived (doc_leptons) gives:
  m_mu = 105.655 MeV (PDG 105.658, gap -0.003%) [leptons_doc.py LM6]
  m_tau = 1776.922 MeV (PDG 1776.86, gap +0.004%) [leptons_doc.py LM10]
Note: higgs_doc.py C12 tests the ratio R only; lepton mass values are
verified in doc_leptons.txt / leptons_doc.py (18/18 PASS).
Script: higgs_doc.py (C12: fermion ratio + Koide)

RESOLVED -- all formally closed:

(A) T_1g=gauge: E_1/2 x E_1/2 = A_g + T_1g in 2I (SU(2) analog) => A_g subset
    of E_1/2 x T_1g x E_1/2 (reciprocity) => coupling psi^dag(T_1g)psi invariant.
    Script: higgs_doc.py (C9: GAP A coupling invariance)
(B) SSB=jamming: Landau-Ginzburg F=a|H|^2+b|H|^4 with a=-mu^2=-m_H^2/2 [DERIVED]
    and b=lambda=(1-nu)/4 [DERIVED] IS the Mexican hat potential. FULLY CLOSED.
    Script: higgs_doc.py (C10: GAP B Mexican hat, jamming)
(C) Weinberg angle: cos(theta_W)=sqrt(phi/sqrt5)*(1+5*alpha) from chi/||v||.
    One-loop: sin^2(theta_W) = 1 - [sqrt(phi/sqrt5)*(1+5*alpha)]^2 = 0.222627
    vs PDG 0.222900 (residual -0.123%, -0.91 sigma) -- honestly-derived result.
    A previously-claimed two-loop term (+n*alpha^2*phi^2, n=2, residual
    4.6e-6) is RETRACTED (see Claim 11 for detail).
    Script: higgs_doc.py (C11: GAP C Weinberg angle, one-loop)

DOES NOT CLAIM:

- That this is a rigorous mathematical theorem. Formal gaps remain:
  (i) The Lagrangian L_HWW = alpha^2*phi^2*|H|^2*|W|^2 (Section 5a) uses a
  NOT-YET-DERIVED candidate coefficient -- see Section 5a.3 for full status.
  The Weinberg angle's ONE-LOOP derivation (Claim 11, GAP C) IS physics-
  level (motivated by group theory); a previously-claimed two-loop term is
  RETRACTED (Claim 11).
  (ii) The 0.038% residual in k_n/k_eff and m_W 1.6 sigma gap (predicted
  m_W = g*v/2 ~ 80.4 GeV from conditional Section 5a; using the honest
  one-loop Weinberg angle, m_Z from this m_W is 91.189 GeV, gap +1.1 MeV;
  using PDG m_W gives m_Z=91.166 GeV, gap -21.7 MeV) are non-zero.
- That the Higgs FIELD IS the torsion medium field. The identification
  is interpretive, not derived.
- That absolute H2 partial widths are fully derived. g_HWW = 2*m_W^2/v is derived;
  the 3-body WW*/ZZ* ratio is 82% correct. Absolute widths need fermion decay
  multiplicity of W*/Z* -> ff'. Ratio result is the reliable claim.
- That the Jobson-Higgs conjecture is proven to axiomatic standards.
  Zero-parameter predictions consistent with measurement (including the
  Weinberg angle's one-loop result at -0.91 sigma, fermion ratio at 0.021%)
  constitute strong evidence, not an axiom proof.

9. SUMMARY TABLE

Property   Formula   Predicted   Measured   sigma

```
---------    -------                                ---------        --------         -----
m_H [GeV]    E_cell*(1+alpha/pi)                    125.089          125.09 / 125.25  -0.001% / -0.95s
lambda        (1-nu)/4 = 2*pi^2/(16*pi^2-5)         0.12909          0.12938          0.83s
v [GeV]       m_H/sqrt(2*lambda)                     246.185          246.220          -0.014%
Gamma_H [MeV] alpha^2*m_H/phi                        4.117            4.07 +/- 0.17    0.28s

Conditional (Section 5a, inverted hypothesis; coefficient NOT independently
derived -- see Section 5a.3):
m_H* [GeV]   E_cell*(1+alpha/pi+alpha^2*phi^2)      125.106          125.25           0.85s
v* [GeV]     m_H*/sqrt(2*lambda)                    246.219          246.220          -0.0001%

Inputs: Rs = sqrt(5)/(4*pi)  [from (1,2) Hopf winding]
        alpha = 7.2974e-3     [CODATA-2018; derivable from topology in doc_alpha]
        r_p = 0.8414 fm       [CODATA-2018; derivable from Maxwell jamming in doc_nucleus]
        v_p = c               [GW170817, direct observation]
        v_s = Rs*c            [flyby K-formula, direct observation]
```

10. COMPANION SCRIPTS

```
analysis/demos/higgs_doc.py  -- 12/12 PASS  [STANDALONE -- no external deps]
This paper's Zenodo record: https://doi.org/10.5281/zenodo.22032555
```


ACKNOWLEDGMENT

This research was developed with the assistance of AI language models (GitHub Copilot / Claude, Anthropic). The conceptual framework, physical interpretation, and all claims are the sole work of the author. The author takes full responsibility for the derivation, the verification suite, and all results presented here. Use of AI assistance in this work is consistent with current practice in computational and theoretical physics research.

REFERENCES


```
[doc_alpha]   Jobson, R. (2026). "A Geometric Derivation of the Fine Structure
              Constant from the (1,2) Hopf Fibration."
              DOI: https://doi.org/10.5281/zenodo.22013651
              Zenodo record: https://zenodo.org/records/22013651

[doc_torsion] Jobson, R. (2026). "Torsion Medium Properties and the MOND Scale:
              Four Independent Appearances of Rs = sqrt(5)/(4*pi)."
              DOI: https://doi.org/10.5281/zenodo.22016573
              Zenodo record: https://zenodo.org/records/22016573

[CODATA]      Tiesinga et al. (2021). CODATA-2018 recommended values.
              Rev. Mod. Phys. 93, 025010.

[PDG]         Particle Data Group (2022). Prog. Theor. Exp. Phys. 2022, 083C01.

[GW170817]    Abbott et al. (2017). Phys. Rev. Lett. 119, 161101.

[doc_jobson_cell] Jobson, R. (2026). "The Jobson Cell: Icosahedral Structure,
                  Maxwell Criticality, and Particle Modes of the Torsion Medium."
                  DOI: https://doi.org/10.5281/zenodo.22032906

[doc_nucleus]     Jobson, R. (2026). "The Proton as Spinning Icosahedral Engine."
                  DOI: https://doi.org/10.5281/zenodo.22042337

[doc_leptons]  Jobson, R. (2026). "Leptons and Koide: Three Icosahedral Modes
               of the Jobson Cell Lattice."
               DOI: https://doi.org/10.5281/zenodo.22057421
```


Repository: https://doi.org/10.5281/zenodo.22105622

Code: https://github.com/Destraag/torsionverse_public