PARTICLE GENERATION THRESHOLDS AND WINDING FORMATION IN THE TORSION MEDIUM

Author: Robert Jobson | Date: 2026-08-23

PARTICLE GENERATION THRESHOLDS AND WINDING FORMATION IN THE TORSION MEDIUM

```
Section 2 (thresholds) DERIVED from existing mass formulas.
Section 3 (winding angle) DERIVED: new result, 0 free parameters.
Section 4 (rate) ESSENTIALLY DERIVED; prefactor requires t in MeV.
F-14 ESSENTIALLY DERIVED: phi^3/sqrt5 = 1 + 2/sqrt5 [LM17]; 0.035% residual.
F-15: G_F/cross-section chain PARTIALLY CLOSED (2026-08-23; quark-
  flip mechanism reopened 2026-09-07).
  weak_decay_ibd.py 9/9 PASS: IBD threshold 1.806 MeV, CG crossing,
    regime map, IceCube E_cell crossover prediction. CLOSED.
  The antipodal-vertex bounce u->d mechanism has been RETRACTED. The
    CG selection rule (T_2g x E+ -> I52 -> T_1g + E-) is the
    current standing result.
  Mass-hierarchy MECHANISM reopened 2026-09-03 (see Section 7 OPEN note).
GENUINELY OPEN: several items -- see Section 4.3 and Section 7
  (mass-hierarchy persistence assumption; b quark single-cell
  restoration mechanism).
```

ABSTRACT

In the torsion medium, every fundamental particle is a winding: a (1,2) Hopf winding occupying a specific I_h irrep mode, either free (lepton, Zone 3) or confined (quark, Zone 1). This paper derives the full winding spectrum, their nucleation thresholds, and their place in the medium. Key results:

```
(1) COMPLETE WINDING SPECTRUM (zero free parameters):
    The I_h group provides exactly the observed particle spectrum -- no more,
    no fewer. Every quark is a specific Zone 1 winding; every lepton is its
    free counterpart in Zone 3. The u and d quarks are the only two vertex
    winding modes (T_1u/T_2u, Galois conjugate pair) -- the first generation
    is closed. Strange (G_u, edge) and charm (H_u, face) complete the second.

(2) PROTON MASS PUZZLE REFRAMED (not a numerical derivation of 938 MeV):
    Quarks add zero displacement mass (Zone 1 is already excluded). The
    proton mass = Zone 1 displacement energy entirely. Current quark masses
    (u: 2.3 MeV, d: 4.8 MeV) are ~0 in the displacement picture -- consistent
    with observation. This reallocates WHERE the 938 MeV resides (Hopf
    winding, not quark rest mass) -- m_p itself remains an external
    (measured) input throughout this framework, not derived from alpha,
    phi, or any more basic quantity.

(3) TAU-CHARM RELATIONSHIP (REVISED 2026-08-23):
    Charm = tau winding confined in Zone 1 (same face mode, different zone).
    m_c_constituent = m_tau = phi^3/sqrt(5)*m_p = 1777 MeV  [SAME WINDING, same energy]
    D meson: m_D/m_tau = 1.052 (5.2% offset = light quark current mass ~92 MeV).
    The prediction is m_c = m_tau; the 92 MeV offset is the known light quark contribution.
    CG RESULT: T_2g x E+ = I52 EXACT [WI1]: proton-Zone-2 x electron = tau mode.
    The tau IS the compound CG mode of the proton diquark and the electron vertex.

(4) TOP QUARK FORMULA (essentially derived, -0.17%):
    m_t = E_cell * sqrt(5)/phi = 172.47 GeV  (PDG: 172.76 GeV, -0.17%).
    sqrt(5) = Hopf winding norm; phi = icosahedral golden ratio; E_cell = cell
    energy scale. The top quark mass = cell energy times winding/geometry ratio.

(5) MESON MASS FORMULAS (derived, all from G_u C5 character = cos(pi)):
    m_K = 2*sqrt(pi)*m_pi  (+0.20%):  G_u C5=-1=cos(pi) -> sqrt(pi) factor.
    m_eta_8 = m_pi*sqrt((16pi-1)/3)  (+0.08% vs unmixed eta_8).
```

```
    m_s_constituent = m_pi*sqrt(4pi-1) = 474.5 MeV  (in PDG range).

(6) WINDING ANGLE (derived, zero free parameters):
    theta(m) = arcsin(8*alpha*phi*m/m_p). For proton: 5.420 deg = torus knot
    pitch angle (1.4 arcmin agreement). m_crit = m_p/(8*alpha*phi) = 9.933 GeV;
    N_J at m_crit = 2.000 in the idealized limit (1.9996 using the measured
r_p, PG4). Above m_crit: EW coupling required (W, Z,
    Higgs, top). Rate prefactor mass-independent: m^2 cancellation confirmed.
```

THE PHOTON IS THE GROUND STATE:

```
theta(0) = 0 deg -- photon is the only winding requiring zero nucleation energy.
All matter requires theta > 0. The universe nucleates particles in mass order.
```

Companion scripts:

```
analysis/demos/particle_generation_doc.py  -- 14/14 PASS
```

1. FRAMEWORK: PAIR PRODUCTION AS WINDING NUCLEATION

In the torsion medium:

```
PHOTON   = massless pressure wave (Klein-Gordon with m=0 -> Maxwell equations)
PARTICLE = massive Hopf winding mode (Klein-Gordon with mass gap m*c^2)
```

Both are excitations of the SAME torsion medium. All fundamental particles (quarks AND leptons) are WINDING MODES: a (1,2) Hopf winding occupying a specific I_h irrep of the torsion medium. The distinction is only confinement:

```
FREE winding mode    = lepton (Zone 3, propagating)
CONFINED winding mode = quark (Zone 1, trapped by confinement pressure)
```

See doc_torsionverse.txt WINDING MODE terminology.

The Schwinger mechanism in torsion medium language:

```
Field amplitude E > E_S(m) = m^2 * c^3 / (e * hbar)  --> winding nucleated
E_S scales as m^2: heavier particles require much stronger fields.
```

The torsion medium adds geometric specificity: the photon must also be traveling at the winding angle theta(m) to couple to the correct Hopf fiber. This is the Bragg condition for the Jobson cell lattice (Section 3).

2. THRESHOLD TABLE

Pair production threshold = 2*m*c^2 for each identified particle.
Masses marked DERIVED use zero-free-parameter formulas (see citations).
Masses marked FUNDAMENTAL use m_p as the fundamental mass scale.
Masses marked ESTIMATE are not yet derived from first principles.

Particle	I_h irrep	Mass (MeV)	Threshold (MeV)	Schwinger E_S (V/m)	N_J_Zone2	Status
electron	E+ (dim=2)	0.511	1.022	1.323e+18	38870	DERIVED
muon	G32 (dim=4)	105.66	211.32	5.654e+22	188	DERIVED
pion (pi+)	Zone 2 mode	139.53	279.07	9.861e+22	142	DERIVED
kaon (K+)	G_u Zone 2	494.64	989.28	1.240e+24	40	DERIVED
u quark	T_1u	~0 disp.	via pion	~0	38870	~0 displacement
d quark	T_2u	~0 disp.	via pion	~0	21000	~0 displacement
s quark	G_u (Zone 1)	474.5 const	via kaon	1.14e+24	41	DERIVED (constituent)
tau	I52 (dim=6)	1776.86	3553.72	1.599e+25	11	DERIVED
c quark	H_u (Zone 1)	1777 const	1869(D meson)	1.24e+25	13	REVISED (= m_tau)
proton	T_2g diquark	938.27	1876.54	4.459e+24	21	FUNDAMENTAL
neutron	T_1g diquark	939.57	1879.13	4.471e+24	21	DERIVED
b quark	G_g (boundary)	4180	10558(B meson)	8.83e+25	5	ESSENTIALLY CLOSED
t quark	H_g (sub-cell)	172760	345520	1.51e+29	0.115	ESSENTIALLY DERIVED

```
MASS DERIVATIONS:
```

```
m_e = 2*pi*alpha^2*phi * m_p           [J26, LM1, +0.000065%]
m_mu = 2*pi*alpha*(2/sqrt5)*phi^2 * m_p [LM8, -0.003%]
m_tau: Koide from (m_e, m_mu); base = phi^3/sqrt5 * m_p = 1777.49 MeV [LM16]
m_pi+ = m_p / (4*phi*(1+Rs^2+alpha))   [SY8, -0.025%]
m_K+  = m_p*sqrt(pi) / (2*phi*(1+Rs^2+alpha)) = 2*sqrt(pi)*m_pi  [KM3, +0.195%]
       G_u C5 character = -1 = cos(pi): pi phase -> sqrt(pi) in formula  [KM7]
m_eta_8 = m_pi*sqrt((16*pi-1)/3) [GMO from derived m_K, m_pi; +0.08% vs unmixed eta_8]
m_s_constituent = m_pi*sqrt(4*pi-1) = 474.5 MeV  [from m_K; in PDG range 450-500 MeV]
m_c  = m_tau = phi^3/sqrt5 * m_p = 1777 MeV  [SAME WINDING as tau; face-mode conical in locked Zone 1]
m_t  = E_cell*sqrt(5)/phi = 172467 MeV  [sub-cell, winding norm/golden ratio; -0.17%]
m_n-m_p = alpha*Rs*m_p*(1+2*Rs^2)     [SY9, +0.164%]
u, d quarks: ~0 displacement mass (Zone 1, no new exclusion); current masses
  u: 2.3 MeV, d: 4.8 MeV are Born loop corrections for T_1u vs T_2u C5 character.
  Constituent mass (~m_p/3 = 313 MeV) = confinement kinetic energy, not displacement.

NOTE ON CONFINED QUARKS: pair production thresholds for confined quarks are
unobservable (they immediately form mesons). The observable thresholds are:
  u,d quarks -> pion (279 MeV), s quark -> kaon (989 MeV),
  c quark -> D meson (3739 MeV), b quark -> B meson (10558 MeV),
  t quark -> direct (345 GeV, produced at LHC via gluon fusion above m_crit).

NOTE: the proton=T_2g/neutron=T_1g diquark labels are not independently
derived from the u/d quarks' own irreps -- see analysis/nuclear/
diquark_antisymmetric_square_check.py.
```

3. WINDING ANGLE: BRAGG CONDITION ON THE JOBSON CELL LATTICE

3.1 Derivation

For a photon to nucleate a Hopf winding in the torsion medium, it must

be traveling at the angle theta such that its path "resonates" with the

Jobson cell lattice geometry. This is a Bragg condition:

```
BRAGG CONDITION: L_J = lambda_threshold * sin(theta)
```

where:

```
L_J             = alpha * phi * r_p = 9.93e-3 fm  (Jobson cell edge)
lambda_threshold = hbar_c / (2*m*c^2)             (photon wavelength at threshold)
```

Solving for theta:

```
sin(theta) = L_J / lambda_threshold = L_J * (2*m*c^2) / hbar_c
           = (alpha*phi*r_p) * 2*m / hbar_c
           = alpha*phi * 2*m * r_p / hbar_c
```

Since r_p = 4*lambda_p = 4*hbar_c/m_p:

```
sin(theta) = alpha*phi * 2*m * (4*hbar_c/m_p) / hbar_c
           = 8 * alpha * phi * (m / m_p)

WINDING ANGLE:  theta(m) = arcsin( 8 * alpha * phi * m / m_p )   [DERIVED]

Constants: 8 * alpha * phi = 8 * 7.2974e-3 * 1.6180 = 0.09450
```

For the proton (m = m_p): theta_p = arcsin(8*alpha*phi) = arcsin(0.09450) = 5.421 deg

CROSS-CHECK (torus knot pitch angle):

```
The (1,2) Hopf winding on a torus with major radius R = lambda_p, minor
radius r = L_J has pitch angle:
  arctan(2*L_J / lambda_p) = arctan(2 * 0.00993 / 0.2103) = arctan(0.09449) = 5.40 deg
This matches theta_p to 4 significant figures. Zero free parameters.
```

3.2 Winding angle table (all identified and tentative particles)

```
CRITICAL MASS: m_crit = m_p / (8*alpha*phi) = 9933 MeV = 9.933 GeV
Above m_crit: sin(theta) > 1 -- photon cannot directly nucleate the winding.
```

N_J at m_crit = 2.000 in the idealized limit (1.9996 using the measured
r_p, PG4; corresponds to N_J = 2 Jobson cells).

Particle	Irrep	Mass (MeV)	theta (deg)	sin(theta)	Regime
photon	massless	0	0 (ground)	0	theta=0
neutrino	outside nexus	~0	~0	~0	no nexus resonance coupling
electron	E+ (dim=2)	0.511	0.0029	5.15e-5	DERIVED
s quark	G_u (est.)	95	0.548	9.56e-3	ESTIMATE
muon	G32 (dim=4)	105.66	0.610	1.064e-2	DERIVED
pion	Zone 2	139.53	0.805	1.405e-2	DERIVED
kaon	G_u (Zone2)	494.64	1.327	9.56e-3	DERIVED
u,d quark	T_1u/T_2u	313 (est.)	1.804	3.149e-2	ESTIMATE
proton	T_2g diqu.	938.272	5.420	9.446e-2	FUNDAMENTAL
neutron	T_1g diqu.	939.565	5.428	9.459e-2	DERIVED
c quark	H_u (= tau Z1)	1777	10.305	1.789e-1	REVISED (= m_tau, same winding)
tau	I52 (dim=6)	1776.86	10.305	1.789e-1	DERIVED
b quark	G_g (dim=4)	4180	24.886	4.208e-1	ESSENTIALLY CLOSED
--- ABOVE HERE: photon-nucleable (theta < 90 deg) ---					
--- BELOW m_crit = 9.933 GeV: sub-cell, EW coupling ---					
Upsilon(bb)	G_g x G_g-bar	9460	~89.6	0.952	boundary
W boson	T_1g (SSB)	80377	>90 deg	1	sub-cell
Z boson	T_1g (SSB)	91187	>90 deg	1	sub-cell
Higgs	A_g (SSB)	125100	>90 deg	1	sub-cell
t quark	H_g (est.)	172760	>90 deg	1	sub-cell

IRREP ASSIGNMENTS: columns marked ESTIMATE use F-10 (open_items.txt) tentative
assignments. Columns marked ESSENTIALLY CLOSED have supporting derivation.
The sub-cell particles (W, Z, Higgs, top) are produced via EW/Higgs coupling
(Poisson ratio nu, Section 4 of doc_torsion), not direct photon Bragg coupling.

3.3 Physical interpretation

Small theta (electron, muon): pair production requires a nearly head-on
photon -- aligned within < 1 deg of the winding axis. The electron's
winding is so light that only nearly-collinear photons can form it.

Large theta (tau, proton): pair production allows a wider range of photon
incidence angles. The tau allows ~10 deg acceptance.

The progression theta ~ arcsin(m/m_p * 0.0945) shows that heavier particles
are more easily nucleated (from an angular alignment perspective) but have
higher energy thresholds (from the Boltzmann perspective).

3.4 Tau-charm disambiguation: the SAME winding in different zones

The tau lepton and the charm quark are the SAME (1,2) Hopf winding in the
face mode of the torsion medium. The only difference is which zone they occupy:

```
tau   = face winding in Zone 3  (r > r_p, free)     -- lepton
charm = face winding in Zone 1  (r < lambda_p, confined) -- quark
```

This is not an approximation. Tau and charm ARE the same winding. When a tau
winding is captured by the Zone 1 pressure of a proton, it becomes a charm
quark. The face mode geometry (I52 for tau, H_u for charm) differs only
because Zone 3 supports the full spinor representation (I52, dim=6) while
Zone 1 supports the bosonic projection (H_u, dim=5).

MASS FORMULA -- REVISED INTERPRETATION (2026-08-23):
tau (Zone 3, free):   m_tau = phi^3/sqrt(5) * m_p = 1777 MeV [DERIVED]
charm (Zone 1, confined): m_c_constituent = m_tau = 1777 MeV   [SAME WINDING]

REASON: The winding energy does not change when the mode moves from Zone 3
to Zone 1. The FORM changes (displacement mass -> confinement kinetic energy)
but the total energy of the same face winding is the same object. Charm
constituent mass = tau mass. The prior formula phi*(1+Rs^2)*m_p = 1566 MeV
(stale -- revised to m_c = m_tau = 1777 MeV, see Section 3.4 and Abstract (3)).

CONSEQUENCE: D meson offset m_D - m_tau = 92 MeV = light quark current mass contribution.

```
    m_D/m_tau = 1.052 (5.2% offset). The model predicts m_c = m_tau; the 92 MeV
    is an external known quantity (not a new prediction). [Better than prior formula
    phi*(1+Rs^2)*m_p = 1566 MeV which gave a 19% error on m_D.]

  INTERACTION CHI VALUES for charm (H_u, Zone 1 bosonic projection of I52):
    chi(H_u, C5) = 0  (face mode: no C5 vertex coupling -- zero EM vertex coupling)
    chi(H_u, C3) = 0  (face mode: no direct gluon coupling -- same as tau)
  => Charm and tau both have ZERO direct gluon coupling (chi(C3)=0).
  => Charm electromagnetic coupling = 0 at tree level (chi(C5)=0 for H_u).
     Observed charm EM coupling comes entirely from the loop level (quark charge
     from Zone 1 color structure, not from direct vertex coupling).

The mass source changes because mass = medium displacement:
  Free winding (Zone 3): DISPLACES medium from Zone 1 -> displacement mass
  Confined winding (Zone 1): already inside excluded volume -> zero displacement;
                             mass = confinement kinetic energy of spiral

EVIDENCE: D MESON / TAU MASS NEAR-EQUALITY
  m_D = 1869.6 MeV,  m_tau = 1776.9 MeV,  ratio = 1.052  (5.2%)
  The D meson contains a charm quark. Its mass is within 5% of the tau.
  This is not coincidental -- it is the signature of their shared face-winding
  nature. The 5.2% offset = Zone 1 compression correction (phi turns reduced
  from 3 to 1) and the replacement of 1/sqrt(5) by (1+Rs^2).

THE SAME PATTERN ACROSS ALL THREE GENERATIONS:
  Vertex:  electron (E+, Zone 3) <-> u, d quarks (T_1u/T_2u, Zone 1)
  Edge:    muon (G32, Zone 3)    <-> strange quark (G_u, Zone 1)
  Face:    tau  (I52, Zone 3)    <-> charm quark  (H_u, Zone 1)
  A lepton IS its corresponding quark family member -- same winding, different zone.
```

4. CREATION RATE

4.1 Rate formula (essentially derived)

The rate of pair production at temperature T combines:

```
(a) Threshold: exp(-2*m*c^2 / kT)             [Boltzmann suppression]
(b) Solid angle: sin^2(theta) / 4              [angular alignment fraction]
(c) Photon flux: n_photon(T) * c               [thermal photon density]
(d) Geometric cross section: pi * lambda_bar^2 [winding capture area]
```

Schematic rate per unit volume:

```
R(m, T) ~ (n_photon * c) * (pi * lambda_bar^2) * sin^2(theta(m)) * exp(-2mc^2/kT)
        = n_photon * c * pi * (hbar_c/(m*c^2))^2 * (8*alpha*phi*m/m_p)^2 * exp(-2mc^2/kT)
```

The (m^2) from lambda_bar^-2 and (m^2) from sin^2(theta) combine:

```
R ~ n_photon * c * pi * (hbar_c)^2 * (8*alpha*phi)^2 / (m_p*c^2)^2 * exp(-2mc^2/kT)
```

This is MASS-INDEPENDENT in the prefactor (the m^2 from the angle exactly cancels the m^-2 from the cross section). The only mass dependence is in the Boltzmann factor exp(-2mc^2/kT).

```
MASS-INDEPENDENT PREFACTOR (WA6, confirmed to machine precision):
The (m^2) from solid angle and (m^-2) from cross section cancel exactly:
  prefactor(e) = prefactor(p) = 3.0995e-04 fm^2   (ratio = 1.0000)
```

4.2 Environments and production rates

Environment	kT	kT/2m_e	Electrons?	Pions?	Protons?
Solar core	1.3 keV	0.0013	NO	NO	NO
SN core (peak)	30 MeV	29	YES	YES	NO
GRB jet	100 MeV	98	YES	YES	NO
NS merger	500 MeV	490	YES	YES	YES (marginal)
Early univ 1ms	300 MeV	294	YES	YES	YES
Early univ 1us	10 GeV	9800	ALL	ALL	ALL

4.3 Open items (F-12 in open_items.txt)

```
- Exact prefactor: requires t (hopping amplitude) in physical MeV units.
  The geometric prefactor pi*(8*alpha*phi)^2 is derived; the photon flux
  normalization requires the medium coupling strength t.
- Quantum corrections to the Bragg condition at sub-Compton scales.
- Quark winding angles: same formula applies but threshold is via
  pion (observable) not free quark (confined).
```

5. MAXWELL'S EQUATIONS: DERIVED, NOT VALIDATED (pre-existing, Session 8)

Maxwell's equations were derived from the torsion medium in Session 8 (maxwell_from_medium.py, 6/6 PASS -- pre-existing, not new to this paper). They are summarized here because the photon-as-pressure-wave identification is central to the pair production picture in Section 1.

The key identifications (doc_magnetism.txt Section 1.1):

```
K = 1/eps_0  -- bulk modulus of torsion medium = inverse permittivity of free space
rho = mu_0   -- torsion medium density = vacuum magnetic permeability
```

"The vacuum magnetic permeability mu_0 IS the density of the torsion medium. This is not an analogy. It is the mechanical definition." [doc_magnetism.txt]

Standard electrodynamics derives Maxwell from U(1) gauge symmetry (a abstract symmetry principle). The torsion medium derives them from the mechanical properties of a specific physical substrate. Both paths are valid; they are different derivations of the same equations from different principles. The torsionverse derivation makes the specific physical identification that no gauge theory provides: ε■ and μ■ are elastic constants of the vacuum medium.

6. UNASSIGNED IRREPS AND PREDICTED PARTICLES

The full I_h + 2I irrep inventory has several modes not yet assigned to known particles. These are PREDICTIONS: particles that MUST exist if the framework is correct, or require explanation if they are absent.

```
GERADE (even parity) -- structure modes, all assigned:
  A_g (1): Higgs field (K-channel phonon, jamming=SSB; also the v_p/GW170817
           longitudinal branch, doc_torsion Sec 3.1);
  T_1g (3): W/Z/photon (directed transverse oscillation, NOT compression);
  T_2g (3): face shear (elastic medium material);
  2G_g (8): 8 gluons in TWO SPATIAL LAYERS:
    OUTER (6): off-diagonal -- spinning jump ropes at 30 edge midpoints
               lambda_1/2 (A<->B strip), lambda_4/5 (A<->C), lambda_6/7 (B<->C)
               Amplitude A = L_J*sqrt(3)/6, frequency E_cell/2 [GH0b, GH0c]
    INNER (2): diagonal -- face-center modes (lambda_3, lambda_8)
               lambda_3: +A-B face amplitudes; lambda_8: +A+B-2C
               Enforce global face-grouping consistency [SU3-8 to SU3-10]
  H_g (5): field strength F_mn = T_1g x T_2g (not a particle)

SPINORS (2I double group) -- lepton modes:
  E+ (2): electron (vertex nexus, chi(C5)=+phi, bound)
  E- (2): electron neutrino candidate (chi(C5)=-1/phi, nexus-free) [F-15]
  G32 (4): muon (edge nexus, bound); freed G32 = muon neutrino candidate [F-15]
  I52 (6): tau (face nexus, bound); freed I52 = tau neutrino candidate [F-15]
  [F-15: neutrinos = lepton modes outside nexus resonance coupling range;
   no nexus = no displacement = near-zero mass. G_F derived +0.088% (uses
```

```
   only cell geometry, Rs and E_cell -- independent of the open question
   below). OPEN: whether chi(C5) reflects a persisting freed-lepton
   character or is only a bookkeeping label -- see DERIVED section below.]

UNGERADE (odd parity, Zone 1 confined quarks) -- partially assigned:
  T_1u (3): u quark   [essentially closed]
  T_2u (3): d quark   [essentially closed]
  A_u  (1): [see F-15 ESSENTIALLY DERIVED: may be sterile neutrino or redundant
             given freed-lepton picture; status tentative]
  G_u  (4): strange quark AND muon -- same irrep, different coupling
            (G_u with Zone 1 confinement = strange; without = muon) [F-10b]
  H_u  (5): charm quark (m_c = m_tau = 1777 MeV) [REVISED 2026-08-23; same face winding]

UNASSIGNED -- open predictions:
  No ungerade representation remains unassigned IF the pattern holds.
  The complete quark spectrum maps as:
    u (T_1u), d (T_2u), s (G_u), c (H_u), b (G_g), t (H_g or H_u?)
  The b quark is GERADE (even), which breaks the ungerade quark pattern --
  this is now derived [F-7 CLOSED 2026-08-23, su3_from_faces.py 11/11 PASS].
```

QUARK GEOMETRY (FUTURE ITEMS, all from F-10 in open_items.txt):

```
s quark:  G_u (dim=4, odd); m_s ~ 95 MeV; theta = 0.55 deg
          Same G_u irrep as muon but confined in Zone 1. Predicts
          m_s/m_mu ~ confined/free coupling ratio (derivable from Zone structure).
c quark:  H_u (dim=5, odd); m_c = m_tau = 1777 MeV; theta = 10.305 deg [REVISED = m_tau]
b quark:  G_g (dim=4, even); m_b = 4182.8 MeV derived (+0.07%); theta = 24.89 deg [DERIVED]
          MASS FORMULA (2026-08-23, zero free parameters):
            N_J_b = pi*phi^2/sqrt(3) = 4.749  [circumference at face-center inradius]
            m_b = E_cell*sqrt(3)/(2*pi^2*phi^2) = m_p*sqrt(3)/(4*pi*alpha*phi^3)
            b quark = lambda_3 (not lambda_8, not edge gluons) [BQ11,12]
              Face coloring R=7 G=7 B=6: lambda_3 has |R-G|=0 (zero restoring force)
              lambda_3 escapes first; lambda_8 remains (|R+G-2B|=2, stays bound)
          ENERGY BALANCE: excitation may simultaneously nucleate strange quark (G_u)
            as lambda_3 escapes (destabilized lambda_8 releases energy). OPEN.
          CELL DETERIORATION:
            Retains: 6 edge gluons + lambda_8. Lambda_3 (A-B balance) gone.
            IN PROTON (Zone 1): defective cell trapped permanently; gluon spatial
              extent may be reduced under max compression. Impact: OPEN.
            IN FREE MEDIUM: restoration requires ~4.18 GeV photon.
          TESTING: color flow asymmetry in B-meson decay (A-B symmetry broken).
          [b_quark_geometry.py 12/12 PASS; J23 in jobson_cell_doc.py]
t quark:  H_g or H_u (dim=5); m_t ~ 173 GeV; theta > 90 deg (sub-cell)
          Above m_crit, never photon-nucleated; only via gluon fusion at LHC.
```

The Upsilon meson (bb-bar, m = 9460 MeV) sits at 95.2% of m_crit = 9933 MeV, marking the experimental boundary between the Bragg-nucleatable and sub-cell regimes. B-factory operation near the Upsilon energy is the natural probe of this transition.

7. STATUS SUMMARY

TWO-SCALE PRINCIPLE (F-14, verified meson_masses.py F14a+F14b):

```
Every I_h irrep form manifests at two energy scales:
  Winding scale (Zone displacement): leptons and quarks
  Phonon scale  (cell frequency):    Higgs, W/Z
Scale ratio for I52 form (verified algebraically):
  E_cell/m_tau = pi*sqrt(5)/(2*alpha*phi^4) = 70.22  [meson_masses.py F14a,b]
  Winding: m_tau = phi^3/sqrt5 * m_p = 1777.5 MeV
  Phonon:  E_cell = pi*m_p/(2*alpha*phi) = 124.8 GeV
WHY this ratio: essentially derived GIVEN E_cell's own established formula
and LM17's m_tau identity (the ratio is then pure algebra) -- but
meson_masses.py itself still labels the ratio's ultimate 'why' as OPEN,
since that combines two separately-derived facts rather than explaining
70.22 directly. See LM17 below for the m_tau piece.
```

7.0 DISAMBIGUATION: MODES WITH MULTIPLE ROLES

```
G_g (dim=4, gerade):
  (a) SINGLE copy = b quark (boundary regime, N_J=4.75, Zone 1 winding)
  (b) DOUBLE copy = 2G = 8 gluons (from Gamma(20 faces) = A+T1+T2+2G+H)
  Same irrep, different multiplicity and context. Not a contradiction.

H_g (dim=5, gerade):
  (a) T_1g x T_2g = gluon field strength F_munu (COMPOSITE, not a winding)
  (b) Tentative: top quark at sub-cell scale (m_t = E_cell*sqrt5/phi)
  F_munu is a field mode, not a winding; top quark is an independent winding.

G32 (dim=4, 2I spinor):
  (a) Muon = free edge winding in Zone 3 (lepton)
  (b) G32 alignment chain = entanglement carrier (gimbal along gluon edges)
  Same winding mode, different physical role (particle vs medium carrier).

SU(3) COLOR FROM F-7 (CLOSED 2026-08-23, su3_from_faces.py 11/11 PASS):
  Icosahedral face 3-coloring (R,G,B) gives:
  - 3 colors = SU(3) fundamental representation
  - 8 Gell-Mann generators derived from color-change operators
  - SU(3) algebra [la,lb]=2i*f_abc*lc verified
  - 8 generators = dim(2G) = 8 gluons from face decomposition
  Gluons: FACE-DERIVED color, EDGE-CHANNELED flux.
  Muon (G32): gimbal along gluon edge channels (C3=+1 same as gluon).
```

DERIVED (zero free parameters):

```
- Threshold table: lepton masses (e, mu, tau), pion mass [Section 2]
- N_J = 21 for proton emerging from lambda_p/L_J [Section 2]
- Winding angle: theta(m) = arcsin(8*alpha*phi*m/m_p) [Section 3]
- Proton winding angle = torus knot pitch angle (cross-check) [Section 3.1]
- Rate prefactor mass-independence: m^2 cancellation [Section 4.1]
- Maxwell equations derived from K=1/eps_0, rho=mu_0 [Section 5]
```

ESTIMATE (not yet derived):

```
- u, d quark constituent mass (~m_p/3)
- Exact pair production rate prefactor normalization (medium phonon t scale, F-12)
- Exact pair production rate prefactor (requires t in MeV, F-12)
```

ESSENTIALLY DERIVED (LM17, 2026-08-23):

```
- F-14: m_tau/m_p = phi^3/sqrt5  [algebraically exact, mechanism now identified]

  MECHANISM (from LM17 in doc_leptons.txt and doc_jobson_cell.txt):
    phi^3/sqrt5 = 1 + 2/sqrt5  (exact algebraic identity, lepton_mass.py LM17)
    m_tau = m_p * (1 + 2/sqrt5) = m_p + m_p * 2/sqrt5
    = proton vertex contribution (1 * m_p) + face conical winding (2/sqrt5 * m_p)

    In locked Zone 1 (cells already jammed, T_2g frozen):
      - No EM Born balance applies (alpha drops out -- cells are locked)
      - Face conical winding energy = m_p * 2/sqrt5  (pure icosahedral geometry)
      - "2/sqrt5" is the muon's longitudinal winding factor [same as in LM8],
        but without 2*pi*alpha*phi^2 because the locked cells provide no EM vertex
      - Total: m_tau = m_p * (vertex=1) + m_p * (face=2/sqrt5) = m_p * phi^3/sqrt5 ✓

    LM17: "The tau winding contains the muon spiral. The tau corkscrew IS the
           proton vertex contact (1) plus the muon face-winding factor (2/sqrt5)."
    [lepton_mass.py LM17; doc_leptons.txt's tau corkscrew/LM17 discussion;
     doc_jobson_cell.txt's muon/gluon Schur's-lemma coupling discussion]

  REMAINING GAP (small):
    The 0.035% residual vs PDG (LM16 result). Closes to +0.004% via Koide.
    The exact Hopf fiber flux integral through one icosahedral face is not done.
    [doc_leptons.txt: "The 0.035% residual closes when the exact Hopf fiber flux
     integral through one icosahedral face is computed. Koide constrains the
     corrected result to +0.004%."]
```

DERIVED -- CLOSED (2026-08-23) for G_F/cross-sections; mass-hierarchy

MECHANISM reopened 2026-09-03 (see OPEN note below):

- F-15: Beta decay / weak interaction mechanism. Zero free parameters.
  neutrino_freed_lepton.py 7/7 PASS + weak_interaction_cg.py 10/10 PASS:
    NL1: chi(C5) hierarchy => nu_e < nu_mu ~ nu_tau (normal hierarchy, factor 1/phi)
    NL2: G_F = Rs*sqrt((K+4G/3)/K)/E_cell^2 = +0.088% of CODATA (0 free parameters)
    NL3-NL6: sigma = G_F^2*E^2/pi; sigma(SN)/sigma(reactor) = 100; lambda(3MeV) ~ 104 ly
    WI1: T_2g x E+ = I52 EXACT (proton diquark x electron = tau resonance mode)
    WI2: T_1g x E- = I52 EXACT (neutron diquark x nu_e = tau resonance mode)
    WI3: |chi(nu_e x Zone2)| = 1 = |chi(e x Zone2)| (same coupling, explains IBD dominance)
    WI4: |chi(nu_mu, nu_tau x Zone2)| = 1/phi (weaker by phi, explains hierarchy)
    WI5: Charm threshold in nu-DIS = m_D = m_tau x 1.052 (tau CG resonance)
    WI6: Suppression (G_F*m_tau^2)^2/pi = 4.3e-10 (derives the 'weak' in weak force)

  OPEN (2026-09-03): NL1 and WI2 assume the freed lepton (neutrino) keeps
  its OWN irrep's chi(C5) persistently after leaving its bound nexus. Real
  neutrino flavor oscillation (experimentally established) requires SOME
  persistent, coherent internal structure -- disfavoring a pure
  "structureless, random contact" picture (see doc_torsion.txt's E=mc^2
  vs E=pc discussion). What remains open is narrower: whether that
  persistent structure IS this same chi(C5) character, or some other
  persistent label. NL2-NL6 (G_F and everything built from it) use only
  cell geometry (Rs, E_cell) and don't depend on this either way, so they
  are unaffected.

  ZONE 2 SHELL AS FIRST COUPLING POINT (2026-08-23, proton_resonance.py PROVEN):
    The proton Zone 2 shell (N_J=21 locked cells) resonates at m_p = 938 MeV.
    Free cells resonate at E_cell = 124.8 GeV.
    Lower resonance = larger displacement amplitude for same driving frequency.
    For a neutrino at E_nu (MeV scale), both sub-resonant but amplitude ratio:
      sigma(proton Zone 2) / sigma(free cell) = (E_cell/m_p)^4 = 3.13e+08
    The proton Zone 2 shell is the FIRST structure the freed lepton couples to.
    Empty space is 3e+08 times more transparent -- proton catches the neutrino first.
    [proton_resonance.py: sigma vs E table, crossover only at E_nu = E_cell = 124.8 GeV]
  When a freed lepton (neutrino) impacts a proton/neutron Zone 1 boundary:
    (1) Freed lepton amplitude couples to Zone 2 boundary (sigma ~ G_F^2 * E_nu^2)
    (2) Impact BREAKS the frozen Zone 2 lock (T_2g jammed state for proton)
        -- the Zone 2 lock is what stabilizes the proton; breaking it costs ~delta_E_lock
        -- this is why the energy threshold is tiny (1.8 MeV) even though W mass is 80 GeV:
           the lock-breaking cost is small; kinematic threshold = mass difference only
    (3) RETRACTED: an antipodal-vertex bounce mechanism previously proposed
        for the u->d quark transition (matching chi(T_1u,C5^2) = -1/phi
        = chi(T_2u,C5)) has been retracted.
    (4) The net transition udd -> uud is representation-theory-consistent:
        Zone 2 re-locks T_1g -> T_2g (CG: WI1+WI2 exact). Electron and
        antineutrino carry off the energy balance (IBD threshold = 1.806
        MeV, WD1) -- this is the current standing result.
    (5) Zone 2 re-locks in the new diquark configuration T_1g (neutron)
        The T_1g lock mode IS the virtual W boson (T_1g = directed transverse
        oscillation = W, not compression -- see doc_torsion Section 3.1)
    (6) Freed lepton gains the released vertex nexus from step (2) -> becomes
        a bound charged lepton (electron, muon, or tau per chi matching)
  W boson is not fundamental -- it is the transient T_1g lock mode during
  Zone 2 diquark restructuring. The 80 GeV W mass = E_cell x (W formula)
  is the energy of the T_1g phonon mode, not the lock-breaking energy itself.
  Range constraint: freed lepton must couple to Zone 1 vertex nexus energy scale.

- Single-cell restoration after b quark escape: energy accounting is
  consistent (released 4.18 GeV > m_b) and the cell we observe IS the new
  formation, but per b_quark_geometry.py's own conclusion the single-cell
  restoration MECHANISM remains OPEN -- only ensemble/statistical
  equilibrium across many cells is argued, not a direct single-cell
  derivation.

CLOSED THIS SESSION (2026-08-23):

- b quark mass DERIVED: m_b = m_p*sqrt(3)/(4*pi*alpha*phi^3) = 4182.8 MeV (+0.07%) [BQ9,BQ10]
- b quark = lambda_3 specifically (not lambda_8; R=G=7 -> zero restoring force) [BQ11,BQ12]
- Gluon C3 Born coupling: g_s_bare ~ alpha (no phi enhancement) [gluon_c3_born.py 9/9]
- Gluon amplitude A = L_J*sqrt(3)/6 DERIVED from face geometry [GH0b,GH0c]
- Tau geometric path: ALL 20 steps = 72 deg (C5), step = 2*phi/3 [GH2-GH5]
- Gluon strip counts: 12 (A<->B) + 9 + 9 = 30, not 10+10+10 [SU3-1 + calculation]
- F-14 algebraic ratio verified: pi*sqrt5/(2*alpha*phi^4) = 70.22 [F14a,b]

```
- Gluon spindle shape: 2 counter-rotating windings = permanent spindle form [doc_jobson_cell]
```

8. COMPANION SCRIPTS

```
analysis/demos/particle_generation_doc.py   -- 14/14 PASS  (STANDALONE; winding angle, F-14/LM17, D meson, CG, G_F
Repository: https://doi.org/10.5281/zenodo.22068557
```


ACKNOWLEDGMENT

This research was developed with the assistance of AI language models (GitHub Copilot / Claude, Anthropic). The conceptual framework, physical interpretation, and all claims are the sole work of the author. The author takes full responsibility for the derivation, the verification suite, and all results presented here. Use of AI assistance in this work is consistent with current practice in computational and theoretical physics research.

Repository: https://doi.org/10.5281/zenodo.22105622
Code: https://github.com/Destraag/torsionverse_public