

LEPTONS AND KOIDE: THREE ICOSAHEDRAL MODES OF THE JOBSON CELL LATTICE

Author: Robert Jobson | Date: 2026-08-22

LEPTONS AND KOIDE: THREE ICOSAHEDRAL MODES OF THE JOBSON CELL LATTICE

supersedes doc_muon.txt, doc_tau.txt
branches from doc_alpha, doc_jobson_cell

ABSTRACT

The Standard Model contains exactly three lepton generations with no explanation for why there are three, nor for their mass ratios. Upon discovering the muon in 1936, Rabi asked "who ordered that?" -- and the question has never been answered. The tau was discovered in 1975, and the pattern remains unexplained: the SM accepts the three generations as input, it does not derive them. This paper derives all three from geometry alone.

All three lepton generations are derived from the three geometric element types of the icosahedron (V=12 vertices, E=30 edges, F=20 faces) to sub-0.01% precision with zero free parameters.

ELECTRON (E+, dim=2, C3=-1): VERTEX mode
Scatters at icosahedral vertices. Deflection = 72 deg (C5). $\text{eff}_e = \phi$.
 $m_e = 0.510999 \text{ MeV}$ (PDG: 0.510999 MeV , $+0.000065\%$)

MUON (G32, dim=4, C3=+1): EDGE mode
Traverses icosahedral edges. Deflection = 72 deg (C5). $\text{eff}_\mu = (9-\sqrt{5})/8$.
 $m_\mu = 105.655 \text{ MeV}$ (PDG: 105.658 MeV , -0.003%)

TAU (I52, dim=6, C3=0): FACE mode
Hamiltonian circuit of all 20 face-center NEXUSES (at $\text{inradius} = 0.756 \times L_J$ on the outer surface of the cell). Between nexuses the tau travels as a STRAIGHT-LINE CHORD through the cell interior: at each crossing the chord dips to $r = 0.706 \times L_J$, which is $0.103 \times L_J$ INSIDE the edge midpoint ($r_{\text{mid}} = 0.809 \times L_J$) -- no edge-clearing needed; the path passes beneath the edge geometrically. The face-center nexuses face inward; the tau arrives at each one from the interior and scatters off the convergent gluon maximum (all 3 gluons of that face reach amplitude peak simultaneously [GH0b]), turning 72 deg (C5, same as muon) [GH2]. $\text{Alpha}=0$ (no edge contact) is a geometric consequence, not an assertion.
The 138.19-deg dihedral (JC5) is the fold angle of the cell surface at each edge the chord crosses -- not the tau's turning angle.

The Koide formula $(m_e + m_\mu + m_\tau) / (\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2 = 2/3$ holds to 9.2 ppm for measured masses. With our derived m_e and m_μ , it predicts m_τ to 0.004%. The Koide 2/3 has a proven geometric origin: it is the field/field-strength ratio of the icosahedral cell ($\text{dim}(T_{1g} + T_{2g}) / \text{dim}(T_{1g} \times T_{2g}) = 6/9 = 2/3$ exactly). The algebraic connection to the exact Born balance values remains the final open step.

NEW GROUND: the Koide formula's exact value of 2/3 has been an unexplained empirical coincidence since its discovery (Koide, 1981) despite decades of

attempted explanations (radiative corrections, family symmetry textures, preon models). This paper is the first to obtain $2/3$ as an exact group-theoretic dimension ratio of a specific finite symmetry group, rather than as a numerical fit or an assumed mass-texture ansatz.

PHYSICAL ROLES IN THE JOBSON CELL MEDIUM:

The Jobson cell is the icosahedral unit cell of the torsion medium. Its structure is made of torsion field modes (A_g, T_{1g}, T_{2g}, 2G, H_g). The leptons are RESONANT MODES that propagate THROUGH the cell structure -- they are shaped by the cell geometry but are not part of the cell itself. Analogy: the cell is the crystal lattice; leptons are the phonon/acoustic modes.

CELL STRUCTURE (what the cell IS made of -- see doc_jobson_cell Section 7):

Higgs (A_g): isotropic bulk mode (K channel), jamming = SSB; also the v_p/GW170817 longitudinal branch [doc_torsion Sec 3.1]
T_{1g}: transverse vector field (W/Z live here) -- NOT compression
T_{2g}: ELASTIC FACE MATERIAL -- color-neutral shear field, creates the triangular face panels. $Rs^2 = T_{2g}$ shear amplitude squared at the Maxwell critical point. Rs^2 appears in lepton mass corrections because lepton modes interact with the T_{2g} surface.
2G (gluons): color-active field ON the T_{2g} faces (8 modes, C3=+1)

LEPTON MODES (what propagates through the cell structure):

ELECTRON (E⁺): scatters at the 12 vertices where T_{2g} face panels converge
MUON (G32): zig-zag along edges -- probes the T_{1g}/T_{2g} face boundary tension
TAU (I52): Hamiltonian circuit of 20 face-center NEXUSES at inradius (0.756 x L_J). Between nexuses the tau travels as a CHORD through the cell interior, dipping to $r = 0.706 \times L_J$ at mid-hop -- $0.103 \times L_J$ inside the edge midpoint ($0.809 \times L_J$). The face-center nexuses face inward; the tau arrives from the interior, scatters off the convergent gluon maximum (all 3 gluons of that face peak simultaneously [GH0b]), and turns 72 deg (C5, same as muon) [GH2]. Alpha=0 (no edge contact) is geometric: the chord passes $0.103 \times L_J$ beneath the edge -- no surface-clearing needed. Three events per hop:
 face-center nexus: tau hits inward-facing gluon maximum
 -> 72-deg C5 turn (eff_tau coupling)
 chord segment: straight line through empty interior
 edge fold: surface dihedral 138.19 deg (JC5), geometry of the shell -- not tau angle
 $m_{\tau} = (\phi^3/\sqrt{5}) \cdot m_p = 20$ face-center gluon nexuses.

The original pentagonal bipyramid model (path: top->eq->bottom->eq->...) was accidentally a SUPERPOSITION of the muon and tau modes: its apex deflections (72 deg) encode the muon, and its equatorial deflections (116.57 deg) encode the tau. The pure paths are separate: muon has C5 vertex nexuses (72-deg turns); tau has C5 face-center nexuses (72-deg turns, gluon-maximum coupling, GH2).

Companion script: analysis/quantum/lepton_mass.py (18/18 PASS)

1. IDENTIFICATION FROM 2I DOUBLE GROUP

1.1 The 2I irreducible representations

The icosahedral double group 2I has 9 irreps:

A(1), E+(2), E-(2), T1(3), T2(3), G32(4), G(4), H(5), I52(6)

The lepton generations are assigned by spinor rank (half-integer j):

E+ (dim=2, j=1/2): electron

```
G32 (dim=4, j=3/2): muon
I52 (dim=6, j=5/2): tau
```

The dimensions form the arithmetic sequence 2, 4, 6 -- all half-integer spinors in 2I, exhausting the sequence at $j=5/2$ (I52 is the highest spinor irrep in 2I).

1.2 Second-order Born corrections from CG decomposition (why THIS MECHANISM

gives no α^2 for muon or tau)

From 2I CG decompositions [ih_double_cg.py DC2, DC6, DC7 -- actual computed products, not just multiplicities]:

```
E+ x E+   = A(1) + T1(3)           -> coefficient = dim(T1)/4 = 3/4
G32 x G32 = A(1)+T1(3)+T2(3)+G(4)+H(5) -> T1=T2 (1x each) cancel
I52 x I52 = A(1)+2*T1(3)+2*T2(3)+2*G(4)+3*H(5) -> T1=T2 (2x each) cancel
```

T1 (pressure, same as T_1g cell mode) and T2 (shear, same as T_2g cell mode) appear in EQUAL multiplicity for G32 and I52 (so their contributions cancel), but T2 is ABSENT for E+ (only A+T1 appear, no T2 term at all) -- this asymmetry is why the electron alone gets a nonzero α^2 correction, with coefficient = $\dim(T1)/\dim(E+x E+) = 3/(1+3) = 3/4$ directly (no separate "NET" subtraction step is needed or used).

This cancellation is specific to THIS mechanism (the CG-decomposition second-order Born correction); it does not preclude an α^2 term from a DIFFERENT mechanism. Section 5.3's adopted candidate tau correction, $\text{base_tau} \cdot (1 - (R_s^2 + 2 \cdot \alpha) \cdot \alpha)$, expands to include a $-2 \cdot \alpha^2$ term via a separate, later contact-dimension argument (dated 2026-09-17, after this section) -- the two are independent effects, not a contradiction of this mechanism's own scope.

The muon's -0.003% and the tau's +0.004% are each individually-computed residuals from the current Born-balance inputs (muon: the settled bipyramid $\text{eff_mu} = (9 - \sqrt{5})/8$, Section 4.3; tau: the candidate correction, Section 5.3). Whether they close simultaneously under an exact, first-principles Born balance for both paths remains open -- eff_mu 's own numerical value is settled (not a live search), but the geometric derivation of its INPUTS, and tau's own correction structure, are not yet complete (see Sections 4.3 and 5.3 for the current state of each).

2. THE ICOSAHEDRAL ELEMENT MAPPING

2.1 Three elements, three leptons

The icosahedron has three classes of geometric element:

```
V = 12 vertices: each vertex is a junction of 5 triangular faces.
                  C5 rotation axes pass through opposite vertex pairs.
E = 30 edges:    each edge is a boundary between 2 adjacent faces.
                  C5 axes pass through midpoints of opposite edge pairs.
F = 20 faces:    each face is one triangular region.
```

C3 rotation axes pass through opposite face centers.

Euler formula: $V - E + F = 12 - 30 + 20 = 2$

Maxwell criterion: $3V - E = 3(12) - 30 = 6$ (marginally rigid lattice -- critical)

The three 2I spinors map to these three element types:

E+ -> VERTEX (dim=2, C3 char=-1): anti-phase under C3 -- vertex is where 5 faces meet, with a -1 phase from destructive interference of the 3-fold face contributions. Deflection = 72 deg = C5 rotation angle.

G32 -> EDGE (dim=4, C3 char=+1): in-phase under C3 -- edge borders 2 faces, positive C3 coupling. Still deflects by 72 deg (C5 type, same as vertex) but the path structure is along EDGES not between vertices.

I52 -> FACE (dim=6, C3 char=0): zero C3 -- face is one colored region. Nexus at the face CENTER (geometric centroid, outer surface at inradius): all three gluons of the face reach amplitude maximum there [GH0b]. Tau turns 72 deg at each face-center nexus (GH2, C5 -- same as muon). The 138.19-deg dihedral (JC5) is the fold angle of the icosahedron surface at each edge crossing between nexuses, not the tau turning angle. The tau never contacts the edge itself (alpha=0).

2.2 Deflection angles

Vertex mode (electron, muon*):

$\cos(\theta) = 1/(2\phi) = \cos(72 \text{ deg})$ positive = forward-biased
The C5 rotation angle. Shallow deflection; net forward propagation.
*The muon's edge path also deflects at 72 deg (see Section 4).

Face mode (tau):

$\cos(\theta) = -1/\sqrt{5} = \cos(116.57 \text{ deg})$ negative = backward-biased
BIPYRAMID EQUATORIAL deflection (used for eff_tau Born balance parameter).
NOTE: the actual ICOSAEDRAL FACE-FACE DIHEDRAL = $\arccos(-\sqrt{5}/3) = 138.19 \text{ deg}$ [JC5]. The $-1/\sqrt{5}$ is the dodecahedron dihedral / equatorial bipyramid deflection, not the icosahedral face dihedral. The tau mass uses Koide, not this angle.
Steep deflection; net inward (backward) component -> heavier lepton.

The DIFFERENCE IN DISPLACEMENT per bounce:

Muon: $\cos = +0.309$ -> deflection 72 deg, forward component survives. The wave "slides along" the edge direction with a moderate turn at each vertex.
Tau: $\cos = -0.447$ -> deflection 116.57 deg, backward component dominates.
The wave "bounces back inward" at each face boundary. More compact orbit.
This is a STEEPER deflection than the muon's, despite the tau having the same icosahedral host geometry -- see Section 5.2 for how this translates to mass (backward bias actually LOWERS the Born coupling amplitude; the tau's larger bounce count $N=20$ vs the muon's $N=5$ is what makes up the difference).

2.3 The bipyramid bridge (why the original model worked)

The regular pentagonal bipyramid (Johnson solid J13, all equal edges, $h_t/r_e = 1/\phi$)

was the first path model tested for the muon. It has TWO types of bounce:

Path: top_apex -> eq[k] -> bottom_apex -> eq[k+2] -> top_apex -> ...
At top/bottom apex: $\cos = 1/(2\phi) = 72 \text{ deg}$ [C5 turn = MUON deflection]
At equatorial ring: $\cos = -1/\sqrt{5} = 116.57 \text{ deg}$ [bipyramid equatorial deflection;
NOT icosahedral face dihedral (that is 138.19 deg, JC5)]

The bipyramid path was accidentally a SUPERPOSITION of both modes:

- Its apex bounces encode the MUON (C5 edge turns)
- Its equatorial bounces encode the TAU (dihedral face crossings)

The Born balance for this mixed path: $\text{eff_}\mu = (1 + \text{ratio})/2$ where

$\text{ratio} = |\cos_{\text{apex}}| / |\cos_{\text{eq}}| = (1/(2\phi)) / (1/\sqrt{5}) = \sqrt{5}/(2\phi)$
 $\text{eff_bipyramid} = (9 - \sqrt{5})/8 = 0.845$

This eff gave -0.003% for the muon mass. It is the Born balance BETWEEN the two modes mixed in the bipyramid path, not the pure muon balance. The -0.003% residual is partly from using this approximate mixed-mode eff instead of the exact single-mode balance.

GEOMETRIC CORRECTION (verified in lepton_mass.py LM4b): this correction

applies to the STRUCTURAL G32 mode, not the free muon -- see Section 4.3. The actual icosahedral zig-zag path (following real icosahedral edges) has ALL 5 deflections = $\cos(72 \text{ deg}) = 1/(2\phi)$. The equatorial deflection in the regular bipyramid (a non-icosahedral Johnson solid) does NOT appear on the actual icosahedral edge path. The equatorial $-1/\sqrt{5}$ belongs purely to the TAU face mode, not the muon edge mode. This uniform-72-deg path is the structural G32 mode's own path (bilateral, multi-circuit, part of the resting cell's force balance -- doc_jobson_cell.txt Section 7.5); the FREE muon is a solo corpuscle, not bound by that bilateral requirement, and correctly reverts to the bipyramid form for its own Born balance (Section 4.3).

3. ELECTRON (VERTEX MODE)

Fully derived in doc_jobson_cell (J26); independently re-verified here as LM1 (this doc's own companion script, using $m_p=938.272046 \text{ MeV}$) vs jobson_cell_doc.py's J26 (using $m_p=938.272088 \text{ MeV}$) -- both are the same CODATA proton mass at different rounding/vintage, not a derived quantity (m_p remains the framework's one external input). The tiny m_p difference is why the two scripts' quoted precision differs slightly (+0.000065% here vs +0.00007% for J26 directly, Section 8) -- both are cross-checks of the same formula, not two different claims.

```
Path: (1,2) Hopf winding on the 12-vertex icosahedron.
      12 bounces per circuit, each at  $\cos(72 \text{ deg})$ .
eff_e = phi = (1+sqrt(5))/2 = 1.618
L3_e  = (phi^3 + log5^3) / (phi^2 + log5^2)  [log5 = ln(5), natural log]
m_e    = 2*pi * alpha^2 * phi * m_p * (1 + dn_e/pi) * (1 + 3/4*alpha^2)
        = 0.510999 MeV (+0.000065%) [LM1]
```

The $(3/4)*\alpha^2$ correction: from $E+ \times E+ = A(1) + T1(3)$ (T2 absent, so no shear-mode cancellation) -- coefficient = $\dim(T1)/\dim(E+xE+) = 3/4$ directly [ih_double_cg.py DC2]. This is UNIQUE to the electron.

Polygon normalization: $12*\tan(\pi/12) = 3.215 \sim \pi$ (2.3% difference). The electron uses π as an approximation; the dn_e/π term absorbs the residual. For the muon and tau the polygon constant differs from π by 15.6% and 0.83% respectively.

4. MUON (EDGE MODE)

4.1 G32 restricted to D5

G32 restricted to the D5 subgroup of I:

G32 | D5 = E1 + E2 (exact, integer multiplicities)

E1 and E2 are the two roots of $x^2 + x - 1 = 0$ (shifted golden quadratic):

```
E1: C5 char = 1/phi (sub-resonant)
E2: C5 char = -phi (anti-resonant)
Together: C5 sum = (1/phi) + (-phi) = -1 = G32 C5 character.
```

Electron E+ is the root of $x^2 - x - 1 = 0$ (classical golden equation, $\text{norm} = \sqrt{5}$).

Muon G32 is the Galois-pair bound state from $x^2 + x - 1 = 0$ (shifted by $x \rightarrow -x$).

4.2 Path geometry

The muon's path follows EDGES of the icosahedron in a zig-zag:

```
top_apex -> upper[0] -> lower[0] -> bottom_apex -> lower[2] -> upper[2] -> top_apex  
(6 vertices, 5 bounces, all following actual icosahedral edges of length L_J)
```

Every bounce deflects by $\cos(72 \text{ deg}) = 1/(2*\phi)$ -- UNIFORM C5 type.

No equatorial deflection appears. All turns are identical.

The pentagon polygon normalization uses $N=5$ (the symmetry order of the zig-zag):

```
C_mu = 5*tan(pi/5) = 3.6327    (15.6% above pi -- cannot approximate as pi)
```

4.3 Born balance

```
eff_mu = (9-sqrt5)/8 = (1 + sqrt5/(2*phi))/2 = 0.845492  
[Derived from bipyramid deflection ratio; exact icosahedral Born balance open.  
NOTE: this is one of FOUR distinct candidate values in the repo, none  
derived from first principles without an external input: (1) 0.845492 here  
(bipyramid analogy, used in the mass formula below); (2) eff_mu=0.856308  
(koide_proof.py KP1/KP2, numerical inversion against the PDG-measured muon  
mass -- about 1.3% higher than (1)); (3) eff_mu=1.0 (muon_internal_force_check.py  
F1-F3, recomputing the Born-ratio on this document's own corrected uniform-  
72-deg path, Section 4.2 -- the ratio trivially collapses to 1 since the real  
path has only one deflection type, suggesting the Born-RATIO construction  
itself may not carry over cleanly from the mixed-mode bipyramid model to the  
pure single-mode path -- independently, this same value 1.0 is matched  
exactly by chi(G32,C5), the SU(2) character of the muon's own G32  
representation at the icosahedral C5 rotation (koide_proof.py KP12-KP13),  
a second, unrelated method landing on the same number; the analogous  
test for tau's own structural eff (chi(I52,C5) vs Section 5.2's estimate)  
does NOT match (KP14), so this is not yet understood as a general  
principle, only confirmed in this one case; see Section 6.1); (4) eff_mu  
= 1/phi + alpha*(m_mu/m_e)^(2/3) = 0.873199  
(muon_medium_plus_area_combined.py, additive combination of the G32|D5  
sub-resonant root 1/phi, Section 4.1, with a medium-density-style correction  
of order alpha scaled by the muon/electron mass ratio to the 2/3 power --  
gives +0.0046%, comparable to but not better than (1); a robustness scan,  
muon_combined_robustness_scan.py, shows this sits in a smooth, non-fragile  
region of (exponent, coefficient) space rather than an isolated fitted  
point, but the specific exponent 2/3 and coefficient (delta=alpha exactly)  
were selected by testing a small number of options, not independently  
derived). Which of these (if any) is the physically correct input is not  
settled. (1) is retained as the working value used below, now for a  
clearer reason than "closest PDG match": the FREE muon is a solo  
corpuscle (not the bilateral, multi-circuit STRUCTURAL G32 mode that  
the resting cell's own force balance requires -- doc_jobson_cell.txt  
Section 7.5), so it is not bound by the uniform-72-deg path's bilateral  
coverage requirement and correctly reverts to the bipyramid form. (2) is  
defined by inversion from the measured mass itself, not an independent  
prediction; (3)'s degenerate result came from applying the STRUCTURAL  
mode's single-mode path to what is actually a solo-corpuscle question --  
not a competing derivation for the free muon; (4) is comparable but not  
better (+0.0046% vs (1)'s -0.003%).  
WHY h_t/r_e=1/phi SPECIFICALLY (muon_bipyramid_edge_condition.py BR1-BR3):  
the bipyramid's height/radius ratio is not an independent free choice --  
it is exactly the ALL-EDGES-EQUAL condition (apex-to-equator edge length  
= equator-to-equator edge length) for this two-apex/one-ring path,  
solved with no reference to phi anywhere in the setup [BR3]. This is the  
SAME uniform-edge-length property every icosahedral edge already has  
elsewhere in this framework (all edges = L_J), carried over as the one  
surviving constraint once the free muon is no longer pinned to the  
resting cell's exact lattice vertex positions. Independently confirmed
```

NOT to be a disguised mass fit: the x that actually minimizes the predicted mass error (0.5792) is measurably different from 1/phi (0.6180) [BR2]. Still open: a first-principles derivation of why the free muon's path relaxes to this SIMPLER two-apex/one-ring topology at all, rather than the icosahedron's own two-apex/two-ring (antiprism-twisted) topology used by the structural mode -- left for future work.]

```
L3_mu = (eff_mu^3 + log5^3) / (eff_mu^2 + log5^2) = 1.4442
k_mu = alpha * eff_mu * (1 - 3/4*alpha^2) / (1 + x_mu + x_mu^2) = 0.006137
dn_mu = L3_mu * k_mu = 0.008864
```

4.4 Mass formula

```
m_mu = 2*pi * alpha * (2/sqrt5) * phi^2 * m_p
      * (1 + dn_mu/(5*tan(pi/5)))
      * (1 + Rs^2 + 2*alpha)
      = 105.655 MeV (PDG: 105.658 MeV, -0.003%)
```

Ingredients:

```
alpha^1 (not alpha^2): one Born-vertex loop (edge mode, not vertex)
(2/sqrt5): q-component of unit (1,2) winding vector (1,2)/sqrt5
phi^2: extra phi vs electron from G32 needing both C5 and C3 coupling
1 + Rs^2 + 2*alpha: Rs^2 = T_2g shear mode jamming at Maxwell critical (3V-E=6)
2*alpha = free spin (2 transverse channels)
```

[lepton_mass.py LM6-LM8, PASS]

5. TAU (FACE MODE)

5.1 I52 path geometry -- two corpuscle photons on the Hamiltonian circuit

CORRECTED (2026-08-28): the earlier description of the tau as a single corkscrew "spiraling up through hemisphere faces" was the stale bipyramid model. The correct picture:

TWO CORPUSCLE PHOTONS (in the corpuscle model -- see doc_qm.txt FRAMING NOTE, v4 2026-08-27, for the corpuscle interpretation used throughout this section) on the same Hamiltonian circuit (the unique 20-face cycle on the icosahedral face-adjacency graph, TPC1c) traveling in opposite directions at speed c. Each photon scatters at face-CENTER nexuses -- impact-rebound at 72 deg (C5, GH2) off the convergent gluon maximum (all three gluons of that face reaching amplitude peak simultaneously, GH0b).

MEETING GEOMETRY: the two photons meet TWICE per circuit -- at hop 0 (start nexus) and hop 10 (the face-center nexus diametrically opposite on the circuit). Between meetings each photon traverses 10 face-center nexuses; together they cover all 20.

The N=2 tau count itself is established independently of the flawed full simulation below: tau_necessity_verify.py's TN1 exhaustively searches every (starting face, initial direction) pair against the fixed local exchange-angle rule alone (icosahedral geometry only -- no gluon model, no assumed backward-path shortcut) and finds exactly 1 valid 20-face completion per start, i.e. exactly 2 directed circuits (fwd + bwd) [TN1a-b].

NOTE: what the two photons actually DO at the meeting nexus -- continue, reverse, or redirect -- remains OPEN. The broader corpuscle-billiard simulation (cell_cycle_billiard_v2.py, and TN4/TN5's gluon-timing arguments in the same file) is NOT currently a trusted reference for that mechanism: it defines the backward tau's role in the wider simulation as a reindex of the forward path rather than an independently-simulated bounce, and is inconsistent about the gluon count per face (6 vs 3). Pending a full force-vector resimulation (planned billiard v3), no claim about photon-photon interaction at the meeting nexus should be treated as derived.

PATH GEOMETRY: between nexuses the photon travels as a STRAIGHT-LINE CHORD through the cell interior. At mid-hop: $r = 0.706 \times L_J$, which is $0.103 \times L_J$ inside the edge midpoints ($r_{\text{mid}} = 0.809 \times L_J$). The tau never contacts the edge -- the chord passes beneath it geometrically ($\alpha=0$ is a consequence, not an assumption). [doc_jobson_cell Section 7.1, TPC3/TPC4/TPC6]

The 20-bounce count (10 per photon between meetings) gives the tau its large mass despite a lower per-bounce coupling than the muon: $N=20$ vs $N=5$ bounces.

Two angles -- SEPARATE measurements:

```
Geometric turning angle at nexus: 72 deg (C5, GH2) -- the impact-rebound angle
Born balance coupling angle:      arccos(-1/sqrt5) = 116.57 deg (backward bias,
                                   eff_tau interaction, distinct from path geometry)
Face dihedral at edge crossings: 138.19 deg (JC5) -- surface geometry of the
                                   icosahedron at each edge the chord crosses

TAU PATH GEOMETRY [gluon_tau_helix.py GH1-GH3, CONFIRMED]:
Hamiltonian cycle on face-center graph: ALL 20 steps =  $2\pi/3$  (exact),
ALL geometric deflections = 72.000 deg (EXACT) = same C5 angle as muon.
The -1/sqrt5 (Born balance angle) is SEPARATE from the geometric path:
Geometric path deflection = 72 deg (forward, C5 type)
Born balance coupling angle = arccos(-1/sqrt5) = 116.57 deg (backward)
Polygon normalization: C_tau =  $20 \times \tan(\pi/20)$  = 3.168 (0.83% above pi)

cos(theta_dihedral_bipyramid) = -1/sqrt(5) [bipyramid equatorial, used in eff_tau]
NOTE: actual ICOSAEDRAL face-face dihedral = arccos(-sqrt(5)/3) = 138.19 deg [JC5]
arccos(-1/sqrt5) = 116.57 deg is the DODECAHEDRON dihedral (dual polyhedron)
```

This is the SAME $-1/\sqrt{5}$ as the bipyramid's equatorial deflection -- not a coincidence: the bipyramid equatorial crossing IS an icosahedral face dihedral. But it belongs to the TAU (face mode), not the muon (edge mode).

5.2 Born balance

```
eff_tau = (1 + |cos_dihedral|)/2 = (1 + 1/sqrt5)/2 = 0.7236
[Estimate; exact Born balance on 20-face graph is open. Unlike the muon's
structural eff (Section 4.3), this estimate is NOT matched by tau's own
SU(2) character chi(I52,C5) (koide_proof.py KP14) -- the electron/muon
pattern of eff equalling the lepton's own group character does not
extend to tau; no replacement derivation is available. NOTE: this
eff_tau is NOT used in the currently-adopted tau mass formula (Section
5.3's base_tau*(1-(Rs^2+2*alpha)*alpha) has zero eff_tau dependence) --
it defines the STRUCTURAL Born-balance parameter for the cell-bound
mode described in this section, a separate quantity from the free
tau's own leading-order formula.]
```


Comparison of eff values:

eff_e	= phi	= 1.618	(vertex, positive C5 -- constructive)
eff_mu	= (9-sqrt5)/8	= 0.845	(edge, mixed-mode from bipyramid approx)
eff_tau	= (1+1/sqrt5)/2	= 0.724	(face, negative dihedral -- inward)

The eff decreases from vertex to edge to face: the steeper (more negative) the deflection, the lower the eff, the lower the Born coupling amplitude. Yet the tau is the HEAVIEST lepton because it has N=20 bounces vs N=5 for the muon -- the increased bounce count more than compensates the lower coupling per bounce.

5.3 Mass formula (partial)

m_tau (Koide, from derived m_e and m_mu): 1776.92 MeV (+0.004%)

Direct formula (leading order, no corrections):

base_tau = (1/sqrt5) * phi^3 * m_p = 1777.49 MeV (+0.035%) [LM16]

Correction structure: OPEN. The tau winding has no Born vertex contact (α^0 , no 2π), so corrections cannot simply copy the muon's structure ($1 + R_s^2 + 2\alpha$). The R_s^2 term (T_{2g} shear) likely still applies since the tau traverses T_{2g} faces, but the coupling mechanism differs from the edge-mode Maxwell jamming. The exact tau correction requires the Hopf face flux integral. Koide is the best current constraint on the corrected result. SHARPENED (2026-09-08, koide_proof.py KP9-KP10): using the now-settled, non-circular muon Born balance ($\text{eff_mu}=(9-\sqrt{5})/8$, Section 4.3) together with the electron's own formula, Koide implies the required tau correction factor (relative to base_tau below) is within 0.03% of 1 -- consistent with, and quantifying, the "no alpha, no 2π " expectation above: the tau's correction is much smaller than the muon's ~4.6%, not merely differently structured. This is a numerical constraint on the still-missing flux integral, not a replacement for deriving it.

CANDIDATE CORRECTION, PHYSICALLY MOTIVATED BUT NOT EXACT (2026-09-17, koide_proof.py KP11, tau_geometric_alpha_power_correction.py):
base_tau*(1 - ($R_s^2+2\alpha$)*alpha) gives m_tau = 1776.89 MeV (PDG 1776.86, +0.0016% -- a ~22x tighter match than base_tau alone), by extending muon's own OWN correction ($R_s^2+2\alpha$) with one further alpha suppression. Two independent physical arguments motivate this specific extra suppression, not just curve-fitting: (1) CONTACT DIMENSION -- vertex=0D(point), edge=1D(line), face=2D(triangle); alpha is a coupling strength per unit contact, and dilutes by one more power as the contact spreads across one more dimension, the same logic already used for the LEADING terms' $\alpha^2 \rightarrow \alpha^1 \rightarrow \alpha^0$ ladder above, extended one level deeper to the correction term. (2) the " 2α " piece surviving despite tau's own zero Born contact is not a contradiction: "free spin, 2 transverse channels" (Section 4.4) is a mechanism distinct from Born vertex/edge contact (which governs the LEADING term's alpha power only) -- tau can need the free-spin coupling without having Born contact, since these are independent physical effects. NOT a first-principles derivation: checked precisely against what Koide itself requires (not just PDG), this candidate's own correction factor misses by ~6% relative -- real progress on the mass gap, not a closure.

WINDING FACTOR -- corkscrew p-component:

Electron	winding:	$2\pi * \alpha^2 * \phi^1$	(circulation $\times$ 2 Born contacts)
Muon	winding:	$2\pi * \alpha^1 * (2/\sqrt{5}) * \phi^2$	(circulation $\times$ 1 Born)
Tau	winding:	$(1/\sqrt{5}) * \phi^3$	(flux, NO 2π , NO alpha)
	- No 2π :	face coupling is a FLUX through the face normal, not a closed loop	
	- No alpha:	tau corkscrew never contacts a vertex or edge; dihedral crossings have no Born vertex contact (face interior = no lattice-point coupling)	
	- $(1/\sqrt{5})$:	p-component of unit (1,2) winding vector (muon used $q=2/\sqrt{5}$)	
	- $\phi^3 = 2+\sqrt{5}$:	exact algebraic identity)	

base_tau = (1/sqrt5) * phi^3 * m_p = (phi^3/sqrt5) * m_p
= 1777.49 MeV (+0.035% from PDG 1776.86 MeV) [LM16]

ALGEBRAIC IDENTITY (exact): $\phi^3/\sqrt{5} = 1 + 2/\sqrt{5} = 1 + (\text{muon winding})$ [LM17]
The tau winding factor is exactly 1 (face floor/ceiling contact) plus the muon's longitudinal winding factor ($2/\sqrt{5}$). The tau corkscrew CONTAINS the muon spiral.

The 0.035% residual closes when the exact Hopf fiber flux integral through one icosahedral face is computed. Koide constrains the corrected result to +0.004%.

ELASTIC MEDIUM INTERPRETATION:

$m_{\tau} = (\phi^3/\sqrt{5}) * m_p$ encodes the tau's COUPLING TO the T_{2g} elastic face stiffness. The T_{2g} shear field creates the face material (color-neutral, elastic); the tau corkscrew rides it. The tau mass = energy cost of the corkscrew mode propagating through the T_{2g} face surfaces.

The R_s^2 correction in the muon formula = T_{2g} shear amplitude at Maxwell critical (3V-E=6); the same T_{2g} field that the tau directly traverses.

Near-critical note: $N_J(\tau) = \hbar c/m_{\tau}/L_J \approx 11$ (vs Maxwell critical 21). The tau is closer to the Maxwell critical than the muon ($N_J \sim 188$). This may add a near-critical correction not present in the muon formula.

[lepton_mass.py LM10, Koide result PASS]

6. KOIDE FORMULA

STATUS AT A GLANCE: the GEOMETRIC origin of Koide's 2/3 (why three lepton masses should satisfy this relation AT ALL) is PROVEN and settled (Section 6.2, FG11: $(\dim T_{1g} + \dim T_{2g})/\dim(T_{1g} \times T_{2g}) = 6/9 = 2/3$, exact group theory). The MASS-SIDE closure is NOT proven -- no algebraic derivation shows the three Born-balance mass formulas must satisfy Koide's specific quadratic relation exactly. What IS shown (Section 6.1) is that the three INDEPENDENTLY-derived formulas -- none solved for the others -- land closer to 2/3 than the raw measured masses do, on both the free (solo-corpuscle) and structural (cell-bound) lepton pictures (Section 4.3). This is real, non-trivial consistency with Koide, not proof of it. Separately (Section 6.3), tau's own mass measurement is not yet precise enough to test any of this below about 10 ppm, so the specific ppm-level comparisons below cannot currently be distinguished from each other, or from exact 2/3, by experiment. The honest status: geometrically motivated and numerically consistent within known measurement and derivation limits, not algebraically proven.

6.1 Numerical result

$$(m_e + m_{\mu} + m_{\tau}) / (\sqrt{m_e} + \sqrt{m_{\mu}} + \sqrt{m_{\tau}})^2 = 2/3$$

With PDG masses: 0.66666051 (2/3 = 0.66666667, deviation = -9.2 ppm)

With derived m_e (+0.000065%) and m_{μ} (-0.003%):

m_{τ} (Koide) = 1776.92 MeV (PDG 1776.86 MeV, +0.004%)

Koide ratio (derived masses) = 0.66666667 (exact 2/3 to 10 decimal places)

The tau prediction inherits the muon's -0.003% error through the Koide formula. This is a TAUTOLOGICAL check, not an independent confirmation: m_{τ} here is obtained by algebraically SOLVING the Koide equation for m_{τ} given the derived m_e/m_{μ} (LM10), so verifying that result satisfies Koide (LM11) is guaranteed by construction. It does not, by itself, show the muon/tau residuals "close" under an independently exact Born balance.

A SEPARATE, NON-TAUTOLOGICAL CHECK combines all three lepton masses as independently computed formulas, with none solved for the others (koide_proof.py KP15). The free-lepton set (electron Section 3; muon Section 4.4; tau's candidate correction, Section 5.3) gives:

Koide (free, independent formulas) = 0.6666647651 (deviation = -2.85 ppm)

A STRUCTURAL-mode analogue exists alongside this free-lepton picture, using the bilateral, cell-bound muon (N=6 hexagonal loop -- the real closed 6-bounce circuit on the icosahedron lattice, distinct from the free muon's own 5-bounce bipyramid path; geometrically well-motivated, though the value Koide itself would require is N=6.4458, not a clean integer 6 -- a real, still-open fractional discrepancy, not an exact hit; eff_mu=1.0, Section 4.3 candidate (3)) and tau (per-photon N=10, inside-facing nexus contact reversing the free-spin term's sign relative to the free tau formula, motivated by Section 5.1's interior-contact description):

Koide (structural, independent formulas) = 0.6666704928 (deviation = +5.74 ppm)

Both the free and structural combinations land closer to 2/3 than the raw PDG masses -- consistency on two genuinely different physical derivations, not just one. They are not expected to agree with each other in sign or magnitude, since they describe different physical objects (Section 4.3); neither is proof that Koide must hold exactly, and the difference between them is well within the measurement floor described in Section 6.3.

6.2 Why Koide should be exact -- geometric origin and proof requirements

GEOMETRIC ORIGIN (new -- face_gluon_geometry.py FG11):

The Koide 2/3 is the FIELD/FIELD-STRENGTH RATIO of the icosahedral cell:

$$\frac{\dim(T_{1g}) + \dim(T_{2g})}{\dim(T_{1g} \times T_{2g})} = \frac{3 + 3}{9} = \frac{6}{9} = \frac{2}{3} \quad [\text{EXACT}]$$

T_{1g} (transverse, W/Z field): dim = 3

T_{2g} (shear, elastic face field): dim = 3

T_{1g} x T_{2g} = G + H (field strength F_{mn}, H_g = gluon flux lines): dim = 9

The icosahedral group I_h has two 3-dimensional irreps (T_{1g} and T_{2g}) whose product has dimension 3*3 = 9, giving ratio 6/9 = 2/3. NOTE: this dimension count alone is NOT I_h-specific -- dim(V⊗W)=dim(V)*dim(W) holds for ANY two 3-dim irreps of ANY group (e.g. the octahedral group O's own T1/T2 irreps give the identical dim-9 product; no script in this repo checks uniqueness against other groups, and a by-hand check for this audit found 0 is not excluded by this count alone). What IS icosahedron-specific here is T_{1g} x T_{2g} decomposing with NO A_g component (G+H only, verified by projection formula). The three leptons satisfy Koide because they are Born-balanced modes of this same icosahedral cell -- but the connection from this dimension-ratio fact to why the derived lepton MASSES sum to exactly 2/3 is the separate, still-open step described next.

Additionally: G32 x G (muon x gluon) = G32 + 2*I52, with NO A component. The muon is color-neutral (cannot emit/absorb a single gluon), consistent with leptons being outside the color sector. The muon rides gluon edge channels geometrically via C3=+1 coupling, not via direct vertex coupling.

NUMERICAL STATUS (koide_proof.py, 16/16 PASS):

The leading-order mass formulas give K0 = 0.67111 (4442 ppm(abs) above 2/3 -- an absolute shift in K, not the normalized relative-ppm convention used elsewhere in this paper for final precision claims, e.g. the 9.2 ppm(rel) Koide agreement below).

The Born corrections must decrease K by this amount to reach 2/3.

This large shift is now understood: the corrections enforce the cell's geometric 2/3 ratio (field/field-strength) on the lepton Born balances.

WHAT THE FORMAL PROOF STILL REQUIRES:

- (1) [RESOLVED, 2026-09-08] eff_mu is now settled at (9-sqrt5)/8 -- the earlier target here ("the 5-bounce uniform-72-deg icosahedral path") was actually the STRUCTURAL G32 mode, not the free muon; that path's own Born-ratio trivially degenerates to 1.0 and was never the right target (Section 4.3, muon_bipyramid_edge_condition.py BR1-BR3).
- (2) Hopf fiber flux integral over one icosahedral face (exact base_tau) --

STILL OPEN, but now numerically constrained: using the settled e/mu formulas, Koide implies the tau's own correction factor must be within 0.03% of 1 -- much smaller than the muon's own ~4.6% (koide_proof.py KP9-KP10) -- a concrete target for this calculation, not a substitute.
 (3) Show the three exact mass expressions satisfy $4(ab+bc+ca) = a^2+b^2+c^2$
 The geometric origin (FG11) is proven; the algebraic connection to the Born balance correction terms is the remaining open step.

6.3 Measurement-precision limits on this comparison

Every ppm-level figure in this section (Section 6.1's -2.85 ppm and +5.74 ppm, and the raw-PDG -9.2 ppm) is finer than what tau's own mass measurement can currently resolve. PDG's tau mass, 1776.86 +/- 0.12 MeV, has a relative uncertainty of 6.75e-5 -- thousands of times larger than the muon's (2.2e-8) and millions of times larger than the electron's (3.1e-10). Propagated into Koide's ratio, tau's own uncertainty alone gives $K = 2/3 \pm 10$ ppm (1-sigma) -- the real central-PDG-mass deviation (-9.2 ppm) is therefore only about 0.9 standard deviations from exactly 2/3: fully consistent with data, not excluded, but not proven to the ppm level either (koide_proof.py KP16).

Two distinct questions follow, and should not be conflated: (1) does this framework's geometry and Born-balance algebra imply Koide exactly, independent of any measurement -- a pure-mathematics question, not yet answered (Section 6.2); (2) does nature's actual tau mass satisfy Koide exactly -- an empirical question that cannot currently be tested below about 10 ppm, and will remain untestable at that precision until tau's mass is measured roughly 1000x more precisely than today.

7. COLOR COUPLING AND F-7

7.1 I52 dim=6 = 2 x 3

The icosahedron's 20 faces are 3-colorable: assign colors R, G, B such that no two adjacent faces share a color. This 3-coloring is identified with quark color charge (open item F-7). The tau (I52) traverses ALL 20 faces, coupling to all 3 colors with equal GROUP-THEORETIC weight ($C_3=0$) -- though the physical face count itself is 7R/7G/6B, not exactly equal (20 is not divisible by 3; su3_from_faces.py SU3-8; face_gluon_geometry.py's own comment: "cannot be perfectly divided into 3 equal color groups"). Since $C_3=0$ in I52, it couples to R, G, B with equal amplitude and zero net color charge:

I52 dim = 6 = 2 (Dirac spinor) x 3 (quark color from face 3-coloring)

7.2 The three C3 characters and color roles

E+ C3 char = -1 (vertex):	5 colored faces meet at each vertex. Anti-phase C3 -> electron is a color SINGLET (destructive interference of all 3 colors at vertex)
G32 C3 char = +1 (edge):	2 colored faces share each edge. In-phase C3 -> muon couples to face BOUNDARY (net color at the interface between two colored regions)
I52 C3 char = 0 (face):	1 colored face = pure color-neutral by C3. Zero C3 -> tau couples to full color content of each face but the full cycle through all 3 colors averages to zero

The muon's $C_3=+1$ is the geometric origin of its sensitivity to the quark-gluon boundary structure (consistent with mu+ capture in nuclei and muon-catalyzed fusion

coupling). The electron's C3=-1 singlet character is why it does not couple to color.

8. COMPARISON TABLE

Property	Electron (E+)	Muon (G32)	Tau (I52)
-----	-----	-----	-----
2I irrep (dim)	E+ (dim=2, j=1/2)	G32 (dim=4, j=3/2)	I52 (dim=6, j=5/2)
C3 character	-1	+1	0
Icosahedral mode	VERTEX	EDGE	FACE
Element count	V = 12	E = 30	F = 20
Deflection	$\cos(72 \text{ deg}) = 1/(2*\phi)$	$\cos(72 \text{ deg}) = 1/(2*\phi)$	GEOMETRIC: 72 deg (C5)
(Born balance)	eff from $\cos(72)$	eff from $\cos(72)$	eff from $\cos(-1/\sqrt{5})$
Deflection bias	forward ($\cos > 0$)	forward ($\cos > 0$)	path: forward (72 deg)
Bounce count N	12	5	20
Polygon N*tan	3.215 (~pi)	3.633 (not pi)	3.168 (~pi)
eff parameter	$\phi = 1.618$	$(9-\sqrt{5})/8 = 0.845$	$(1+1/\sqrt{5})/2 = 0.724$
Alpha power	α^2	α^1	α^0 (no Born contact)
phi power	ϕ^1	ϕ^2	ϕ^3
Free-spin corr	$3/4 * \alpha^2$	$Rs^2 + 2*\alpha$	$(Rs^2+2*\alpha)*\alpha$ [candidate, ~6% off exact, 2026-09-17]
T2 cancellation	no (T2 absent)	yes (T1=T2 in G32xG32)	yes (T1=T2 in I52xI52)
Color role	SINGLET (intersection)	BOUNDARY (edge)	NEUTRAL (face=T_2g surface)
Mass derived	0.510999 MeV (+0.00007%)	105.655 MeV (-0.003%)	1776.92 MeV (+0.004%)
PDG value	0.510999 MeV	105.658 MeV	1776.86 MeV
Path formula	J26 in doc_jobson_cell	Section 4 above	corkscrew: $\phi^3/\sqrt{5} * m_p$

The bipyramid model bridges muon and tau:

Bipyramid apex (72 deg) = muon deflection -> pure muon path uses ONLY these
 Bipyramid equatorial (116.57 deg) = tau deflection -> pure tau path uses ONLY these
 Mixed bipyramid eff = $(9-\sqrt{5})/8$ = Born balance BETWEEN the two modes

9. STATUS

DERIVED (zero free parameters, all verified by companion scripts):

- All three lepton = 2I spinors, assigned by spinor rank [exact, group theory]
- Electron = VERTEX, muon = EDGE, tau = FACE (icosahedral element mapping) [LM12-LM13]
- Bipyramid was superposition of muon (apex) and tau (equatorial) modes [Section 2.3]
- $m_e = 0.510999 \text{ MeV}$ (+0.000065%) J26 formula [LM1]
- $m_\mu = 105.655 \text{ MeV}$ (-0.003%) bipyramid Born balance, $\text{eff}_\mu = (9-\sqrt{5})/8$
 now SETTLED (not a fitted approximation) -- the free muon is a solo
 corpuscle correctly reverting to the bipyramid form, with the height
 ratio $h_t/r_e = 1/\phi$ proven to be the all-edges-equal condition
 independent of any mass fit (Section 4.3, muon_bipyramid_edge_condition.py
 BR1-BR3). The numerically-inverted value 0.8563 (koide_proof.py KP1) is
 a PDG-mass fit only, not a competing geometric target. [LM6-LM8]
- $m_\tau = 1776.92 \text{ MeV}$ (+0.004%) Koide from derived m_e, m_μ [LM10]
- Tau has a nexus with the gluon at each face boundary (20 per circuit) [LM14]
- Born balance deflection = $\arccos(-1/\sqrt{5})$; actual face-face dihedral 138.19 deg [JC5]
- I52 dim=6 = 2×3 = spinor x quark colors [LM15]
- Tau corkscrew base = $(1/\sqrt{5})*\phi^3*m_p = 1777.49 \text{ MeV}$ (+0.035%) [LM16]
- $\phi^3/\sqrt{5} = 1 + 2/\sqrt{5}$ (exact algebraic identity) [LM17]
- Tau winding: α^0 (no Born vertex contact), no $2*\pi$ (face flux not loop)
- T_2g = color-neutral elastic face material; $Rs^2 = T_2g$ shear amplitude [FG4,FG5]
- 2G (8-dim, C3=+1) = 8 gluons from Gamma(20 face centers) decomp [FG3]
- Tau rides T_2g surface; does not constitute it [FG7]
- $H_g = T_{1g} \times T_{2g}$ (field strength F_{mn} , not a new particle) [FG8]
- Muon rides gluon edge channels; 72-deg = gluon flux bending at C5 vertex [FG9]
- $G32 \times G = G32 + 2*I52$ (no A): muon is color-neutral in gluon sector [DG15]
- KOIDE 2/3 geometric origin: $\text{dim}(T_{1g}+T_{2g})/\text{dim}(T_{1g} \times T_{2g}) = 6/9 = 2/3$ [FG11]
- Koide holds to 9.2 ppm for raw PDG masses [LM9]; the derived-mass check
 solving Koide for m_τ given m_e/m_μ [LM10-LM11] is TAUTOLOGICAL, not
 independent confirmation. A separate, non-tautological check combining
 all three INDEPENDENTLY-derived formulas (none solved for the others)
 gives -2.85 ppm (free-lepton) and +5.74 ppm (structural-mode) deviation
 from 2/3 -- both closer than the raw-PDG figure, on two different
 physical derivations (Section 6.1). This is consistency with Koide,
 not algebraic proof of it; tau's own mass measurement (+/-0.12 MeV)
 limits any such comparison to about 10 ppm resolution (Section 6.3).

- F-7: SU(3) from icosahedral face 3-coloring [CLOSED 2026-08-23, su3_from_faces.py 11/11]

ESSENTIALLY CLOSED:

- No α^2 correction FROM THE CG-DECOMPOSITION MECHANISM for muon or tau ($T_1=T_2$ in $G_{32} \times G_{32}$, $I_{52} \times I_{52}$) [Section 1.2] -- a separate, later mechanism (Section 5.3's candidate tau correction) does introduce an α^2 term via a different, independent argument; this does not contradict THIS mechanism's own scope.
- Color coupling from C3 characters ($E+=-1$, $G_{32}=+1$, $I_{52}=0$) [Section 7]
- Euler $V-E+F=2$ and Maxwell $3V-E=6$ constrain all three path geometries [LM12-LM13]
- Lepton generations exhaust 2I spinors (3 independent spinor irreps)
- KOIDE GEOMETRIC ORIGIN: $2/3 = (\dim T_{1g}+T_{2g})/\dim(T_{1g} \times T_{2g}) = 6/9$ [FG11, proven]
- Muon color-neutrality: $G_{32} \times G$ has no A component (color-neutral by group theory) [DG15]

OPEN:

- Exact Hopf fiber flux integral through one icosahedral face (closes tau +0.035%)
- Tau correction structure (no Born vertex contact -- differs from muon's $R_s^2+2\alpha$; now numerically constrained to within 0.03% of 1, much smaller than the muon's ~4.6% -- koide_proof.py KP9-KP10). CANDIDATE (2026-09-17, koide_proof.py KP11): $(R_s^2+2\alpha)\alpha$, physically motivated by the vertex/edge/face contact-dimension ladder and the free-spin/Born-contact distinction (Section 5.3) -- gives m_τ within +0.0016% of PDG but still ~6% off Koide's own exact requirement, not yet a full derivation.
- Born balance algebraic closure: show 4442 ppm correction enforces 6/9 ratio [KP7]
- Near-critical tau correction ($N_J=11$, Maxwell critical ~21)
- Tau meeting-nexus mechanism at hop 0/10: what the two photons actually do when they coincide (continue, reverse, redirect) is OPEN. (The $N=2$ tau count itself is NOT part of this open item -- it is independently established by TN_1 's exchange-angle-rule enumeration, Section 5.1.) The broader corpuscle-billiard simulation is NOT a trusted reference for the meeting mechanism (undisclosed tau-backward shortcut in the wider sim, inconsistent gluon-per-face count); needs a full force-vector resimulation (planned billiard v3) before any claim here is derived.
- Section 4.1's $G_{32}|D_5=E_1+E_2$ branching (C5 character sum = -1) and Section 4.3/6.1's $\chi(G_{32}, C_5)=+1$ cross-check are both individually correct computations, evaluated at DIFFERENT classes of 2I's four order-5 conjugacy classes ($\chi(G_{32})$ is +1 at C_5/C_5^3 , -1 at C_5^2/C_5^4 -- all four are genuine values of the same character formula). Which class the D_5 subgroup's own generator actually corresponds to once embedded in 2I has not been traced through the explicit geometric embedding; until it is, the two results should be read as complementary data points, not as a resolved contradiction in either direction.

WHY THREE GENERATIONS (no more, no fewer):

The 2I double group has exactly 4 spinor irreps: $E+(2)$, $E-(2)$, $G_{32}(4)$, $I_{52}(6)$. $E+/E-$ are a Galois-conjugate pair distinguished by vertex-nexus coupling strength: $\chi(C_5)=+\phi$ for $E+$ (strongly constructive -> binds at the vertex nexus -> the electron) vs $\chi(C_5)=-1/\phi$ for $E-$ (not constructive -> no nexus binding -> propagates freely -> electron neutrino candidate) -- the same bound/freed-at-its-own-nexus pattern already used for G_{32}/μ on and I_{52}/τ au, not a particle/antiparticle relationship [ih_double_group.py DG11-DG14; neutrino_freed_lepton.py NL1-NL6, which independently derives G_F and the Fermi cross-section from this same picture (the mass-hierarchy PATTERN also matches; whether a freed lepton actually keeps a persistent $\chi(C_5)$ character to produce it, or $\chi(C_5)$ is only a bookkeeping label, is open)]. Counting $E+/E-$ as ONE generation (bound lepton + its own freed/neutrino partner), there are exactly 3 independent generations. These ARE the 3 lepton generations. No 4th exists in 2I.

10. COMPANION SCRIPTS

```
analysis/demos/leptons_doc.py          -- 18/18 PASS [STANDALONE -- no project imports; uses numpy for vector geom
analysis/quantum/lepton_mass.py        -- 18/18 PASS (main: all three leptons, exact formulas)
analysis/quantum/face_gluon_geometry.py -- 14/14 PASS (T_2g, 2G, H_g, Koide origin, lattice geometry)
analysis/quantum/koide_proof.py         -- 16/16 PASS (Koide gap, eff_mu inversion, geometric origin
                                         2/3=6/9 KP1-KP8; settled-eff_mu tau constraint KP9-KP10;
                                         candidate tau correction, speculative, KP11; chi(G32,C5)/
```

chi(I52,C5) cross-checks KP12-14; non-tautological free vs structural check KP15; measurement-precision KP16)

analysis/nuclear/ih_double_group.py -- 15/15 PASS (2I character table, spinor irreps, G32xG=G32+2*I52 DG15)

analysis/quantum/gluon_tau_helix.py -- 10/10 PASS (tau path geometry, GH0-GH0c, GH6-GH7)

analysis/quantum/tau_pair_configuration.py -- 11/11 PASS (TPC1-TPC6: tau circuit uniqueness, two-photon meeting geometry -- NOT currently trusted, see Section 9 OPEN item)

analysis/corpuscle/tau_necessity_verify.py -- 12/12 PASS (TN1 only: N=2 tau count independently established by exchange-angle-rule enumeration, trusted; TN4/TN5 gluon-timing claims NOT trusted, see Section 9 OPEN item)

analysis/quantum/su3_from_faces.py -- 11/11 PASS (F-7: SU(3) from face 3-coloring)

LM1-2: Electron J26 reference (+0.000065%)

LM3-5: Pentagonal bipyramid deflection geometry (exact, $h_t/r_e = 1/\phi$)

LM4b: Actual icosahedral path: all deflections = 72 deg (uniform C5)

LM6-8: Muon mass from pentagon Born balance (-0.003%)

LM9: Koide formula (9.2 ppm)

LM10-11: Tau from Koide (derived m_e, m_μ , +0.004%)

LM12-13: Icosahedral Euler/Maxwell constraints (V=12, E=30, F=20)

LM14: Tau dihedral = $\arccos(-1/\sqrt{5})$ (exact)

LM15: I52 dim=6 = 2^3 (spinor x quark colors)

LM16: Tau corkscrew base = $\phi^3/\sqrt{5} * m_p$ (+0.035%, leading order; correction open)

LM17: $\phi^3/\sqrt{5} = 1 + 2/\sqrt{5}$ (tau contains muon winding, exact)

FG1-3: Gamma(20 face centers) = A+T1+T2+2G+H; 2G=8 gluons

FG4-5: $T_{2g} C5 = -1/\phi$ (shear), $Rs^2 = T_{2g}$ shear amplitude at Maxwell critical

FG6-7: G C3=+1 (gluon color-active), T2 C3=0 (face elastic, color-neutral)

FG8: $T1g \times T2g = G + H$ (H_g = field strength, not a new particle)

FG9: Muon 72-deg deflection = gluon edge-channel bending at vertex (exact)

FG10: G32 (muon) and G (gluon) share C3=+1 -- C3 coupling locks muon to gluon channels

FG11: Koide 2/3 = (dim T1g + dim T2g) / dim(T1g x T2g) = 6/9 [EXACT]

FG12-14: V=12 from spinor sum 2+4+6; E=30 from C5; F=20 from Euler V-E+F=2

DG15: G32 x G = G32 + 2*I52, no A (muon color-neutral; ih_double_group.py 15/15 PASS)

KP1-2: Exact $eff_\mu = 0.8563$ by inversion (bipyramid approx 0.8455 = 1.3% off)

KP3: base_tau/ m_p scenario B vs $\phi^3/\sqrt{5}$ (0.28% off: needs Hopf flux)

KP4-5: Koide holds to 9.2 ppm; algebraic form $a^2+b^2+c^2=4(ab+bc+ca)$ to 0.003%

KP6: Koide texture angle $\theta = 132.73$ deg; m_e reconstruction to 0.001%

KP7: Leading-order $K0 = 4442$ ppm above 2/3 (proof needs exact Born balances)

KP8: Geometric origin: $2/3 = (\dim T1g+T2g)/\dim(T1g \times T2g) = 6/9$ exactly [FG11]

KP9-10: Settled $eff_\mu \rightarrow$ required tau correction factor within 0.03% of 1

KP11: Candidate tau correction ($Rs^2+2*\alpha$)* α [speculative, ~6% off] combined with muon's own analogous test; both physically motivated, neither confirmed -- see Section 5.3

KP12-14: chi(G32,C5)/chi(I52,C5) group-character cross-check vs the free and structural eff_μ/eff_τ values -- see Section 4.3, 5.2

KP15: Non-tautological Koide check, free and structural masses each independently computed -- see Section 6.1

KP16: Measurement-precision propagation (tau's own PDG uncertainty limits any Koide comparison to ~10 ppm) -- see Section 6.3

REFERENCES

[doc_alpha]	https://doi.org/10.5281/zenodo.22013651	[alpha, Born balance]
[doc_jobson_cell]	https://doi.org/10.5281/zenodo.22032906	[J26 electron mass]
[doc_nucleus]	https://doi.org/10.5281/zenodo.22042337	[2I CG, irrep assignments]
[doc_leptons]	https://doi.org/10.5281/zenodo.22057421	[this document]
[doc_muon.txt]	superseded by this document	
[doc_tau.txt]	superseded by this document	

ACKNOWLEDGMENT

This research was developed with the assistance of AI language models (GitHub Copilot / Claude, Anthropic). The conceptual framework, physical interpretation, and all claims are the sole work of the author.

Repository: <https://doi.org/10.5281/zenodo.22105622>

Code: https://github.com/Destraag/torsionverse_public