

**Bachelor Level / First Year/ First Semester/ Science
Computer Science and Information Technology (MTH112)
(Mathematics I)
(OLD COURSE)**

**Full Marks: 80
Pass Marks: 32
Time: 3 hours.**

Candidates are required to give their answers in their own words as far as practicable. The figures in the margin indicate full marks.

The figures in the margin indicate full marks.
Group A $(3 \times 10 = 30)$

Group A (3 × 10 = 30).

Attempt any THREE questions:

1. (a) If $f(x) = 3x^2 - x + 2$, then find $f(2)$, $f(-2)$, $f(a)$, $f(-a)$ and $f(0)$. [5]

(b) Find $\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$ [5]

2. (a) Find the derivative of $y = \frac{x^2 + x - 2}{x^3 + 6}$. [5]

(b) Estimate the area between the parabola $y = x^2$ and the line $y = x$. [5]

3. (a) Verify the Mean Value Theorem for the function $f(x) = 3x^2 + 2x + 5$ $x \in [-1, 1]$. [4]

(b) Define initial value problem. Solve the equation $xy' + 2y = 3x$, $y(1) = 2$. [4]

(c) Find the volume of a sphere of radius a . [2]

4. (a) Find the local maximum and minimum values and saddle points of $f(x, y) = x^4 + y^4 - 4xy + 1$. [5]

(b) Find the curvature of the parabola $y = x^2$ at the points $(0, 0)$, $(0, 1)$ and $(2, 4)$. [5]

GUPTA TUTORIAL

Group B $(10 \times 5 = 50)$.

Attempt any TEN questions:

5. Verify Rolle's theorem for $f(x) = x^2 - 4$, $x \in [-2, 2]$. [5]

6. Find the Maclaurin series expansion of $\sin x$ at $x = 0$. [5]

7. If $f(x) = \sqrt{x-2}$ and $g(x) = \sqrt{x}$, find $(f \circ g)(x)$ and $(g \circ f)(x)$. [5]

8. Show that the absolute value function $f(x) = |x|$ is continuous everywhere. [5]

9. Find $\int_0^{\frac{1}{2}} \frac{x}{\sqrt{1-4x^2}} dx$. [5]

10. Sketch the curve $y = x^3$. [5]

11. Find the solution of $y'' + 4y' + 4 = 0$, $y(0) = 2$, $y'(0) = 1$. [5]

12. Test the series $\sum_{n=1}^{\infty} \frac{n^2}{5n^2 + 4}$ diverges or converges. [5]

GUPTA TUTORIAL

13. Define cross product of two vectors. If $\vec{a} = \vec{i} + 2\vec{j} + 3\vec{k}$ and $\vec{b} = 2\vec{i} + 3\vec{j} - 4\vec{k}$, find $\vec{a} \times \vec{b}$ and $\vec{b} \times \vec{a}$. [5]
14. Find the second partial derivatives of $f(x, y) = x^3 + x^2y^3 - 2y^2$. [5]
15. Find the length of the arc of the semi-cubical parabola $y^2 = x^3$ between the points $(1, 1)$ and $(4, 8)$. [5]

2080 (odd Syllable) Questions Solution

Subj:- Mathematics - I

- Pravin Gupta

GOOD MORNING
PAGE NO. _____
DATE: _____

1(a)

Given

$$f(n) = 3n^2 - n + 2 \quad \dots \dots \dots (1)$$

Find $f(2) = ?$ $f(-2) = ?$ $f(a) = ?$
 $f(-a) = ?$ $f(0) = ?$

Now,

for $f(2) = ?$

if we put $n = 2$ in eq¹(1)

$$\begin{aligned} f(2) &= 3n^2 - n + 2 \\ &= 3 \times 2^2 - 2 + 2 \end{aligned}$$

$$\therefore f(2) = 12$$

GUPTA TUTORIAL

for $f(-2)$; put $n = -2$ then,

$$\begin{aligned} f(-2) &= 3 \times (-2)^2 - (-2) + 2 \\ &= 12 + 2 + 2 \end{aligned}$$

$$\therefore f(-2) = 16$$

for $f(a)$; put $n = a$ then

$$f(a) = 3a^2 - a + 2$$

$$\therefore f(a) = 3a^2 - a + 2$$

for $f(-a)$; put $n = -a$ then

$$f(-a) = 3 \times (-a)^2 - (-a) + 2$$

$$\therefore f(-a) = 3a^2 + a + 2$$

for $f(0)$; put $n = 0$ then,

$$f(0) = 3 \times 0 - 0 + 2$$

$$\therefore f(0) = 2$$

Therefore,

$$f(2) = 12$$

$$f(-a) = 3a^2 + a + 2$$

$$f(-2) = 16$$

$$f(0) = 2$$

$$f(a) = 3a^2 - a + 2$$

AM



b Given,

$$\lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x}$$

$$= \frac{x(-2)^3 + 2x(-2)^2 - 1}{5 - 3x(-2)}$$

$$= \frac{-8 + 8 - 1}{5 + 6}$$

$$= \frac{-1}{11}$$

$$\therefore \lim_{x \rightarrow -2} \frac{x^3 + 2x^2 - 1}{5 - 3x} = \frac{-1}{11} \quad \text{Ans}$$

20) Given,

GUPTA TUTORIAL

$$y = \frac{x^2 + x - 2}{x^3 + 6} \quad \dots \quad (1)$$

Diff' eq (1) w.r.t 'x'

$$\frac{d}{dx} \left(\frac{x^2 + x - 2}{x^3 + 6} \right) = \frac{(x^3 + 6) \frac{d}{dx}(x^2 + x - 2) - (x^2 + x - 2) \frac{d}{dx}(x^3 + 6)}{(x^3 + 6)^2}$$

$$= \frac{(x^3 + 6)(2x + 1) - (x^2 + x - 2)(3x)}{(x^3 + 6)^2}$$

$$= \frac{2x^4 + x^3 + 12x + 6 - 3x^3 - 3x^2 + 6x}{(x^3 + 6)^2}$$

$$= \frac{2x^4 - 2x^3 - 3x^2 + 18x + 6}{(x^3 + 6)^2}$$

$$\therefore \text{The derivative of } y = \frac{x^2 + x - 2}{x^3 + 6} \text{ is } \frac{(2x^4 - 2x^3 - 3x^2 + 18x + 6)}{(x^3 + 6)^2}$$

2(b)

Given,

the parabola is $y = x^2 - 1$ and the line is $y = x + b$

Solving eq(1) & (2), we get limits of integration

$$x^2 = x$$

$$\text{or, } x^2 - x = 0$$

$$\text{or, } x(x-1) = 0$$

$$\therefore x = 0 \text{ and } x = 1$$

The area between the curves is

$$A = \int_a^b (f(x) - g(x)) dx$$

$$= \int_0^1 (x^2 - x) dx$$

$$= \left[\frac{x^3}{3} - \frac{x^2}{2} \right]_0^1$$

$$= \frac{1}{3} - \frac{1}{2}$$

$$= \frac{2-3}{6}$$

$$= -\frac{1}{6} \text{ square unit}$$

\therefore The area can't be negative so $\frac{1}{6}$ square unit

3(a) Statement of Mean Value Theorem

Let f be a function that satisfies the following conditions.

- (i) f is continuous on the closed interval $[a, b]$
- (ii) f is differentiable on the open interval (a, b)

Then there is a number c in (a, b) such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

GUPTA TUTORIAL

Given,

$$f(x) = 3x^2 + 2x + 5 \quad x \in [-1, 1]$$

- (i) Since $F(x)$ is a polynomial function. So, it is continuous on the closed interval $[-1, 1]$

- (ii) $f'(x) = 6x + 2$ which exists for all $x \in (-1, 1)$
 $\therefore f(x)$ is differentiable on the open interval

All the condition of Mean Value Theorem are satisfied hence, there exist a number c in $(-1, 1)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f'(x) = 6x + 2$$

$$f'(c) = 6c + 2$$

$$f(x) = 3x^2 + 2x + 5$$

$$f(a) = f(1) = 3 \times 1^2 + 2 \times 1 + 5 = 10$$

$$f(b) = f(-1) = 3 \times (-1)^2 + 2 \times (-1) + 5 = 3 - 2 + 5 = 6$$

Then,



$$6C + 2 = \frac{10}{6}$$

$$\text{or, } 2(3C+1) = \frac{10}{3}$$

$$\text{or, } 18C + 6 = 5$$

$$\text{or, } 18C = 5 - 6$$

$$\text{or, } 18C = -1$$

$$\therefore C = \frac{-1}{18} \in (-1, 1)$$

$$\text{Hence, } C = \frac{-1}{18} \in (-1, 1)$$

∴ Mean Value theorem verify.

so?

1st part.

Q.N;3(b)

A differential equation together with initial condition(s) is called the initial value.

For example:-

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 0, \quad y(0) = -1, \quad y'(0) = 1$$

Here, $y(0) = -1$ and $y'(0) = 1$ is an initial condition.

36

2nd part,

Given

$$xy' + 2y = 3x \quad , \quad y(1) = 2$$

$$\frac{dy}{dx} + \frac{2}{x}y = 3$$

dividing x on both sides, we get

$$\frac{dy}{dx} + \frac{2}{x}y = 3 \quad \dots \dots \dots \textcircled{1}$$

Comparing eq² ① with $\frac{dy}{dx} + Py = Q$ where, $P = \frac{2}{x}$, $Q = 3$

Now,

$$\text{Integrating factor (I.F)} = e^{\int P dx} = e^{\int \frac{2}{x} dx} = e^{2 \int \frac{1}{x} dx} = e^{2 \log x} = x^2$$

Multiplying eq² ① by I.F

$$x^2 \left[\frac{dy}{dx} + \frac{2}{x}y \right] = 3x^2$$



$$d(y \cdot n^2) = 3n^2$$

on integrating both sides

$$\int d(y \cdot n^2) = \int 3n^2$$

$$\text{or, } y \cdot n^2 = 3n^2 + C$$

$$\text{or, } y \cdot n^2 = \frac{3n^3}{3}$$

$$y = \frac{n^3 + C}{n^2} \quad \dots \text{(i)}$$

SINCE,

$$y(1) = 2$$

$$y = 2, n = 1$$

then eq(i) becomes

$$2 = 1 + C$$

$$2 = 1 + C$$

$$\therefore C = 1$$

Substituting the value of C in eq(i)

$$y = \frac{n^3 + 1}{n^2}$$

$$\text{or, } y = \frac{n^3}{n^2} + \frac{1}{n^2}$$

$$\therefore y = n + \frac{1}{n^2} \quad \text{Ans}$$

3 (c) Find the volume of sphere of radius 'a'
Sol Since, we know that intersection of sphere of radius 'a' and the plane is of a circle of radius 'a'

Then circle is

$$x^2 + y^2 = a^2 \\ y^2 = a^2 - x^2 \quad \dots \dots \text{①}$$

clearly the circle has end $y = -a$ to $y = a$. Now,
 the volume of the solid that is generated by revolving the circle ① about x -axis is (i.e. $y = 0$)

GUPTA TUTORIAL

$$\begin{aligned} \text{Volume} &= \pi \int_{-a}^a y^2 dx \\ &= \pi \int_{-a}^a (a^2 - x^2) dx \\ &= \pi \left[x^2 - \frac{x^3}{3} \right]_{-a}^a \\ &= \pi \left\{ \left(a^2 \cdot a - \frac{a^3}{3} \right) - \left(a^2 \cdot (-a) - \frac{(-a)^3}{3} \right) \right\} \\ &= \pi \left\{ \left(a^3 - \frac{a^3}{3} \right) - \left(-a^3 + \frac{a^3}{3} \right) \right\} \\ &= \pi \left\{ \frac{3a^3 - a^3}{3} - \left(-3a^3 + a^3 \right) \right\} \\ &= \pi \left(\frac{2a^3}{3} + \frac{2a^3}{3} \right) = \pi \times \frac{4a^3}{3} = \frac{4\pi a^3}{3} \text{ Ans} \end{aligned}$$

\therefore The volume of the sphere of radius 'a' is $\frac{4\pi a^3}{3}$ Ans

Q.N;4(a)

Given function,

$$f(x,y) = x^4 + y^4 - 4xy + 1$$

so,

$$f_x = 4x^3 - 4y$$

$$f_y = 4y^3 - 4x$$

$$f_{xy} = -4$$

$$f_x = 4y^3 - 4x$$

$$f_{yy} = 12y^2$$

$$f_{yx} = -4$$

for stationary point

$$f_x = 0$$

$$4x^3 - 4y = 0$$

$$x^3 = y \quad \dots \textcircled{i}$$

$$f_y = 0$$

$$4y^3 - 4x = 0$$

$$y^3 = x \quad \dots \textcircled{ii}$$

GUPTA TUTORIAL

from eqⁿ \textcircled{i} and \textcircled{ii}

$$(x^3)^3 = x \quad (\therefore y = x^3)$$

$$\text{or, } x^9 = x$$

$$\text{or, } x^9 - x = 0$$

$$\text{or, } x(x^8 - 1) = 0$$

$$\therefore x = 0, x = 1$$

Substitute the point value of y in \textcircled{i}

$$\text{when } x=0, y=0$$

$$\text{" } \quad x=1, y=1$$

At point (0,0)

$$\begin{aligned}f_{xx} &= 0 \\f_{yy} &= 0\end{aligned}$$

Then,

$$\begin{aligned}D &= \begin{vmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{vmatrix} \\&= \begin{vmatrix} 0 & -4 \\ -4 & 0 \end{vmatrix} = 0 - 16 = -16 < 0\end{aligned}$$

so, it has saddle point at (0,0)

GUPTA TUTORIAL

At point (1,1)

$$f_{xx} = 12, f_{yy} = 12$$

$$D = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{vmatrix} = \begin{vmatrix} 12 & -4 \\ -4 & 12 \end{vmatrix} = 144 - 16 = 128 > 0$$

so, it has minimum value at ~~(1,1)~~ (1,1)

∴ Minimum value is

$$\begin{aligned}f(1,1) &= 1^4 + 1^4 - 8 \times 1 \times 1 + 1 \\&= 1 + 1 - 8 + 1 \\&= -1\end{aligned}$$

4b Given,

Parabola is $y = x^2$ --- (1)
and the points are $(0,0), (0,1)$ & $(2,4)$

For curvature we are using formula as;

$$K = \frac{1}{|\vec{r}|} \left| \frac{d\vec{r}}{dt} \right| \quad \text{--- (ii)}$$

diff' eq (1) w.r.t 'x'

i.e

$$\vec{r} = \frac{\partial \vec{y}}{\partial x} = 2x \quad \text{--- (iii)}$$

$ \vec{r} = \sqrt{2^2} = 2$	at $(0,0)$ $ \vec{r} = 0$	at $(0,1)$ $ \vec{r} = 0$	at $(2,4)$ $ \vec{r} = \sqrt{4^2} = 4$
------------------------------	-------------------------------	-------------------------------	--

GUPTA TUTORIAL

Again, diff' eq (1) w.r.t 'x'

$$a(x) = 2$$

Then,

$$\text{at } (2,4) \quad K = \frac{1 \times 2}{4} = \frac{1}{2}$$

at $(0,0)$ & $(0,1)$

$$K = \frac{1 \times 2}{0}$$



5. Given,

$$f(x) = x^2 - 4, \quad x \in [-2, 2]$$

(i) Since $f(x)$ is a polynomial function. So, it is continuous on the closed interval $[-2, 2]$

(ii) $f'(x) = 2x$ which exists for all $x \in (-2, 2)$
 $\therefore f(x)$ is differentiable in $(-2, 2)$

GUPTA TUTORIAL

(iii) $f(-2) = (-2)^2 - 4 = 0$
 $f(2) = 2^2 - 4 = 0$
 $\therefore f(-2) = f(2)$

\therefore All the condition of Rolle's theorem are satisfied Hence.
 There exists at least a point $c \in (0, 3)$
 such that

$$\begin{aligned} f'(c) &= 0 \\ \text{or, } 2c &= 0 \\ \therefore c &= 0 \in (-2, 2) \end{aligned}$$

Hence, Rolle's theorem verified an $0 \in (-2, 2)$ Ans



Sol

The given function is

$$f(n) = \sin n \quad \text{--- (1)}$$

Q.N;6

We have to find the Taylor's series generated by the function $f(n) = \sin n$ at $n=0$.

• diff eq⁷ O w.r.t. 'n' successively we get.

$$f(n) = \sin n$$

$$f'(n) = \cos n$$

$$f''(n) = -\sin n$$

$$f'''(n) = -\cos n$$

$$f^{IV}(n) = \sin n$$

$$f^V(n) = \cos n$$

$$f^VI(n) = -\sin n$$

rearrangement all derivative as below.

$$f'(n) = \cos n$$

$$f''(n) = -\sin n$$

$$f'''(n) = -\cos n$$

$$f^{IV}(n) = \sin n$$

$$f^V(n) = \cos n$$

$$f^VI(n) = -\sin n$$

GUPTA TUTORIAL

$$f^{2n+1}(n) = (-1)^n \cos n$$

$$f^{2n}(n) = (-1)^n \sin n$$

at $n=0$

$$f(0) = \cos 0 = 1$$

$$f''(0) = -\sin 0 = 0$$

$$f'''(0) = -\cos 0 = -1$$

$$f^{IV}(0) = \sin 0 = 0$$

$$f^V(0) = \cos 0 = 1$$

$$f^VI(0) = -\sin 0 = 0$$

$$f^{2n+1}(0) = (-1)^n$$

$$f^{2n}(0) = 0$$

we know,

$$f(n) = f(a) + f'(a)(n-a)^1 + f''(a)(n-a)^2 + f'''(a)(n-a)^3 + \dots + n^m \text{ term} + \dots$$

$$\sin n = 0 + \frac{1}{1!}(n-0)^1 + 0 + \frac{(-1)}{2!}n^3 + \dots - \frac{n^5}{3!} + \dots + \frac{n^m}{m!} + \dots$$

$$\sin n = \frac{n}{1!} - \frac{1}{3!} \frac{n^3}{3!} + \frac{n^5}{5!} - \frac{n^7}{7!} - \dots - \frac{3!}{(-1)^{\frac{n-1}{2}} \frac{(n+1)!}{(n+1)!}} + \dots \neq$$

7 Given,

$$f(x) = \sqrt{x-2} \quad \& \quad g(x) = \sqrt{x}$$

then,

$$fog(x) = ? \quad gof(x) = ?$$

we know,

$$\begin{aligned} fog(x) &= f(g(x)) \\ &= f(\sqrt{x}) \\ &= \sqrt{\sqrt{x}-2} \end{aligned}$$

$$\therefore fog(x) = \sqrt{\sqrt{x}-2}$$

Again for $gof(x)$

GUPTA TUTORIAL

$$\begin{aligned} gof(x) &= g(f(x)) \\ &= g(\sqrt{x-2}) \\ &= \sqrt{\sqrt{x-2}} \end{aligned}$$

$$\therefore gof(x) = \sqrt{\sqrt{x-2}}$$

$$\therefore fog(x) = \sqrt{\sqrt{x-2}} \quad \left. \right\} \text{Ans}$$

$$gof(x) = \sqrt{\sqrt{x-2}}$$



8

A function $f(x)$ is continuous at a point $x=a$ if the following conditions are satisfied.

- (i) $f(a)$ is defined
- (ii) $\lim_{n \rightarrow a} f(n)$ exists
- (iii) $\lim_{n \rightarrow a} f(n) = f(a)$

GUPTA TUTORIAL

For the absolute value function $f(x)=|x|$

- (i) $f(a)$ is defined for all real numbers
- (ii) The limit of the function as n approaches a exists for all real numbers a . Since the function is piecewise defined as $f(n)=n$ for $n \geq 0$ and $f(n)=-n$ for $n < 0$, the limit from both the right and the left as n approaches a is a .
- (iii) The limit of the function as n approaches a is equal to the value of the function at $n=a$ for all real numbers a . This is true because the function's value at $n=a$ is either a or $-a$, depending on the sign of a .

∴ The absolute value satisfies all three conditions for continuity at every point so, it is continuous everywhere on the real number.

∴ $f(x)=|x|$ is continuous everywhere.

9. Given, $\int_0^{1/2} \frac{x}{\sqrt{1-4x^2}} dx$

let,

$$\begin{aligned} y &= 1-4x^2 \\ dy &= -8x dx \\ -\frac{1}{8} dy &= x dx \end{aligned}$$

then,

$$\int_0^{1/2} \frac{-y}{\sqrt{y}} dy = -\frac{1}{8} \int_0^{1/2} \frac{dy}{\sqrt{y}} = -\frac{1}{8} \int_0^{1/2} y^{-1/2} dy$$

$$= -\frac{1}{8} \left[\frac{y^{1/2}}{\frac{1}{2}+1} \right]_0^{1/2}$$

GUPTA TUTORIAL

$$\begin{aligned} &= -\frac{1}{8} \left[\frac{y^{1/2}}{\frac{3}{2}} \right]_0^{1/2} \\ &= -\frac{1}{4} \left[(\sqrt{y})^{1/2} \right]_0^{1/2} \\ &= -\frac{1}{4} \left[\sqrt{1-4x^2} \right]_0^{1/2} \\ &= -\frac{1}{4} \left[(\sqrt{1-4 \times (\frac{1}{2})^2}) - \sqrt{1-0} \right] \\ &= -\frac{1}{4} (\sqrt{0} - 1) \\ &= \frac{1}{4} \end{aligned}$$



To

Given,

$$f(x) = y = x^3 \quad \dots \text{ (1)}$$

(i) Domain:-

For domain; The function exists for all x . So, the domain is

$$\text{Domain} = (-\infty, \infty)$$

diff' eq (1) w.r.t 'x'

$$f'(x) = 3x^2$$

for critical point

$$f'(x) = 0$$

$$3x^2 = 0$$

$$\therefore x = 0$$

GUPTA TUTORIAL

Interval	$(-\infty, 0)$	$(0, \infty)$
Sign of $f'(x)$	+ve	+ve
Nature of f	increasing	increasing

Thus, the ~~maxima~~ ^{minima} occur at $x = 0$

$f(0) = 0$ is ^{minimum} ~~maximum~~ point

Again,

$$f''(x) = 6x$$

for Point of inflection,

$$f''(x) = 0$$

$$6x = 0$$

$$\therefore x = 0$$



Interval	$(-\infty, 0)$	$(0, \infty)$
Sign of f''	-ve	+ve
Nature of f	Concave down	Concave up


Thus, Point of inflection occur at $x=0$.

So,

$f(0) = 0$ is the point of inflection.

Summarizing above two tables,

$(-\infty, 0)$	$(0, \infty)$
+ve Increasing	increasing
Concave down	Concave up



11 The given 2nd order homogenous linear eqn is.

$$y'' + 7y' + 14y = 0 \quad \text{--- (1)}$$

$$y'' + 7y' + 14y = 0 \quad \text{--- (1)}$$

Let

$y = e^{mx}$ be the reqⁿ solⁿ of eqⁿ (1)

We get an auxiliarily eqⁿ of (1) by replacing y'' by m^2 , y' by m and y by 1.

Then,

$$m^2 + 7m + 14 = 0$$

$$(m+2)^2 = 0$$

$m = -2$ which is real & equal.

The reqⁿ solⁿ is

$$y = e^{mx} (C_1 + C_2 x)$$

$$y = e^{-2x} (C_1 + C_2 x) \quad \text{--- (II)}$$

Since, $y(0) = 2$ **GUPTA TUTORIAL**

if $y=2$, $x=0$ then eqⁿ (II) becomes

$$y = e^{-2x} (C_1 + C_2 x)$$

$$2 = e^0 (C_1 + C_2 \times 0)$$

$$C_1 = 2 \quad \text{--- (III)}$$

Again,

diff' eqⁿ (II) w.r.t 'x'

$$y' = e^{-2x} (0 + C_2) + (C_1 + C_2 x) (-2) \times e^{-2x}$$

$$y' = e^{-2x} \times C_2 - 2 e^{-2x} (C_1 + C_2 x) \quad \text{--- (IV)}$$



Since, $y'(0)=1$

If $y'=1$, $n=0$ then eqⁿ(v) becomes,

$$y' = e^{-2x} C_2 - 2e^{-2x} (C_1 + C_2 n)$$

$$\text{or, } 1 = e^0 \times C_2 - 2 \times e^0 (C_1 + C_2 \times 0)$$

$$\text{or, } 1 = C_2 - 2 \times C_1 \quad (\text{from eq } \text{viii})$$

$$\text{or, } 1 = C_2 - 2 \times 2$$

$$\text{or, } C_2 = 1+4$$

$$\therefore C_2 = 5$$

Substituting the value of C_2 & C_1
then eqⁿ(v) becomes.

$$y = e^{-2x} (2 + 5n) \quad \underline{\text{Ans}}$$



Given Series is

Q.N:12

$$\sum_{n=1}^{\infty} \frac{n^2}{5n^2 + 1}$$

i.e. $\sum_{n=1}^{\infty} a_n$

where,

$$a_n = \frac{n^2}{5n^2 + 1}$$

Now,

$$\lim_{n \rightarrow \infty} \frac{n^2}{5n^2 + 1}$$

$$= \lim_{n \rightarrow \infty} \frac{n^2}{n^2(5 + \frac{1}{n^2})}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{5 + \frac{1}{n^2}}$$

$$= \frac{1}{5 + 0}$$

$$= \frac{1}{5}$$

GUPTA TUTORIAL

\therefore The Given Series $\sum_{n=1}^{\infty} \frac{n^2}{5n^2 + 1}$ is divergent by
 n^{th} Root Test.



sol

Ist part

Q.N;13

let \vec{a} and \vec{b} are two non-zero vectors.
Then the cross product of \vec{a} and \vec{b} is denoted
by $\vec{a} \times \vec{b}$ and is defined as

$$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin\theta \hat{n}$$

where θ be the angle between \vec{a} and \vec{b} and \hat{n} is
unit vector along $(\vec{a} \times \vec{b})$

and

13

2nd Part,

Given,

$$\vec{a} = \vec{i} + 2\vec{j} + 3\vec{k}$$

$$\vec{b} = 2\vec{i} + 3\vec{j} - \vec{k}$$

Then,

$$\vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 2 & 3 \\ 2 & 3 & -1 \end{vmatrix}$$

$$= \vec{i}(-8-9) - \vec{j}(1-6) + \vec{k}(3-4)$$

$$= -17\vec{i} + 10\vec{j} - \vec{k}$$

GUPTA TUTORIAL

And we know,

$$(\vec{a} \times \vec{b}) = -(\vec{b} \times \vec{a})$$

So,

$$\vec{b} \times \vec{a} = -(\vec{a} \times \vec{b})$$

$$= -\{-17\vec{i} + 10\vec{j} - \vec{k}\}$$

$$= 17\vec{i} - 10\vec{j} + \vec{k}$$

$$\therefore (\vec{a} \times \vec{b}) = (-17\vec{i} + 10\vec{j} - \vec{k})$$

$$(\vec{b} \times \vec{a}) = (17\vec{i} - 10\vec{j} + \vec{k})$$



14 Given.

$$f(x,y) = x^3 + x^2y^3 - 2y^2 \quad \dots \quad (1)$$

diff' eq'(1) w.r.to 'n' partially,

$$\frac{\delta f}{\delta x} = 3x^2 + 2xy^3 \quad \text{Ans} - - - (11)$$

diff eq¹ w.r. to 'y' Partially

$$\frac{\delta f}{\delta y} = 3y^2n^2 - 4y \quad \text{Ans} \quad \text{--- (11)}$$

diff' eq⁽ⁱⁱ⁾ w.r. to 'x' Partially

$$\frac{\partial^2 f}{\partial x^2} = 6x + 2y^3 \quad \text{Ans}$$

GUPTA TUTORIAL

diff' eq' (iii) w.r. to 'y' Partially

$$\frac{dy}{dx} = 6yx^2 - 4 \quad \text{Ap}$$

diff' eq'(ii) w.r.to 'y' partially

$$\frac{\partial f}{\partial x} = 6xy^2 \quad \text{Ans}$$

diff' eg? (iii) w.r. to 'n' Partially

$$\frac{\delta f}{\delta y} = 6xy^2 \quad \text{Ans}$$

Sol Given $y^2 = x^3$

$$y^2 = x^3$$

S.taking root on both sides

$$\sqrt{y^2} = \sqrt{x^3}$$

or, $y = \sqrt{x^3}$

$$\therefore y = x^{3/2} \quad \dots \text{--- (1)}$$

diff eq (1) w.r.t x

$$\frac{dy}{dx} = \frac{3}{2} x^{3/2 - 1}$$

$$\therefore \frac{dy}{dx} = \frac{3}{2} x^{1/2}$$

Now, using arc length formula,

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

GUPTA TUTORIAL

$$L = \int_1^4 \sqrt{1 + \left(\frac{3}{2} x^{1/2}\right)^2} dx$$

$$L = \int_1^4 \sqrt{1 + \frac{9}{4} x^{3/2}} dx$$

$$L = \int_1^4 \sqrt{1 + \frac{9}{4} x} dx$$

Date:

Page:

Let

$$z = I + \frac{9}{4} n$$

$$dz = \frac{9}{4} dn \Rightarrow \frac{4}{9} dz = dn$$

changing limit 4

$$\Rightarrow \text{if } n=1, z = I + \frac{9}{4} = \frac{13}{4}$$

$$\text{if } n=4, z = I + \frac{9 \times 4}{4} = 10$$

Now,

$$I = \int_{\frac{13}{4}}^{10} \sqrt{z} \times \frac{4}{9} dz = \int_{\frac{13}{4}}^{10} (z)^{\frac{1}{2}} \times \frac{4}{9} dz$$

$$= \frac{4}{9} \left[\frac{z^{\frac{3}{2}}}{\frac{3}{2} + 1} \right]_{\frac{13}{4}}^{10}$$

$$= \frac{4 \times 2}{9 \times 3} [z^{\frac{3}{2}}]_{\frac{13}{4}}^{10}$$

$$= \frac{8}{27} [10^{\frac{3}{2}} - (\frac{13}{4})^{\frac{3}{2}}]$$

$$= \frac{7}{27} (8 \times \sqrt{10^3} - 8 \times \sqrt{(\frac{13}{4})^3})$$

$$= \frac{7}{27} (8 \times 10 \sqrt{10} - 8 \times \frac{13}{4} \sqrt{\frac{13}{4}})$$

$$= \frac{7}{27} (80\sqrt{10} - 26\sqrt{\frac{13}{4}}) \quad \cancel{4}$$

or,

$$= \frac{7}{27} (80\sqrt{10} - 26 \times \frac{1}{2} \sqrt{13}) \Rightarrow \frac{7}{27} (80\sqrt{10} - 13\sqrt{13})$$