

Stability Report · Dissolution Data Summary

This report summarizes dissolution testing data collected during stability studies conducted under ICH Q1A(R2) conditions for the drug product. Dissolution testing was performed per USP <711> using the paddle method (Apparatus 2, 50 rpm, 900 mL phosphate buffer pH 6.8, 37°C).

Dissolution Specification

The release specification requires that the dissolution result meets the criterion of NMT 80% at 30 minutes (see Dissolution Test Procedures, Table A). This criterion represents the Q value established during development and approved in the regulatory filing. The criterion applies to each individual unit tested as well as the calculated mean across the sampled units.

Any sample with a total average dissolution rate below the individual minimum values reported in Table A of the Dissolution Test Procedures document would represent an arithmetic inconsistency requiring review.

Cross-Reference

For detailed dissolution measurements, refer to the Dissolution Test Procedures document (doc_y.docx), specifically:

- Table A: Individual Unit Dissolution Results and Total Average
- Table B: Batch Release Data and Maximum Release Value

These tables contain the measured dissolution values compared against the acceptance criteria NMT 80% at 30 minutes stated in this Stability Report.

ICH References

- ICH Q1A(R2): Stability Testing of New Drug Substances and Products
- ICH Q1E: Evaluation for Stability Data
- USP <711>: Dissolution