


SPhOH

SAUDI PHYSICS OLYMPIAD HUB



Trigonometry I

Olympiad Problem Set

Olympiad Training • Math

SPhOH Team

FIRST EDITION • 2026

Contents

About This Problem Set	ii
1 Trigonometry I	1
1.1 Trigonometric Identities	1
1.2 Right-Triangle Trigonometry	3
1.3 Special Triangles and Geometry	5
1.4 Similarity and Trigonometric Ratios	6
1.5 Trigonometry in Physics	7
1.6 Geometric Applications	9
1.7 Vectors and Trigonometric Synthesis	10
2 Hints	13

About This Problem Set

This problem set is part of the training resources of the **Saudi Physics Olympiad Hub (SPhOH)**.

It is designed primarily for students preparing for the Saudi Physics Olympiad and related advanced physics competitions. The purpose of the handout is to provide a focused collection of problems that develops trigonometric fluency and its application to physical problems.

The handout is **problem-centered**. Explanations are intentionally brief. The main purpose is to provide deliberate problem-solving practice.

How to Use the Problem Set

Students are encouraged to attempt each problem independently before looking at its hint. A hint should be treated as a way to overcome a specific obstacle, not as a substitute for solving the problem.

For some problems, multiple hints may be provided. Try to use the least amount of assistance necessary before returning to the problem.

Difficulty Levels

The normal problems are divided into three training levels:

- **Level I** — Fundamental Olympiad problem-solving practice.
- **Level II** — Problems requiring combinations of ideas.
- **Level III** — More serious problems that require deeper analysis.


Trigonometry I

Topic Note.

Trigonometry provides a language for describing angles, triangles, directions, and components. In physics, the same ideas appear naturally in vectors, forces, motion, geometry, and many other areas.

Here are some important definitions:

For a right triangle ABC where $C = 90^\circ$:



- **Pythagorean Theorem:**

$$AC^2 + BC^2 = AB^2$$

- **Sine:**

$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}}$$

- **Cosine:**

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}}$$

- **Tangent:**

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}} = \frac{\sin \theta}{\cos \theta}$$

1.1 Trigonometric Identities

PROBLEM 1

LEVEL I

The Fundamental Identity

Show that

$$\sin^2 x + \cos^2 x = 1.$$

PROBLEM 2

LEVEL II

A Tangent Identity

Show that


$$1 + \tan^2 x = \frac{1}{\cos^2 x}.$$

PROBLEM 3

LEVEL II

Complementary Angles

A right triangle ABC has $C = 90^\circ$.



(a) Show that

$$A + B = 90^\circ.$$

(b) Show that

$$\cos B = \sin A.$$

(c) Show that

$$\cos B = \sin(90^\circ - B).$$

(d) Explain geometrically why the two expressions are equal.


1.2 Right-Triangle Trigonometry

PROBLEM 4

LEVEL I

The 3–4–5 Triangle

Without using a calculator, find all six trigonometric ratios for a right triangle whose sides are 3, 4, and 5.



PROBLEM 5

LEVEL I

A Right Triangle with Known Sides

Triangle ABC is right-angled at C . The hypotenuse is 16 cm, and $AC = 12$ cm.

- (a) Find BC .
- (b) Find $\sin B$.
- (c) Find $\cos B$.
- (d) Find $\tan C$.
- (e) Explain why $\tan C$ has that result.

PROBLEM 6

LEVEL II

Working Backwards from Sine

Triangle XYZ is right-angled at Z . $XY = 10$ cm and

$$\sin X = \frac{3}{5}.$$


- (a) Find XZ .
- (b) Find YZ .
- (c) Find $\cos X$.
- (d) Find $\tan Y$.

PROBLEM 7

LEVEL II

The 30–60–90 Triangle

A right triangle has one acute angle of 30° and a hypotenuse of 12 cm.



- (a) Find the two shorter sides.
- (b) Find the area of the triangle.
- (c) Explain why a 30° - 60° - 90° triangle always has the same side ratio.

PROBLEM 8

LEVEL I

Exact Trigonometric Values

Without using a calculator, find:

- (a) $\sin 30^\circ$
- (b) $\cos 60^\circ$
- (c) $\sin 45^\circ$
- (d) $\cos 45^\circ$


(e) $\tan 45^\circ$

PROBLEM 9

LEVEL II

The Right Isosceles Triangle

A right isosceles triangle has two equal legs of length a .



- (a) Show that its hypotenuse is $a\sqrt{2}$.
- (b) Determine its two acute angles.
- (c) Find $\sin 45^\circ$, $\cos 45^\circ$, and $\tan 45^\circ$.

1.3 Special Triangles and Geometry


PROBLEM 10

LEVEL II

Area of an Equilateral Triangle

An equilateral triangle has side length a .

Derive an expression for its area in terms of a .



PROBLEM 11

LEVEL II

Dividing an Equilateral Triangle

An equilateral triangle has side length 10 cm.

- (a) Find its height.
- (b) Find its area.
- (c) Find the sine and cosine of the 30° and 60° angles formed when the triangle is divided into two right triangles.


1.4 Similarity and Trigonometric Ratios

PROBLEM 12

LEVEL II

Similar Right Triangles

Two similar right triangles ABC and XYZ are shown. C and Z are right angles. $AB = 6$ m and $AC = 4$ m.



- (a) Find BC .
- (b) Find $\cos A$.
- (c) Find $\cos X$ if $A = X$.
- (d) Explain why the size of a triangle does not affect its trigonometric ratios.

PROBLEM 13

LEVEL II

Same Angle, Different Size

Two right triangles have the same acute angle θ but different side lengths.

Triangle 1 has opposite and adjacent sides of 3 cm and 4 cm.


Triangle 2 has an adjacent side of 10 cm.

- (a) Find the opposite side of Triangle 2.
- (b) Explain why you can determine it without finding θ .

1.5 Trigonometry in Physics

In physics, vectors are often resolved into perpendicular components. Trigonometry provides the connection between the magnitude of a vector, its direction, and its components.

For Vector A, where θ is taken from positive x axis:



- **Pythagorean Theorem:**

$$A_x^2 + A_y^2 = A^2$$

- **Components:**

$$A_y = A \sin \theta, A_x = A \cos \theta$$

- **Angle Identity:**


$$\theta = \tan^{-1}\left(\frac{A_y}{A_x}\right)$$

PROBLEM 14

LEVEL II

Resolving a Force

A 10 N force acts at an angle of 30° above the horizontal.



- (a) Find the horizontal component of the force.
- (b) Find the vertical component.
- (c) Show that the original force can be reconstructed from its two components.

PROBLEM 15

LEVEL II

Velocity Components

A vector has a magnitude of 20 m/s and makes an angle of 60° with the horizontal.

- (a) Find its horizontal component.
- (b) Find its vertical component.
- (c) Which component is larger, and why?

PROBLEM 16

LEVEL II

Recovering an Unknown Direction

A 50 N force acts at an unknown angle θ . Its horizontal component is 30 N.

- (a) Write an equation involving $\cos \theta$.
- (b) Find θ .
- (c) Find the vertical component.

PROBLEM 17

LEVEL II

Velocity of a Ball

A ball is moving with a velocity of 12 m/s at an angle of 40° above the horizontal.


- (a) Find v_x .
- (b) Find v_y .
- (c) Explain why the horizontal and vertical components are not both 12 m/s.

1.6 Geometric Applications**PROBLEM 18**

LEVEL II

A Ladder Against a Wall

A ladder of length 5 m leans against a vertical wall. Its base is 3 m from the wall.



- (a) Find the height reached by the ladder.
- (b) Find the angle between the ladder and the ground.
- (c) Find the angle between the ladder and the wall.

PROBLEM 19

LEVEL II

A Sloping Ramp

A ramp is 6 m long and rises 1.5 m.




- (a) Find the angle of the ramp above the horizontal.
- (b) Find the horizontal distance covered by the ramp.
- (c) Which trigonometric ratio gives the most direct solution?

PROBLEM 20

LEVEL II

Angle of Elevation

From a point on the ground, the angle of elevation to the top of a 20 m tower is 35° .



- (a) Find the distance from the observer to the base of the tower.
- (b) Find the distance from the observer to the top of the tower.
- (c) Draw the triangle and identify the opposite, adjacent, and hypotenuse sides.

1.7 Vectors and Trigonometric Synthesis

PROBLEM 21

LEVEL II

Components and the Fundamental Identity

A vector has components A_x and A_y . Its magnitude is A .

Show that

$$A^2 = A_x^2 + A_y^2.$$

Then, using

$$A_x = A \cos \theta, \quad A_y = A \sin \theta,$$

recover the identity


$$\sin^2 \theta + \cos^2 \theta = 1.$$

PROBLEM 22

LEVEL II

Perpendicular Forces

Two forces of 6 N and 8 N act perpendicular to each other.



- (a) Find the magnitude of their resultant.
- (b) Find the angle that the resultant makes with the 8 N force.
- (c) Explain why the Pythagorean theorem and trigonometry can be used to describe the resultant.

PROBLEM 23

LEVEL II

A Vector with Related Components

A vector has magnitude 10 and makes an angle θ with the positive x -axis.

Its x -component is twice its y -component.

- (a) Write an equation relating $\sin \theta$ and $\cos \theta$.
- (b) Determine θ .
- (c) Find both components.

PROBLEM 24**LEVEL III****Area and Hypotenuse**

A right triangle has an area of 24 cm^2 and a hypotenuse of 10 cm.

- (a) Find the two possible leg lengths.
- (b) Find the two acute angles.
- (c) Explain why knowing the area and hypotenuse is enough to determine the triangle.

CHAPTER 2

Hints

The hints below correspond to the problems in the handout. They are provided separately so that students can work through the problem collection without immediately seeing the associated guidance.

@hintlist