

Informed Consent Form

Study: Phase 3 Xanomeline-Trospium in Alzheimer's Disease

IRB Protocol Number: CT-2024-001-ICF

Purpose of the Study

You are being asked to take part in a research study to evaluate whether the combination of two study medications, xanomeline and trospium, can improve cognitive function in patients with Alzheimer's disease. Xanomeline is an investigational drug that targets muscarinic receptors in the brain, potentially improving memory and thinking. Trospium is an FDA-approved medication for overactive bladder that may help reduce certain side effects of xanomeline.

Study Procedures

If you agree to participate, the study will last approximately 52 weeks (24 weeks of double-blind treatment plus 28 weeks of open-label extension). You will be asked to attend 12-14 clinic visits. At each visit, you will undergo cognitive assessments, physical examinations, vital sign measurements, ECGs, and blood laboratory tests. Study medication will be taken orally twice daily with food.

Risks and Discomforts

The most common side effects of xanomeline include gastrointestinal symptoms such as nausea, vomiting, diarrhea, and abdominal pain. Trospium may cause dry mouth, constipation, and urinary retention. There is a risk of falls, particularly in elderly patients. All medications will be monitored by the study physician. You should report any side effects to the study team immediately.

Confidentiality

Your study records will be kept confidential to the extent permitted by law. Your identity will be coded using a unique patient identification number. Data will be stored in secure, password-protected databases accessible only to authorized study personnel. Results may be published in medical journals but your identity will not be disclosed.