

Studying Development of Psychopathology Across the Lifespan

Isaac T. Petersen, Zachary Demko, and Megan Schumer

Department of Psychological and Brain Sciences, University of Iowa

Author Note

Isaac T. Petersen  <https://orcid.org/0000-0003-3072-6673>

Zachary Demko  <https://orcid.org/0000-0002-5717-6205>

Megan Schumer  <https://orcid.org/0009-0009-5488-6541>

The data for the present study are published online: [LINK]. We have no conflicts of interest to disclose. The present study was approved by the University of Iowa Institutional Review Board (Study #: INSERT). The authors complied with APA ethical standards in the treatment of participants. I.P. was funded by Grant HD098235 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Institutes of Health. We thank Jordan L. Harris and Alexis Hosch for their help identifying measures of externalizing problems and transcribing and coding their content. Author roles were classified using the Contributor Role Taxonomy (CRediT; <https://credit.niso.org>) as follows: Isaac T. Petersen: conceptualization, data curation, analysis, funding, investigation, methodology, project administration, resources, software, supervision, validation, visualization, writing, editing; Zachary Demko: conceptualization, data curation, analysis, investigation, methodology, software, editing; Megan Schumer: editing

Correspondence concerning this article should be addressed to Isaac T. Petersen, University of Iowa, G60 Psychological and Brain Sciences Building, Iowa City, IA 52242. Email: isaac-t-petersen@uiowa.edu

Abstract

The field has not adequately studied individuals' development of psychopathology across the lifespan, instead taking a piecewise approach to development, limiting our understanding of lifespan development and the origins of psychopathology. We describe why this is the case—beyond logistical challenges. First, diagnostic and taxonomic systems do not define psychopathology constructs from a developmental perspective. A developmental perspective is crucial because psychopathology constructs often have earlier origins than their full clinical presentation, and forms of psychopathology frequently show changes in behavioral manifestation across development (heterotypic continuity). Second, current assessments do not adequately capture these changes in behavioral manifestation. Third, traditional scoring systems do not allow charting individuals' change over time in the face of changing behavioral manifestation. Fourth, assessments are not well-suited for capturing individual differences, despite the dimensional nature of psychopathology. We propose solutions to these challenges by introducing a developmental scaling framework for psychopathology. Using externalizing psychopathology as an example, we draw on our content analysis of externalizing measures, as well as lessons from the study of other constructs across lengthy developmental periods. Addressing these challenges will allow charting individuals' trajectories of psychopathology across the lifespan and identifying risk and protective factors that influence developmental courses.

Keywords: psychopathology, lifespan development, longitudinal trajectories, heterotypic continuity, externalizing behavior problems

Studying Development of Psychopathology Across the Lifespan

NOTE: this manuscript is in progress of being drafted—it is incomplete.

When entrenched, psychopathology is costly and difficult to treat (Insel, 2008; Wakschlag et al., 2019). Understanding its developmental origins is therefore crucial for preventing later, more severe psychopathology. Developmental science seeks to build a bridge that spans the lifespan—from infancy to older adulthood—and not just transitory periods in people's lives. However, the field has not adequately studied individuals' development of psychopathology across the lifespan, instead taking a piecewise approach to development that focuses on narrower age ranges. The focus on narrow age ranges restricts our ability to understand development across important transitions, such as puberty, school entry, and entry into parenthood, thus limiting our understanding of lifespan development and the origins of psychopathology. In this paper, we describe reasons why this is the case—beyond logistical challenges of time and money. And we propose solutions to these challenges by introducing a developmental scaling framework for psychopathology. Using externalizing psychopathology as an example, we draw on our content analysis of over 270 externalizing measures, as well as lessons from the study of other constructs such as academic and cognitive skills across lengthy developmental periods.

The Goal

A key goal of developmental science is to understand how people develop across the lifespan. A variety of methods, including cross-sectional and longitudinal designs, could be leveraged for this purpose. In a cross-sectional design, people are assessed at one timepoint. In a longitudinal design, the same individuals are followed across time.

Cross-sectional designs provide the ability to cheaply and quickly assess a wide age range, providing a snapshot at a particular point in time. Cross-sectional designs may inform our understanding of general normative patterns of change—such as the typical pattern of cognitive growth and decline across the lifespan. Thus, cross-sectional designs provide an important starting point.


However, cross-sectional designs have key limitations. First, cross-sectional designs do

not allow examining individuals' change over time. Second, cross-sectional designs may obscure diverse patterns of change. Individuals show considerable heterogeneity in level and change across time in many constructs. As an example, despite the normative pattern of steep cognitive decline in episodic memory with age, there are a subset of individuals who do not show this same pattern. For instance, superagers are individuals who are older adults (age 80+) who have episodic memory at or above the normative level of middle-aged adults (50–65 years old) and who may be resilient to the typical aging process (Godoy et al., 2021). Theoretical depictions of the heterogeneity of cognitive decline are depicted in Figure 1.


Figure 1

Panel A: Hypothetical models for different cognitive trajectories of aging. MCI = mild cognitive impairment. (Figure in Panel B reprinted from Borelli et al. (2018), Figure 1, p. 222. Borelli, W. V., Carmona, K. C., Studart-Neto, A., Nitrini, R., Caramelli, P., & Costa, J. C. d. (2018). Operationalized definition of older adults with high cognitive performance. *Dementia & Neuropsychologia*, 12(3), 221–227. <https://doi.org/10.1590/1980-57642018dn12-030001>). **Panel B:** Theoretical illustration of inter-individual variability in cognitive aging trajectories. Each grey line represents one individual hypothetical cognitive health trajectory. The orange line marks the threshold for functional independence; once an individual's cognitive health drops below this level, autonomous daily living is compromised. (Figure in Panel B reprinted from Abrous et al. (2026), Figure 1, p. 2. Abrous, D. N., Blin, N., Boraxbekk, C.-J., Catheline, G., Fitzsimons, C. P., Hilscher, M., Lemoine, M., Lopes, L. V., Maass, A., Nilsson, M., Nyberg, L., Rasmussen, L. J., Remondes, M., Sauvage, M., Schreiber, S., & Wolbers, T. (2026). Hallmarks of healthy cognitive aging: Inter-individual differences in aging trajectories. *Ageing Research Reviews*, 119, 103102. <https://doi.org/10.1016/j.arr.2026.103102>)

(A) Hypothetical models for different cognitive trajectories of aging.



(B) Theoretical illustration of inter-individual variability in cognitive aging trajectories.



Heterogeneity also applies to psychopathology trajectories including externalizing and

criminal behavior. The age-crime curve depicts the well-known strong association between age and engagement in crime, with crime rates peaking markedly (in many Western countries) in adolescence around ages 15–19. An example of the age-crime curve is depicted in Figure 2A.

However, Moffitt's (1993) developmental taxonomy of externalizing ("antisocial") behavior posits that there are heterogeneous subgroups such that one subgroup exhibits low levels of externalizing behavior across the lifespan, an "adolescence limited" subgroup exhibits high levels of externalizing behavior during adolescence but low levels during childhood and adulthood, and a "life-course persistent" subgroup that exhibits high levels of externalizing behavior across the lifespan; in addition, more recent work has identified a "childhood limited" subgroup that exhibits high levels of externalizing behavior during childhood but low levels after childhood (Fairchild et al., 2013; Odgers et al., 2008). The developmental taxonomy is depicted in Figure 2B. Moffitt's general hypothesis is that life-course persistent externalizing behavior results from early disruptions in brain development either in utero (resulting from, for example, maternal substance use, poor prenatal nutrition, genetics, or exposure to toxins), during delivery (e.g., delivery complications), or after birth (e.g., child abuse or neglect, or exposure to toxins) that lead to impaired cognitive abilities and difficult temperament, in perhaps conjunction with coercive parent-child interactions or other adverse, criminogenic environments. Adolescence-limited externalizing behavior, by contrast, is thought to reflect a more normative—even adaptive—pattern of social behavior that is driven by exposure to deviant peers and a maturity gap. Childhood-limited externalizing behavior, though not hypothesized by the original taxonomy, may in some cases reflect recovery and in other cases reflect a shift from externalizing problems into problems in other domains (Odgers et al., 2008). It has been hypothesized that both childhood-limited and life-course persistent externalizing behavior may be characterized by high levels of individual-level risk (e.g., genetic and neuropsychological risk), whereas life-course persistent externalizing behavior may be characterized by greater environmental risk (e.g., adversity and trauma; Fairchild et al., 2013). A cross-sectional study that does not capture the subgroups' different developmental histories would not be well-positioned to adjudicate

whether a particular individual is part of the adolescence-limited versus life-course persistent developmental course ([Moffitt, 1993](#)).

The age-crime curve that is based on official police records has key problems ([Moffitt, 1993](#)). First, it misses the developmental course of externalizing behavior that leads up to the commission of crime. Second, it is based on only those (criminal) behaviors that are known to official arrest records and does not capture other behaviors the person engaged in that are known to the individual or collateral report by caregivers, teachers, siblings, or peers. Longitudinal work examining informant report of children's behavior allows estimating the prototypical trajectory of externalizing problems at ages younger than those encompassed by the age-crime curve, thus providing important developmental context to the origins of externalizing behavior. For instance, in a longitudinal study examining the development of externalizing behavior from childhood to adulthood that incorporated ratings of the child's externalizing behavior from mothers, fathers, teachers, peers, and self-report (at relevant ages), Petersen et al. ([2015](#)) found that externalizing problems normatively tended to decrease from early childhood to preadolescence, increased during adolescence (peaking around age 17), and decreased from late adolescence to adulthood.

Since Moffitt ([1993](#)) proposed the developmental taxonomy of externalizing behavior, work has examined individual differences in externalizing trajectories to determine whether they are best represented by categories—i.e., taxa such as adolescence-limited, life-course persistent, etc.—or dimensionally. Findings have consistently shown that variation in externalizing trajectories is better represented dimensionally than with categories ([Fairchild et al., 2013](#); [Walters, 2011, 2012, 2015](#); [Walters & Ruscio, 2013](#)), suggesting that differences between people in their developmental course of externalizing behavior appear to reflect differences in degree rather than kind (i.e., quantitative differences rather than qualitative differences). For example, the trajectories in Petersen et al. ([2015](#)) were characterized by marked individual differences—some individuals stayed low, some increased, some decreased, some increased then decreased, and seemingly everything in between. The heterogeneous trajectories of externalizing problems are depicted in Figure [2C](#).

In sum, if we rely solely on normative age-related patterns of cognitive functioning or psychopathology from cross-sectional research, we would miss the great individual differences in people's developmental courses. And these individual differences are crucial to identify the risk and protective mechanisms that explain these differing developmental courses.

Third, in a cross-sectional study, any apparent age-related differences in a construct could reflect cohort differences instead of developmental changes. For example, average differences in stress between 5-year-olds and 10-year-olds could reflect the differing experiences that the 10-year-olds shared—such as going to school during the Coronavirus pandemic. Fourth, findings from cross-sectional studies can differ from findings examining the same people over time, which violates the convergence assumption. In cross-sectional studies, the convergence assumption is the assumption that cross-sectional age differences and longitudinal age changes converge onto a common trajectory. An example where the convergence assumption has been shown to be violated is in studies of cognitive decline. In particular, cross-sectional studies overestimate age-related cognitive declines compared to following the same people over time in a longitudinal study, which may reflect cohort effects such as the Flynn effect (where population intelligence scores at a given age tend to increase across generations; [Ackerman, 2013](#)).

Because longitudinal designs follow the same person(s) over time, such a design allows examining individuals' change across time. However, as we describe [later](#), longitudinal designs also have potential confounds of change. Moreover, compared to cross-sectional designs, longitudinal designs take more time and tend to be more costly and to require more personnel. Thus, longitudinal studies that span the lifespan are not common. However, even cross-sectional work spanning the lifespan is not often conducted.

The Problem/Obstacle


The focal problem examined in this paper is that, despite calls to do so ([De Los Reyes, 2026](#)), the field has not adequately studied individuals' development of psychopathology across the lifespan, instead taking a piecewise approach to development. The piecewise approach to development limits our understanding of lifespan development and the origins of psychopathology.

Figure 2

Panel A: The age-crime participation curve from age 9 through 64 in cohort members registered for antisocial and/or criminal behavior in a birth cohort recruited in Stockholm, Sweden ($N = 4825$), based on official records. The steep peak in crime rate in the adolescent years is similar to the age-crime curve in the United States; however, the steepness of the age-crime curve differs within country across birth cohorts (Steffensmeier et al., 2025). (Figure adapted from Sivertsson et al. (2024), Figure 1, p. 7. Sivertsson, F., Carlsson, C., Almquist, Y. B., & Brännström, L. (2024). Offending trajectories from childhood to retirement age: Findings from the Stockholm birth cohort study. *Journal of Criminal Justice*, 91, 102155. <https://doi.org/10.1016/j.jcrimjus.2024.102155>).


Panel B: A stylized depiction of Moffitt's (1993) developmental taxonomy of externalizing behavior, adapted to include a childhood limited subgroup. **Panel C:** Individuals' ($N = 585$) longitudinal trajectories of externalizing problems from childhood to adulthood based on annual assessments of mother-, father-, teacher-, peer-, and self-report (at relevant ages). Average trajectory in red. (Figure adapted from Petersen et al. (2015), Petersen, I. T., Bates, J. E., Dodge, K. A., Lansford, J. E., & Pettit, G. S. (2015). Describing and predicting developmental profiles of externalizing problems from childhood to adulthood. *Development and Psychopathology*, 27(3), 791–818. <https://doi.org/10.1017/S0954579414000789>).

(A) Age-crime curve (among those who have a record of antisocial or criminal behavior).



(C) Heterogeneity of individuals' trajectories of externalizing behavior.

(B) A stylized depiction of Moffitt's (1993) developmental taxonomy of externalizing behavior, adapted to include a childhood limited subgroup.



A modest number of longitudinal studies have followed people for lengthy developmental periods.

General Longitudinal Studies

The Fels Longitudinal Study ([Roche, 1992](#)) was a cohort-sequential study lasting from 1929 to 2018 that recruited 10–20 newborn infants each year and followed them over time, eventually following some until age 76 ([Chumlea et al., 2009](#)). The study collected data on participants' body composition, including height and weight, which had key impacts and formed the basis of pediatric growth charts. In addition, the study collected information on the participants' behavior—and other constructs—in childhood (based on observation) and adulthood (based on interview). These rich data allowed researchers to examine the extent of personality continuity and change from childhood to adulthood, and the extent to which mothers may influence the personality development of the child ([Moss & Kagan, 1964](#)). On the basis of predictive associations from childhood to adulthood in the Fels Longitudinal Study, Kagan and Moss ([1962](#)) noted that girls who had frequent tantrums at 6 to 10 years of age tended to become women who were more motivated in school, less dependent on others, and more masculine in their interests than women who had fewer tantrums as children. They interpreted the different behaviors at different ages as stemming from the same psychological process: a tendency to avoid adopting “female sex-role standards” (p. 200). Kagan described such findings, when psychological processes remain the same but the form of behavior changes, as examples of heterotypic continuity ([Kagan, 1969, 1971, 1980](#)).

Longitudinal Studies of Psychopathology

Longitudinal studies such as these have had immense impacts. However, as we will describe, these lengthy longitudinal studies of psychopathology (e.g., [Odgers et al., 2008](#))—including our own ([Petersen et al., 2015](#))—have important limitations in (a) how they assessed psychopathology constructs across development and (b) how they estimated people's scores on those constructs over time that prevent strong inferences regarding individuals' change over time.

Reasons for the Problem

Beyond logistical challenges of time and money, there are several key reasons why the field has failed to study people's development of psychopathology, longitudinally, across the lifespan. First, diagnostic and taxonomic systems do not define psychopathology constructs from a developmental perspective. A developmental perspective is crucial because psychopathology constructs often have earlier origins than their full clinical presentation, and forms of psychopathology frequently show changes in behavioral manifestation across development (heterotypic continuity). Second, current assessments do not adequately capture these changes in behavioral manifestation. Third, traditional scoring systems do not allow charting individuals' change over time in the face of changing behavioral manifestation. Fourth, current assessments are not well-suited for capturing the full range of individual differences, despite the dimensional nature of psychopathology. Fifth, as a result of these limitations, current assessments are not well-suited for capturing individuals' change over time.

Below, we describe these reasons in detail. To ground the discussion, we provide examples relating to externalizing psychopathology; however, these issues are not specific to externalizing.

Problems of Constructs

The first reason the field has failed to adequately study individuals' development of psychopathology across the lifespan represents a theory problem—namely, problems with constructs, and the codifying of those constructs in diagnostic and taxonomic systems. In particular, our theories do not fully specify how constructs manifest behaviorally across the lifespan. For instance, theories do not specify the features (e.g., cognitive, affective, or biological processes) or behaviors that characterize a given construct in each developmental period. In addition, theories do not specify how those features or behaviors change across development with respect to how strongly they reflect the construct—as indicated by changes in factor loadings (in factor analysis) or discrimination (in item response theory)—or how they change in their level on the construct—as indicated by changes in intercepts (factor analysis) or difficulty/severity (item response theory).

These gaps are important for several reasons. First, psychopathology does not come from nowhere; psychopathology constructs often have earlier origins than their full clinical presentation. In many cases, psychopathology may arise from the interaction of genetic and environmental factors, and their effects on biology, cognition, emotion, and behavior. Such effects may accumulate and transpire over years. Second, forms of psychopathology frequently show changes in behavioral manifestation across development—a phenomenon called heterotypic continuity (Caspi & Shiner, 2006; Cicchetti & Rogosch, 2002; Petersen et al., 2020; Petersen, 2024).

Considerable theoretical work attempts to characterize constructs across development. For instance, Patterson (1993) characterized externalizing (“antisocial”) behavior as a chimera. A chimera is a mythical creature with the body of a goat that, with development, grows the head a lion and then a tail of a snake. Patterson (1993) and Moffitt (1993) argued that externalizing behavior changes in manifestation across development such that underlying antisociality is maintained while more mature features are added across development. In particular, Patterson (1993) noted that some externalizing behaviors wane with age, such as temper tantrums, whereas other externalizing behaviors emerge with age, including the ability to inflict serious damage or harm with violence, burglary, and shoplifting, in addition to problems such as academic failure, peer rejection, substance problems, and arrest. Moffitt (1993) noted that externalizing behaviors may manifest as “biting and hitting at age 4, shoplifting and truancy at age 10, selling drugs and stealing cars at age 16, robbery and rape at age 22, and fraud and child abuse at age 30”, and that the prognosis for individuals who engage in externalizing behavior throughout the life-course includes “drug and alcohol addiction; unsatisfactory employment; unpaid debts; homelessness; drunk driving; violent assault; multiple and unstable relationships; spouse battery; abandoned, neglected, or abused children; and psychiatric illness” (p. 679). In general, externalizing behaviors are often expressed as overt acts in early childhood, such as physical aggression and temper tantrums; whereas externalizing behaviors are more often expressed later in development as covert and indirect or relational forms of aggression, rule breaking, and substance use (Miller et al., 2009; Petersen et al., 2026). These age-differing behaviors show strong stability of

individual differences, indicating a consistent underlying construct (Moffitt, 1993; Patterson, 1993). Consistent with the Developmental Issues Framework (Sroufe, 2016), the changing expression of externalizing behavior likely reflects a combination of time-varying genetic and environmental factors, such as school entry transition, in combination with varying developmental tasks, different opportunities, and greater experience-dependent capacity (Petersen et al., 2020).

In addition to externalizing problems, both internalizing (Petersen et al., 2018; Weems, 2008; Weiss & Garber, 2003) and thought-disordered (Rutter et al., 2006) problems are thought to change in manifestation across development.

However, more work is needed to developmentally parameterize constructs—to define constructs developmentally in terms of their features. Recent empirical work suggests that it is uncommon for symptoms to change from one symptom to another for a given person at the subsequent measurement occasion (Applegate & Lahey, in press). Nevertheless, the authors found that adding new symptoms and subtracting symptoms—other ways that a construct can change in manifestation—were both relatively common. Moreover, just because a symptom persists does not mean that the behavioral manifestation of that symptom stays the same for that individual. For instance, even if a symptom (e.g., “fighting”) persists for an individual, their manifestation of that symptom could differ in form, function, or mechanism (Petersen, 2024). Thus, as noted by Applegate and Lahey (in press), “future causal theories will need to rest on a more detailed and nuanced description of persistence and change at the level of specific psychological problems.”

Moreover, where such theories do indicate changes in behavioral manifestation across development, diagnostic and taxonomic systems do not incorporate these developmental changes.

Lack of a Developmental Perspective

Diagnostic and taxonomic systems do not define psychopathology constructs from a developmental perspective. The diagnostic systems, including the Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR; American Psychiatric Association, 2022) and the International Classification of Diseases (ICD-11; World Health Organization, 2022), specify various externalizing-related disorders, including attention-deficit/hyperactivity disorder (ADHD),

oppositional defiant disorder (ODD), conduct disorder (CD), and antisocial personality disorder (ASPD). Among externalizing-related disorders—with few exceptions¹—the diagnostic criteria do not differ across development, even though diagnostic manuals acknowledge that developmental factors impact symptom presentation². In addition to diagnostic systems, emerging taxonomies and frameworks of psychopathology—including the hierarchical taxonomy of psychopathology (HiTOP) and research domain criteria (RDoC)—do not incorporate a developmental perspective (Tackett & Hallquist, 2022).

Problems of Existing Assessments and Traditional Scoring Systems

Other key reasons the field has failed to adequately study individuals' development of psychopathology across the lifespan represent method problems—namely, problems with existing

¹ In the DSM-5-TR (American Psychiatric Association, 2022), the minimum frequency for considering a symptom of ODD to be present is “most days” (for a period of six months) for children younger than 5 years, whereas it is “at least once per week” (for at least six months) for children 5 years or older. For the diagnosis of ADHD, six or more symptoms are required for children and younger adolescents (up to age 16), whereas five or more symptoms are required for older adolescents and adults (aged 17 or over). However, the symptoms for defining ODD and ADHD do not differ across development.

² In the DSM-5-TR (American Psychiatric Association, 2022), some disorders have a “Development and Course” subsection that describes when disorders tend to emerge and their rates across ages, in addition to describing the types of symptoms that may be more or less common at various points in development. However, the symptoms that define the disorders do not differ across development. For instance, in the section on major depressive disorder, it is noted that “hypersomnia and hyperphagia are more likely in younger individuals, and melancholic symptoms, particularly psychomotor disturbances, are more common in older individuals. Depressions with earlier ages at onset are more familial and more likely to involve personality disturbances.” (p. 189). In the section on ADHD, it is noted that, “In preschool, the main manifestation is hyperactivity. Inattention becomes more prominent during elementary school. During adolescence, signs of hyperactivity (e.g., running and climbing) are less common and may be confined to fidgetiness or an inner feeling of jitteriness, restlessness, or impatience. In adulthood, along with inattention and restlessness, impulsivity may remain problematic even when hyperactivity has diminished.” (p. 71). In the section on CD, it is noted that “Symptoms of the disorder vary with age as the individual develops increased physical strength, cognitive abilities, and sexual maturity. Symptom behaviors that emerge first tend to be less serious (e.g., lying, shoplifting), whereas conduct problems that emerge last tend to be more severe (e.g., rape, theft while confronting a victim)... When individuals with conduct disorder reach adulthood, symptoms of aggression, property destruction, deceitfulness, and rule violation, including violence against co-workers, partners, and children, may be exhibited in the workplace and the home” (p. 534). Moreover, some disorders in the DSM-5-TR note that, to be considered a symptom, the frequency, intensity, or persistence, of the behavior must be inconsistent with the level expected for their developmental level. For instance, in the diagnosis of ADHD, it is noted that “the symptoms have persisted for at least 6 months to a degree that is inconsistent with developmental level” (p. 68). In the diagnosis of ODD, it is noted that “other factors [besides a minimum frequency] should also be considered, such as whether the frequency and intensity of the behaviors are outside a range that is normative for the individual’s developmental level, gender, and culture.” (p. 522). In addition, some disorders require that some symptoms onset before or after a particular age. For instance, in the diagnosis of ADHD, several symptoms must be present prior to age 12. In the diagnosis of CD, the symptoms “often stays out at night despite parental prohibitions” and “if often truant from school” must begin before age 13 (p. 531). In the diagnosis of ASPD, symptoms must be persistent since age 15 but have onset prior to age 15.

assessments and how those assessments are used, based on traditional scoring systems. There is a mismatch between theory and method. In particular, there is a mismatch between theory of constructs—including how a construct manifests behaviorally—and the methods used to assess the construct and to evaluate people's level and change in that construct.

Do Not Adequately Capture Constructs' Change in Behavioral Manifestation Across Development

Given the substantial changes in behavioral manifestation of psychopathology across development, it is important for measures to capture the constructs' changing behavioral manifestation, to stay developmentally relevant and construct valid. As noted by Moffitt (1993), "Measures of antisocial behavior should be sensitive to developmental heterogeneity to tap individual differences while allowing for the emergence of new forms of antisocial behavior (e.g., automobile theft) or for the forsaking of old forms (e.g., tantrums)." (p. 694).


However, measures show insufficient differences in which items are assessed at which ages to account for heterotypic continuity. For example, the same version of the Child Behavior Checklist (CBCL) spans ages 6–18. Considerable development occurs during that span.

There are two primary ways that researchers have examined people's change in psychopathology across development Figure 3. The "common items" approach removes items that are age specific. For example, in a study of externalizing behavior from early childhood to adolescence, a study might remove some substance use items that might not be developmentally relevant at all ages. The second approach, the "upward/downward extension" approach, takes items that are valid at a given age and uses the items across ages when the item may not be valid or useful—i.e., all items are assessed at all ages. For instance, the upward extension approach might take the item "disobedient to authority figures" and apply it to older individuals for whom such a behavior may no longer validly reflect externalizing behavior—and may instead reflect prosocial functions including protesting against societally unjust actions. Both the "common items" (e.g., Odgers et al., 2008; Sterba et al., 2007) and "upward/downward extension" (e.g., Briggs-Gowan et al., 2016; Broeren et al., 2013) approaches are widely used, likely because they

result in the same items assessed across ages, but both have key problems. As an example of the common items approach, Odgers et al. (2008, p. 676) examined children's development of DSM-IV symptoms of conduct disorder from ages 7 to 26 years, but they dropped age-specific symptoms (e.g., running away, staying out late) "because [these symptoms] did not cover the study's age span". As an example of the upward extension approach, Broeren et al. (2013, p. 84) examined children's development of anxiety from ages 4 to 11 years; they noted, "Although the questionnaire was originally developed to measure anxiety in preschoolers, the current study also employed the scale with older children to promote uniformity in measures".

Figure 3

Item set A refers to items that are construct-valid at only timepoint 1 (T1); item set B is construct-valid at both T1 and T2; item set C is construct-valid at only T2. A dash indicates that the item set was not assessed at a given timepoint. A white box indicates invalid assessment in terms of either a content gap (i.e., important missing items) or intrusion (i.e., invalid items at a given timepoint). In a study of externalizing behavior from early childhood to adulthood, "biting others" may be in item set A; "noncompliant" in B; "drug use" in C. The three approaches are: (1) common items: B at T1 and T2, (2) upward/downward extension: ABC at T1 and T2, or (3) construct-valid items: AB at T1; BC at T2. The "common items" and "upward/downward extension" approaches are by far the most widely used in the literature, even though they likely lead to inaccurate scores and thus inaccurate trajectories. (Figure reprinted from Petersen et al. (2026), Figure 1, p. 113. Petersen, I. T., Demko, Z., Lee, W.-C., & Oleson, J. J. (2026). Studying development of psychopathology using changing measures to account for heterotypic continuity. JAACAP Open, 4(1), 111–123. <https://doi.org/10.1016/j.jaacop.2025.10.008>)



Forcing use of the same measure across time naturally limits the ages a study can span, which prevents charting wider age spans. The common items approach yields low content validity

because it does not assess all construct facets, especially age-specific manifestations. The upward/downward extension approach violates construct validity because it assesses developmentally inappropriate items. Due to these problems, both simulation (Petersen et al., 2021) and empirical work (Chen & Jaffee, 2015; Petersen et al., 2018, 2026) has shown that the “common items” and “upward/downward extension” approaches lead to inaccurate developmental inferences. In sum, despite these approaches being the most widely used for assessing individuals’ development, they likely lead to inaccurate scores and thus inaccurate trajectories.

Do Not Allow Charting Individuals’ Change Over Time When the Construct Changes in Behavioral Manifestation

Current tools fail to implement the construct-valid item approach—existing measures show insufficient differences in which items are assessed at which ages. Moreover, even in cases where the items of a measure differ across ages (e.g., CBCL: ages 1.5–5 vs. ages 6–18), the scoring systems do not link scores across ages to be on the same scale while preserving change in level. Thus, the measures do not yield scores that are comparable across time and that enable change in an individual’s level to be observed across a lengthy developmental span. Age norms and standardized scores (*T*- or *z*-scores) do not allow observing individuals’ absolute change because scores have a fixed mean and variance, and they do not equal developmental understanding. Absolute change is necessary to identify true growth for an individual, and is thus important for treatment monitoring and longitudinal growth curves.

Not Well-Suited For Capturing the Full Range of Individual Differences

Many measures use coarse, subjective response scales such as Likert-type response scales (e.g., “rarely”, “sometimes”, “often”, “very often”) leading to cultural bias and insensitivity to change (Paalman et al., 2013; Schaeffer, 1991; Schwarz, 1999). Moreover, Likert-type scales tend to focus on frequency of misbehavior and ignore intensity and severity.

In addition, the measures focus on extreme or clinical-range behaviors; scores are thus positively skewed with floor effects and have unacceptably low reliability (including the CBCL; Kaat et al., 2019; Pavlovich et al., 2026; Tiego et al., 2023), making the CBCL and existing

measures unsuitable for dimensional research (Pavlovich et al., 2026; Tiego et al., 2023).

Considerable evidence indicates that externalizing problems (and psychopathology, more generally; Haslam et al., 2012; Haslam et al., 2020; Kotov et al., 2017; Krueger & DeYoung, 2016; Krueger & Markon, 2011; Krueger & Piasecki, 2002; Markon et al., 2011; Wright et al., 2013) and their trajectories (Burt, 2012; Fairchild et al., 2013; Walters, 2011, 2012, 2015; Walters & Ruscio, 2013) and risk factors (Walters, 2014) are best conceptualized dimensionally (or multi-dimensionally; Bolhuis et al., 2017; Wakschlag et al., 2014), not categorically (Barry et al., 2013; Coghill & Sonuga-Barke, 2012; Forbes et al., 2016; Kliem et al., 2018, 2022; Krueger et al., 2004, 2005, 2007, 2021; Markon & Krueger, 2005; Sellbom, 2016; Walton et al., 2011). Thus, the existing measures are impoverished for studying the full range of individual differences on the externalizing spectrum, which prevents subthreshold children who are at risk for impairment from being identified. Better identification of subthreshold externalizing is critical, in both research and practice, for identifying at-risk individuals in the early-stage trajectory of psychopathology.

Not Well-Suited for Capturing Individuals' Change Over Time

In sum, although they have been helpful, existing measures are poorly suited to classify individuals' development of psychopathology across a lengthy span, which is necessary for tracking individuals' change and identifying mechanisms in the development of psychopathology.


Example: Externalizing Psychopathology

Solution: A Developmental Scaling Framework

A conceptual depiction of an example construct-valid item design spanning infancy to adulthood is in Figure 4. In the construct-valid item design, some items represent the core of the phenotype and are assessed across all ages, whereas other items are age-specific. Despite some age-specific items, some common items are assessed at adjacent ages, which help serve as anchor items for linking the scores across the adjacent ages. The use of overlapping yet distinct items is widely used in educational testing to help link individuals' scores across grades, and is called vertical scaling because the item sets appear to form a line moving upward (see Figure 4). For example, for a standardized test of math ability, Kindergartners may receive items assessing

Figure 4

Conceptual depiction of a construct-valid item design from infancy to adulthood. The age(s) at which an item set is assessed are represented by the letter x and a colored box. Some items (“Core”, in green) represent the core of the phenotype and are assessed across all ages. Some items (in varying shades of brown) are age-specific and are assessed at only one or a few ages. However, some of the items assessed at a given age are shared with those assessed at adjacent ages, to facilitate linking the scores across ages. For instance, item set f can be used to help link scores from age 6 to those at age 5, and item set g can be used to help link scores from age 6 to those at age 7. Hypothetically, items could be assessed for as many or as few years as desired—as long as the items are construct valid for those ages and there enough common items between adjacent ages for linking scores across ages.



Other Key Challenges to Address

Potential Confounds of Change

Practice/Retest Effects

Period and Cohort Effects

Measurement Error

Change in Meaning of the Scores

Factorial Invariance Across Time

Integration Across Multiple Sources

Obsolescence of Items and Measures

Conclusion

Addressing these challenges will allow charting individuals' trajectories of psychopathology across the lifespan and identifying risk and protective factors that influence their developmental courses. Doing so is critical for understanding how psychopathology develops, when to intervene, and how to most effectively prevent psychopathology. Although this manuscript focuses on development of psychopathology, the principles discussed likely apply to many other developmental domains.

References

- Abrous, D. N., Blin, N., Boraxbekk, C.-J., Catheline, G., Fitzsimons, C. P., Hilscher, M., Lemoine, M., Lopes, L. V., Maass, A., Nilsson, M., Nyberg, L., Rasmussen, L. J., Remondes, M., Sauvage, M., Schreiber, S., & Wolbers, T. (2026). Hallmarks of healthy cognitive aging: Inter-individual differences in aging trajectories. *Ageing Research Reviews*, 119, 103102. <https://doi.org/10.1016/j.arr.2026.103102>
- Ackerman, P. L. (2013). Assessment of intellectual functioning in adults. In K. F. Geisinger, J. F. Carlson, J.-I. C. Hansen, N. R. Kuncel, S. P. Reise, & M. C. Rodriguez (Eds.), *APA handbook of testing and assessment in psychology: Vols. 2: Testing and Assessment in Clinical and Counseling Psychology* (pp. 119–132). American Psychological Association.

- American Psychiatric Association. (2022). *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR)*. American Psychiatric Publishing.
- Applegate, B., & Lahey, B. B. (in press). Mapping persistence and change in psychological problems during the transition to adolescence: Adding, subtracting, shifting, and persisting. *JCPP Advances*, e70043. <https://doi.org/10.1002/jcv2.70043>
- Barry, T. D., Marcus, D. K., Barry, C. T., & Coccaro, E. F. (2013). The latent structure of oppositional defiant disorder in children and adults. *Journal of Psychiatric Research*, 47(12), 1932–1939. <https://doi.org/10.1016/j.jpsychires.2013.08.016>
- Bolhuis, K., Lubke, G. H., Ende, J. van der, Bartels, M., Beijsterveldt, C. E. M. van, Lichtenstein, P., Larsson, H., Jaddoe, V. W. V., Kushner, S. A., Verhulst, F. C., Boomsma, D. I., & Tiemeier, H. (2017). Disentangling heterogeneity of childhood disruptive behavior problems into dimensions and subgroups. *Journal of the American Academy of Child & Adolescent Psychiatry*, 56(8), 678–686. <https://doi.org/10.1016/j.jaac.2017.05.019>
- Borelli, W. V., Carmona, K. C., Studart-Neto, A., Nitrini, R., Caramelli, P., & Costa, J. C. da. (2018). Operationalized definition of older adults with high cognitive performance. *Dementia & Neuropsychologia*, 12(3), 221–227. <https://doi.org/10.1590/1980-57642018dn12-030001>
- Briggs-Gowan, M. J., Godoy, L., Heberle, A., & Carter, A. S. (2016). Assessment of psychopathology in young children. In D. Cicchetti (Ed.), *Developmental psychopathology: Vols. 1: Theory and Method* (pp. 1–45).
- Broeren, S., Muris, P., Diamantopoulou, S., & Baker, J. R. (2013). The course of childhood anxiety symptoms: Developmental trajectories and child-related factors in normal children. *Journal of Abnormal Child Psychology*, 41(1), 81–95. <https://doi.org/10.1007/s10802-012-9669-9>
- Burt, S. A. (2012). How do we optimally conceptualize the heterogeneity within antisocial behavior? An argument for aggressive versus non-aggressive behavioral dimensions. *Clinical Psychology Review*, 32(4), 263–279. <https://doi.org/10.1016/j.cpr.2012.02.006>

- Caspi, A., & Shiner, R. L. (2006). Personality development. In N. Eisenberg, W. Damon, & R. M. Lerner (Eds.), *Handbook of child psychology* (6th ed., Vol. 3, pp. 300–365). John Wiley & Sons, Inc.
- Chen, F. R., & Jaffee, S. R. (2015). The heterogeneity in the development of homotypic and heterotypic antisocial behavior. *Journal of Developmental and Life-Course Criminology*, 1(3), 269–288. <https://doi.org/10.1007/s40865-015-0012-3>
- Chumlea, W. M. C., Choh, A., Lee, M., Towne, B., Sherwood, R. J., Duren, D., Czerwinski, S., & Siervogel, R. M. (2009). The first serial study into old age for weight, stature and BMI: The Fels Longitudinal Study. *The Journal of Nutrition, Health and Aging*, 13(1), 3–5. <https://doi.org/10.1007/s12603-009-0001-5>
- Cicchetti, D., & Rogosch, F. A. (2002). A developmental psychopathology perspective on adolescence. *Journal of Consulting and Clinical Psychology*, 70(1), 6–20. <https://doi.org/10.1037/0022-006X.70.1.6>
- Coghill, D., & Sonuga-Barke, E. J. S. (2012). Annual research review: Categories versus dimensions in the classification and conceptualisation of child and adolescent mental disorders: Implications of recent empirical study. *Journal of Child Psychology and Psychiatry*, 53(5), 469–489. <https://doi.org/10.1111/j.1469-7610.2011.02511.x>
- De Los Reyes, A. (2026). Toward integrative, multisource, and multimodal approaches to assessing psychopathology intergenerationally and across the lifespan: Commentary on Caspi et al. (2026). *Journal of Psychopathology and Clinical Science*, 135(4), 495–497. <https://doi.org/10.1037/abn0001084>
- Fairchild, G., Goozen, S. H. M. van, Calder, A. J., & Goodyer, I. M. (2013). Research review: Evaluating and reformulating the developmental taxonomic theory of antisocial behaviour. *Journal of Child Psychology and Psychiatry*, 54(9), 924–940. <https://doi.org/10.1111/jcpp.12102>
- Forbes, M. K., Tackett, J. L., Markon, K. E., & Krueger, R. F. (2016). Beyond comorbidity: Toward a dimensional and hierarchical approach to understanding psychopathology across

- the life span. *Development and Psychopathology*, 28(4pt1), 971–986.
<https://doi.org/10.1017/S0954579416000651>
- Godoy, L. L. de, Alves, C. A. P. F., Saavedra, J. S. M., Studart-Neto, A., Nitrini, R., Costa Leite, C. da, & Bisdas, S. (2021). Understanding brain resilience in superagers: A systematic review. *Neuroradiology*, 63(5), 663–683. <https://doi.org/10.1007/s00234-020-02562-1>
- Haslam, N., Holland, E., & Kuppens, P. (2012). Categories versus dimensions in personality and psychopathology: A quantitative review of taxometric research. *Psychological Medicine*, 42(05), 903–920. <https://doi.org/10.1017/S0033291711001966>
- Haslam, N., McGrath, M. J., Viechtbauer, W., & Kuppens, P. (2020). Dimensions over categories: A meta-analysis of taxometric research. *Psychological Medicine*, 50(9), 1418–1432.
<https://doi.org/10.1017/S003329172000183X>
- Insel, T. R. (2008). Assessing the economic costs of serious mental illness. *American Journal of Psychiatry*, 165(6), 663–665. <https://doi.org/10.1176/appi.ajp.2008.08030366>
- Kaat, A. J., Blackwell, C. K., Estabrook, R., Burns, J. L., Petitcherc, A., Briggs-Gowan, M. J., Gershon, R. C., Cella, D., Perlman, S. B., & Wakschlag, L. S. (2019). Linking the child behavior checklist (CBCL) with the multidimensional assessment profile of disruptive behavior (MAP-DB): Advancing a dimensional spectrum approach to disruptive behavior. *Journal of Child and Family Studies*, 28(2), 343–353.
<https://doi.org/10.1007/s10826-018-1272-4>
- Kagan, J. (1969). The three faces of continuity in human development. In D. A. Goslin (Ed.), *Handbook of socialization theory and research* (pp. 983–1002). Rand McNally.
- Kagan, J. (1971). *Change and continuity in infancy*. Wiley.
- Kagan, J. (1980). Perspectives on continuity. In O. G. Brim Jr. & J. Kagan (Eds.), *Constancy and change in human development* (pp. 26–74). Harvard University Press.
- Kagan, J., & Moss, H. A. (1962). *Birth to maturity: A study in psychological development*. Wiley.
- Kliem, S., Heinrichs, N., Lohmann, A., Bussing, R., Schwarzer, G., & Briegel, W. (2018). Dimensional latent structure of early disruptive behavior disorders: A taxometric analysis

- in preschoolers. *Journal of Abnormal Child Psychology*, 46, 1385–1394.
<https://doi.org/10.1007/s10802-017-0383-5>
- Kliem, S., Krieg, Y., Klatt, T., & Baier, D. (2022). Dimensional latent structure of callous-unemotional traits in german adolescents: Results from taxometric analyses. *Research on Child and Adolescent Psychopathology*, 50(6), 771–780.
<https://doi.org/10.1007/s10802-021-00885-y>
- Kotov, R., Krueger, R. F., Watson, D., Achenbach, T. M., Althoff, R. R., Bagby, R. M., Brown, T. A., Carpenter, W. T., Caspi, A., Clark, L. A., Eaton, N. R., Forbes, M. K., Forbush, K. T., Goldberg, D., Hasin, D., Hyman, S. E., Ivanova, M. Y., Lynam, D. R., Markon, K., ... Zimmerman, M. (2017). The hierarchical taxonomy of psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *Journal of Abnormal Psychology*, 126(4), 454–477. <https://doi.org/10.1037/abn0000258>
- Krueger, R. F., & DeYoung, C. G. (2016). The RDoC initiative and the structure of psychopathology. *Psychophysiology*, 53(3), 351–354. <https://doi.org/10.1111/psyp.12551>
- Krueger, R. F., Hobbs, K. A., Conway, C. C., Dick, D. M., Dretsch, M. N., Eaton, N. R., Forbes, M. K., Forbush, K. T., Keyes, K. M., Latzman, R. D., Michelini, G., Patrick, C. J., Sellbom, M., Slade, T., South, S. C., Sunderland, M., Tackett, J. L., Waldman, I., Waszczuk, M. A., ... Workgroup, H. U. (2021). Validity and utility of hierarchical taxonomy of psychopathology (HiTOP): II. Externalizing superspectrum. *World Psychiatry*, 20(2), 171–193. <https://doi.org/10.1002/wps.20844>
- Krueger, R. F., & Markon, K. E. (2011). A dimensional-spectrum model of psychopathology: Progress and opportunities. *Archives of General Psychiatry*, 68(1), 10–11.
<https://doi.org/10.1001/archgenpsychiatry.2010.188>
- Krueger, R. F., Markon, K. E., Patrick, C. J., Benning, S. D., & Kramer, M. D. (2007). Linking antisocial behavior, substance use, and personality: An integrative quantitative model of the adult externalizing spectrum. *Journal of Abnormal Psychology*, 116(4), 645–666.
<https://doi.org/10.1037/0021-843X.116.4.645>

- Krueger, R. F., Markon, K. E., Patrick, C. J., & Iacono, W. G. (2005). Externalizing psychopathology in adulthood: A dimensional-spectrum conceptualization and its implications for DSM-v. *Journal of Abnormal Psychology, 114*(4), 537–550. <https://doi.org/10.1037/0021-843x.114.4.537>
- Krueger, R. F., Nichol, P. E., Hicks, B. M., Markon, K. E., Patrick, C. J., Iacono, W. G., & McGue, M. (2004). Using latent trait modeling to conceptualize an alcohol problems continuum. *Psychological Assessment, 16*(2), 107–119. <https://doi.org/10.1037/1040-3590.16.2.107>
- Krueger, R. F., & Piasecki, T. M. (2002). Toward a dimensional and psychometrically-informed approach to conceptualizing psychopathology. *Behaviour Research and Therapy, 40*(5), 485–499. [https://doi.org/10.1016/S0005-7967\(02\)00016-5](https://doi.org/10.1016/S0005-7967(02)00016-5)
- Markon, K. E., Chmielewski, M., & Miller, C. J. (2011). The reliability and validity of discrete and continuous measures of psychopathology: A quantitative review. *Psychological Bulletin, 137*(5), 856–879. <https://doi.org/10.1037/a0023678>
- Markon, K. E., & Krueger, R. F. (2005). Categorical and continuous models of liability to externalizing disorders: A direct comparison in NESARC. *Archives of General Psychiatry, 62*(12), 1352–1359. <https://doi.org/10.1001/archpsyc.62.12.1352>
- Miller, J. L., Vaillancourt, T., & Boyle, M. H. (2009). Examining the heterotypic continuity of aggression using teacher reports: Results from a national canadian study. *Social Development, 18*(1), 164–180. <https://doi.org/10.1111/j.1467-9507.2008.00480.x>
- Moffitt, T. E. (1993). Adolescence-limited and life-course-persistent antisocial behavior: A developmental taxonomy. *Psychological Review, 100*(4), 674–701. <https://doi.org/10.1037/0033-295X.100.4.674>
- Moss, H. A., & Kagan, J. (1964). Report on personality consistency and change from the Fels Longitudinal Study. *Vita Humana, 7*(2), 127–138. <http://www.jstor.org/stable/26761430>
- Odgers, C. L., Moffitt, T. E., Broadbent, J. M., Dickson, N., Hancox, R. J., Harrington, H., Poulton, R., Sears, M. R., Thomson, W. M., & Caspi, A. (2008). Female and male antisocial trajectories: From childhood origins to adult outcomes. *Development and*

- Psychopathology*, 20(2), 673–716. <https://doi.org/10.1017/S0954579408000333>
- Paalman, C. H., Terwee, C. B., Jansma, E. P., & Jansen, L. M. C. (2013). Instruments measuring externalizing mental health problems in immigrant ethnic minority youths: A systematic review of measurement properties. *PLOS ONE*, 8(5), e63109. <https://doi.org/10.1371/journal.pone.0063109>
- Patterson, G. R. (1993). Orderly change in a stable world: The antisocial trait as a chimera. *Journal of Consulting and Clinical Psychology*, 61(6), 911–919. <https://doi.org/10.1037/0022-006X.61.6.911>
- Pavlovich, K., Tiego, J., Constable, T., Fornito, A., & Imaging, M. B. (2026). Assessing the psychometric properties of the child behavior checklist in the ABCD study. *Developmental Science*, 29(4), e70216. <https://doi.org/10.1111/desc.70216>
- Petersen, I. T. (2024). Reexamining developmental continuity and discontinuity in the 21st century: Better aligning behaviors, functions, and mechanisms. *Developmental Psychology*, 60(11), 1992–2007. <https://doi.org/10.1037/dev0001657>
- Petersen, I. T., Bates, J. E., Dodge, K. A., Lansford, J. E., & Pettit, G. S. (2015). Describing and predicting developmental profiles of externalizing problems from childhood to adulthood. *Development and Psychopathology*, 27(3), 791–818. <https://doi.org/10.1017/S0954579414000789>
- Petersen, I. T., Choe, D. E., & LeBeau, B. (2020). Studying a moving target in development: The challenge and opportunity of heterotypic continuity. *Developmental Review*, 58, 100935. <https://doi.org/10.1016/j.dr.2020.100935>
- Petersen, I. T., Demko, Z., Lee, W.-C., & Oleson, J. J. (2026). Studying development of psychopathology using changing measures to account for heterotypic continuity. *JAACAP Open*, 4(1), 111–123. <https://doi.org/10.1016/j.jaacop.2025.10.008>
- Petersen, I. T., LeBeau, B., & Choe, D. E. (2021). Creating a developmental scale to account for heterotypic continuity in development: A simulation study. *Child Development*, 92(1), e1–e19. <https://doi.org/10.1111/cdev.13433>

- Petersen, I. T., Lindhiem, O., LeBeau, B., Bates, J. E., Pettit, G. S., Lansford, J. E., & Dodge, K. A. (2018). Development of internalizing problems from adolescence to emerging adulthood: Accounting for heterotypic continuity with vertical scaling. *Developmental Psychology*, 54(3), 586–599. <https://doi.org/10.1037/dev0000449>
- Roche, A. F. (1992). *Growth, maturation, and body composition: The Fels Longitudinal Study 1929–1991*. Cambridge University Press.
- Rutter, M., Kim-Cohen, J., & Maughan, B. (2006). Continuities and discontinuities in psychopathology between childhood and adult life. *Journal of Child Psychology and Psychiatry*, 47(3-4), 276–295. <https://doi.org/10.1111/j.1469-7610.2006.01614.x>
- Schaeffer, N. C. (1991). Hardly ever or constantly? Group comparisons using vague quantifiers. *Public Opinion Quarterly*, 55(3), 395–423. <https://doi.org/10.1086/269270>
- Schwarz, N. (1999). Self-reports: How the questions shape the answers. *American Psychologist*, 54(2), 93–105. <https://doi.org/10.1037/0003-066X.54.2.93>
- Sellbom, M. (2016). Elucidating the validity of the externalizing spectrum of psychopathology in correctional, forensic, and community samples. *Journal of Abnormal Psychology*, 125(8), 1027–1038. <https://doi.org/10.1037/abn0000171>
- Sivertsson, F., Carlsson, C., Almquist, Y. B., & Brännström, L. (2024). Offending trajectories from childhood to retirement age: Findings from the Stockholm birth cohort study. *Journal of Criminal Justice*, 91, 102155. <https://doi.org/10.1016/j.jcrimjus.2024.102155>
- Sroufe, L. A. (2016). The place of attachment in development. In J. Cassidy & P. Shaver (Eds.), *Handbook of attachment: Theory, research, and clinical applications* (Vol. 3, pp. 997–1011). Guilford.
- Steffensmeier, D., Slepicka, J., & Schwartz, J. (2025). International and historical variation in the age–crime curve. *Annual Review of Criminology*, 8, 239–268. <https://doi.org/10.1146/annurev-criminol-111523-122451>
- Sterba, S. K., Prinstein, M. J., & Cox, M. J. (2007). Trajectories of internalizing problems across childhood: Heterogeneity, external validity, and gender differences. *Development and*

- Psychopathology*, 19(2), 345–366. <https://doi.org/10.1017/S0954579407070174>
- Tackett, J. L., & Hallquist, M. (2022). The need to grow: Developmental considerations and challenges for modern psychiatric taxonomies. *Journal of Psychopathology and Clinical Science*, 131(6), 660–663. <https://doi.org/10.1037/abn0000751>
- Tiego, J., Martin, E. A., DeYoung, C. G., Hagan, K., Cooper, S. E., Pasion, R., Satchell, L., Shackman, A. J., Bellgrove, M. A., Fornito, A., Abend, R., Goulter, N., Eaton, N. R., Kaczkurkin, A. N., Nusslock, R., & the HiTOP Neurobiological Foundations Work Group. (2023). Precision behavioral phenotyping as a strategy for uncovering the biological correlates of psychopathology. *Nature Mental Health*, 1(5), 304–315. <https://doi.org/10.1038/s44220-023-00057-5>
- Wakschlag, L. S., Briggs-Gowan, M. J., Choi, S. W., Nichols, S. R., Kestler, J., Burns, J. L., Carter, A. S., & Henry, D. (2014). Advancing a multidimensional, developmental spectrum approach to preschool disruptive behavior. *Journal of the American Academy of Child & Adolescent Psychiatry*, 53(1), 82–96.e3. <https://doi.org/10.1016/j.jaac.2013.10.011>
- Wakschlag, L. S., Roberts, M. Y., Flynn, R. M., Smith, J. D., Krogh-Jespersen, S., Kaat, A. J., Gray, L., Walkup, J., Marino, B. S., Norton, E. S., & Davis, M. M. (2019). Future directions for early childhood prevention of mental disorders: A road map to mental health, earlier. *Journal of Clinical Child & Adolescent Psychology*, 48(3), 539–554. <https://doi.org/10.1080/15374416.2018.1561296>
- Walters, G. D. (2011). The latent structure of life-course-persistent antisocial behavior: Is moffitt's developmental taxonomy a true taxonomy? *Journal of Consulting and Clinical Psychology*, 79(1), 96–105. <https://doi.org/10.1037/a0021519>
- Walters, G. D. (2012). Developmental trajectories of delinquent behavior: One pattern or several? *Criminal Justice and Behavior*, 39(9), 1192–1203. <https://doi.org/10.1177/0093854812443645>
- Walters, G. D. (2014). Continuous versus categorical models of delinquency risk. *American Journal of Criminal Justice*, 39(3), 395–410. <https://doi.org/10.1007/s12103-013-9235-1>

- Walters, G. D. (2015). The latent structure of criminal persistence: A taxometric analysis of offending behavior from late adolescence to early adulthood in adjudicated male delinquents. *American Journal of Criminal Justice*, 40(3), 542–559.
<https://doi.org/10.1007/s12103-014-9272-4>
- Walters, G. D., & Ruscio, J. (2013). Trajectories of youthful antisocial behavior: Categories or continua? *Journal of Abnormal Child Psychology*, 41(4), 653–666.
<https://doi.org/10.1007/s10802-012-9700-1>
- Walton, K. E., Ormel, J., & Krueger, R. F. (2011). The dimensional nature of externalizing behaviors in adolescence: Evidence from a direct comparison of categorical, dimensional, and hybrid models. *Journal of Abnormal Child Psychology*, 39(4), 553–561.
<https://doi.org/10.1007/s10802-010-9478-y>
- Weems, C. F. (2008). Developmental trajectories of childhood anxiety: Identifying continuity and change in anxious emotion. *Developmental Review*, 28(4), 488–502.
<https://doi.org/10.1016/j.dr.2008.01.001>
- Weiss, B., & Garber, J. (2003). Developmental differences in the phenomenology of depression. *Development and Psychopathology*, 15(2), 403–430.
<https://doi.org/10.1017/S0954579403000221>
- World Health Organization. (2022). *International Statistical Classification of Diseases and Related Health Problems 11th revision (ICD-11)*. <https://icd.who.int>
- Wright, A. G. C., Krueger, R. F., Hobbs, M. J., Markon, K. E., Eaton, N. R., & Slade, T. (2013). The structure of psychopathology: Toward an expanded quantitative empirical model. *Journal of Abnormal Psychology*, 122(1), 281–294. <https://doi.org/10.1037/a0030133>

Appendix A

This Section Is an Appendix

Appendix B

Another Appendix