Paediatric Urology Rapid Revision

Campbell-Walsh-Wein · Chapters 41, 49–53, 55–57, 59, 61, 63, 64

For MCh Urology theory & viva · NEET-SS / competitive MCQs

How to use (total under 7 h even at a slow, careful pace)

- Each chapter reads top-to-bottom in the order you would write an essay answer: definition → pathology → diagnosis → management → complications → follow-up. The same material serves theory and viva.
- Boxed **ALGORITHM** and shaded **KEY POINT** panels carry the reasoning examiners ask about ("why do we…?").
- **MCQ one-liners** end each chapter: re-read these the day before the paper.
- The last section puts the big treatment choices side by side for essay questions.

Conventions: **underlined bold** = number to memorise · MCQ = exam trivia only · → *Ch 53* = cross-reference · † = not in the source chapter notes (added because it is essential in practice). Abbreviations are spelled out at first use in each chapter.

Chapter	Topic	Slow-reader time
41	Vesicoureteral Reflux	33 min
49	Anomalies of the Kidney & Collecting System	31 min
50	Renal Dysgenesis & Cystic Disease	31 min
51	Pathophysiology of Urinary Tract Obstruction	18 min
52	Ectopic Ureter, Ureterocele & Ureteral Anomalies	37 min
53	Surgery of the Ureter in Children (PUJ, megaureter, VUR surgery)	40 min
55	External Genitalia in Males & Acute Scrotum	37 min
56	Hypospadias	34 min
57	Undescended Testis	35 min
59	Differences of Sexual Development	33 min
61	Paediatric Renal Transplantation	18 min
63	Renal & Adrenal Tumours (Wilms, Neuroblastoma)	33 min
64	Other Paediatric Urological Tumours	24 min
★	Putting it together: common paediatric decisions	8 min
	Total (28794 words, at 70 words/min)	**6h 52min**

41 Management Strategies for Vesicoureteral Reflux

Reading time ≈ 33 min (slow pace)

1 · DEFINITION AND CONCEPT

- **Vesicoureteral reflux (VUR)**: retrograde flow of urine from bladder to upper tract. Associated with **acquired scarring** and with **congenital renal dysplasia**.
- Now seen as a **spectrum of severity** needing risk-stratified management, not a yes/no disease.
- **Infection, not reflux, is the injury.** VUR does not cause urinary tract infection (UTI); it is an accelerant that carries infected bladder urine to the kidney. Sterile urine and good emptying are the targets of treatment.

2 · EMBRYOLOGY AND ANTIREFLUX MECHANISM

- Ureteric bud arises from the mesonephric (wolffian) duct at weeks 4–6; the common duct is absorbed into the urogenital sinus (UGS). **Mackie–Stephens**: bud position sets both orifice and kidney.
  - **Early bud** → over-rotation → orifice **high and lateral**, short tunnel → VUR.
  - **Late bud** → orifice **distal and medial** → ectopic, often obstructed ureter.
  - Abnormal bud–metanephros contact → dysplasia/hypoplasia. Hence high-grade VUR and dysplasia travel together and some "scars" are congenital.
- VUR = imbalance of three factors: ureteric peristalsis, ureterovesical junction (UVJ) anatomy, bladder dynamics.
- **Passive flap-valve**: intramural ureter compressed against detrusor backing during filling. **Paquin**: tunnel length : ureteric diameter **5:1** in non-refluxing vs 1.4:1 in refluxing UVJs; orifice configuration matters as much.
- Opening is active (longitudinal tunnel muscle shortens/widens the tunnel); closure is passive.

3 · PRIMARY VS SECONDARY VUR

	Primary	Secondary
Fault	Congenital UVJ deficiency, normal low-pressure bladder	Normal-ish UVJ overwhelmed by pressure, or abnormal UVJ
Causes	Short tunnel, weak detrusor backing, dilated/stiff ureter	Outlet obstruction (posterior urethral valves (PUV), stricture); neurogenic dyssynergia; dysfunctional voiding; ureterocele, duplication, paraureteral diverticulum, iatrogenic
Treat	Wait for resolution / correct	Treat the cause first

- Spina bifida: VUR in up to **30%** → early clean intermittent catheterisation (CIC), anticholinergic or β3-agonist, serial imaging. Look for occult dysraphism (sacral dimple, hairy patch, gluteal cleft anomaly, refractory constipation).
- **Infant male VUR**: high voiding pressures from poor sphincter relaxation; resolves as voiding matures — explains infant male predominance and high male resolution.
- Normal bladders void at ~100 cm H_2O without reflux, so structural UVJ failure is key. **Reflux at low pressure / early filling = poor prognosis for resolution.**
- Detrusor overactivity is the commonest urodynamic finding; treating it (timed voiding, biofeedback, anticholinergics, α-blockers) improves resolution.

> **KEY POINT** Primary and secondary are not exclusive: "primary" reflux often has a treatable secondary component (overactivity, bladder and bowel dysfunction). Removing it leaves only the true primary VUR to manage.

4 · EPIDEMIOLOGY (HEADLINE)

- VUR in **~30%** of children with UTI; ~17% of those imaged for hydronephrosis. Younger child with UTI → likelier VUR (**38.6%** under 2 y vs 7.6% at 12–21 y).
- Girls ~2× overall, but **infant males predominate** (up to 3:1 at 0–6 months; uncircumcised boys have 10–12× more UTI). Less common in Black children.

5 · GRADING — INTERNATIONAL REFLUX STUDY (1981)

Grade	Ureter	Pelvis and calyces
I	Into non-dilated ureter only	Not reached
II	Not dilated	Reached, no dilatation
III	Mild–moderate dilatation	Mild–moderate dilatation, minimal forniceal blunting
IV	Moderate tortuosity	Moderate dilatation
V	Gross dilatation and tortuosity	Gross dilatation; **papillary impressions lost**

- I–II non-dilating; III–V dilating. Interobserver agreement poor (all three readers in RIVUR (Randomized Intervention for Children with Vesicoureteral Reflux) agreed in only ~60%, worst II–IV).
- Radionuclide cystography (RNC) has 3 grades: 1 = I; 2 = II–III; 3 = IV–V.
- **Ureteral diameter ratio (UDR)** = largest ureteric diameter in false pelvis ÷ L1–L3 vertebral distance. Predicts resolution and breakthrough UTI (BUTI) better than grade.

6 · DIAGNOSIS

When to image after UTI

- Accurate urine collection and UTI confirmation first — major decisions (including surgery) rest on it.
- American Academy of Pediatrics (AAP) 2011, febrile UTI 2–24 months: **ultrasound (US) first**; voiding cystourethrogram (VCUG) only if ultrasound abnormal or **second febrile UTI**.
- National Institute for Health and Care Excellence (NICE, UK): US; VCUG mainly for atypical/recurrent UTI in younger children; none routinely >3 y. Misses much VUR including grade IV–V.
- **Top-down approach**: dimercaptosuccinic acid (DMSA) scan first; VCUG only if abnormal. Normal DMSA → **<1%** chance of high-grade VUR; misses ~1/3 of (mostly low-grade) VUR.

Cystographic methods

Method	Strength	Weakness
VCUG (gold standard; gravity ≤70 cm H_2O, filling + voiding fluoroscopy)	Anatomy: grade, intrarenal reflux, diverticula, ureterocele, urethra, postvoid drainage	Catheter, radiation
Direct RNC (technetium-99m pertechnetate)	More sensitive for II–V; low dose; ideal for screening and follow-up	Misses grade I; little anatomy
Indirect RNC	No catheter	Needs toilet-trained child; sensitivity 70–90%
Contrast-enhanced voiding urosonography	No radiation; comparable to VCUG; detects intrarenal reflux	Grading not standardised

- **Cyclic VCUG** (2–3 fill–void cycles) detects 12–20% more VUR.
- **Filling-phase reflux** = poor sign for resolution; voiding-only reflux is more favourable.
- Timing: once urine sterile and child improved, early VCUG (~8 days) gives same VUR prevalence as delayed.

- **Postvoid films**: retained pelvic contrast → suspect ureteropelvic junction obstruction (UPJO).
- Suspect coexisting **UVJ obstruction**: dilated proximal ureter, distal aperistaltic segment, dilute contrast, delayed drainage → **relative contraindication to endoscopic injection** (may worsen obstruction).
- Routine cystoscopy has no role (orifice shape correlates poorly with VUR); use just before open surgery.
- Non-invasive uroflowmetry + postvoid residual (PVR) ± electromyography (EMG) is useful; invasive urodynamics only for refractory cases.

Upper tract

- **Renal and bladder ultrasound (RBUS)**: initial and serial mainstay (size, growth, echogenicity). Cannot diagnose or exclude VUR.
- **DMSA**: gold standard for cortical defects (sensitivity 98%, specificity 92%). Acute photopenia may resolve; persistence = scar.

7 · CONGENITAL DYSPLASIA VS ACQUIRED SCAR

	Reflux-associated dysplasia	Acquired scar
Cause	Abnormal bud–blastema interaction	Pyelonephritis: infected urine + VUR
Typical patient	Male infant, grade IV–V, antenatal hydronephrosis, no UTI	Girl, recurrent febrile UTI, bladder and bowel dysfunction
Imaging	Small kidney, globally reduced function	Focal photopenic defects after infection
Preventable?	No	Yes — keep urine sterile

- Prenatally detected boys: damage confined to grades IV–V (nearly half of grade V severely damaged) before any infection.
- **Ransley–Risdon** (pig): reflux caused nephropathy only with infected urine. Sterile VUR damages only with high pressure (PUV, Hodson).
- **Compound (concave) papillae** at poles → intrarenal reflux → **polar scars**. Intrarenal reflux occurs at only 2 mm Hg in neonates.
- Scarring risk highest in first year. "Big bang" theory (most scarring with first pyelonephritis) is now challenged.
- Long-term: hypertension in **34–38%** of scarred children. **Correcting VUR does not lower blood pressure** (parenchymal mechanism); nephrectomy/segmental resection of a small kidney may.

8 · BLADDER AND BOWEL DYSFUNCTION (BBD)

- Symptoms: frequency, holding, urgency, incontinence, constipation; curtsying (girls), penile squeezing (boys). **Treat constipation first** — it causes co-contraction of anal and urinary sphincters.
- American Urological Association (AUA) guideline: BBD is one of the most critical modifiable variables. BBD → more UTI on prophylaxis and after surgery, less resolution, **lower endoscopic success, but no reduction in open surgical success**.
- Off prophylaxis, recurrent UTI: ~1/2 with BBD + VUR, ~1/5 VUR alone, ~1/3 BBD alone.
- **Phimosis**: in boys with VUR number needed to treat (NNT) by circumcision to prevent one UTI falls from ~111 to **~4**; topical steroid is equivalent.

9 · NATURAL HISTORY AND RESOLUTION

Grade	Spontaneous resolution
I–II	Most; grade II 63–85%
III	~50%; ~90% of resolutions within 4 years

Grade	Spontaneous resolution
IV–V, bilateral III	Few (but infants with IV–V resolve well in year 1)

- Mechanism: UVJ remodelling (tunnel elongation) + stabilising bladder dynamics. Persistent BBD explains failure.
- Predictors of resolution: **low grade, young age, unilateral, no BBD, voiding-only reflux, higher bladder volume at onset, low UDR**. Age at presentation matters as much as grade (AUA 1997 nomograms).

10 · MEDICAL MANAGEMENT — CONTINUOUS ANTIBIOTIC PROPHYLAXIS (CAP)

- Principle: **manage infection risk, not VUR**; prevent UTI while awaiting resolution.
- Daily low-dose oral CAP, **at night** — longest bladder concentration during longest retention.

Agent	Age	Note
Trimethoprim / amoxicillin	<2 months	Avoids sulfonamide in neonate
Trimethoprim-sulfamethoxazole (TMP-SMX)	>2 months	RIVUR agent; sulfonamide displaces bilirubin → neonatal jaundice; rare reversible leukopenia
Nitrofurantoin	>2 months	Least faecal-flora resistance; pulmonary fibrosis, hepatic toxicity (rare at low dose)

- **Smellie's BUTI rule**: organism sensitive to the prophylactic → non-adherence/low dose; resistant → residual volume too high or dose too high.
- **Breakthrough febrile UTI on CAP despite good bladder and bowel habits → stop medical management and correct VUR** (confirm proper urine collection first).
- Stopping CAP after toilet training/BBD resolution: ~8–10% get febrile UTI → reassess, possibly correct.

Risk stratification (Hidas)

Factor	Low risk	High risk
Sex / age	Male; >5 y	Female; <1 y
Laterality / grade	Unilateral; I–II	Bilateral; III–V
BBD / scars	No	Yes
Presentation	Prenatal hydronephrosis	Multiple febrile UTIs

- 2-year BUTI on CAP: **8.6% low, 26% intermediate, 62.5% high**. About two-thirds of children are low risk.

11 · PRACTICAL ALGORITHM (AUA-BASED, RISK-STRATIFIED)

ALGORITHM

History of number/severity of true febrile UTIs + RBUS → VCUG if multiple febrile UTIs, admissions or upper-tract findings (afebrile, doubtful UTI with normal US → conservative)

All: treat BBD (constipation first), hygiene, timed voiding

Low risk (grade I–II, no BBD, male, older) → observation ± CAP; treat febrile UTI promptly

High risk (grade ≥III, girl, BBD, febrile presentation, scars, infant) → CAP

Breakthrough febrile UTI, new scars, high grade with poor function, persistent VUR with low tolerance for infection → correction (endoscopic or reimplant)

- Evidence summary: little CAP benefit at grade ≤II without BBD; benefit at grade ≥III, at least in girls.
- Extensive scarring, globally reduced function or dysplasia lower tolerance for another infection → favour correction.

- Unresolved: asymptomatic bilateral high-grade VUR in an older boy; correction of persistent VUR in girls before pregnancy (unproven).

12 · SURGICAL CORRECTION

- Open correction succeeds in **>95%**; robotic correction is probably less morbid. No superiority of surgery over medical therapy for scarring or renal growth (International Reflux Study in Children (IRSC), Birmingham).
- After correction, almost no new scars despite asymptomatic bacteriuria — but adult UTI risk persists.
- **Endoscopic injection** of dextranomer/hyaluronic acid copolymer (Dx/HA): BBD lowers success; avoid with UVJ obstruction or obstructive refluxing megaureter (may obstruct). Mounds calcify in over half and **mimic ureteric stones** later.
- Surgical techniques — endoscopic injection, intravesical/extravesical reimplantation, complications → *Ch 53*.
- After cross-trigonal reimplant, **retrograde catheterisation and ureteroscopy are difficult** — know the surgical history.

13 · ASSOCIATED CONDITIONS

- **UPJO + VUR**: VCUG signs — dilated ureter with little pelvic filling, dilute pelvic contrast, ureter–pelvis discrepancy, pelvis not draining postvoid. **Correct obstruction first**; simultaneous open repair of both risks ureteric blood supply.
- **Duplex**: VUR usually into **lower pole** (Weigert–Meyer: lateral, proximal orifice, short tunnel). Duplication alone does not increase risk → conservative management reasonable.
- **Paraureteral diverticulum**: resolves like primary VUR. **Ureter entering diverticulum**: no muscle backing → will not resolve.
- Multicystic dysplastic kidney (MCDK): contralateral VUR ~1/4, usually resolves.
- **Megacystis–megaureter**: massive bilateral VUR recycles urine → large thin bladder; normal posterior urethra. Treat by reimplant (not outlet surgery).
- **Obstructive refluxing megaureter**: ureteric contrast more dilute than bladder; avoid injection; reimplant ± tapering.

14 · SIBLING SCREENING

- Sibling prevalence ~**30%** (up to 50%), higher in younger siblings; identical twins near-complete concordance.
- Target is **renal damage, not VUR**. AUA: **VCUG for siblings with cortical abnormality or renal asymmetry on US, or UTI history**; otherwise observe. Older sibling with normal kidneys and no BBD gains little — treat any febrile UTI promptly.

15 · FOLLOW-UP AND ADULTHOOD

- Lifelong follow-up if scarred: blood pressure, proteinuria, renal function; transition programmes.
- **Pregnancy**: VUR → pyelonephritis in ~1/3; renal scars → ~3× hypertension, **~8× pre-eclampsia**. **Scarring, not VUR presence, drives pregnancy morbidity.** Reimplanted women still at UTI risk but no excess miscarriage.

16 · LANDMARK TRIALS

Trial	Design	Result	Take-home
IRSC	<9 y, high grade; CAP vs open surgery	UTI equal (38%); surgery fewer pyelonephritis; new scars equal; 10 y renal function equal	Surgery ≈ medical for scarring

Trial	Design	Result	Take-home
Birmingham Reflux Study	High grade; medical vs surgical, 5 y	New scars equal, all within 2 y	Equal; supports "big bang"
Swedish Reflux Trial	1–2 y, grade III–IV; CAP vs endoscopic vs surveillance	Girls febrile UTI: 19% CAP, 23% endoscopic, **57%** surveillance; new scars mostly on surveillance, none on CAP; boys no difference	Treat girls with dilating VUR
RIVUR	607 children, grade I–IV after first/second UTI; TMP-SMX vs placebo	Recurrent UTI **12.9% vs 23.6%**; scarring unchanged; benefit greatest with febrile index UTI/BBD	CAP prevents UTI, not scars
PRIVENT	576 children with UTI; CAP vs placebo	Absolute UTI reduction 6% (19% → 13%)	Modest benefit
French / low-risk trials	Grade I–III; CAP vs none	No significant difference	No benefit at low risk

KEY POINT Why do the trials disagree? Population risk. RIVUR was diluted by low-risk children (only ~8% grade IV); reanalysis gives NNT 5 in high-risk vs 18 in low-risk. Prophylaxis reduces infection in proportion to baseline risk; neither CAP nor surgery prevents scarring better than the other.

17 · MCQ ONE-LINERS

1. Early ureteric budding → lateral, high orifice → short tunnel → VUR.
2. Paquin ratio of tunnel length to ureteric diameter in non-refluxing UVJ: 5:1.
3. VUR prevalence in children with UTI: ~30%.
4. Grade V VUR: loss of papillary impressions.
5. First dilating grade on international classification: III.
6. Direct RNC misses grade I VUR.
7. Cyclic VCUG detects 12–20% more VUR.
8. Reflux during filling phase predicts poor spontaneous resolution.
9. Normal DMSA: <1% chance of high-grade VUR.
0. Compound papillae at poles allow intrarenal reflux and polar scars.
1. Ransley–Risdon pig model: scarring needs infected urine.
2. Congenital reflux nephropathy: male infant, grade IV–V, no UTI.
3. VUR correction does not cure reflux-related hypertension.
14. BBD lowers endoscopic but not open reimplant success.
15. Circumcision NNT to prevent one UTI in boys with VUR: ~4.
16. RIVUR agent: trimethoprim-sulfamethoxazole; halved recurrent UTI, no effect on scarring.
17. Swedish trial: surveillance girls had 57% febrile UTI recurrence.
18. Avoid sulfonamides under 2 months: bilirubin displacement → neonatal jaundice.
19. Breakthrough febrile UTI on CAP: indication for surgical correction.
20. Hidas high-risk group: 62.5% BUTI at 2 years.
21. Duplex VUR usually affects the lower pole.
22. Ureter entering diverticulum: VUR will not resolve.
23. UPJO with VUR: correct obstruction first.
24. Megacystis–megaureter: treat by ureteric reimplantation, not outlet surgery.
25. Calcified Dx/HA mound may mimic distal ureteric stone.
26. Pregnancy morbidity is driven by renal scarring, not VUR.

49 Anomalies of the Kidney and Collecting System

Reading time ≈ 31 min (slow pace)

1 · EMBRYOLOGY THAT EXPLAINS EVERYTHING

- Ureteric bud (UB) arises from the caudal wolffian duct (WD) at **5–7 wk**, invades the metanephric mesenchyme (MM) and branches into the collecting system; UB tips induce nephrons (**reciprocal induction**).
- Ascent with 90° ventromedial rotation: weeks 6–9; kidney ends below the adrenal, calyces lateral, pelvis medial.
- Renal and genital development are integrated → renal anomalies travel with WD (male) and müllerian duct (MD, female) anomalies. Adrenal (cortex from mesoderm medial to the ridge, medulla from neural crest) is independent of the kidney.

> **KEY POINT** Every anomaly of number is a failure of UB–MM induction: timing and extent set the phenotype. An early insult drags the WD/MD with it (genital anomalies). Arrested ascent leaves an unrotated kidney with anterior pelvis and fetal blood supply; fusion freezes kidneys where they meet.

2 · BILATERAL RENAL AGENESIS (BRA)

- **≈1 in 5000** births; **75% male**. First-degree relatives ≈1000× population risk → **ultrasound (US) parents and siblings** of any infant with unilateral renal agenesis (URA), BRA or dysgenesis.
- **Pathology**: kidneys absent; ureters atretic; bladder absent (≈½) or hypoplastic. Adrenal present but flattened — **"lying down" sign** on US. Bladder fails because UB/WD entry into the cloaca initiates bladder development (favoured over "lack of urine").
- **Potter facies** (sine qua non of absent amniotic fluid): senile look, semicircular fold below inner canthus, blunted nose, low-set-appearing ears; claw hands, bowed clubbed legs. These are **deformations**, not malformations (absent in a BRA twin sharing a normal sac).
- **Pulmonary hypoplasia** kills. Peters two-step model: early "renal factor" plus amniotic fluid in the canalicular phase (16–26 wk).
- **Diagnosis**: prenatal US — severe oligohydramnios, no kidneys, small chest. Neonate: **anuria >24 h without bladder distension** = BRA until proved otherwise. Postnatal US + power Doppler; equivocal → technetium-99m dimercaptosuccinic acid (DMSA) scan: no uptake anywhere confirms.
- **Prognosis**: ≈40% stillborn; rest die in 24–48 h of respiratory failure. Amnioinfusion is experimental (RAFT trial, §9).

> **KEY POINT** BRA is lethal through the lungs, not the kidneys. Amnioinfusion replaces only the amniotic step, so it converts pulmonary deaths into neonatal end-stage renal disease (ESRD).

3 · UNILATERAL RENAL AGENESIS AND THE SOLITARY FUNCTIONING KIDNEY (SFK)

- **1 in 1200**; male 1.8:1; left slightly more. Many apparent URAs are **involuted multicystic dysplastic kidneys (MCDK)** (clue: curvilinear calcification).
- Associations: VACTERL (vertebral, anal, cardiac, tracheo-oesophageal, renal, limb; ≈30% have URA), MURCS, Turner, Kallmann, branchio-oto-renal, DiGeorge (22q11). **Maternal diabetes triples risk**. Isolated preauricular pit/tag or single umbilical artery → no routine renal US; ≥2 anomalies → US; recognised complex → image all systems incl. spine.
- **Ipsilateral ureter absent in ≈60%**.
- **Contralateral tract**: vesicoureteric reflux (VUR) **≈30%**, pelviureteric junction (PUJ) obstruction ≈11%, vesicoureteric junction obstruction ≈7%.

- **Genital anomalies 20–40%**: females 25–50%, males 10–15% (although URA is commoner in males). Gonads usually normal; adrenal agenesis <10%.

	Male (WD derivatives)	Female (MD derivatives)
Always present	Testis, head of epididymis	Fimbriated end of tube
Commonest	Absent vas/epididymal body-tail/seminal vesicle (≈50%). **79% of men with absent vas lack the ipsilateral kidney**	Unicornuate/bicornuate, didelphys or septate uterus; vaginal atresia or duplication
Named syndrome	**Zinner**: seminal vesicle cyst (ejaculatory duct atresia) + ipsilateral agenesis	**OHVIRA** (obstructed hemivagina, ipsilateral renal anomaly; Herlyn–Werner–Wunderlich): pubertal haematocolpos, pelvic mass, pain
Trigger to image kidneys	Impalpable vas in infertility work-up, or at herniotomy/orchidopexy	Any uterine anomaly; pelvic US after thelarche, before menarche

- **Magee classification** (URA + müllerian anomaly, by timing): I (before wk 4, whole ridge) → unicornuate uterus; II (early wk 4, WD + UB) → didelphys with obstructed hemivagina on agenesis side; III (after wk 4, UB only) → isolated URA, normal genitalia.
- **Mayer–Rokitansky–Küster–Hauser (MRKH)**: 46,XX, upper vaginal atresia to total müllerian agenesis (1 in 4500 females). Type I (typical): symmetrical buds, renal anomalies uncommon. **Type II (atypical, commoner): renal anomalies 40–60%** (URA, ectopia, horseshoe). MURCS = most severe end (C5–T1 vertebral dysplasia).
- **Work-up**: compensatory hypertrophy from 20 wk in ≈90%. US + colour Doppler; DMSA finds a missed pelvic or crossed ectopic kidney. Voiding cystourethrogram (VCUG) for 28% contralateral VUR is **not routine** — no reduction in febrile urinary tract infection (UTI); shared decision with parents.
- **Sport**: a solitary kidney does not preclude sport (individual assessment, padding).

Prognosis and surveillance of the SFK

- **Brenner hyperfiltration**: reduced nephron mass → glomerular hypertension → albuminuria → glomerulosclerosis. Risk highest when the SFK itself has congenital anomalies of the kidney and urinary tract (CAKUT).
- KIMONO study: ≈⅓ show renal injury (hypertension, albuminuria, medication) by ≈10 y; population data (Israeli cohort) ≈1% ESRD over 30 y, ≈4× controls. Verdict: lifelong chronic kidney disease (CKD) risk → lifelong follow-up.

Situation	blood pressure (BP) + first-morning urine albumin	Creatinine
SFK without CAKUT	Annually	Every 5 y if glomerular filtration rate (GFR) >60 and no medication
SFK with CAKUT	Twice yearly	As indicated
GFR <60 or on antihypertensive/antiproteinuric	BP, albumin, creatinine 2–4×/year	

Warning signs: hypertension, microalbuminuria (>30 mg/24 h). Treat with angiotensin-converting enzyme (ACE) inhibitor, salt limitation, avoid excess protein. Renal hypoplasia and dysplasia → *Ch 50*.

4 · SUPERNUMERARY KIDNEY

- True accessory organ: **own capsule, own collecting system and own (separate) blood supply** — vs duplex kidney (one parenchyma, one capsule, shared supply). Rare; left; ≈60% caudal to the dominant kidney.
- Second UB; ureters converge (≈50%) or are separate (≈50%, Weigert–Meyer usually obeyed).

- ≈½ are hydronephrotic; symptoms (pain, fever, mass) in early adulthood from obstruction/infection/stone. Magnetic resonance urography (MRU) or retrograde study for anatomy.

5 · ECTOPIA (ANOMALIES OF ASCENT)

Simple ectopia

- **1 in 900** (autopsy); left slightly more; bilateral ≈10%. Types: **pelvic (commonest)**, lumbar/iliac, abdominal, cephalad, thoracic. "Vascular barrier" theory rejected — aberrant vessels are a consequence, not cause.
- Kidney small, fetally lobulated, malrotated (pelvis anterior). **Hydronephrosis common**: ½ obstruction (PUJ ≈70% > VUJ), ¼ VUR ≥ grade III, ¼ malrotation alone. **VUR 30–50%** (often into the orthotopic kidney).
- Blood supply anomalous (distal aorta, iliac, inferior mesenteric) → **map by CT/MRI angiography before surgery**.
- Genital anomalies 15–45% (females 20–66%); ectopic kidney in ≈14% of cloacal malformations.
- Diagnosis: empty renal fossa; DMSA/MRU for small non-functioning kidney mistaken for URA. A girl with continuous incontinence → look for ectopic ureter of the ectopic kidney.
- Prognosis: normal except hydronephrosis, stones, blunt-trauma vulnerability and reduced differential function. **Never remove a "pelvic mass" without imaging** — solitary pelvic kidneys have been excised as tumours.

Cephalad and thoracic kidney

- Cephalad (near T10): classic with **omphalocele**.
- Thoracic: ≈1 in 13,000. Lies in the **foramen of Bochdalek** within a thin membrane (not in pleural space); rotation complete; usually asymptomatic. Chest X-ray mass → CT/MRU (avoid thoracotomy). Operate only for respiratory distress.

6 · FUSION ANOMALIES

Crossed renal ectopia

- Kidney lies opposite its ureteric orifice; **90% fused** — second commonest fusion anomaly. Left-to-right 3:1. Crossed ureter always crosses midline at the pelvic brim; orifice usually orthotopic.
- **McDonald–McClellan types**: with fusion (90%), without fusion, solitary crossed (imperforate anus ≈20%), bilateral crossed (rarest).

Crossed fused form	Key feature
Inferior ectopia (≈⅔, commonest)	Crossed upper pole fused to normal lower pole; both pelves anterior (early fusion)
Sigmoid (S-shaped)	Second commonest; pelves face opposite ways (late fusion)
Lump (cake)	Lobulated pelvic mass, ureters leave below
L-shaped (tandem)	Crossed kidney transverse, anterior to L4
Disc (pancake)	Medial fusion, reniform outline kept
Superior ectopia	Least common; crossed lower pole to normal upper pole

- Pelvis orientation times fusion: **anterior = early** (before rotation); medial = late. No rotation after fusion; midline structures stop ascent.
- VUR ≈20% (71% if bilateral crossed). Stone in the crossed ureter hurts on the **anephric side** (the ureter's embryonic side). Imaging: US, renography, MRU/MR angiography (MRA) before surgery.

Horseshoe kidney

- **Commonest fusion anomaly**: **1 in 400** (0.25%); male 2:1. **Lower-pole fusion in 95%**, before rotation.

- **Isthmus**: bulky parenchyma (own supply) or fibrous band at L3/L4, **just below the inferior mesenteric artery (IMA)**, which blocks ascent; usually anterior to aorta and inferior vena cava.
- Calyces normal in number but **point posteriorly**; lowest calyces point medially and may overlie the spine; pelvic axis vertical or outward; high lateral ureteric insertion; ureter bends over the isthmus.
- **Blood supply highly variable** (aorta above/below isthmus, IMA, iliac, sacral) → image (MRA/CT) before surgery.
- **Associations** (≈½ of children): **Turner syndrome — ≈60% have horseshoe** (commonest syndrome), caudal regression, neural tube defects (3%), anorectal malformations/VATER association (vertebral, anal, tracheo-oesophageal, renal/radial anomalies); hypospadias/undescended testis ≈4%.
- **PUJ obstruction 13–34%**: (1) high ureteric insertion, (2) ureter anterior to isthmus, (3) crossing vessels. Many hydronephroses are non-obstructed on renography. VUR 8–32% (managed as usual).
- **Stones ≈⅓ of adults**: stasis *plus* usual metabolic risk (hypercalciuria, hypocitraturia) → still do metabolic work-up.
- **Tumours**: **Wilms 1.76–7.9×** expected (favourable histology; preoperative chemotherapy preserved ≈75% of parenchyma — National Wilms Tumor Study); renal pelvic tumours increased (infection, stones, obstruction); renal cell carcinoma (RCC) not increased. A surprising share arise in the isthmus.
- Most stay asymptomatic.

> **KEY POINT** † PUJ obstruction in a horseshoe kidney: dismembered pyeloplasty with a dependent, anterior anastomosis is standard; division of the isthmus (symphysiotomy) is not routinely needed.

7 · MALROTATION AND VASCULAR ANOMALIES

- Malrotation: ≈1 in 939; male 2:1; frequent in Turner. **Weyrauch positions**: **ventral (no rotation, commonest)**; ventromedial (incomplete); dorsal (180°, rarest; vessels pass behind); lateral (hyper- or reverse rotation). **Vessel course is the only clue** to direction of rotation.
- Hilar fibrosis distorts the PUJ → intermittent obstruction, stones; function unaffected. Bilateral malrotation mimics horseshoe (look for the isthmus).
- **Vessel terminology**: *multiple* = >1 aortic vessel (preferred term); *aberrant* = from vessels other than aorta/main renal; *accessory* = ≥2 branches to one segment. Single artery in 71–85%. Segmental arteries are **end-arteries**.
- Crossing vessels occur in ≈¼ of normal unobstructed PUJs → a vessel alone does not prove obstruction.
- **Renal artery aneurysm** (≈0.1%): saccular ≈93%; ring ("wreath") calcification ≈30%; mostly silent; hypertension renin-mediated. Rupture only >3 cm. **Treat** (endovascular/open): rupture, rapid growth, pregnancy (or planned), **>2 cm**, symptoms. Otherwise observe.
- **Arteriovenous fistula**: congenital (cirsoid, <25%) vs acquired (biopsy, trauma — may close spontaneously). Congenital: bruit, haematuria, hypertension, high-output failure → embolisation.

8 · COLLECTING SYSTEM ANOMALIES

- **Bifid pelvis**: ≈10% of normal pelves; normal variant.
- † **Extrarenal pelvis**: a normal variant in which the pelvis lies outside the renal sinus; it looks dilated on US but drains normally — confirm with diuretic renography before calling it obstruction.
- **Calyceal diverticulum**: transitional-lined cavity joined to a calyx (type I, upper pole, small) or pelvis (type II, larger, symptomatic) via a narrow neck; ≈4.5/1000. **Stones in ≈40%**; milk of calcium layers on US. Diagnosis: **delayed CT/MRU contrast pooling**. Asymptomatic → observe; pain, infection, stones → ureteroscopic neck dilatation/ablation, **percutaneous ablation + fulguration** (best long-term), or laparoscopic marsupialisation.

- **Hydrocalycosis**: dilated major calyx with demonstrable connection; infundibular obstruction (vessel, stenosis, achalasia). Treat for symptoms.
- **Megacalycosis**: **non-obstructive** calyceal enlargement from malformed (falciform) papillae; calyces increased (12–20); pelvis and PUJ normal; **normal diuretic renogram**. Male 6:1, bilateral in males. Non-progressive → **no surgery**. May coexist with segmental distal megaureter (pitfall: mimics obstruction).
- **Infundibulopelvic stenosis**: usually bilateral, with VUR; spectrum toward cystic dysplasia. Function normal early, but **>⅓ develop renal insufficiency** (bilateral cases; hyperfiltration). Manage: yearly creatinine/estimated GFR, ACE inhibitor; surgery (endoscopic, calicocalicostomy, ureterocalicostomy) only if obstruction harms function.

> **KEY POINT** Collecting-system anomalies split on one question — is there obstruction? Dilatation without obstruction (bifid, extrarenal pelvis, megacalycosis) needs nothing; stasis (diverticulum, hydrocalycosis) is treated for symptoms; infundibulopelvic stenosis threatens function through bilateral dysplasia.

9 · LANDMARK TRIALS

Trial	Design	Result	Take-home
RAFT (Renal Anhydramnios Fetal Therapy)	Non-randomised multicentre; isolated BRA, serial amnioinfusion before 26 wk	94% live births; 82% survived ≥14 d with dialysis access; only 35% discharged on peritoneal dialysis; stopped early	Amnioinfusion prevents lethal lung hypoplasia but leaves neonatal ESRD — experimental, research setting only

10 · MCQ ONE-LINERS

1. Flattened adrenal lying along the spine on US ("lying down" sign) = absent ipsilateral kidney.
2. Potter facies are deformations from oligohydramnios, not primary malformations.
3. URA incidence 1 in 1200; ipsilateral ureter absent in ≈60%.
4. Commonest associated urological anomaly in URA: contralateral VUR (≈30%).
5. 79% of men with absent vas have ipsilateral renal agenesis.
6. MRKH type II carries renal anomalies in 40–60%.
7. Maternal diabetes triples the risk of renal agenesis.
8. Supernumerary kidney requires its own capsule and separate blood supply.
9. Thoracic kidney lies in the foramen of Bochdalek, outside the pleural space.
0. Cephalad ectopia is classically seen with omphalocele.
1. 90% of crossed ectopic kidneys are fused; left-to-right is 3× commoner.
2. Commonest crossed fused form: inferior ectopia.
13. Anterior-facing pelvis in fused kidneys indicates fusion before rotation.
14. Horseshoe kidney: 1 in 400; isthmus lies below the IMA at L3/L4.
15. About 60% of Turner syndrome patients have a horseshoe kidney.
16. Horseshoe kidney raises Wilms tumour risk, not RCC risk.
17. PUJ obstruction in horseshoe kidney: 13–34%.
18. Commonest malrotation: ventral (no rotation).
19. Direction of malrotation is told by the course of the renal vessels.
20. Treat renal artery aneurysm >2 cm, enlarging, symptomatic or with pregnancy.
21. Congenital renal AVF may cause high-output cardiac failure.
22. Bifid pelvis occurs in ≈10% of normal kidneys.
23. Calyceal diverticulum: stones in ≈40%; delayed contrast pooling confirms.
24. Megacalycosis: increased calyces, normal pelvis, normal diuretic renogram — no surgery.
25. Infundibulopelvic stenosis is usually bilateral and can progress to ESRD.

50 Renal Dysgenesis and Cystic Disease of the Kidney

Reading time ≈ 31 min (slow pace)

1 · GENETICS IN ONE TABLE

Gene	Renal phenotype
PAX2	Renal hypodysplasia or aplasia
EYA1	Branchio-oto-renal (BOR) syndrome
HNF1B	Renal cysts and diabetes syndrome
PKD1, PKD2 (primary cilium)	Polycystic kidney disease

2 · DYSGENESIS: DYSPLASIA VS HYPOPLASIA

- **Dysgenesis** = maldevelopment of size, shape or structure; subtypes dysplasia and hypoplasia. Agenesis → *Ch 49*.
- **Dysplasia** is a **histological** diagnosis: arrested metanephric differentiation from a wolffian duct–ureteric bud (UB)–blastema error. Hallmark (sine qua non): **primitive ducts** — tubules ringed by fibromuscular collars; also cartilage, immature glomeruli. Cysts optional: not all cysts are dysplastic, not all dysplasia is cystic.
- Cause: primary UB failure (genetic) or fetal outflow obstruction. **Dysplasia is the leading cause of childhood end-stage renal disease (ESRD).** Ultrasound (US): small, echogenic (brighter than liver) kidney, lost corticomedullary differentiation ± cysts.
- **Hypoplasia** = reduced renal mass from too few cells; simple hypoplasia has normal nephron number.

Entity	Defining feature	Course / management
Oligomeganephronia	Few, hypertrophied nephrons; usually bilateral	Polyuria, short stature; **proteinuria is first sign**; ESRD in second decade. Fluids, correct salt loss/acidosis, protein 1.5 g/kg/d, angiotensin-converting enzyme (ACE) inhibitor; living related donor acceptable (not familial)
Ask-Upmark (segmental hypoplasia)	Mostly acquired reflux nephropathy; deep midzone "slit-scars", thyroid-like tubules	Girls ≥10 y with **severe hypertension**; partial/total nephrectomy cures blood pressure (BP); reflux correction does not
Hypodysplasia	With ectopic orifice, megaureter, ureterocele, posterior urethral valves (PUV), prune-belly	**Degree of dysplasia ∝ degree of ureteric ectopia**

> **KEY POINT** The ureteric orifice predicts the kidney: a bud from the wrong place on the wolffian duct meets the wrong blastema, so ectopia, obstruction and dysplasia travel together.

3 · CYSTIC DISEASE: CONCEPTS AND CLASSIFICATION

- Cysts arise from tubules by three processes: **epithelial proliferation** (after a "second hit" on the normal allele), **fluid secretion** (net NaCl secretion when Na^+ absorption falls), and **abnormal extracellular matrix**. The **primary cilium** is central.
- **Multicystic** (multicystic dysplastic kidney, MCDK) = dysplastic, no normal nephrons; **polycystic** = normally developed kidney with nephrons throughout, cysts progressively replace them.

Inheritable	Non-heritable
Autosomal recessive polycystic kidney disease (ARPKD)	Multicystic dysplastic kidney (MCDK)

Inheritable	Non-heritable
Autosomal dominant polycystic kidney disease (ADPKD)	Benign multilocular cyst (cystic nephroma)
Juvenile nephronophthisis (recessive) / medullary cystic disease (dominant)	Simple cysts
Congenital nephrosis; familial hypoplastic glomerulocystic disease	Medullary sponge kidney; sporadic glomerulocystic disease
Syndromes: tuberous sclerosis, von Hippel–Lindau (VHL)	Acquired renal cystic disease; calyceal diverticulum

Extrarenal clue decides: hepatic fibrosis → ARPKD; berry aneurysm → ADPKD; retinitis pigmentosa → nephronophthisis; angiofibroma → tuberous sclerosis; haemangioblastoma → VHL. Chromosome 16 carries both PKD1 and TSC2.

4 · ARPKD

- **PKHD1, chromosome 6** († protein fibrocystin/polyductin); collecting-duct ectasia; **all have congenital hepatic fibrosis**. 1 in 10,000–50,000. "Infantile" is a misnomer — the earlier it presents, the more severe.
- **Fetus/neonate**: huge, smooth, reniform, non-transilluminating flank masses; oligohydramnios → Potter phenotype, pulmonary hypoplasia. **30–50% die** neonatally (respiratory failure, uraemia); ≈½ of neonatal survivors alive at 10 y.
- **US**: bilaterally enlarged, **homogeneously hyperechoic** kidneys (innumerable microcysts creating interfaces); pyramids merge with cortex. Discrete cysts appear with age.
- Older children: hypertension, chronic kidney disease (growth failure, anaemia, osteodystrophy) and liver — **portal hypertension, varices, splenomegaly with normal liver enzymes**. No neoplasm risk.
- Differential: ADPKD in a neonate (can be identical → **scan the parents**), bilateral hydronephrosis, mesoblastic nephroma, Wilms, renal vein thrombosis.
- **Management**: supportive — ventilation, antihypertensives, portosystemic shunt/sclerotherapy, dialysis and transplantation.
- **Nephrectomy** (respiratory/feeding compromise, BP, space for peritoneal dialysis) — limited evidence. ARegPKD registry: **very early bilateral nephrectomy (<3 months)** and hypotensive episodes independently predicted severe neurological injury. Prefer unilateral; buy time.

5 · ADPKD — PAEDIATRIC FOCUS

- **Commonest inheritable cystic disease** (1 in 400–1000); 50% of offspring; ≈10% de novo; 7–15% of haemodialysis patients. Usually presents 30–50 y, but in utero/infantile onset is aggressive.

	PKD1	PKD2
Locus / protein	16p / polycystin-1	4q / polycystin-2
Share	≈85%	≈15%
Cysts / ESRD	Most by 10 y, nearly all by 20; ESRD 50s	Later; ESRD 70s

- **Contiguous gene syndrome**: deletion of PKD1 + adjacent TSC2 → tuberous sclerosis with severe early polycystic kidneys.
- In children ≈17% look unilateral/asymmetric; the other kidney follows.
- **Clinical**: pain (commonest adult symptom — haemorrhage, infection, stone), haematuria ≈50%, stones 20–30%, urinary tract infection (UTI). **Hypertension** (renin-mediated by cyst stretch) in ≈50% at 20–35 y with normal function — the key paediatric finding.
- **Extrarenal**: liver cysts (commonest; rare in children, ≈all by 50, worse in females), **intracranial berry aneurysm 10–30%**, seminal vesicle cysts 40%, mitral valve prolapse, colonic diverticula.
- Renal cell carcinoma (RCC) **not increased** (but when present: younger, bilateral, sarcomatoid).

Diagnosis

Age (at 50% risk)	Ravine US criterion
<30 y	≥2 cysts, uni- or bilateral
30–59 y	≥2 cysts in each kidney
≥60 y	≥4 cysts in each kidney

- Negative US in a young person does **not** exclude PKD2 (sensitivity ≈⅔ <30 y) → genetic testing when certainty matters (e.g., living donor). Counsel before testing (insurance, psychological harms).
- CT: haemorrhagic cyst 50–90 Hounsfield units (HU); MRI: bright T1/dark T2.

Management

- **BP**: ACE inhibitor/angiotensin receptor blocker (ARB) first line; target <130/80. HALT-PKD: tight BP slowed kidney growth.
- **Tolvaptan** (vasopressin V2 antagonist; ↓ cyclic AMP) slows total kidney volume (TKV) growth and glomerular filtration rate (GFR) decline (TEMPO, REPRISE); first approved adult therapy (2018); costs — aquaresis, hepatotoxicity, discontinuation.
- **Urological issues**: pain → exclude infection, stone, tumour; avoid non-steroidal anti-inflammatory drugs; aspiration or **laparoscopic unroofing** (relieves pain only, no effect on renal decline). Infected cyst unresponsive to antibiotics → **lipophilic antibiotics** (trimethoprim–sulfamethoxazole, fluoroquinolones) → drainage. Stones: alkalinisation, shock wave lithotripsy.
- Aneurysm screening (MR angiography) only for family history of aneurysm/subarachnoid haemorrhage, prior rupture, high-risk job, major surgery, anxiety. Coil/clip >7–10 mm or symptomatic. Adult management → *kidney book*.

> **KEY POINT** ADPKD is a volume disease: cyst growth stretches vessels (hypertension), compresses nephrons and hides stones and tumours. Every therapy is judged by whether it slows total kidney volume.

6 · JUVENILE NEPHRONOPHTHISIS (NPH) / MEDULLARY CYSTIC KIDNEY DISEASE (MCKD)

- Ciliopathies; shared triad: **tubular basement membrane thickening/disintegration, tubular atrophy with corticomedullary cysts, interstitial fibrosis**. Small-to-normal kidneys.
- Shared clinical: **polyuria/polydipsia >80%** (vasopressin-resistant, **salt wasting**); ESRD 5–10 y after presentation.

	Juvenile NPH	MCKD
Inheritance	Autosomal recessive	Autosomal dominant
ESRD	Mean 13 y; **10–20% of childhood renal failure**	Third–fourth decade
Hypertension	Absent	Present
Extrarenal	20%: **retinitis pigmentosa (Senior–Løken)**, Joubert, Cogan apraxia, hepatic fibrosis, Bardet–Biedl	None (hyperuricaemia/gout)

- US often misses cysts (echogenic small kidneys); CT/MRI better; absence of cysts does not exclude.
- Treatment: **sodium replacement; avoid salt restriction and diuretics**; dialysis/transplant — does not recur in grafts.

7 · SYNDROMES WITH RENAL CYSTS

Tuberous sclerosis complex (TSC)

- Autosomal dominant; **TSC1 (chr 9), TSC2 (chr 16)**. Triad: epilepsy ≈80%, intellectual disability ≈60%, adenoma sebaceum ≈75%; earliest skin sign ash-leaf macule (Wood lamp). Diagnosis: 2

major or 1 major + 2 minor features (angiomyolipoma, AML = major; multiple renal cysts = minor).

- Kidney: cysts (even in utero; mimic ADPKD), **AML 40–80%** (bilateral, multiple, after 10 y; never metastasise; risk = haemorrhage), RCC ≈2% (younger, female, bilateral — suspect in a growing fat-free or calcified lesion).
- Imaging: AML echogenic on US, fat (≈−50 HU) on CT; **MRI preferred** (25–30% fat-poor AMLs missed on US).
- **AML >3 cm and growing → mammalian target of rapamycin (mTOR) inhibitor (everolimus) first line**; embolisation then corticosteroids for acute bleed; kidney-sparing surgery second line; avoid nephrectomy. Imaging every 1–3 y; BP and GFR yearly; ACE inhibitor/ARB for hypertension.

Von Hippel–Lindau disease

- Autosomal dominant, **VHL gene 3p25**; pVHL degrades hypoxia-inducible factor (HIF) → loss upregulates angiogenic factors (vascular endothelial growth factor foremost).
- Renal cysts 76% (earliest), **clear cell RCC ≈50%**, phaeochromocytoma 10–17%, cerebellar/retinal haemangioblastoma, pancreatic cysts, epididymal cystadenoma 10–26%. Metastatic RCC and haemangioblastoma are leading causes of death.
- Surveillance (VHL Alliance 2020): exam yearly from 1 y, BP from 2, eye from <1, metanephrines from 5, brain/spine MRI 2-yearly from 11, **abdominal MRI 2-yearly from 15**.
- Goal: cancer control while preserving parenchyma — **nephron-sparing surgery** (local recurrence 75–85% but good survival); partial adrenalectomy for phaeochromocytoma.

8 · MULTICYSTIC DYSPLASTIC KIDNEY (MCDK)

- Multiple non-communicating cysts, **no normal parenchyma, no function**, atretic ureter ("bunch of grapes"). Types: infundibulopelvic (commoner, pelvis + ureter atretic) and hydronephrotic.
- **Commonest renal cystic disease of the newborn; second commonest neonatal abdominal mass**; 1 in 1000–4000; male 55–70%. Bilateral = lethal. UB-gene theory explains contralateral anomalies.
- **Contralateral kidney**: compensatory hypertrophy; **vesicoureteric reflux (VUR) 18–43%, pelviureteric junction (PUJ) obstruction 3–12%**.
- **Natural history**: ≈40% involute (later imaged as "agenesis"); hypertension and malignancy (Wilms) not increased in large series.

US feature	MCDK	Severe PUJ obstruction
Cysts	Haphazard, varied sizes, non-communicating	Calyces arranged around a central pelvis, communicating
Central large cyst	Absent	Present (pelvis)
Isotope uptake	None	Usually some

ALGORITHM
Prenatal MCDK → US day 2–3 and at 1 month (contralateral kidney, bladder) → technetium-99m dimercaptosuccinic acid (DMSA) or mercaptoacetyltriglycine (MAG3) scan if doubt → voiding cystourethrogram (VCUG) only if UTI or abnormal contralateral kidney/bladder → primary care: BP, mass, UTI

- **Nephrectomy** only for mass effect (respiration, feeding), hypertension, cyst rupture, or **increasing solid tissue**. Routine US surveillance has no proven benefit.

> **KEY POINT** MCDK removes itself; management looks away from it to the solitary contralateral kidney and its reflux or obstruction. Mistaking a non-functioning PUJ kidney for MCDK costs little.

9 · BENIGN MULTILOCULAR CYST (CYSTIC NEPHROMA)

- Spectrum (no lesion converts): benign multilocular cyst → cystic partially differentiated Wilms tumour (CPDN) → multilocular cyst with Wilms nodules → cystic Wilms tumour. Adult counterpart: multilocular cystic RCC.
- **Bimodal: boys <4 y (M 2:1, flank mass) and women >30 y (F 8:1, haematuria from herniation into pelvis)**.
- Imaging cannot exclude malignancy → **excision is diagnosis and treatment** (nephrectomy in children; partial if localised). Wilms nodules → treat per National Wilms Tumor Study staging (favourable outlook).

10 · SIMPLE CYSTS IN CHILDREN AND BOSNIAK

- **US criteria (all four)**: anechoic; thin smooth wall; posterior acoustic enhancement; round/oval. CT: −10 to +20 HU, no enhancement. Enhancement = vascular tissue = suspicion.
- **Children**: simple cyst <15 mm, asymptomatic → repeat US at 2 y, discharge if stable. Continue surveillance if ≥15 mm, complex, family history of cystic disease, or genitourinary anomalies. Many bilateral cysts without liver cysts — do not label ADPKD without family history/genetics.

Bosniak	Features	Management
I	Simple	None
II	Fine septa, minimal calcification, hyperdense ≤3 cm	None
IIF	More septa/calcification, no measurable enhancement	Follow-up (≈11% progress)
III	Thick/irregular enhancing walls or septa	Excision (paediatric III–IV: >90% intermediate/malignant)
IV	Enhancing solid components	Excision

Symptomatic cyst: aspiration ± sclerosant, or laparoscopic unroofing; infected → drainage + antibiotics. Bosniak 2019 adds MRI and down-classifies to curb over-treatment.

11 · MEDULLARY SPONGE KIDNEY, GLOMERULOCYSTIC AND ACQUIRED CYSTIC DISEASE

- **Medullary sponge kidney**: ectatic medullary collecting ducts ("brush bristles", "bouquet of flowers" on urography); non-heritable; presents >20 y with colic 50–60%, UTI, haematuria. Hypercalciuria ⅓–½, incomplete distal renal tubular acidosis 30–40% (hypocitraturia). Prevention: **potassium citrate first**, thiazide, fluids, low sodium.
- **Sporadic glomerulocystic disease**: bilateral enlarged kidneys, Bowman-space cysts; diagnosis of exclusion (no family history).
- **Acquired renal cystic disease**: ≥3 cysts per kidney in ESRD; caused by **uraemia**, not dialysis; 10% pre-dialysis → >90% at 10 y; children on dialysis 20–45%. Small kidneys (vs large in ADPKD). Haemorrhage commonest; adenomas common; RCC risk ↑ (aggressive after transplantation, mostly in native kidneys) → screen long-term dialysis and transplant patients.

12 · LANDMARK TRIALS

Trial	Design	Result	Take-home
HALT-PKD	Randomised: lisinopril + telmisartan vs lisinopril; rigorous vs standard BP	Dual blockade no benefit; low BP slowed TKV growth, ↓ left ventricular mass and albuminuria, GFR unchanged	Single renin–angiotensin blocker; tight BP early

Trial	Design	Result	Take-home
TEMPO / REPRISE	Randomised, tolvaptan vs placebo	Slower TKV growth and GFR decline	First disease-modifying drug; aquaresis, hepatotoxicity
ARegPKD registry	Observational ARPKD cohort	Bilateral nephrectomy <3 months → worst neurological outcome	Avoid early bilateral nephrectomy

13 · MCQ ONE-LINERS

1. Sine qua non of renal dysplasia: primitive ducts with fibromuscular collars.
2. Leading cause of childhood ESRD: renal dysplasia.
3. Degree of hypodysplasia correlates with degree of ureteric orifice ectopia.
4. Oligomeganephronia: fewer, hypertrophied nephrons; proteinuria first sign.
5. Ask-Upmark kidney: segmental hypoplasia, severe hypertension, cured by nephrectomy.
6. ARPKD gene: PKHD1 on chromosome 6 (fibrocystin).
7. ARPKD is always accompanied by congenital hepatic fibrosis.
8. Neonatal ARPKD US: large homogeneously hyperechoic kidneys.
9. Neonatal ADPKD may mimic ARPKD — scan the parents.
0. PKD1 (chr 16) ≈85%; PKD2 (chr 4) milder, later ESRD.
1. PKD1–TSC2 contiguous deletion gives TSC with polycystic kidneys.
2. Commonest extrarenal ADPKD manifestation: hepatic cysts.
3. Berry aneurysm in ADPKD: 10–30%.
4. Tolvaptan is a vasopressin V2 antagonist lowering renal cAMP.
15. Infected ADPKD cyst: use lipophilic antibiotics.
16. NPH: salt wasting, no hypertension, ESRD around 13 years.
17. Senior–Løken syndrome: nephronophthisis plus retinitis pigmentosa.
18. TSC AML >3 cm and growing: everolimus first line.
19. Fat-poor AML is missed on US in 25–30%; MRI preferred.
20. VHL gene: chromosome 3p25; RCC is clear cell type.
21. Commonest renal cystic disease in newborns: MCDK.
22. Contralateral VUR in MCDK: 18–43%.
23. About 40% of MCDKs involute spontaneously.
24. MCDK nephrectomy indication: increasing solid tissue or mass effect.
25. Multilocular cyst: boys <4 y and women >30 y.
26. Bosniak IIF: follow-up; III and IV: excision.
27. Paediatric simple cyst <15 mm: repeat US at 2 years, then discharge.
28. Medullary sponge kidney stone prevention: potassium citrate first.
29. Acquired cystic disease is caused by uraemia, not dialysis.

51 Pathophysiology of Urinary Tract Obstruction

Reading time ≈ 18 min (slow pace)

1 · DEFINITION AND CLASSIFICATION

- **Obstruction**: any anatomic or functional abnormality impeding urine flow, anywhere from calyces to meatus, that injures the kidney or lower tract. Classify: **acute/chronic · partial/complete · intrinsic/extrinsic · unilateral/bilateral**.
- Defined by effect on **flow and function, not dilation**: a dilated system may be unobstructed and resolve alone, so the partial vs high-grade distinction, not the image, decides treatment.
- Incidence ≈ **1 in 500** live births. High-grade obstruction: lifetime end-stage renal disease (ESRD) risk up to **30%**.

	Lower urinary tract obstruction (LUTO)	Upper tract obstruction
Commonest cause	Posterior urethral valves (PUV) and urethral atresia ≈ 50%	Ureteropelvic junction obstruction (UPJO), 1 in 1500; M:F 2:1
Others	Anterior urethral valves, megalourethra, prolapsing ureterocele, cloaca, megacystis–microcolon–intestinal hypoperistalsis syndrome, trisomy 18/21	Ureterovesical junction obstruction (UVJO), ectopic ureter, ureterocele, fibroepithelial polyp, stones, crossing vessel (extrinsic)
Outcome driver	ESRD + pulmonary hypoplasia (bilateral, whole-fetus disease)	Progressive unilateral function loss; normal contralateral kidney buffers

Presentation depends on site, severity, age at onset and duration: prenatal hydronephrosis on anomaly scan (most at 18–20 weeks); postnatally mass, fever/recurrent infection, flank pain, polyuria or retention, failure to thrive, electrolyte imbalance, or respiratory failure from pulmonary hypoplasia.

> **KEY POINT** Trap: **detrusor hypertrophy** is a lower-tract cause of upper-tract (UVJ) obstruction — the thickened wall narrows the intramural ureter. Neurogenic bladder obstructs without an anatomical lesion.

2 · HAEMODYNAMIC CHANGES

Urine production continues against the block → proximal pressure rises → net filtration pressure falls → glomerular filtration rate (GFR) and renal blood flow (RBF) fall → ischaemia. Ischaemic injury (inflammation, tubular atrophy, interstitial fibrosis) begins within **2 h** of complete obstruction. Bilateral obstruction or a solitary kidney → more profound disease.

† Phase (complete obstruction)	Unilateral	Bilateral / solitary kidney
I (0–1.5 h)	RBF ↑, ureteric pressure ↑ (afferent vasodilatation: prostaglandin E2, nitric oxide)	Modest RBF ↑
II (1.5–5 h)	RBF ↓, ureteric pressure still ↑ (efferent constriction)	RBF ↓, pressure ↑
III (> 5 h)	RBF ↓, ureteric pressure ↓ (afferent constriction: angiotensin II, thromboxane)	Pressure stays high (retained solute, atrial natriuretic peptide) — underlies post-obstructive diuresis

- **Hypertension**: ↑ intrarenal pressure → ↓ perfusion → renin–angiotensin–aldosterone system activation → vasoconstriction + Na retention. Bilateral ≈ **77%** hypertensive; unilateral ≈ **20%** (renin lateralises to affected kidney).

3 · TUBULAR FUNCTION

Obstruction **downregulates aquaporins (AQP1–4)** and Na transporters along the nephron → concentrating defect (hyposthenuria), salt wasting, polyuria; impaired H^+ transport → **distal renal tubular acidosis**.

Early obstruction — wasting	Prolonged obstruction — retention
Na, K, HCO_3, Mg, Ca, phosphate	K, H^+, Cl, ammonium

- K is **wasted early, retained late**.
- Fractional excretion of sodium (FENa): low (< 1%) in acute obstruction (filtration falls, tubular avidity kept); high (> 1%) with established tubular injury (salt wasting).

4 · PATHOLOGY AND FIBROSIS MEDIATORS

Immature kidney	Mature kidney
Apoptosis (pressure, ischaemia); impaired nephrogenesis and abnormal differentiation — nephron number permanently capped	Compensatory tubular proliferation → if persistent, collagen deposition → scarring

Renal fibrosis (hallmark ↑ α-smooth muscle actin and interstitial extracellular matrix [ECM]) is the final common path to ESRD.

Pathway	Mechanism
Mechanotransduction	Stretch-sensitive receptors convert stretch into fibrotic signals (initiator)
Inflammation	Tumour necrosis factor-α (TNF-α), interleukins → immune cell recruitment, ECM deposition
Transforming growth factor-β (TGF-β)	TGF-β superfamily → **Smad** signalling → fibrogenesis; **angiotensin II** contributes (amplifier)
MicroRNAs	miR-34a ↑ → epithelial-to-mesenchymal transition; possible therapeutic target

Bladder (outlet obstruction): hypertrophy + hyperplasia → compensation → decompensation. Interleukin-6, TNF-α, monocyte chemoattractant protein-1 → collagen/fibronectin → poor compliance. Storage pressure > **40 cm H_2O** transmits injury to kidneys.

5 · RENAL RECOVERY AND COMPENSATORY GROWTH

- Prolonged obstruction → scarring and atony → renal failure over days–months. Relief can reverse damage, but **incomplete recovery is common**.
- Acute obstruction on chronic kidney disease (CKD) → rapid acute kidney injury; obstruction erodes **renal reserve**.
- **Compensatory growth** of the contralateral/remaining nephrons: insulin-like growth factor-1, TGF-β1, hepatocyte growth factor → glomerular enlargement and tubular hypertrophy (**glomerulomegaly, not new nephrons**).

6 · POST-OBSTRUCTIVE DIURESIS (POD)

- Polyuric phase after relief; unilateral ≈ **2%**, bilateral ≈ **50%** (no normal kidney to regulate volume).
- **Mechanism**: aquaporin downregulation, loss of medullary tonicity, tubular transporter dysfunction (apoptosis); physiological excretion of retained salt and water.
- **Risk after valve ablation**: oligohydramnios, preterm birth (low AQP2), raised creatinine.

ALGORITHM
Monitor intake/output, serum and urine electrolytes → replace losses, correct electrolytes → **do not over-replace** (perpetuates POD) → usually resolves in **24 h**, seldom > 48 h

7 · FETUS AND CHILD

- Fetal kidneys visible from 9 weeks; bladder by 12 weeks (not seen by 15 weeks = abnormal); visible fetal ureter = dilated.
- Urinary tract dilation classification; anteroposterior diameter (APD) most reliable: obstruction considered at **≥ 7 mm** (16–27 weeks), **≥ 10 mm** (≥ 28 weeks).
- LUTO: raised pressure impairs glomerular development; anhydramnios → pulmonary hypoplasia. Second-trimester oligo/anhydramnios → mortality **60–80%**.
- **Fetal urine** (vesicocentesis, three serial samples; last reflects current function) favourable if: Na < 100 mEq/L, Cl < 90, Ca < 8 mg/dL, osmolality < 200 mOsm/L, β2-microglobulin < 6 mg/L — the healthy tubule makes hypotonic urine.
- Fetal therapy: vesicoamniotic shunt/fetal cystoscopy (LUTO, preserved function) treats obstruction; amnioinfusion only replaces fluid.
- Postnatal: ultrasound (US) first; mercaptoacetyltriglycine (MAG3) for drainage; dimercaptosuccinic acid (DMSA) for scar; voiding cystourethrogram (VCUG) for outlet and vesicoureteral reflux (VUR). Whitaker test: obstructed if pelvic pressure > **12 cm H_2O** or continuous rise.
- Cystatin C–based GFR is the most accurate paediatric estimate.

8 · TREATMENT PRINCIPLES

Logic is **temporise, then define**: drainage buys renal function and stability; definitive reconstruction follows; complications of relief (POD, electrolytes) are managed as carefully as the obstruction.

Option	Indication / key point
Clean intermittent catheterisation (CIC)	Neurogenic bladder; ↓ urinary tract infection (UTI), stones, retention
Continuous overnight catheter drainage	Valve bladder with nocturnal polyuria and progressive CKD; delays progression
Vesicostomy	Infant with complex LUTO, unstable, or unable to do CIC; temporising
Percutaneous nephrostomy	Upper tract obstruction; needs some hydronephrosis
Ureteral stent / cutaneous ureterostomy	Temporising or with definitive repair; stent excellent for extrinsic obstruction
Pyeloplasty / reimplantation	Correct all intrinsic and extrinsic causes; aberrant vessel → anastomose anterior to it; non-refluxing reimplant in children

- Pain: non-steroidal anti-inflammatory drugs only if renal function is preserved.
- Oliguria/acute kidney injury: treat cause, optimise fluids and perfusion. CKD: blood pressure, diet; ESRD → dialysis or transplant → *Ch 61*.

9 · MCQ ONE-LINERS

1. Commonest cause of LUTO: posterior urethral valves.
2. Commonest upper tract obstruction: UPJO; male:female 2:1.
3. Ischaemic injury starts within 2 h of complete obstruction.
4. Unilateral obstruction late phase: RBF ↓ and ureteric pressure ↓.
5. Bilateral obstruction: ureteric pressure stays high.
6. Characteristic acidification defect: distal renal tubular acidosis.
7. Potassium: wasted early, retained in prolonged obstruction.
8. Aquaporins downregulated in obstruction: AQP1–4.
9. Key fibrotic signalling: TGF-β → Smad.
10. MicroRNA linked to obstructive fibrosis: miR-34a.
11. Compensatory growth = glomerulomegaly, not nephrogenesis.

2. POD: unilateral ≈ 2%, bilateral ≈ 50%.
3. POD usually resolves within 24 h; over-replacement perpetuates it.
4. Preterm infants: low AQP2 → POD risk after valve ablation.
5. Hypertension: bilateral ≈ 77%, unilateral ≈ 20%.
6. Dangerous bladder storage pressure: > 40 cm H_2O.
7. Prenatal APD threshold: ≥ 7 mm (16–27 wk), ≥ 10 mm (≥ 28 wk).
18. Favourable fetal urine Na < 100 mEq/L; β2-microglobulin < 6 mg/L.
19. Fetal bladder not seen by 15 weeks: abnormal.
20. Whitaker test positive: pelvic pressure > 12 cm H_2O.
21. Megacystis: fetal bladder failing to empty over 40 min at ≥ 18 weeks.
22. FENa < 1% in acute obstruction; > 1% with tubular injury.

52 Ectopic Ureter, Ureterocele and Ureteral Anomalies

Reading time ≈ 37 min (slow pace)

1 · DEFINITIONS AND CLASSIFICATION

- Ectopic ureter and ureterocele are one developmental defect of **ureterotrigonal connection** expressed along a continuum; appearance changes the technique, not the strategy.
- **Ectopic ureter**: any ureter, single or duplex, that does not enter the trigone. Ectopia requires entry **at or distal to the bladder neck**; a ureter merely near the bladder neck is not ectopic.
- **Ureterocele**: ectopic-ureter variant with **cystic dilation of the distal ureter**, lying within the bladder or spanning bladder neck and urethra. Never forms entirely within the urethra; never attaches to wolffian structures.
- In a **duplex system, both lesions always involve the upper pole** ureter.

Ureterocele classification

System	Basis	Categories	Clinical value
Glassberg (most useful)	Position vs bladder neck	**Intravesical** (entirely above bladder neck) vs **ectopic/extravesical** (part permanently at bladder neck or urethra)	Predicts outcome of endoscopic incision
Stephens	Morphology	Stenotic, sphincteric, sphincterostenotic, **cecoureterocele**, blind, non-obstructed	Identifies the surgically dangerous cecoureterocele
Churchill	Functional impact	Upper pole only → whole ipsilateral kidney → contralateral too (reflux or bladder outlet obstruction (BOO))	Extent of threat

- **Cecoureterocele**: orifice inside the bladder (large, incompetent), but the cavity tunnels submucosally beyond the bladder neck into the urethra. Often missed preoperatively; troublesome at incision; risk factor for incontinence.
- Distinguish ectopic ureterocele from an intravesical ureterocele that **prolapses** into the urethra only during voiding.
- **Ureterocele disproportion**: small dysplastic, non-dilated upper pole invisible on imaging, with a typical ureterocele and apparently normal kidney.

2 · EMBRYOLOGY ESSENTIALS

- Ureteric bud arises from the mesonephric (wolffian) duct. In a duplex, the upper pole bud arises **later and more cephalad**, so it is incorporated into the urogenital sinus later and its orifice is carried **caudally** (ectopic).
- **Weigert–Meyer rule**: inverse relationship of duplex orifices — the **upper pole orifice is caudal (and medial) to the lower pole orifice**. Upper pole → ectopia/ureterocele (obstruction); lower pole → short tunnel → reflux.
- **Chwalle membrane**: failure of rupture of a membrane at the distal ureteric orifice → ureterocele (classic theory).
- Renal dysplasia is **downstream** of the connection defect, not a parallel event.
- Bud derives from the wolffian duct, so an ectopic ureter never inserts directly into müllerian structures; it reaches the vagina/uterus via the **Gartner duct** (wolffian remnant alongside the vagina).

3 · ECTOPIC URETER — SITES AND SEX ASYMMETRY

	Female	Male
Sites	Bladder neck → perineum; vagina, uterus, rectum (via Gartner duct)	Always **above the external sphincter**: bladder neck, prostatic urethra, vas, seminal vesicle, ejaculatory duct
Continence	**Continuous incontinence** (classic)	**Never incontinent**
Presentation	Continuous dribbling, perineal (purulent) discharge, urinary tract infection (UTI)	Epididymitis/orchitis, UTI, hydronephrosis, flank pain
Association	Gartner duct cyst	**Zinner syndrome**: ipsilateral renal agenesis + seminal vesicle cyst

> **KEY POINT** Why does only the girl leak? Everything follows from where the ureter opens relative to the external sphincter. The female ureter may open distal to it and leaks continuously; the male ureter, confined to wolffian derivatives above it, cannot leak and presents as infection or obstruction.

- **Single-system ectopic ureter** may appear as an "absent kidney" (small, poorly functioning unit missed on ultrasound) → computed tomography (CT) or renal function study.
- **Bilateral single-system ectopic ureters**: hypoplastic bladder and bladder neck, bilateral dysplasia.

4 · CLINICAL PRESENTATION

Prenatal

- Most detected on prenatal ultrasound as hydronephrosis; fetal upper pole "cyst" = upper pole hydronephrosis.
- **Diagnosis is made at the bladder**: upper tract appearances are identical. A large ureterocele can fill and mask the bladder; await filling. Record upper pole parenchymal thickness and echogenicity.

Infection

- Urosepsis → ultrasound in all. Unresponsive sepsis → **urgent transurethral incision (TUI)** of ureterocele, or percutaneous/open drainage (end ureterostomy) of ectopic ureter.
- Ectopic ureter: low-grade fever with spikes. **Negative urine culture does not exclude infection** — the infected system may not drain into the bladder.
- Any young male with bacterial epididymitis → upper tract ultrasound.

Incontinence

- Toilet-trained girl with verified **continuous low-volume dampness day and night**, despite normal voiding → evaluate for ectopic ureter. Untreated ureteroceles do not cause incontinence.
- Discriminating question: **can she be dry for 30–60 minutes?** With ectopic ureter — no.
- The leaking moiety may be non-dilated and missed on imaging; suspicion is history-driven.

Ureterocele prolapse

- **Smooth, congested, mucosa-covered interlabial mass** from the urethra, distinct from the vagina, often with voiding difficulty.
- Lobulated, grape-like mass suggests **sarcoma botryoides** instead.

Late presentation

- Adult single-system non-obstructing ureterocele, frequently containing a **stone**.

5 · EVALUATION

Test	Answers	Note
Ultrasound	Anatomy; inferred function; bladder	First in every case; cannot separate ectopic ureter from ureterocele above the bladder
Voiding cystourethrogram (VCUG)	Reflux (lower pole, contralateral, into ureterocele); bladder base, urethra	Optional if obstruction alone dictates incision; repeat after incision
Dimercaptosuccinic acid (DMSA)	Differential function — gold standard	Only "some vs none"; no normal upper pole criteria
Mercaptoacetyltriglycine (MAG3) diuretic renogram	Function + drainage	For observation candidates; generally after 2 months
Magnetic resonance (MR) urography	Most detailed anatomy + function	Complex/undefined anatomy only; needs sedation
Intravenous pyelogram (IVP)	Anatomy + qualitative function	Non-dilated ectopic ureter with incontinence

- **Ureterocele on ultrasound**: thin-walled cyst within the bladder, not beyond its wall. **Pseudoureterocele**: dilated ectopic ureter indenting the bladder — **bilaminar, thicker wall, lumen extending outside the bladder**.
- VCUG: lower pole reflux shows the **"drooping lily"** (lower pole displaced down/out by dilated upper pole).
- † On excretory urography a single-system intravesical ureterocele gives the **"cobra head" (spring-onion)** appearance — contrast-filled dilated distal ureter with a radiolucent halo (ureterocele wall).

VCUG terms

Term	Meaning	Significance
Effacement	Ureterocele flattens with filling	Mimics ectopic ureter on full-bladder imaging
Prolapse	Ureterocele moves into urethra	Outlet obstruction; predicts secondary surgery
Eversion	Diverticulum-like protrusion of bladder base with filling/voiding	**Weak trigonal floor → likely needs open repair**

Endoscopy

- A **site-selection** exercise before incision: start with little filling and increase slowly — the lowest portion of the ureterocele (best incision site) is seen with limited filling.
- Look for urethral extension (cecoureterocele), other orifices, patulous bladder neck.
- Patulous bladder neck / cecoureterocele → risk of later incontinence; consider bladder neck tightening at definitive surgery (videourodynamics may help).

6 · MANAGEMENT PRINCIPLES

- Goals: **preserve renal function; eliminate infection, obstruction and reflux; maintain continence; minimise total procedural morbidity**.
- Early daily prophylactic antibiotic in significant hydronephrosis (agreed).
- Relieving obstruction may **create** reflux or allow existing lower pole reflux to resolve — the central trade-off.
- Function cannot be predicted before or after decompression; upper pole contributes little and recovery after early decompression is limited. Decisions are empirical and individual.

Neonatal algorithm

ALGORITHM

Prenatal hydronephrosis → postnatal ultrasound → antibiotic prophylaxis if significant hydroureteronephrosis → VCUG (optional if it will not change decompression) → renal scan after 2 months

Small ureterocele, minimal hydronephrosis, no obstruction → observe

Significant obstruction, parenchyma at risk → TUI (or cutaneous distal ureterostomy if endoscopy impossible)

Sepsis not responding → TUI (ureterocele) / nephrostomy or end ureterostomy (ectopic ureter); after stabilisation an elective drainage procedure is still needed

Observation (selected ureteroceles)

- Criteria: no obstruction of ipsilateral lower pole or contralateral kidney, lower pole reflux ≤ grade III; non-functioning upper pole (e.g. multicystic dysplastic upper segment); or no obstruction on diuretic renogram.

7 · OPERATIVE OPTIONS

Transurethral incision (TUI) of ureterocele

- Simplest decompression; relieves obstruction in **78–97%**. Procedure of choice in **sepsis** and in the **infant**.
- Technique: **low transverse full-thickness incision** with cutting current, **as distal and as close to the bladder floor as possible** to lessen reflux. Angled-tip wire preferred; incise deeply.
- Adequacy: jet of urine or view of urothelium inside. Outpatient; ultrasound and VCUG at **4–6 weeks**.
- Ectopic ureterocele: longitudinal incision from intravesical to urethral portion, or separate intravesical and urethral punctures (avoid leaving an obstructing urethral flap).
- **"Watering-can"** technique: multiple holmium laser punctures — equal decompression, less de novo reflux.
- New reflux after TUI **0–50%**. **Intravesical**: decompression without reflux in 70–80%. **Extravesical**: secondary surgery up to 70%. Reoperation relative risk 1.74 with lower pole reflux, 2.78 if extravesical.
- Persisting reflux into either pole → typically ureterocele excision and reimplantation.

> **KEY POINT** Why offer TUI first even if half need more surgery? It is an outpatient procedure that may be definitive; the second operation can be deferred to an older child; and decompressing the ureter often avoids tapering at later reimplantation. Less appealing in the older child with a massive non-functioning upper pole.

Upper pole partial nephrectomy (heminephrectomy)

- For a clearly non-functioning upper pole or doubtful drainage because of massive dilation. Open (flank or dorsal lumbotomy), laparoscopic or robotic — same steps.
- Steps: transect upper pole ureter, use as handle → pass it **behind the renal vessels** → ligate upper pole vessels (usually 2–3) → demarcation → resect upper pole → close defect over a fat pedicle (mattress sutures, 3-0 polydioxanone (PDS)/Vicryl).
- **Dissect on the wall of the upper pole ureter**; leave its back wall attached to the lower pole ureter near the bladder — the shared blood supply runs between the two ureters.
- Distal stump: ligate and leave if small; if obstructed and refluxing, excise as low as possible (second Gibson incision in open surgery; protect vas).
- Postoperative Doppler ultrasound: anatomy, urinoma, lower pole flow.
- Outcomes: ectopic ureter — usually curative (stump rarely problematic). Ureterocele — pre-existing lower pole reflux resolves in up to 20%; new reflux **15–50%**; secondary surgery **40–50%**.

- Laparoscopic/robotic: less pain, quicker discharge, avoids second incision.

Ureteroureterostomy / pyeloureterostomy

- Upper pole drained into lower pole system (end-to-side). For functioning upper poles and, increasingly, non-functioning ones (hypertension concern unsubstantiated); small incision.
- **Distal** (inguinal): **stent the lower pole ureter cystoscopically first** to identify the recipient. **Proximal** preferred for dilated upper pole/size mismatch (less stasis, "yo-yo" concern).
- Minimal medial dissection; generous recipient ureterotomy; decompress and resect distal upper pole ureter staying on its wall. Stent the recipient only, never across the anastomosis.

Lower tract reconstruction

- **Ureterocele excision + common sheath reimplantation**: relieves obstruction and corrects reflux in one operation. Risks: bladder neck, sphincter, vagina.
- Steps: incise ureterocele → dissect it off the bladder to its junction with the lower pole ureter → mobilise **both ureters as a unit** (never separate them) → taper upper pole ureter if needed → excise distal ureterocele to bladder neck → **plicate the detrusor defect** → submucosal reimplant → mucosal flaps.
- Cecoureterocele remnant can act as an obstructing **urethral windsock**: excise, close in two layers, or fulgurate.
- **Marsupialisation**: excise thin intravesical roof and oversew edges; for small ureteroceles without detrusor defect.
- Persisting reflux **5–10%**, more with tapering.
- **Total reconstruction** (heminephrectomy + excision + lower pole reimplant): definitive, two incisions; best for **older child with massive ureterocele, non-functioning upper pole and significant lower pole reflux**.

8 · DECISION SUMMARY

Ectopic ureter (duplex)

ALGORITHM

Keep upper pole? → Yes + lower pole reflux → common sheath reimplant, or lower pole reimplant + distal ureteroureterostomy

Yes + no lower pole reflux → proximal or distal ureteroureterostomy

Function unclear / massive dilation / sepsis → temporary end ureterostomy (no ureteric trimming; stoma at lateral end of future Pfannenstiel) → DMSA at ~4–6 months → reconstruct or remove

Remove → upper pole nephrectomy with ureterectomy

Ureterocele (Table 52.1)

Procedure	Ideal indication	Advantage	Limitation
TUI	Infant; large ureterocele with vesicoureteral reflux (VUR)	Outpatient; effective decompression; occasionally definitive	Reflux into ureterocele moiety → bladder surgery
Upper pole nephrectomy	Older; large non-functioning upper pole; no VUR	Removes pathology; avoids bladder	Lower pole risk; may still need bladder surgery
Ureteroureterostomy / pyelostomy	Older; functioning upper pole; no VUR	Drains with little risk	Ureterocele left; may reflux
Excision + common sheath reimplant	VUR; functioning upper pole without major dilation	Eliminates obstruction and reflux; no renal risk	Complex; vagina/bladder neck risk; tapering

> **KEY POINT** VUR is the pivot: no VUR → upper tract solution; VUR → bladder-level solution — except in the infant, where TUI is chosen precisely to defer bladder surgery. Upper pole function then chooses nephrectomy vs ureteroureterostomy.

9 · COMPLICATIONS AND THEIR MANAGEMENT

- **Loss of lower pole function** after heminephrectomy (most significant; fever, pain, haematuria in first week) — prevented by preserving the inter-ureteric blood supply and gentle traction.
- **Urinoma**: up to 20% after laparoscopic/robotic cases, mostly when the defect is not closed; asymptomatic non-expanding → observe.
- **De novo reflux** after TUI or heminephrectomy → observe if no infection; excision and reimplantation if persistent/infecting.
- **Voiding dysfunction after repair**: due to inadequate trigonal support (ballooning trigone, poor emptying, infection, upper tract dilation), also after TUI of cecoureterocele. Assess with VCUG **including lateral voiding views** ± videourodynamics; treat by trigonal/bladder neck reconstruction, clean intermittent catheterisation (CIC) or bulking injection for bladder neck incompetence.

10 · OTHER URETERAL ANOMALIES

Duplication

- ~**1 in 125** (0.8%) at autopsy; F:M 1.6:1; unilateral 6× bilateral. **Look for contralateral duplication** — an occult second ectopic ureter explains persistent wetting.
- **Bifid pelvis** = single ureter; **bifid ureter** = partial duplication joining below the ureteropelvic junction (UPJ).
- **Lower pole gets single-system disease** (UPJ obstruction, VUR); **upper pole gets ureteral-formation disease** (ectopia, ureterocele).
- Lower pole UPJ obstruction: manage as single system; option of pyeloureterostomy to the normal upper pole ureter. Lower pole VUR: often high grade, slower resolution; common sheath reimplant, endoscopic injection, or low lower-to-upper ureteroureterostomy.

Triplication (Smith)

- Type 1 three separate ureters (**35%**, commonest); 2 incomplete, two orifices (21%); 3 trifid, one orifice (31%); 4 two ureters, three orifices (13%). Weigert–Meyer usually holds; violated in quadruplication.

Fibroepithelial polyp

- Flank pain, haematuria, hydronephrosis; commonest at **UPJ**; **89% male, 78% left**; only 22% seen preoperatively (filling defect, mistaken for clot).
- At UPJ → pyeloplasty; elsewhere → **ureteroscopic removal**. Sleeve resection/nephrectomy unjustified.

Preureteral vena cava (retrocaval/circumcaval ureter)

- A **venous** anomaly: lumbar **subcardinal vein persists** as the cava, trapping the ureter dorsally. Always **right**-sided. ~1 in 1500; male 3:1; presents 3rd–4th decade.
- Ureter passes behind the cava at L3–L4 and crosses in front medial-to-lateral. **Type I** (commoner): low crossing, **fishhook/J** pelvis, obstruction at iliopsoas edge. **Type II**: high crossing, smooth horizontal curve, little/no obstruction.
- Retrograde pyelogram: **S curve**; **CT urography** is test of choice; MRI shows the cava. Classic mimic of UPJ obstruction — a pyeloplasty will fail.
- Treatment: divide ureter, relocate anterior to cava, reanastomose with **excision/bypass of the aperistaltic retrocaval segment**; laparoscopic/robotic possible. Move the ureter, not the vessel.

Other position anomalies

- **Preureteral iliac artery**: either side; obstruction at L5–S1; associated vasal/mesonephric duct anomalies and ureteral ectopia.

11 · MCQ ONE-LINERS

1. Ectopic ureter in a duplex system always drains the upper pole.
2. Weigert–Meyer: upper pole orifice lies caudal and medial to the lower pole orifice.
3. Female ectopic ureter: continuous dribbling with normal voiding.
4. Male ectopic ureter never causes incontinence; always above external sphincter.
5. Commonest male presentation of ectopic ureter: epididymitis or UTI.
6. Zinner syndrome: ectopic ureter, ipsilateral renal agenesis, seminal vesicle cyst.
7. Most useful ureterocele classification: intravesical versus ectopic (extravesical).
8. Cecoureterocele: intravesical orifice, submucosal cavity extending into urethra.
9. Smooth interlabial mass: prolapsed ureterocele; grape-like: sarcoma botryoides.
10. "Drooping lily": lower pole displaced by dilated upper pole.
11. Pseudoureterocele: bilaminar thick wall, lumen outside bladder.
12. Eversion on VCUG: weak trigone, likely needs open repair.
13. Negative urine culture does not exclude infected ectopic system.
14. TUI incision: low, transverse, close to bladder floor.
15. Watering-can puncture: equal decompression, less reflux.
16. Intravesical ureterocele after TUI: 70–80% decompressed without reflux.
17. Extravesical ureterocele: secondary surgery after TUI up to 70%.
18. Secondary surgery after upper pole nephrectomy for ureterocele: 40–50%.
19. Major heminephrectomy risk: loss of lower pole function.
20. Never separate duplex ureters intravesically: shared blood supply.
21. Distal ureteroureterostomy: stent lower pole ureter first.
22. Duplication incidence: ~1 in 125 (0.8%).
23. Lower pole: UPJ obstruction and VUR; upper pole: ectopia and ureterocele.
24. Fibroepithelial polyp: male, left, at UPJ.
25. Retrocaval ureter: persistent right subcardinal vein; fishhook deformity at L3–L4.

53 Surgery of the Ureter in Children

Reading time ≈ 40 min (slow pace)

1 · PELVIURETERIC JUNCTION (PUJ) OBSTRUCTION — DEFINITION AND AETIOLOGY

- Paediatric hydronephrosis: renal pelvic anteroposterior diameter (APD) **12 mm**; hydronephrosis is **not synonymous with obstruction**.
- **Koff's definition**: obstruction is "any restriction to urinary outflow that, if left untreated, will lead to progressive kidney damage." It is retrospective — so every surgical indication is an attempt to predict harm before function is lost.
- Male and left predominance; bilateral in **10–40%**.

	Intrinsic PUJ obstruction	Extrinsic (crossing vessel)
Age	Newborns, infants (commonest at this age)	Older children, adolescents
Lesion	Narrow segment with interrupted circular muscle → functional discontinuity of peristalsis → poor pelvic emptying	Accessory lower-pole vessel crossing anteriorly; ureter kinks over it
Presentation	Antenatal or incidental dilatation	Intermittent loin pain, nausea, vomiting

- After mobilisation off a crossing vessel there is often no intrinsic narrowing, yet **dismembering and transposing the ureter anterior to the vessel prevents recurrence**.
- **Secondary PUJ obstruction**: high-grade vesicoureteral reflux (VUR) with tortuous ureter kinks the PUJ. Avoid simultaneous reimplant + dismembered pyeloplasty (devascularises ureter from both ends) → conservative or **two-stage** repair.
- **Lower-moiety PUJ obstruction (duplex)**: incomplete duplication → upper-pole ureter to lower-pole pelvis (ureteropyelostomy); complete → standard pyeloplasty; moiety ≤ **5–10%** function → partial nephrectomy.
- **Associated anomalies** in up to 50%: contralateral PUJ obstruction (commonest, 10–40%) → renal dysplasia → multicystic dysplastic kidney → agenesis. VUR in up to 40% (high spontaneous resolution). Horseshoe kidney and VATER association.

2 · DIAGNOSIS

- Antenatal ultrasound (US) at ≈ 20 weeks; postnatal US (incidental or after urinary tract infection [UTI]). Febrile UTI usually responds to antibiotics; few need nephrostomy.
- † **Society for Fetal Urology (SFU) grading**: 0 none; 1 renal pelvis only; 2 pelvis + few calyces; 3 all calyces dilated, parenchyma normal; 4 as 3 + parenchymal thinning.
- † **Mercaptoacetyltriglycine (MAG3) diuretic renogram**: drainage half-time (T½) < 10 min unobstructed, 10–20 min equivocal, > 20 min obstructed; differential renal function (DRF) < 40% or falling is the more reliable marker.
- Neonatal washout curves are confounded by renal immaturity, hydration, diuretic timing, posture and pelvic volume/compliance — false positives up to **44%**, so decide on DRF trend and dilatation, not T½ alone.

3 · INDICATIONS FOR PYELOPLASTY

About **one-third** need surgery; most resolve.

- Increasing APD on US
- Low or falling DRF
- Breakthrough infection on prophylaxis

- Symptoms: pain (older child); failure to thrive, feeding difficulty in infants; haematuria (ruptured mucosal vessels); rarely hypertension (renin–angiotensin activation)

4 · ANDERSON–HYNES DISMEMBERED PYELOPLASTY

Gold standard whatever the access, success ≈ **95%**; originally devised for retrocaval ureter. Dismembering preserves a crossing vessel (ureter transposed anterior), excises the adynamic segment and allows pelvic reduction.

ALGORITHM
Open Gerota fascia → expose PUJ → free lower pole (do not miss an accessory vessel) → sparing bipolar diathermy (protect pelvic/ureteric blood supply) → stay sutures on pelvis and ureter → decompress tense pelvis with 21-G needle → dismember ± reduce pelvis → **spatulate laterally** well into healthy ureter, stenotic segment kept as handle ("no-touch") → apex of spatulation to inferior pelvis → tension-free, watertight 6-0/5-0 absorbable anastomosis → excise stenotic segment just before completion → irrigate clot → cover with perinephric fat

- Long gap: full renal mobilisation; horizontal lower-pelvis incision ("bypass" side-to-side configuration).
- Pitfall: excess stay-suture tension kinks the ureter when released. Gradual taper; drain usually unnecessary.

Alternative	Best indication	Point
Foley Y-V-plasty	High insertion, no vessel	Base of V on dependent medial pelvis, apex at PUJ; stem down lateral ureter
Spiral flap	Long proximal stricture, dependent PUJ	Length:width ≤ 3:1
Vertical flap	Dependent PUJ, large box-shaped extrarenal pelvis	Flap length must match gap
Vascular hitch (Hellström)	Anterior lower-pole vessel, **no intrinsic stenosis**: moderate hydronephrosis, no calyceal dilatation, good cortex, preserved split function; normal PUJ with peristalsis at surgery	Vessel moved cranially, fixed in pelvic wall; no hypertension seen
Uretero-calicostomy	Failed repair, short scarred ureter, dilated lower calyx with thin cortex	Lower-pole nephrectomy to expose calyx widely → spatulated ureter sutured over stent ± nephrostomy; capsule not closed near anastomosis (compression); cover with fat/omentum

Flaps are reserved for severe hydronephrosis with a long stenotic segment where tension-free primary anastomosis is impossible. Non-dismembered (Heineke–Mikulicz) repair is inferior.

KEY POINT Every variant answers two questions: is there a crossing vessel (dismember), and is the gap too long for tension-free anastomosis (flap, mobilisation)?

KEY POINT Repair has one fixed principle and many routes: dismember, excise, spatulate, anastomose without tension. Access, stent and robot are choices of convenience; failure is managed by repeating the same principle.

5 · ACCESS: OPEN VS LAPAROSCOPIC/ROBOTIC

- **Infants**: open preferred (incision barely larger than ports). **Posterior lumbotomy** (prone, lumbodorsal fascia) only in infants/toddlers; flank approach at 11th–12th rib tip at any age; open route extraperitoneal.
- Laparoscopic outcome = open (meta-analysis); robotic: shorter learning curve (3D, dexterity), faster resolution of hydronephrosis vs open.

	Transperitoneal	Retroperitoneal
For	Easier, familiar landmarks; left side transmesocolic	No bowel injury; leak stays extraperitoneal; shorter stay; mirrors open
Against	Bowel risk; urinary ascites → ileus	Learning curve; small space; orientation
Note	Preferred for horseshoe kidney, retrocaval ureter	One randomised trial favoured it (time, stay, feeding); 80–90° flank, table **not flexed**, balloon space 200–300 mL

- **Endopyelotomy**/balloon: poor long-term in < 18 years; niche is **failed pyeloplasty** (anatomy known, no vessel) with double-J for 6 weeks.
- Stones found at pyeloplasty → flexible scope through a port.
- Lap/robotic set-up: transperitoneal 45–60° flank, Hasson camera port, instrument ports more medial for large pelvis or small infant; stay suture on PUJ (transabdominal hitch stitch) for traction. Retroperitoneal: gel roll under contralateral iliac crest, extra 5-mm assistant port for suction and sutures.

6 · STENTS AND DRAINAGE

Stent	Removal	Point
Internal double-J	Second anaesthetic unless string/magnet	Bladder catheter 24–48 h (stent transmits reflux pressure); string → more febrile UTI (girls)
Externalised nephro-ureteric	Pulled at 7–14 days, no anaesthesia	Fewer complications and shorter stay in infants; can be unknotted to drain as nephrostomy
Stentless	—	Needs flawless technique

- Stent duration 1–4 weeks does not change outcome. Antegrade double-J may not pass the narrow infant ureterovesical junction (UVJ) — confirm bladder position with methylene blue in bladder.
- Emergency diversion before repair: **nephrostomy, not retrograde stent** (stent inflames the PUJ).

7 · OUTCOMES AND FAILED PYELOPLASTY

- Success 90–100%, but "success" variably defined (strictest: no reintervention). Commonest reason for conversion: bleeding.
- **Early** failure (infection, pain, ↑ dilatation): usually clot/oedema → settles, ± temporary nephrostomy.
- **Late** failure (falling function, ↑ dilatation): balloon dilatation/endopyelotomy with stent 4–6 weeks, or — usually best — **redo pyeloplasty, transperitoneal** (retroperitoneum scarred); short scarred ureter → uretero-calicostomy.

8 · PRIMARY MEGAURETER

- Descriptive: dilated ureter **≥ 7 mm** (normal ≤ 5–6.5 mm to 16 years). Primary (intrinsic) vs secondary (neurogenic bladder, outlet obstruction).
- **Four types**: obstructed · refluxing · non-obstructed non-refluxing · refluxing + obstructed. Hard part: separating obstructed from non-obstructed non-refluxing — dilatation does not prove obstruction.
- **Primary obstructive megaureter (POM)**: adynamic juxtavesical segment. Ureteric muscle matures **craniocaudally**, distal segment last (after birth) → transient dysfunction and spontaneous resolution. Also excess collagen, abnormal interstitial cells of Cajal.
- 10–23% of antenatal upper-tract dilatation; males, left; bilateral 25%. Symptoms: UTI, pain, haematuria, stones.

Ureter diameter	< 10 mm	≥ 10 mm
Course	Few febrile UTIs; usually resolves	Febrile UTI, stones, pain; minority resolve

Ureter diameter	< 10 mm	≥ 10 mm
Surgery	Virtually none	Substantial minority

Indications: recurrent febrile UTI · progressive dilatation · DRF **< 40%** · DRF fall **≥ 5%** on comparable scans. Follow into adulthood (some worsen at puberty).

> **KEY POINT** Why not rely on the renogram? Neonatal washout depends on immature function, hydration and the capacity of a huge ureter — so serial DRF, diameter and infection history drive surgery. Megaureter is the PUJ problem moved to the bottom of the ureter: an immature adynamic segment that usually matures on its own.

Temporising in infancy

Small bladder and massive ureter are incongruent — a safe non-refluxing tunnel is impossible, so definitive repair is delayed.

Option	Point
Distal (loop) cutaneous ureterostomy	Oblique inguinal incision, ureter brought out as loop, fixed to fascia, opened and everted to skin; rapid decompression of infected system; stomal stenosis rare (wide ureter)
Refluxing reimplant	Converts dangerous obstruction into the "lesser evil" of reflux
Endoscopic balloon dilatation + double-J 2–6 months	Dilate with high-pressure balloon under fluoroscopy until the waist disappears; failure → re-stent or reimplant. Least invasive; success ≈ 70–80% (weak, mostly retrospective evidence); complications up to one-third (migration, infection, VUR, stones)
Nephrostomy	Hard to maintain in infants

Definitive repair

ALGORITHM
Excise stenotic distal segment → straighten → taper → non-refluxing reimplant at **5:1** tunnel length:diameter → stent (feeding tube 7 days or double-J 4 weeks)

	Starr plication	Kalicinski folding	Hendren excisional tapering
Principle	Imbricate lateral wall inward	Exclude lateral lumen, fold over	Excise lateral wall over 8–10 Fr catheter
Blood supply	Preserved — less ischaemia		At risk → ischaemic stricture
Use	Moderate dilatation (< 1.75 cm); bulky if larger		Massively dilated ureter

- A refinement of Hendren excises only mucosa and muscle, leaving vascular adventitia.
- Hendren: running locking suture proximal two-thirds, interrupted distal third (shortened at reimplant). Taper gradually — an abrupt calibre change is itself obstructive.
- Approach: Pfannenstiel, mobilise as for intravesical reimplant (Cohen or Politano–Leadbetter tunnel).
- Very large ureter: extravesical approach avoids a huge trigonal hiatus. Laparoscopic/robotic results comparable short–medium term.
- Success > **90%**; POM better than refluxing megaureter (stiffer collagen-rich UVJ). Worse with valves/neurogenic bladder.
- Complications: obstruction → stent first (oedema); ischaemic stricture → redo reimplant. Reflux → observe/subureteric injection → revision. Transureteroureterostomy for unilateral severe scarring.

9 · SURGERY FOR VESICOURETERAL REFLUX

Aim: one-way, unobstructed antegrade flow by reimplanting the ureter in an antireflux tunnel. Open, laparoscopic/robotic and endoscopic options.

KEY POINT **Principles**: exclude or treat secondary (bladder) causes first · handle the ureter gently (blood supply) and the bladder gently (haematuria, spasm) · mobilise enough ureter for a tension-free repair · tunnel length:diameter **5:1** (Paquin) · keep the course straight at both empty and full bladder (kinking).

- Classified by access (open/laparoscopic), approach (intravesical/extravesical), and tunnel relative to the original hiatus (suprahiatal/infrahiatal).
- Pfannenstiel 2 cm above symphysis; vertical midline for redo (more cranial ureteric mobilisation).

Technique	Approach / tunnel	Strength	Weakness
Politano–Leadbetter	Intravesical; suprahiatal — new superomedial hiatus, vertical tunnel	Long tunnel for high grade	Blind neo-hiatus (bowel, vas, vagina); kink if too lateral/anterior
Glenn–Anderson	Intravesical; infrahiatal, towards bladder neck	Least kinking	Short tunnel
Cohen	Intravesical; transhiatal cross-trigonal	Commonest; success > **98%**; thick trabeculated bladder; with bladder-neck reconstruction	Difficult later retrograde catheterisation
Lich–Gregoir	Extravesical detrusorotomy, ≈ 5 cm	No haematuria or spasm; short catheter time	Transient voiding dysfunction/retention up to **10%**, especially bilateral
Robotic extravesical (RALUR-EV)	Y-shaped detrusorotomy 4–5 cm	Less pain, spasm, stay	Early series: leak, obstruction, ureteric injury

Cohen — key steps

ALGORITHM
Open bladder vertically to 2 cm from neck → pack dome, ring retractor; stay sutures on lateral walls → 5-Fr feeding tube in orifice as handle → circumscribe orifice 1–2 mm with fine diathermy → free ureter in correct plane (preserve distal blood supply) until it reaches the contralateral wall without tension → submucosal tunnel across trigone → new orifice just above contralateral orifice → mucosa-to-mucosa 6-0, detrusor hitch at 6 o'clock

Later access to a cross-trigonal orifice: flexible ureteroscope, or suprapubic needle/cannula aimed at the orifice to pass a guidewire.

Lich–Gregoir — key steps

ALGORITHM
Bladder one-third full → divide obliterated umbilical artery → free ureter **hugging it**, only down to the UVJ → mark 5-cm tunnel along natural course with bladder moderately distended → detrusorotomy to intact mucosa (uniform bulge) → raise flaps wide enough not to constrict → lay ureter in → close detrusor caudocranially, catching ureteric adventitia at apex (prevents slippage) → Foley 12–48 h

KEY POINT Why voiding dysfunction? Pelvic plexus lies ≈ 1.5–2 cm dorsomedial to the UVJ, with branches along the medial ureter. Stay on the ureter, avoid the dorsomedial UVJ, stop at the UVJ. The Zaontz modification (detrusorotomy around the UVJ) encroaches on these nerves and is now avoided.

- Open benchmark: American Urological Association (AUA) meta-analysis — success **95.9%**. Minimally invasive results converging with experience.
- **Follow-up**: US at 6–12 weeks for all. Voiding cystourethrogram (VCUG) may be omitted after low-grade reflux if US normal and no voiding dysfunction or UTI; do it for high grade, new hydronephrosis, UTI or dysfunctional bladder.

Complications of reimplantation

	Early	Long-term
Reflux	Ipsilateral low grade (bladder inflammation) — usually resolves by 1 year. Contralateral ≈ 1 in 10 after unilateral repair, mostly resolves → observe; prophylaxis < 4–5 years; intervene for pyelonephritis. Routine bilateral repair not justified	Rare after low grade; due to high grade, short tunnel/untapered ureter, or **unrecognised secondary reflux** (neurogenic or valve bladder) → treat bladder first (anticholinergic + clean intermittent catheterisation)
Obstruction	Oedema, clot, submucosal haematoma, kink; within 2 weeks with pain, vomiting → double-J or nephrostomy (internalise early); most resolve	Distal ischaemia, angulation (hiatus too lateral/anterior), tight tunnel, orifice stenosis → balloon + stent or unroofing → re-reimplant

- Mild early hydronephrosis is common; it should not exceed the maximal dilatation the reflux caused.
- Contralateral reflux risk: age < 6 years, capacity < 50% expected, refluxing duplex; technique irrelevant.
- **Reoperation**: combined intra/extravesical mobilisation; excise ischaemic ureter to bleeding, peristaltic tissue; short ureter → **psoas hitch** (non-absorbable, before tunnelling), Boari flap, nipple valve (dilated ureter), or reconfigured (Monti) bowel ureter — no tapering, no metabolic effects.

> **KEY POINT** Early complications are tissue reactions — drain and wait. Late ones are design errors (blood supply, hiatus position, tunnel ratio) or an untreated bladder. Reimplanting into an abnormal bladder only worsens dilatation.

Endoscopic injection

- Dextranomer/hyaluronic acid (Dx/HA, Deflux) is the standard agent; ≈ **90%** initial resolution of low-grade reflux after one injection. Teflon (STING, subureteric Teflon injection) abandoned — particle migration, never approved by the US Food and Drug Administration.
- Technique: cystoscopy first (abandon if inflamed) → needle bevel up at 6 o'clock → mound should appear within **0.1–0.2 mL**, else reposition → aim for a **"volcano"** mound with slit/crescent orifice. Bleeding: press with scope, **do not cauterise** (extrusion). Day case, no catheter.

Technique	Needle entry	Result
STING	2–3 mm below orifice, 4–5 mm submucosal	Baseline
HIT (hydrodistension implantation)	Inside hydrodistended orifice, intramural ureter	Better than STING for all grades
Double HIT	Two intraureteric punctures (proximal + distal)	≈ 90%, durable

- Follow-up: antibiotics 3 months; US at 2–6 weeks (acute obstruction); US ± VCUG at 3 months. Failure → 1–2 repeat injections → reimplant (not harder after Deflux; implant encapsulated).
- **Late obstruction** reported (Dx/HA, especially polyacrylate polyalcohol [Vantris]) → long-term US. Deflux loses ≈ 23% volume; Swedish Reflux Trial: **20%** recurrence at 2 years.
- Migration: particles < 50 µm pass lungs to systemic circulation; < 80 µm phagocytosed. Deflux 80–250 µm.

10 · OTHER URETERIC LESIONS

- † **Retrocaval ureter**: right side, ureter passes behind inferior vena cava (persistent posterior cardinal vein); "fish-hook/S" deformity → dismember and re-anastomose anterior to the cava.
- **Congenital mid-ureteric stricture**: misread as megaureter; excise (1–3 cm) + ureteroureterostomy.
- **Fibroepithelial polyp**: upper third, male, left; pain + haematuria in later childhood; confirm by retrograde pyelography or ureteroscopy; endoscopic polypectomy, or resection with

pyeloplasty/ureteroureterostomy.

- **Retroperitoneal fibrosis**: steroids → azathioprine → double-J → ureterolysis.
- **Short ureter in reoperation**: psoas hitch (one side only), Boari flap, transureteroureterostomy, reconfigured (Monti) bowel ureter.

11 · MCQ ONE-LINERS

1. Gold-standard PUJ repair: Anderson–Hynes dismembered pyeloplasty (≈ 95%).
2. Anderson–Hynes was originally devised for retrocaval ureter.
3. Commonest associated anomaly with PUJ obstruction: contralateral PUJ obstruction.
4. Crossing vessel PUJ obstruction typically presents in older children with intermittent pain.
5. Foley Y-V-plasty is contraindicated with a crossing vessel.
6. Spiral flap length:width ratio ≤ 3:1.
7. Vascular hitch needs no intrinsic stenosis and visible peristalsis.
8. Posterior lumbotomy: infants and toddlers only.
9. Retroperitoneoscopic positioning: table not flexed.
10. Emergency drainage before pyeloplasty: nephrostomy, not stent.
11. Stent duration 1–4 weeks does not affect pyeloplasty outcome.
12. Redo pyeloplasty: transperitoneal route preferred.
13. Endopyelotomy niche: failed primary pyeloplasty.
14. Diuretic renogram false-positive rate in neonates up to 44%.
15. Megaureter: diameter ≥ 7 mm.
16. Ureteric muscle matures craniocaudally; juxtavesical segment last.
17. Megaureter < 10 mm virtually never needs surgery.
18. Megaureter surgery if DRF < 40% or falls ≥ 5%.
19. Tunnel length:diameter for reimplant: 5:1 (Paquin).
20. Folding/plication limit: ureter < 1.75 cm.
21. Hendren excisional tapering: risk of ischaemic stricture.
22. POM outcomes better than refluxing megaureter.
23. Balloon dilatation of POM: double-J 2–6 months, complications up to one-third.
24. Duplex lower-moiety with ≤ 5–10% function: partial nephrectomy.
25. Fibroepithelial polyp: upper ureter, left, males.
26. Psoas hitch: unilateral only; second short ureter → transureteroureterostomy.
27. Externalised nephro-ureteric stent: removed at 7–14 days without anaesthesia.
28. Secondary PUJ obstruction from high-grade VUR: two-stage repair, not simultaneous.
29. Commonest reason for conversion of minimally invasive pyeloplasty: bleeding.
30. Commonest intravesical reimplant: Cohen cross-trigonal; drawback is difficult retrograde access.
31. Glenn–Anderson infrahiatal tunnel: least kinking, length limited by bladder neck.
32. Lich–Gregoir voiding dysfunction: pelvic plexus injury dorsomedial to the UVJ.
33. Open reimplant success (AUA meta-analysis): 95.9%.
34. Endoscopic injection: mound within 0.1–0.2 mL; volcano shape predicts success.
35. Injection-site bleeding: compress, never cauterise.

55 Management of Abnormalities of the External Genitalia in Males

Reading time ≈ 37 min (slow pace)

1 · NORMAL DEVELOPMENT AND PENILE SIZE

- Male differentiation (weeks 9–13) needs testosterone (fetal Leydig cells) → dihydrotestosterone (DHT, via 5α-reductase) → functional androgen receptor. Any break → undervirilisation; female is the passive default.
- **Stretched penile length (SPL)**: pubic symphysis (fat pad fully compressed) to glans tip. Term neonate **3.5 ± 0.7 cm**; adult 13.3 ± 1.6 cm. Micropenis cut-off (mean − 2.5 SD): **1.9 cm** at term–5 months; 1.5 cm at 30 weeks, 2.0 cm at 34 weeks; adult 9.3 cm.

2 · PREPUCE: PHIMOSIS AND PARAPHIMOSIS

- **Physiological phimosis** at birth (glanular adhesions ± preputial ring). Smegma accumulation and intermittent erections separate the prepuce: ≈ **90% retractable by 3 years**, < 1% phimotic at 17 years.
- **Secondary (pathological) phimosis**: scarring after forceful retraction, or balanitis xerotica obliterans (BXO). Forceful retraction is discouraged — it converts a physiological problem into a cicatricial one.
- Indications to improve retractility: persistent primary or secondary phimosis, balanitis, posthitis, BXO, urinary tract infection (UTI).
- **Topical steroid**: betamethasone 0.05% twice daily × 30 days (or thrice daily × 21 days) — equally effective; response within 2 weeks; ≈ 1 in 6 fails (more collagen, less elastic fibre).
- **Paraphimosis** (retracted prepuce trapped behind the glans) — oedema builds within hours and can progress to gangrene. Escalation: manual glans compression with distal traction → oedema reduction (iced glove 5 min, granulated sugar 1–2 h, multiple punctures) → dorsal slit → circumcision.

3 · CIRCUMCISION

Policy, technique, anaesthesia

- American Academy of Pediatrics (AAP) 2012: benefits (less UTI, heterosexual human immunodeficiency virus [HIV] acquisition, other sexually transmitted infections) insufficient to recommend routine neonatal circumcision, but sufficient to justify access and third-party payment.
- Devices: Gomco, Mogen, Plastibell — first separate the prepuce fully and inspect meatus/corona to exclude hypospadias. Older children: freehand under general anaesthesia.
- Analgesia is mandatory: **dorsal penile nerve block superior to EMLA** (eutectic mixture of local anaesthetics; methaemoglobinaemia risk); ring block alternative.

Defer / avoid neonatal circumcision	Reason
Hypospadias, penile curvature, torsion, dorsal hood, buried or webbed penis	Prepuce needed for reconstruction; devices unsuitable
Large hydrocele / inguinal hernia	Push skin forward → secondary phimosis, buried or trapped penis
Coagulopathy (contraindication); hyperbilirubinaemia (some)	Haemorrhage
Neonatal intensive care unit setting	More complications

Benefits: penile cancer almost exclusively in men not circumcised at birth; febrile infant UTI risk lower (colonisation of prepuce by uropathogens), but **111** circumcisions prevent one UTI; ≈ **60%**

protection against HIV acquisition in men; no proven sexual dysfunction.

Complications (overall 0.2–5%; rises 10–20-fold after infancy)

Complication	Management
Bleeding — commonest (≈ 0.1%), usually frenular	Compression, cautery, silver nitrate, suture; occasionally haematoma evacuation
Penile degloving (excess skin removed)	Heals by re-epithelialisation: antibiotic ointment + warm baths; no immediate graft
Glanular adhesions / skin bridges	Adhesions separate spontaneously (rare after 9 y); office lysis with EMLA; **steroids do not help**. Skin bridge (can cause chordee/torsion) → excise
Trapped penis (cicatrix)	Betamethasone 0.05% + manual retraction → vertical relaxing incision → formal repair (flaps/Z-plasty)
Amputated glans	Re-attach within **8 h** (often without microscope)
Penile necrosis (cautery touching clamp/ring)	Phallic reconstruction vs female reassignment — latter problematic (male identity from androgen imprinting)

Meatal stenosis

- Almost exclusively in the **circumcised** penis: meatitis from diaper irritation of the exposed meatus, frenular devascularisation, BXO.
- Normal calibre (French, Fr): < 4 months 6–10 Fr; 1–12 months 10–12 Fr; 1–10 years 12–14 Fr; > 10 years 14–18 Fr.
- Symptoms: upward-deflected stream (ventral web), thin forceful stream with straining/frequency, dysuria at start. Urethrography not needed unless UTI/incontinence.
- Treatment: ventral meatotomy/meatoplasty (office with EMLA, or general anaesthesia); suturing mucosa to glans reduces recurrence.

Balanitis xerotica obliterans (lichen sclerosus)

Chronic cicatrising dermatosis → pathological phimosis, may involve glans, meatus, urethra; typically at puberty, rare < 5 years. Steroids only for mild disease; **circumcision is preferred** ± meatotomy; ≈ 1/5 with meatal disease need further meatal surgery; buccal mucosa urethroplasty for severe disease.

4 · INCONSPICUOUS PENIS VERSUS MICROPENIS

> KEY POINT **Stretched length is the single discriminator.** Inconspicuous penis (buried, webbed, trapped) = normal SPL hidden by skin, fat or scar → anatomical treatment. Micropenis = small but normally formed organ → endocrine work-up.

Buried penis type	Mechanism	Treatment
Congenital (poor penopubic/penoscrotal fixation)	Inelastic dartos; skin not anchored to deep fascia	Surgery
Obesity	Suprapubic fat hides shaft	Weight loss first; liposuction only in adolescents
Trapped (after circumcision)	Cicatrix embeds penis in fat	Office dilation or betamethasone + retraction → repair at ≥ 6 months

- **Repair sequence**: excise dysgenic dartos bands (proximal dorsal shaft) → unfurl prepuce for ventral cover → fix dorsal subcutaneous tissue to pubic fascia → fix scrotal subcutaneous tissue to ventral penile base (spare neurovascular bundles).
- **Webbed penis** (penoscrotal fusion): scrotal skin on ventral shaft; congenital or post-circumcision. Repair: transverse incision of web + vertical closure, or Byars' flap transfer; avoid neonatal circumcision.

Micropenis

- Normally formed penis ≥ 2.5 SD below mean SPL (< **1.9 cm** at term). Testes often small/cryptorchid; scrotum small.
- Embryology: differentiation complete by week 12 under placental human chorionic gonadotrophin (hCG)-driven testosterone (defect → small penis *with hypospadias*); later growth needs fetal luteinising hormone (LH) → gonadotrophin deficiency after 14 weeks gives a normally formed micropenis.

Cause	Examples	Hormones
Hypogonadotrophic hypogonadism (commonest)	Kallmann, Prader–Willi, Laurence–Moon–Biedl, CHARGE, hypopituitarism (hypoglycaemia), midline brain defects	Low LH/FSH (follicle-stimulating hormone)
Hypergonadotrophic (primary testicular failure)	Rudimentary testes, gonadal dysgenesis, Robinow	High LH/FSH; no testosterone rise after hCG
Idiopathic	Delayed gonadotrophin timing	Normal axis
Partial androgen insensitivity (rare)	—	Ambiguity commoner

ALGORITHM

History, examination, **karyotype**, SPL, corpora, cryptorchidism → paediatric endocrinologist → testosterone before/after hCG stimulation (± gonadotrophin-releasing hormone [GnRH] test) → anterior pituitary screen (glucose, Na/K, cortisol, thyroid) → magnetic resonance imaging (MRI) brain → **test end-organ response: intramuscular testosterone enanthate for 3 months** (short courses do not affect height); repeat at puberty.

Outcome: male identity and satisfactory sexual function in most.

5 · ABNORMAL ORIENTATION: CURVATURE (CHORDEE WITHOUT HYPOSPADIAS) AND TORSION

Type	Associations	Repair (stepwise)
Ventral (commonest)	Hypospadias; isolated chordee with dorsal hood/deficient ventral skin; frenular chordee	Deglove → excise fibrous tissue superficial to Buck's fascia → Byars' flaps → persistent: dorsal plication/Nesbit/corporal rotation → short urethra: urethral reconstruction
Dorsal	Isolated, epispadias	Deglove → release → ventral plication; protect dorsal neurovascular bundles
Lateral	Corporal asymmetry; often noticed only on erection, later childhood	Plication/ellipse of the corpus **opposite** the maximal curvature

Principle: superficial → deep; release skin and tethers first, plicate the convex side only if curvature persists. **Artificial erection** (saline ± tourniquet) locates the apex and confirms correction. → *Ch 56* for hypospadias.

Penile torsion: usually **counter-clockwise (to the left)**, < 60°; raphe spirals; corpora at the base normally oriented — a skin-level deformity. Avoid neonatal circumcision if repair planned. < 90°: deglove and rotate glans opposite (clockwise) and redrape. ≥ 90°: mobilise base, divide bands, non-absorbable anchoring sutures at corporal base ± dorsal dartos flap.

6 · ABNORMAL NUMBER, MASSES AND LYMPHOEDEMA

Feature	Aphallia (penile agenesis)	Diphallia
Incidence / defect	1 in 10–30 million; genital tubercle fails; 46,XY, scrotum + testes normal	1 in 5 million; accessory penis to complete duplication

Feature	Aphallia (penile agenesis)	Diphallia
Associations	Urethra at anal verge (postsphincteric — commonest, best survival) or rectum; more proximal = higher mortality; vesicoureteral reflux (VUR), renal	Bladder duplication, bifid scrotum, renal
Work-up / decision	Karyotype, anomaly screen, MRI → neophallus vs reassignment (multidisciplinary; reassigned children may develop male identity)	Renal ultrasound (US) + voiding cystourethrogram (VCUG) → individualised

- Penile masses: cysts commonest (excise if symptomatic); **juvenile xanthogranuloma — observe 1 year**.
- Genital lymphoedema (Milroy, praecox, Crohn disease): excisional surgery — all tissue superficial to Buck's fascia; scrotum except posterior skin.

7 · URETHRAL ANOMALIES

- **Congenital urethral fistula** (coronal/subcoronal, normal meatus): multilayer closure, or Thiersch–Duplay if glans bridge thin.
- **Urethral duplication**: usually sagittal (≈ 94%); the **ventral channel is the functional urethra** (sphincter, verumontanum). **Effmann**: type I incomplete (blind), II complete, III with bladder duplication. Presentation: double stream, incontinence (more proximal accessory origin → more incontinence), UTI. VUR commonest association. Evaluate with VCUG, retrograde urethrogram, cystoscopy. Treat if symptomatic; **never use the hypoplastic dorsal accessory urethra as the main channel** — excise, fulgurate/sclerose, septotomy or urethro-urethrostomy.
- Meatal stenosis — section 3.

8 · PRIAPISM

Type	Cause in children	Corporal gas	Treatment
Ischaemic (low-flow)	**Sickle cell disease** (commonest), leukaemia	Dark, hypoxic, acidotic	Hydration, oxygen, alkalinisation, analgesia ± transfusion → aspiration/irrigation ± phenylephrine → shunt (Winter distal; Quackels proximal)
Non-ischaemic (high-flow)	Straddle injury → arteriocavernous fistula (Doppler)	Bright red	Observe → superselective embolisation
Stuttering	Sickle cell	—	Bedtime pseudoephedrine; long-term phosphodiesterase-5 inhibitor

9 · SCROTAL ANOMALIES

- **Penoscrotal transposition**: failed inferomedial migration of labioscrotal swellings; partial forms — shawl, doughnut, prepenile scrotum. Associated with proximal hypospadias + chordee; with complete transposition, **up to 75%** have significant upper-tract anomalies → renal US + VCUG. Repair usually with hypospadias repair: **rotation-flap scrotoplasty** (circumscribe upper hemiscrotums, rotate wings medially under penis) or buttonhole penile relocation.
- **Bifid scrotum**: unfused labioscrotal folds, no raphe; with proximal hypospadias; Z-plasty/scrotoplasty with repair.
- **Ectopic scrotum** (suprainguinal commonest): ipsilateral renal agenesis/dysplasia, ectopic ureter → upper-tract US; scrotoplasty + orchidopexy at 6–12 months.

10 · INGUINAL HERNIA AND HYDROCELE

Entity	Processus vaginalis	Plan
Indirect hernia	Patent; abdominal contents enter	Repair soon after diagnosis

Entity	Processus vaginalis	Plan
Communicating hydrocele	Patent to testis, fluid only	Repair if persists > 1–2 y or fluctuates
Hydrocele of cord (encysted)	Isolated patent segment	Repair after 1 year
Scrotal (non-communicating)	Closed	Observe — most resolve by 1 y
Abdominoscrotal	Closed; extends through ring	Excise/drain; ring tightening unnecessary

- Hernia in 1–5% of children; prematurity (≈ 30% if < 1 kg); right 3:1 (later right descent). Bilaterality risk: female, premature, < 1 year, cryptorchidism.
- **Intermittent bulge** = patent processus → surgery; fixed collection → observe. Parent photographs are reliable; neonatal bowel also transilluminates.
- Timing: reducible → elective within weeks; irreducible → immediate.
- **Repair**: high ligation of the sac at the internal ring (Mitchell-Banks in infants — external oblique not opened). Peripubertal non-communicating hydrocele (> 10–12 y) → scrotal approach.
- Recurrence 0.5–1% (2% preterm, 3–6% after incarceration). Others: iatrogenic cryptorchidism (check testis position at end), atrophy, vas injury.
- Laparoscopic repair now matches open; advantage = sees and closes contralateral ring (fewer metachronous hernias). Contralateral patency ≈ 40% infants → 20% after 10 years, but metachronous hernia only ≈ **7%** — argument against routine contralateral exploration.

11 · ACUTE SCROTUM

Feature	Cord torsion	Appendage torsion	Epididymitis
Frequency	20–30%	**40–60% (commonest)**	5–15%
Peak age	Perinatal; 12–16 y	7–12 y	Infancy, puberty
Onset / vomiting	Acute severe; nausea/vomiting common	Sudden or insidious; uncommon	Insidious; rare
Cremasteric reflex	Usually absent	Present	Present
Signs	High-riding, horizontal lie, diffuse tenderness	**Blue dot**, tender upper-pole nodule	Enlarged tender epididymis; pyuria 20–40%
Colour Doppler US (CDUS)	Absent/reduced flow, heterogeneity; **whirlpool/snail sign**	Avascular upper-pole lesion, epididymal hyperaemia	Increased epididymal flow
Management	**Emergency exploration**	Ice, non-steroidal anti-inflammatory drugs, rest	Supportive; antibiotics only if UTI

Intravaginal torsion

- **Bell-clapper** deformity (high tunica attachment, ≈ 12% at autopsy); familial cases; left-sided predominance; incidence ≈ 1 in 4000.
- Cremasteric reflex may be present in up to 10% of proven torsions — presence does not exclude torsion.
- Predictors: absent reflex, nausea/vomiting, scrotal skin changes. **TWIST score** (validated) stratifies risk: † testis swelling 2, hard testis 2, absent cremasteric reflex 1, nausea/vomiting 1, high-riding testis 1; ≥ 5 high risk → theatre without imaging; ≤ 2 low risk.
- CDUS: absent flow is highly specific, but sensitivity only 63–90% (residual flow in partial torsion; operator-dependent). **Do not delay surgery for imaging when suspicion is high.**

Hours from onset	0–6	7–12	13–18	19–24	> 24	> 48
Orchidectomy risk	**≈ 5%**	20%	40%	60%	80%	90%

ALGORITHM
Suspected torsion → scrotal (hemiscrotal or raphe) exploration → affected side first: open tunica, count twists → detorse, warm packs → **meanwhile fix contralateral testis with non-absorbable suture** → reassess: viable → dartos pouch fixation (non-absorbable); non-viable → orchidectomy (cord ligated in segments); prosthesis later.

- Manual detorsion fails in ≈ 1/3 (twist opposite to that assumed); still needs orchidopexy.
- Late: atrophy even after salvage; subtle semen defects from ischaemia–reperfusion (not antisperm antibodies).
- **Intermittent torsion**: precedes acute torsion in 30–50%; abrupt self-limiting pain → **elective bilateral orchidopexy**.

Perinatal (extravaginal) torsion

Whole cord + tunica twist before scrotal fixation, usually prenatal → hard painless testis with discoloured fixed skin, or vanished testis/nubbin; rarely salvageable; bilateral 5–22%. Elective vs immediate exploration is debated (contralateral asynchronous torsion argues for early surgery). Explore **urgently** if the change follows a documented normal postnatal examination; inguinal approach allows tumour exclusion.

Appendage torsion, epididymitis, others

- Appendix testis (Müllerian, ≈ 80% of testes) and appendix epididymis (Wolffian). Explore only if cord torsion cannot be excluded.
- Epididymitis: prepubertal boy with positive urine culture → renal US + VCUG (look for ectopic ureter, urethral or ejaculatory duct anomaly); recurrent → endoscopy.
- Idiopathic scrotal oedema (fountain sign; self-limiting); Henoch–Schönlein purpura (may cause torsion).

12 · ADOLESCENT VARICOCELE

- Prevalence ≈ **15%** (as in adults); appears after 10 years, peaks **Tanner 3**; left-sided (high left renal vein pressure, incompetent valves, nutcracker); tall thin boys; familial.
- **Isolated right or non-decompressing varicocele** → abdominal/pelvic imaging for a mass.
- Grading: 0 subclinical (CDUS only); 1 palpable with Valsalva; 2 palpable at rest; 3 visible.
- Testicular volume by **US (L × W × H × 0.71)** — most sensitive for asymmetry; volume doubles Tanner 2→3, so serial measures.
- Surrogates (hypotrophy, grade, hormones, histology) do not reliably predict adult semen; catch-up growth after repair may also occur spontaneously. Hence **observation is the default**.

KEY POINT **Indications for repair**: left testis > **20%** smaller (or bilateral hypotrophy), pain, abnormal semen analysis (reliable at Tanner 5 / ≥ 18 y), intratesticular varicocele with asymmetry.

Technique	Recurrence	Hydrocele	Note
Open retroperitoneal (Palomo, mass ligation)	2–4%	Up to 30%	Simple; hydrocele = lymphatics ligated
Laparoscopic, non-sparing	0–9%	11–32%	—
Laparoscopic artery/lymphatic-sparing	1–7%	0–4%	Lymphatic sparing cuts hydrocele
Microscopic subinguinal	0–10%	0–6%	Best balance; rare atrophy; microscope > loupes
Non-microscopic inguinal	7–33%	8–14%	—
Sclerotherapy (retrograde/antegrade)	6–35% / 4–12%	Occasional	Mainly for recurrence; maps collaterals

Mass ligation trades hydrocele for certainty; sparing trades recurrence for lymphatics. Late hydroceles (≥ 2 y) often need surgery. Genitofemoral nerve injury → transient thigh paraesthesia.

13 · EPIDIDYMAL AND VASAL ANOMALIES

- Congenital absence of vas: bilateral → **CFTR** (cystic fibrosis transmembrane conductance regulator) mutations; unilateral → **renal anomalies** (Wolffian origin).
- Testicular microlithiasis → *Ch 57*; undescended testis → *Ch 57*.

14 · MCQ ONE-LINERS

1. Micropenis in a term neonate: stretched length < 1.9 cm (−2.5 SD).
2. Commonest cause of micropenis: hypogonadotrophic hypogonadism.
3. First therapeutic step in micropenis: intramuscular testosterone enanthate for 3 months to test response.
4. Phimosis topical therapy: betamethasone 0.05%; ≈ 1 in 6 fail.
5. Commonest circumcision complication: bleeding, usually frenular.
6. Meatal stenosis occurs almost exclusively after circumcision.
7. Best neonatal circumcision analgesia: dorsal penile nerve block (better than EMLA).
8. Penile torsion is usually counter-clockwise; corpora at base normal.
9. Urethral duplication: ventral urethra is functional; Effmann classification.
0. Aphallia: commonest type has postsphincteric opening at anal verge, best survival.
1. Complete penoscrotal transposition: up to 75% have upper-tract anomalies.
12. Suprainguinal ectopic scrotum: ipsilateral renal agenesis — image upper tracts.
13. Commonest cause of paediatric priapism: sickle cell disease (ischaemic).
14. High-flow priapism follows straddle injury; embolise if persistent.
15. Commonest acute scrotum cause overall (prepubertal): torsion of appendix testis.
16. Present cremasteric reflex does not exclude testicular torsion.
17. Whirlpool/snail sign on high-resolution US = twisted spermatic cord.
18. Torsion < 6 hours: orchidectomy ≈ 5%; > 24 hours ≈ 80%.
19. Always fix the contralateral testis with non-absorbable suture in torsion.
20. Perinatal torsion is extravaginal; bilateral in 5–22%.
21. Varicocele peaks at Tanner 3; repair if > 20% volume discrepancy.
22. Lowest hydrocele + recurrence: microscopic subinguinal varicocelectomy.
23. Bilateral absent vas = CFTR mutation; unilateral absent vas = renal anomaly.

56 Hypospadias

Reading time ≈ 34 min (slow pace)

1 · DEFINITION AND DIAGNOSIS

- **Classic triad**: proximal (ventral) ectopic meatus, ventral penile curvature, dorsally hooded foreskin — all from arrested penile development. Curvature and hood are not always present.
- Incidence ≈ **1 in 150–300** male births; one of the commonest congenital anomalies. Diagnosis is clinical.
- **Megameatus with intact prepuce (MIP)**, ≈ 5%: wide distal meatus, normal foreskin, no severe curvature — hidden until the foreskin is retracted. Retract before every newborn circumcision.
- Avoid circumcision with curvature or ventral skin deficiency (foreskin may be needed); curvature with orthotopic meatus → defer circumcision until curvature is repaired.

2 · EMBRYOLOGY

- Critical window **weeks 8–16**; urethral closure proceeds from the anus towards the glans tip.
- Sex-determining region Y (SRY) at 7–8 weeks → testis → Leydig testosterone → **5α-reductase type 2** → dihydrotestosterone (DHT) locally. **DHT at 8–12 weeks** is the key mediator. Sonic hedgehog drives tubercle and plate formation.
- **Endodermal "two-zipper" theory** (Kurzrock, current): plate continuous through the glans; "opening zipper" canalises plate into a groove (both sexes), then androgen-dependent "closing zipper" (male only). Older **ectodermal ingrowth theory** (Glenister): glanular urethra from ectodermal ingrowth fusing at the corona. MCQ
- Prepuce fuses dorsally first; ventral fusion awaits urethral fold fusion (≈ 20 weeks), so ectopic meatus → incomplete foreskin in **95%**.

> **KEY POINT** Hypospadias is an arrest of the androgen-dependent closing zipper: the meatus stops where closure stopped, the prepuce cannot close ventrally over an open urethra, and unfused dysplastic ventral tissue tethers the shaft into curvature.

3 · AETIOLOGY

- **Multifactorial**; every thread converges on insufficient androgen action in weeks 8–16.
- **Fetal/placental**: prematurity, small for gestational age, intrauterine growth restriction, placental insufficiency (≈ 2× risk); assisted reproduction (independent risk).
- **Genetic**: twin concordance 9–27%; first-degree relative 9–15%. Chromosomal abnormality in up to 30%; no single gene. Gene groups: penile development (HOX, fibroblast growth factor), gonadal (WT1, SOX9, DMRT1, GATA4), androgen cascade (LH receptor, 5α-reductase, androgen receptor). ≈ 200 syndromes: WAGR syndrome (Wilms, aniridia, genitourinary anomalies, intellectual disability) (Wilms tumour, aniridia, genitourinary anomalies, developmental delay), Denys–Drash, Smith–Lemli–Opitz.
- **Endocrine-disrupting chemicals (EDCs)**: pesticides, oestrogens/progestins, diethylstilbestrol, phthalates, phytoestrogens — compete at the androgen receptor; proven in animals, unproven in humans.
- **Testicular dysgenesis syndrome**: hypospadias, cryptorchidism, poor semen quality and testis cancer share one fetal androgen-disrupting exposure.
- **Anogenital distance (AGD)** (anus to penoscrotal junction): marker of androgen action; shorter AGD = more severe hypospadias.

4 · CLASSIFICATION AND SEVERITY

- By meatal position: distal (glanular, coronal, distal shaft) **70–85%**; proximal (proximal shaft, penoscrotal, scrotal, perineal) 10–25%.

- Meatus is the **least reliable** marker: a "distal" meatus may overlie absent spongiosum to the penoscrotal junction with severe curvature. **Classify after degloving**: meatus + curvature + plate.

Score	Glans (G)	Meatus (M)	Shaft curvature (S)
1	Good size, deep grooved plate	Glanular	None
2	Adequate, grooved	Coronal sulcus	<30°
3	Small; narrow/flat plate	Mid or distal shaft	30–60°
4	Very small; indistinct plate	Proximal shaft/penoscrotal	>60°

GMS score (Merriman 2013): total 3–12, scored in theatre; higher → more complications. A coronal meatus with a small flat glans and 45° curvature scores worse than its position suggests.

Plate objective scoring tool (POST): ratio of neomeatus length (AB) to glans fusion length (BC); low ratio = poor plate → favour inlay graft over plain tubularised incised plate (TIP) repair.

5 · ASSOCIATED ANOMALIES AND DSD WORK-UP

- Inguinal hernia/hydrocele 9–16%; undescended testis (UDT) ≈ 7%, ≈ 10% in proximal cases.
- **Disorder of sex development (DSD) work-up** when: UDT (uni- or bilateral) + proximal hypospadias; perineal hypospadias; atypical genitalia. Since 2006 severe hypospadias counts as 46,XY DSD. DSD risk ≈ **50%** with non-palpable vs 15% palpable testis; 64% proximal vs 7% distal.
- Work-up: ultrasound (US) for Müllerian remnants, karyotype, testosterone, follicle-stimulating hormone (FSH), luteinising hormone (LH), anti-Müllerian hormone (AMH). A 46,XY karyotype does not exclude 5α-reductase deficiency, androgen insensitivity, testicular dysgenesis or ovotesticular DSD → *Ch 59*.
- **Prostatic utricle**: commoner in proximal cases; large ones → stasis, urinary tract infection (UTI), difficult catheterisation. Most asymptomatic; excision (endoscopic/robotic) rare because the vas is at risk.
- No routine voiding cystourethrogram (VCUG) or renal US; image only with other systemic anomalies.

> **KEY POINT** Hypospadias + cryptorchidism signals a systemic androgen problem, not a local one — hence karyotype before surgery.

6 · PRE-OPERATIVE ASSESSMENT

- **Goals**: straight erection, urethra to the tip, acceptable cosmesis — fewest procedures, durable into adulthood. A straight penis voiding standing may gain little; counsel honestly.
- History: previous surgery, UTI, erectile curvature (parents' photographs), bleeding disorders. Examine: stretched penile length (exclude micropenis), both groins, AGD, penoscrotal transposition, torsion, scrotal cleft, meatal stenosis ("laser-beam" stream — may justify surgery even in distal cases).
- **Timing**: genital awareness ≈ 18 months. American Academy of Pediatrics (AAP, 1996): 6–12 months; authors start at 6 months within **6–18 months**. Systematic review (>16,000) → fewer complications when younger; repair by 18 months. One short anaesthetic appears low-risk for neurodevelopment.
- **Androgen stimulation**: glans width **<14 mm** → more complications. Testosterone, DHT or human chorionic gonadotrophin (hCG) enlarge glans, reducing glansplasty tension. Authors: testosterone cypionate **2 mg/kg intramuscular (IM) at 5 and 2 weeks** pre-op if glans <15 mm (≈ 4 mm gain). Two randomised trials → fewer complications; meta-analyses inconclusive. Side effects (aggression, erections, pigmentation) resolve within 6 months; concerns: bleeding, healing; severe cases may be partly androgen-insensitive.

7 · INTRA-OPERATIVE PRINCIPLES

- Caudal block (most popular; fistula link disputed), penile or pudendal block; ketorolac 0.5 mg/kg once. IV antibiotics at induction. Tourniquet (release every 30 min) or epinephrine.
- Cystoscopy only for severe proximal/redo (utricle may block catheter).
- Sutures: rapidly absorbed (chromic, polyglactin) beat polydioxanone; 6-0–8-0 in infants. Slow sutures leave epithelialised tracks → fistula.
- Magnification, fine handling, **tension-free** urethra/glans/skin (tension → ischaemia → fibrosis, breakdown), non-overlapping suture lines, haemostasis (haematoma kills flaps).
- **Never use hair-bearing skin** for urethroplasty: postpubertal hair → stones, recurrent UTI.
- Higher surgeon volume → fewer complications.

8 · CURVATURE: ASSESSMENT AND CORRECTION

Curvature, not meatal position, is the main determinant of **one-stage vs staged repair**. Causes: tethered ventral skin, dysplastic dartos/Buck's fascia, short urethra, corporal disproportion.

ALGORITHM
Circumscribe → deglove to penoscrotal junction, excise dysplastic dartos → **artificial erection** (Gittes–McLaughlin tourniquet + saline; or corporal compression; or prostaglandin) → straight → urethroplasty · residual <30° → dorsal plication · >30° → divide urethral plate → still >30° → ventral corporal lengthening (usually staged)

Grading: mild <30°, moderate 30–45°, severe ≥45°. Eyeballing underestimates; a tourniquet can hide basal curvature (compress corpora against pubis instead).

Feature	Dorsal plication (shortening)	Ventral corporotomy ± graft (lengthening)
Technique	Nesbit ellipses; Baskin single midline 12 o'clock plication (nerve-free zone); Heineke–Mikulicz dorsal incision	Transverse ungrafted corporotomies (staged tubularised autograft, STAG) or single 9-to-3 o'clock corporotomy + graft (dermis 20% larger, tunica vaginalis, small intestinal submucosa)
Indication	≤30° (authors ≤15°)	Moderate/severe after degloving and plate division
Risks	Neurovascular injury off midline; recurrence at puberty; shortens	Complex; limits later ventral skin; graft contraction

Recurrence: plication 28% vs grafting 9% in one series; not dividing the plate is an independent risk. Correct fully first time — under-correction generates repeated surgery through puberty.

9 · URETHROPLASTY TECHNIQUES

Tenets: correct curvature → advance urethra or urethroplasty → glansplasty → skin cover ± preputioplasty. TIP is used for >90% of distal repairs.

Technique	Indication	Key point / pitfall
MAGPI (meatal advancement and glanuloplasty; Duckett 1981)	Glanular meatus, no curvature, mobile meatus	No tubularisation; used more proximally → retrusive meatus, flat glans
Urethromeatoplasty	Stenotic distal meatus, blind pit	Heineke–Mikulicz
GAP (glans approximation)	Wide deep groove, fish-mouth meatus (MIP)	Deep glans wings, two-layer neourethra
Thiersch–Duplay	Wide plate, adequate glans	U-incision, tubularise over 12 Fr, dartos flap
TIP / Snodgrass (1994)	Distal and midshaft; plate too narrow to tubularise; mild curvature	Deep midline plate relaxing incision; stop 2–3 mm short → slit meatus; risk meatal stenosis; add inlay graft if poor plate

Technique	Indication	Key point / pitfall
Mathieu (flip-flap)	Distal, shallow/deficient plate	Parameatal flap; narrow base → ischaemia, fistula; fish-mouth meatus
Onlay island flap	Plate kept but too narrow/thin; curvature corrected by degloving	Inner preputial island flap; diverticulum if flap too wide; less stenosis
Duckett tube (transverse preputial island flap)	Proximal; plate divided for curvature	Single-stage tube; higher complications; anchor flap edge to avoid diverticulum
Two-stage Bracka (free graft)	Proximal with severe curvature, redo, poor plate	Stage 1: correct curvature, divide plate, quilted inner preputial/buccal graft; stage 2 (≥6 months): tubularise
Two-stage Byars (pedicle flaps)	Proximal with severe curvature	Split dorsal prepuce transposed ventrally; highest staged complication rates (fistula, diverticulum)

- Variants: **STAG** (ungrafted corporotomies + graft; lowest reported proximal complications, ≈ 23%); STAC (no graft at stage 1, graft stage 2, tubularise stage 3 — authors' choice); distal-first/Ulaanbaatar.
- **Why stage**: a urethroplasty laid over a fresh corporotomy/graft fails more often; stage 1 creates supple ventral tissue. Always repeat artificial erection at stage 2.
- Proximal TIP abandoned: plate mobilisation devascularises and more recurrent curvature.

> **KEY POINT** Why proximal repairs fail — Poiseuille: resistance ∝ length / radius⁴. A long, rigid, spongiosum-free skin tube ending in a fixed non-distensible glans raises proximal pressure → fistula, diverticulum, stricture. 25–75% need extra procedures after planned two-stage repair.

Glansplasty and coverage

- Deep glans wing mobilisation, tension-free **two-layer glans closure**, mucosal collar. Glans tension → dehiscence.
- **Barrier layer** of vascularised non-epithelialised tissue over the neourethra reduces fistula: de-epithelialised dorsal preputial dartos (Belman flap), ventral dartos, **tunica vaginalis** (mobilise to external ring to avoid tethering), spongiosum. Offset suture lines; must not compress glans.

10 · DRESSING, STENTS AND POST-OPERATIVE CARE

- Compressive dressing (circumferential Tegaderm including glans, or "sandwich" dressing) for ≈ **72 h**; then petroleum jelly.
- Diversion **3–21 days** by complexity; soft dripping stent; suprapubic tube for older/redo. Short distal repairs can be unstented.
- Oxybutynin until stent removal (bladder spasm → voiding around stent). Stiff stents/laterally pulled Foley → glans dehiscence.
- Paracetamol + ibuprofen regularly (non-steroidal anti-inflammatory drugs do not increase bleeding). Antibiotic prophylaxis: evidence mixed.

11 · COMPLICATIONS AND MANAGEMENT

Complication	Standard definition
Fistula	Urethral leak anywhere below meatus
Glans dehiscence	Wing separation → coronal/proximal meatus; glans fusion <2 mm
Meatal stenosis	Obstructive symptoms + calibration **<8 Fr** prepubertal / <12 Fr postpubertal
Diverticulum	Segmental sacculation on voiding
Recurrent curvature	**>30°** on erection

- **Urethrocutaneous fistula** — commonest (just under 10%), usually at original meatus or coronal margin. Causes: ischaemia, haematoma, infection, overlapping suture lines and especially **distal obstruction**. Rarely closes spontaneously. Repair ≥ **6–12 months** after surgery: calibrate (bougie à boule) ± urethroscopy, stress test with saline/methylene blue → excise tract, close in offset layers with dartos/tunica vaginalis interposition. Coronal fistula → redo urethroplasty and glans closure, not simple excision. Recurrence ≈ 20%.
- **Meatal stenosis**: dilatation ± topical steroid → meatotomy or redo urethroplasty. Commoner after TIP than Mathieu.
- **Glans dehiscence**: reoperate only for voiding symptoms; inlay graft avoids glans tension.
- **Urethral stricture** (often late): cystoscopy; dilatation/urethrotomy rarely durable → revision urethroplasty with buccal mucosa, often staged.
- **Diverticulum** (after flaps, staged, proximal): excise redundant epithelium, reduce to uniform calibre, keep dartos as extra layer; always exclude distal obstruction.
- **Balanitis xerotica obliterans (BXO)/lichen sclerosus**: white scarring of meatus/neourethra. Topical clobetasol/betamethasone usually fails → excise all involved tissue (histology), staged buccal or inner preputial graft. Recurrence up to **40%**; BXO can invade buccal grafts.
- **Recurrent curvature**: under-correction, graft contraction, skin tethering, pubertal disproportion. Full degloving + artificial erection; staged ventral lengthening, buccal graft, Cecil modification if skin short.
- **Hairy urethra**: hair + narrowing → stones, UTI → laser epilation, endoscopic stone removal, or resect and replace with buccal graft.

> **KEY POINT** Nearly every urethroplasty complication traces to distal obstruction, ischaemia or tension — so fistula/diverticulum repair begins by calibrating the distal urethra.

Reoperation

- Each redo degrades tissue; Devine: avoid "small operations" — wide scar excision and tissue transfer instead. Always cystoscope and do artificial erection.
- **Buccal mucosa** = graft of choice (hairless, wet-adapted, little contraction); **lip mucosa** preferred within the glans (thinner). Bladder mucosa abandoned (meatal prolapse). Graft 10% larger, quilted, on preserved dartos; urethroplasty ≥6 months later. Dorsal inlay beats onlay or tube in redo.
- **Cecil–Culp**: bury repaired penis in scrotum to parasitise its blood supply; modified release at 9–12 months.
- Adjuncts (unproven): topical nitroglycerin, hyperbaric oxygen, tissue expanders.

12 · OUTCOMES AND FOLLOW-UP

- TIP meta-analysis (Pfistermuller 2015, 4,675 boys): complications 10.6%, reoperation 4.5%; fistula > stenosis > stricture; dartos layer improves results.
- Distal success 83–95%; proximal complications 23–68%.
- Only ≈ 50% of complications appear in year 1, ≈ 75% by 3 years; fistula early, stricture/curvature late → follow proximal cases **through puberty** (stream after toilet training; curvature, sexual function at puberty).
- Recurrent UTI after repair (≈ 2%) → image for diverticulum, stricture, utricle.
- Instruments: Hypospadias Objective Scoring Evaluation (HOSE; surgeon), Penile Perception Score (patient/parent), Hypospadias Objective Penile Evaluation (HOPE; photo panels; cosmetic only).

Adult issues

- Untreated mild distal hypospadias: minimal sexual, fertility or voiding problems.
- Proximal repairs: worse satisfaction, sensation and fertility; ejaculatory problems; concern about penile length (chief adult complaint); sexual inhibition and lower self-esteem but no major psychopathology. Even 20–30° curvature can impair intercourse.

13 · MCQ ONE-LINERS

1. Hypospadias triad: ventral ectopic meatus, ventral curvature, dorsal hooded foreskin.
2. Critical window of urethral development: weeks 8–16; DHT at 8–12 weeks is key.
3. Current embryological model: endodermal "two-zipper" theory (Kurzrock).
4. Megameatus intact prepuce: ≈ 5%; found on foreskin retraction.
5. Distal hypospadias forms 70–85% of cases.
6. GMS score ranges 3–12; higher predicts complications.
7. Hypospadias + undescended testis → karyotype; DSD ≈ 50% if testis non-palpable.
8. Glans width <14 mm increases complication risk.
9. Pre-op testosterone cypionate 2 mg/kg at 5 and 2 weeks before surgery.
0. Recommended repair age: 6–18 months.
1. Residual curvature <30° after degloving → dorsal plication.
2. Curvature >30° after plate division → ventral corporotomy ± graft, usually staged.
3. MAGPI suits a glanular meatus with no curvature.
14. TIP: midline plate incision; commonest late problem meatal stenosis.
15. Mathieu flip-flap: fewer meatal stenoses than TIP, similar fistula rate.
16. Onlay island flap complication typical: diverticulum.
17. Duckett tube = transverse preputial island flap tube.
18. STAG has the lowest reported proximal complication rate (≈ 23%).
19. Byars two-stage flaps: highest complication rate among staged options.
20. Poiseuille: resistance ∝ length / radius⁴ — explains proximal failures.
21. Commonest complication: urethrocutaneous fistula.
22. Coronal fistula → redo urethroplasty, not simple closure.
23. Meatal stenosis: symptoms + <8 Fr prepubertal.
24. Recurrent curvature defined as >30°.
25. BXO recurrence after excision and buccal graft: up to 40%.
26. Graft of choice for redo: buccal mucosa; lip mucosa within glans.
27. Only half of complications appear within 1 year.

57 Etiology, Diagnosis and Management of the Undescended Testis

Reading time ≈ 35 min (slow pace)

1 · DEFINITIONS

Term	Definition / key point
Undescended testis (UDT) / cryptorchidism	Testis not in normal scrotal position (midpoint at or below mid-scrotum); clinically palpable vs non-palpable
Ectopic testis	Off the line of descent; commonest site **superficial inguinal pouch (SIP)**; others perineal, femoral, prepubic, penile, transverse/contralateral scrotal
Retractile testis	Retracts but can be brought into the scrotum and stays there at least temporarily; may progress to acquired UDT
Acquired (ascending)	Previously documented scrotal; usually diagnosed at 8–11 years
Recurrent	Undescended at birth → descended spontaneously → re-ascended (10–22%, at 1–5 years)
Secondary	Suprascrotal after inguinal hernia repair
Vanishing testis	Formed, then lost (perinatal torsion/vascular accident): monorchia; bilateral = anorchia
Agenesis	Never formed; associated with ipsilateral Müllerian persistence (extremely rare)

2 · EMBRYOLOGY OF DESCENT

- SRY (sex-determining region Y) + steroidogenic factor 1 (SF1) → SOX9 → Sertoli cells → testis cords. Fetal Leydig cells active from 6–7 weeks.
- Two Leydig hormones control descent: **insulin-like 3 (INSL3)** via its receptor RXFP2 (relaxin family peptide receptor 2), peak 15–17 weeks, and **androgens** via the androgen receptor, testosterone peak 14–16 weeks (placental human chorionic gonadotrophin [hCG]-driven). Anti-Müllerian hormone (AMH) from Sertoli cells regresses Müllerian ducts at 9–10 weeks.
- The gubernaculum attaches to the epididymis; it does not pull the testis.

Phase	Timing	Event	Control
Transabdominal (relative)	10–15 weeks	Testis held near internal ring while gubernaculum swells (outgrowth); cranial mesonephric ligament regresses	Classically INSL3 ("non-androgenic factor"); now INSL3 + androgen synergistic
Inguinoscrotal	**20–28 weeks**	Gubernaculum widens canal, processus vaginalis forms; testis passes rapidly through canal, then gubernaculum regresses; most scrotal after 27 weeks	Androgens acting via **genitofemoral nerve (GFN)**, released **calcitonin gene-related peptide (CGRP)**; cremaster myogenesis

> **KEY POINT** Outgrowth creates the canal; regression lets the testis follow — so a defect in either produces the same phenotype. INSL3 and androgen act on one target (gubernacular/cremaster myogenesis), hence one clinical picture whichever fails.

Minipuberty (hypothalamic–pituitary–gonadal [HPG] axis surge, peak 1–3 months): luteinising hormone (LH), follicle-stimulating hormone (FSH), testosterone, AMH and inhibin B (INHB) rise; gonocytes transform into **adult dark (Ad) spermatogonia** (stem-cell pool). This is the window in which a positional defect becomes a germ cell deficit.

3 · EPIDEMIOLOGY AND RISK FACTORS

- Prevalence **2–4% at birth** (term) → **1–2% at 3 months**; up to 45% in preterm. Spontaneous descent mainly in first 3–6 months; rarely after 6 months.
- 80–85% isolated (non-syndromic). Risk factors: **prematurity / low birth weight**, family history (twins > brothers > sons; maternal half-brothers > paternal), maternal diabetes, smoking; diethylstilbestrol (DES) exposure doubles risk; endocrine-disrupting chemicals (EDCs) — mixture effects, unproven singly.
- Genetics: polygenic threshold trait; rare INSL3, RXFP2, NR5A1 variants.
- Testicular dysgenesis syndrome: cryptorchidism + hypospadias + infertility + germ cell tumour from impaired fetal testis function. Reduced anogenital distance marks low prenatal androgen.
- Syndromic: androgen synthesis/action defects, Kallmann, Klinefelter (> 50% of prepubertal Klinefelter boys cryptorchid), persistent Müllerian duct syndrome (PMDS — AMH or AMHR2 mutation; transverse testicular ectopia), prune-belly (all), posterior urethral valves, cerebral palsy (bilateral), abdominal wall defects. **Gastroschisis — 50–55% descend spontaneously: observe.** Caudal-field and diaphragmatic hernia associations are ipsilateral.

Congenital vs acquired

Acquired UDT is probably a never-fully-descended testis unmasked by growth: lower position (often SIP), usually closed processus, reduced Ad spermatogonia like congenital cases, and similar semen outcomes. Retractile testes carry a risk of ascent (prospective ≈ 2–3%) → **annual examination through childhood**.

4 · HISTOLOGY AND FERTILITY

- Germ cell loss begins **after infancy** and progresses; contralateral descended testis also affected to a lesser degree — evidence of an intrinsic defect.
- Normal spermatogonia per tubule (S/T): 2.5 at birth, nadir 1.2 at 2–3 years — needs age-specific reference.
- Delayed gonocyte disappearance and **Ad spermatogonia** formation; Ad absent in 7–85%; primary spermatocytes delayed. Sertoli cell number falls after 4 months.
- Higher germ cell counts with younger age at surgery and palpable testis.
- Associated anomalies: **patent processus** in 87% of unilateral UDT; aberrant lateral gubernacular attachment 66–75%; epididymal non-fusion/elongation (commoner in congenital than acquired). An undescended testis is also at higher risk of torsion, which can mimic an incarcerated hernia.

Trial	Design	Result	Take-home
Kollin et al.	Randomised controlled trial: orchidopexy at 9 months vs 3 years	Early group: higher Sertoli and germ cell counts, better testicular growth (to 4 years)	Operate in first year; age at surgery, not hormone levels, predicts germ cells

Fertility outcome

Outcome	Former unilateral	Former bilateral
Oligospermia / azoospermia	31% / 14%	42% / 31%
Paternity	**≈ 90% of those attempting — same as controls**	≈ 65% of those attempting (≈ 1/3 overall)

- Unilateral: the normal testis carries fertility; semen overstates infertility. Bilateral: semen reflects true infertility.
- Ad spermatogonia at biopsy predict adult counts (present in both → normal count in 84%; absent → subnormal in all).
- Population study: surgery **before 18 months** reduced need for assisted reproduction; each 6-month delay increased it by 5%.

- Orchidopexy before 19 years improves testicular sperm retrieval in azoospermia.

5 · MALIGNANCY

- Testicular germ cell tumour (TGCT) risk **2–5×** (pooled odds ratio ≈ 4); ≈ 1% of cryptorchid men; 5–10% of TGCT patients have a history of UDT. Contralateral testis: odds ratio ≈ 1.5.
- **Swedish population study**: cryptorchidism odds ratio 3.2 — ≈ 2.8 if diagnosed before puberty vs ≈ 5.0 after (hypospadias 2.3, inguinal hernia 1.3). Meta-analysis (Walsh 2007): orchidopexy after 10–11 years → relative risk **5.8**.
- Dose–response by age at orchidopexy: < 9 y odds ratio 1.1–2.35; 13–15 y 5.1; ≥ 16 y 6.2; untreated adults 14.4. Risk rises with each 6-month delay beyond 18 months.
- Persistently cryptorchid → **seminoma** (74%); after orchidopexy → mostly non-seminoma (63%).
- Precursor germ cell neoplasia in situ (GCNIS, formerly intratubular germ cell neoplasia/carcinoma in situ): 2–3% of formerly cryptorchid men, ≤ 0.4% of non-syndromic children. Higher risk with disorders of sex development (DSD)/chromosomal defects.
- Postpubertal UDT with normal contralateral testis: **orchidectomy** preferred (up to age 50).
- **Testicular microlithiasis**: 3–6% of boys, not clearly an independent TGCT risk factor, no added risk with UDT → routine ultrasound (US) surveillance not recommended.
- Counsel teaching testicular self-examination.

† The Swedish data (Pettersson 2007): relative risk 2.23 if operated before 13 years vs 5.40 at ≥ 13 years.

6 · DIAGNOSIS

Examination — the gold standard

ALGORITHM
Warm room, warm hands, supine (frog-leg/cross-legged or squatting if needed), lubricant on fingers → sweep from anterior superior iliac spine along the canal towards the scrotum with one hand → palpate with the other for the lowest attainable position → while compressing the canal, invaginate scrotum to feel SIP/distal testis → sustained gentle traction fatigues cremaster: **stays in scrotum = retractile; springs back = undescended**.

- Document palpability, position, size, mobility, hernia/hydrocele, penile size, meatus.
- ≈ **80% palpable**; 60–70% unilateral. Non-palpable at surgery: abdominal/peeping 25–50%, vanished 15–40%, extra-abdominal missed 10–30%.
- Contralateral hypertrophy (volume > 2 mL, length ≥ 1.8–2 cm) suggests monorchia but does not replace exploration.
- Interobserver agreement is poor; examination under anaesthesia (EUA) is the final arbiter.

Imaging — not recommended

American Urological Association (AUA) 2014: **US or magnetic resonance imaging (MRI) is not recommended**. US sensitivity ≈ 45%, specificity 78%; MRI 65%/100%. A positive scan does not avoid surgery and a negative scan does not exclude a testis — the only effect is a delay of ≈ 3 months. Possible MRI use: ectopic abdominal testis not found at laparoscopy.

Bilateral non-palpable or UDT with hypospadias → DSD work-up

- Bilateral non-palpable newborn (especially with abnormal phallus): **urgent karyotype and hormones** to exclude **congenital adrenal hyperplasia (CAH)** (undiagnosed salt-wasting); pelvic US for Müllerian structures; defer circumcision.
- Proximal hypospadias + UDT → chromosomal abnormality in 32–47% → *Ch 56*.
- Micropenis + bilateral UDT: hypogonadotrophic hypogonadism or anorchia — measure LH, FSH, testosterone during minipuberty.

Marker	Anorchia	Bilateral abdominal UDT
FSH, LH	Raised	Normal
AMH, INHB (best prepubertal Sertoli markers)	Undetectable/low	Present
Testosterone after hCG stimulation	No rise	Rises

hCG stimulation is poorly standardised and is no longer the test of choice; gonadotrophins may be normal in mid-childhood anorchia. If hormones prove testicular tissue, it still has to be found: **laparoscopy remains the confirmatory test**. In a genetic male with bilateral non-palpable testes, ≥ 95% are abdominal.

7 · MANAGEMENT PRINCIPLES AND TIMING

- Aims: optimise fertility, reduce/allow detection of malignancy, cosmesis, prevent torsion and hernia.
- **Timing**: observe to **6 months** (corrected age) → refer → orchidopexy **6–18 months**, preferably by 12 months. Preterm: observe to 6 months beyond due date.
- Retractile testis: no surgery; examine yearly.
- Acquired UDT: operate (observation not recommended, especially bilateral).

Hormonal therapy — not recommended (AUA)

- Luteinising hormone-releasing hormone (LHRH) intranasal: descent ≈ 20%; hCG: limited, never placebo-tested. Neither distinguishes retractile from undescended.
- hCG may increase germ cell apoptosis and reduce adult volume (infertility relative risk 4.7 in Lee series).
- Adjuvant LHRH (buserelin 10 µg intranasally alternate days × 6 months; Hadziselimovic) for bilateral UDT with absent Ad spermatogonia: pilot randomised data only; not widely used.

8 · SURGERY FOR THE PALPABLE TESTIS

Inguinal orchidopexy — key steps

ALGORITHM

EUA → skin-crease groin incision → look for testis in SIP → open external oblique sparing ilioinguinal nerve → free testis and divide gubernaculum distal to the sac (protect a long-looping vas) → **separate and high-ligate the patent processus at internal ring** (the main length-gaining step) → divide lateral spermatic fascial bands → retroperitoneal dissection → **Prentiss manoeuvre** (route medial to inferior epigastric vessels) → transverse scrotal incision, **subdartos pouch** → pass testis without twist → excise appendages, record volume → close dartos around cord (transalbugineal sutures unnecessary — 98.7% success without).

- Cannot reach scrotum → staged procedure; orchidectomy only for atrophic/abnormal testis, postpubertal patient, or insufficient vas.
- Biopsy: not routine; indicated in DSD/ambiguity, consider > 12–13 years at orchidopexy.

Scrotal (Bianchi) orchidopexy

Single high scrotal-border incision (Bianchi–Squire) → deliver testis → dissect sac and cremaster high, ligate processus through the scrotal wound → subdartos pouch. Inguinal conversion 3.5–4.4%; atrophy 0.3–0.6%; less pain than inguinal; increasingly first choice for palpable (especially low) testes.

9 · THE NON-PALPABLE TESTIS

ALGORITHM

EUA — testis now palpable? → inguinal/scrotal orchidopexy. **Still non-palpable → diagnostic laparoscopy** (umbilical 5-mm port; CO_2 8–12 mmHg) — inspect internal ring, vessels, vas:

• **Blind-ending spermatic vessels** (± vas) above the ring = vanished testis → stop (only unequivocal endpoint).

• **Vessels + vas entering a closed or open ring** → inguinal/scrotal exploration; excise the nubbin (viable tubules in 5–15%).

• **Intra-abdominal testis** → laparoscopic orchidopexy: primary (vessels long enough) or Fowler–Stephens (one or two stage); orchidectomy if postpubertal or dysgenetic.

• Peeping testis at ring → laparoscopic or inguinal orchidopexy.

• Neither vas nor vessels seen → search perivesical area and retroperitoneum to the kidney (agenesis extremely rare).

- Pitfalls: an open ring/hernia often (not always) means a viable distal testis; apparently blind-ending vessels may coexist with a distal or ectopic testis — if the view is equivocal, keep exploring (add ports).
- Palpable scrotal nubbin + contralateral hypertrophy (≥ 1.8 cm) → scrotal exploration first is acceptable.
- Contralateral fixation of a solitary testis: debated — some fix empirically; AUA algorithm does not require it.

Laparoscopic orchidopexy

Incise peritoneum lateral to vessels and distal to vas, mobilise vessels proximally **preserving the peritoneal strip between vas and vessels** (keeps Fowler–Stephens possible) → bring testis to scrotum via new hiatus medial to inferior epigastric vessels (Prentiss-like). Ring closure unnecessary. Success > 90–97% (primary).

Fowler–Stephens (FS)

- Principle: divide/clip the testicular vessels; testis survives on collaterals from the **deferential (vasal) artery** and cremasteric/gubernacular vessels. For testes > 2–4 cm from the internal ring with short vessels.
- **Single-stage**: clip and bring down at once. **Two-stage**: clip vessels high (1.5–3 cm above testis) → wait ≈ **6 months** for collaterals → mobilise on a wide peritoneal pedicle. Two-stage slightly favoured (better position, similar atrophy).
- Success: FS overall 60–97%; primary procedures more reliable — avoid vessel division when possible (it reduces germ cells, especially in young boys). Gubernacular sparing reduces atrophy.
- Alternatives: **Shehata traction** (testis fixed near contralateral anterior superior iliac spine; orchidopexy after 12 weeks; success 84–85%); **microvascular autotransplantation** (success ≈ 96%; for the solitary abdominal testis; needs microsurgical skill and time).
- Bilateral high testes: stage sides ≈ 6 months apart to avoid bilateral atrophy.

Approach	Success
Primary laparoscopic	> 90–97%
Primary open (Jones/Bagley transabdominal)	89–95%
One-stage / two-stage FS	63% / 68% (series range 60–97%)
Microvascular autotransplant	≈ 96%

10 · COMPLICATIONS AND FOLLOW-UP

Complication	Cause / prevention	Management
Atrophy (≈ 2% in modern series; atrophy/failure 24% for abdominal/peeping historically vs 10% distal)	Vessel injury, skeletonisation, FS, excess tension	No further surgery; offer **prosthesis** ≥ 6 months later or after puberty, inguinal route
Retraction / non-scrotal position (≈ 2–10%)	Inadequate retroperitoneal mobilisation, unligated processus	Redo orchidopexy: approach cord laterally through unscarred plane (Redman) or with external spermatic fascia patch (Cartwright); retroperitoneal mobilisation
Vas injury	Long-looping vas cut with gubernaculum; dissecting sac off vas	Divide gubernaculum distal to sac; avoid skeletonising
Torsion after orchidopexy	Rare	Prevented by extravaginal subdartos fixation
Laparoscopic	Bladder/vessel injury, hypercapnia, bowel obstruction	Open (Hasson) entry, bladder drainage

Follow-up: examine at ≥ **6 months** for position and size; long-term counselling on fertility and cancer; teach self-examination from puberty.

11 · MCQ ONE-LINERS

1. Cryptorchidism prevalence: 2–4% at term birth, 1–2% at 3 months.
2. Commonest ectopic site: superficial inguinal pouch.
3. Retractile testis stays in scrotum after cremaster fatigue; observe yearly.
4. INSL3 (Leydig cell) acts on gubernacular RXFP2 receptor.
5. Inguinoscrotal descent: 20–28 weeks, androgen via genitofemoral nerve and CGRP.
6. Gubernaculum attaches to the epididymis, not the testis.
7. Gonocyte → Ad spermatogonia transformation occurs during minipuberty (1–3 months).
8. Kollin RCT: orchidopexy at 9 months beats 3 years for testicular growth.
9. About 80% of UDTs are palpable.
10. Imaging (US/MRI) not recommended for UDT (AUA 2014).
11. Bilateral non-palpable testes in newborn: exclude CAH urgently.
12. Anorchia: high FSH/LH, undetectable AMH and INHB, no hCG response.
13. Orchidopexy timing: 6–18 months, ideally by 12 months.
14. Hormonal therapy for descent: not recommended; LHRH success ≈ 20%.
15. Laparoscopy blind-ending vessels: stop, vanished testis confirmed.
16. Vessels entering internal ring: inguinal exploration and nubbin excision.
17. Fowler–Stephens testis survives on deferential artery collaterals.
18. Second stage of Fowler–Stephens at about 6 months.
19. Prentiss manoeuvre: pass cord medial to inferior epigastric vessels.
20. Gastroschisis-associated UDT: observe; 50–55% descend.
21. Paternity after unilateral UDT ≈ controls; bilateral markedly reduced.
22. TGCT risk 2–5×; uncorrected UDT tumour is usually seminoma.
23. Orchidopexy after puberty: TGCT risk roughly double that of prepubertal surgery.
24. Postpubertal UDT with normal contralateral testis: orchidectomy.
25. Microlithiasis is not an independent TGCT risk; no routine US surveillance.
26. PMDS: AMH or AMHR2 mutation; transverse testicular ectopia.
27. Agenesis is associated with ipsilateral Müllerian duct persistence.

59 Differences of Sexual Development

Reading time ≈ 33 min (slow pace)

1 · DEFINITION AND NOMENCLATURE

- **Differences (disorders) of sex development (DSD)**: chromosomal, gonadal or anatomical sex not typically male or female. Examples: congenital adrenal hyperplasia (CAH), ovotesticular DSD (OVT DSD), androgen insensitivity syndrome (AIS).
- **2006 Chicago consensus** (Lee et al.) replaced "intersex", "hermaphrodite" and "sex reversal" with karyotype-based terms; "atypical" preferred to "ambiguous" genitalia.

Old term	2006 term
Male pseudohermaphrodite	46,XY DSD
Female pseudohermaphrodite	46,XX DSD
True hermaphrodite	Ovotesticular DSD
XX male / XX sex reversal	46,XX testicular DSD
XY sex reversal	46,XY complete gonadal dysgenesis (not AIS) MCQ

Offer endocrine and genetic testing for **proximal hypospadias even with descended testes** — a named diagnosis predicts gender identity data, fertility, tumour risk, comorbidities and recurrence.

2 · NORMAL SEX DEVELOPMENT

Three steps: **chromosomal sex** (fixed at fertilisation) → **gonadal sex** (bipotential gonad until ≈ 6 weeks) → **phenotypic sex** (bipotential ducts and external genitalia until ≈ 8 weeks).

- **SRY** (sex-determining region Y): earliest testis-determining transcription factor; upregulates **SOX9** → Sertoli differentiation.
- Genes act elsewhere too, explaining syndromic DSD: **WT1** (kidney — Wilms, nephropathy), **SF1** (NR5A1) and **DAX1** (adrenal — adrenal failure), SOX9 (bone — campomelic dysplasia). DAX1 duplication (Xp21) causes 46,XY dysgenesis.
- Ovary needs RSPO1/WNT4/β-catenin and FOXL2; ovarian germ cells enter meiosis in fetal life, testicular ones only at puberty.
- **Sertoli → anti-Müllerian hormone (AMH)** at 7–8 weeks → Müllerian regression by ≈ 10 weeks.
- **Leydig → testosterone** from ≈ 9 weeks (autonomous, then placental human chorionic gonadotrophin [hCG], then fetal luteinising hormone [LH]) → Wolffian ducts → epididymis, vas, seminal vesicles.
- **5α-reductase type 2 → dihydrotestosterone (DHT)**, higher receptor affinity; DHT is the active androgen in prostate, urogenital sinus and external genitalia. External virilisation complete by 12–13 weeks; penile growth and descent in third trimester (androgen, INSL3).
- Female: no AMH → tubes, uterus, upper vagina; distal one-third of vagina from urogenital sinus; no androgen → clitoris, labia.

> **KEY POINT** The testis uses two separate hormones: AMH removes Müllerian ducts, testosterone/DHT builds Wolffian and external structures. Any DSD phenotype can be reasoned from which hormone was missing, where and when — e.g. a uterus in 46,XY means Sertoli (AMH) failure, i.e. dysgenesis, not a synthesis or receptor defect.

3 · CLASSIFICATION (2006 / CHICAGO)

Category	Conditions
Sex chromosome DSD	47,XXY Klinefelter; 45,X Turner; 45,X/46,XY mixed gonadal dysgenesis; 46,XX/46,XY chimerism

Category	Conditions
46,XX DSD — gonadal	Gonadal dysgenesis; ovotesticular DSD; testicular DSD (SRY+, SOX9 duplication)
46,XX — androgen excess	Fetal: CAH (21-hydroxylase, 11β-hydroxylase, 3β-HSD, P450 oxidoreductase); fetoplacental: aromatase deficiency; maternal: luteoma, androgenic drugs
46,XX — other	Cloacal anomalies; Müllerian agenesis (Mayer–Rokitansky–Küster–Hauser, MRKH); vaginal atresia
46,XY DSD — gonadal	Complete/partial gonadal dysgenesis (SRY, SOX9, SF1, WT1, DHH, MAP3K1); ovotesticular DSD; testis regression
46,XY — androgen synthesis	LH receptor; Smith–Lemli–Opitz; StAR; 3β-HSD; 17α-hydroxylase; 17β-HSD3; 5α-reductase 2
46,XY — androgen action	Complete/partial AIS (CAIS, PAIS)
46,XY — other	Persistent Müllerian duct syndrome (PMDS); vanishing testis; isolated hypospadias; hypogonadotropic hypogonadism; cryptorchidism

3β-hydroxysteroid dehydrogenase (3β-HSD) and P450 oxidoreductase virilise 46,XX yet undervirilise 46,XY. 21-hydroxylase deficiency causes no genital difference in 46,XY.

4 · 46,XX DSD

Congenital adrenal hyperplasia — 21-hydroxylase deficiency

- CAH = commonest 46,XX DSD and **commonest cause of neonatal atypical genitalia**. 21-hydroxylase deficiency (21-OHD) = **95%** of CAH; autosomal recessive, ≈ 1 in 15,000.
- Mechanism: cortisol deficiency → ↑adrenocorticotropic hormone (ACTH) → adrenal androgen excess → virilisation; aldosterone deficiency → **salt wasting**.
- Infant presentation: variable virilisation, risk of adrenal crisis in the first weeks; infant death in the family suggests CAH.
- 46,XX: normal ovaries and Müllerian structures; clitoromegaly to fused labioscrotal folds, often a **common urogenital sinus**; may look like a boy with **bilateral impalpable gonads**. Graded by **Prader I–V** (increasing masculinisation; † V = male-appearing phallus with meatus at tip and fused empty scrotum).
- Diagnosis: ↑**17-hydroxyprogesterone (17-OHP)**, measured after **36 h** (physiological neonatal surge). Also ↑androstenedione, testosterone, renin; ↓cortisol; cosyntropin test.
- **Salt-wasting crisis**: hyponatraemia, hyperkalaemia, acidosis, hypotension, shock — usually not before **day 5–7**, so normal early electrolytes do not exclude CAH. Treatment: glucocorticoid + mineralocorticoid + salt — lifesaving.
- 46,XY boys have normal genitalia, so **newborn screening** matters most for them.
- Long-term: lifelong replacement balancing androgen suppression against steroid excess (growth, obesity, bone). 46,XY: testicular adrenal rest tumours (benign; ↑glucocorticoid; bank sperm).
- Prenatal dexamethasone must start by 6–7 weeks; ≥ 4 fetuses treated per girl helped — experimental (ethics-board-approved protocol only).

Form	Features
Classic salt-wasting	Cortisol + aldosterone deficient; crisis in first weeks
Simple virilising	Androgen excess, no infant shock
Nonclassic	20–50% enzyme activity; premature adrenarche, hirsutism, oligomenorrhoea; may need no treatment

Enzyme	Share	blood pressure (BP) / K^+	46,XY	Signature
21-hydroxylase	95%	Salt wasting; ↓BP, ↑K^+	Normal	↑17-OHP
11β-hydroxylase	≈ 5%	Hypertension, ↓K^+	Normal	—
3β-HSD	<1%	Salt wasting	Undervirilised	↑dehydroepiandrosterone (DHEA)
17α-hydroxylase	Rare	Hypertension, ↓K^+	Female/atypical; delayed puberty	↓androgens

Other 46,XX DSD

- **46,XX testicular DSD**: SRY translocated to X → testes, male genitalia, no Müllerian structures; azoospermia (no AZF region).
- Maternal androgens / aromatase deficiency: exposure ends at birth → no postnatal medical therapy.

5 · 46,XY DSD

Condition	Key features
Complete gonadal dysgenesis (Swyer)	Bilateral streaks, female genitalia, uterus present; absent puberty; tumour 30–45% → prophylactic gonadectomy; lifelong oestrogen; pregnancy with donor ovum
Partial gonadal dysgenesis	Dysgenetic testes, atypical genitalia, variable Müllerian structures; identity variable; WT1 → Denys–Drash (congenital nephrotic syndrome, Wilms 90%) / Frasier (late focal segmental glomerulosclerosis, gonadoblastoma 60%)
CAIS	X-linked receptor defect; female genitalia, short blind vagina, no uterus, inguinal/abdominal testes; breasts by aromatisation; ≈ 50% have inguinal hernia; 1–2% of girls with hernia are CAIS
PAIS	Atypical genitalia, residual receptor function; identity mostly male; pubertal breast growth
5α-reductase 2 deficiency	Autosomal recessive; atypical/female at birth, blind vaginal pouch; marked **virilisation at puberty**, fertility reported; testosterone:DHT **>20:1**; Dominican "guevedoces" (Imperato-McGinley 1979)
17β-HSD3 deficiency	↑androstenedione:testosterone; virilises at puberty
LH receptor defect	Leydig hypoplasia; no hCG response
PMDS	AMH or AMH-receptor mutation (autosomal recessive); normal male genitalia + uterus/tubes; found at hernia repair/orchidopexy; 30–50% of transverse testicular ectopia
Vanishing testes	Undetectable testosterone and AMH, ↑gonadotropins at mini-puberty; testosterone from adolescence

- **Synthesis/action defects**: AMH normal → no uterus; adrenal failure only if enzyme shared with cortisol pathway.
- **PAIS management**: raised male — high-dose testosterone may improve genital development; female identity — gonadotropin-releasing hormone agonist or gonadectomy, then oestrogen. CAIS: vaginal dilation usually suffices; vaginoplasty rarely needed.
- **CAIS hernia**: in a girl at herniorrhaphy, vaginoscopy for cervix or look for a testis in the sac.
- **PMDS (Clarnette)**: 1 bilateral intra-abdominal (60–70%); 2 one testis in hernia sac/scrotum + contralateral hernia (20–30%, "hernia uteri inguinale"); 3 transverse testicular ectopia (10%). Orchidopexy; vasa hug the uterus, so remnants traditionally kept; remnant malignancy 3–8% → if removing, laparoscopic/robotic to spare vasa.

6 · SEX CHROMOSOME DSD AND OVOTESTICULAR DSD

Feature	Klinefelter (47,XXY)	Turner (45,X and variants)
Frequency	1 in 600 males; commonest aneuploidy	25–50 per 100,000 females
Phenotype	Tall; small firm testes <3.5 cm; gynaecomastia; neurodevelopmental issues	Short; female genitalia; absent puberty; webbed neck; coarctation, bicuspid valve; horseshoe kidney (renal anomaly ⅓–½)
Hormones	↑gonadotropins, ↓testosterone	Ovarian insufficiency ≈ 95%
Treatment	Testosterone; sperm extraction + intracytoplasmic sperm injection	Growth hormone; oestrogen + progesterone; oocyte donation
Tumour	Extragonadal germ cell; male breast	None without Y; **Turner + Y** 12–20% (gonadoblastoma → dysgerminoma); 2017 guideline: gonadectomy at diagnosis (low-level evidence, now challenged)

- **Mixed gonadal dysgenesis (MGD)**, usually 45,X/46,XY: dysgenetic testis + contralateral streak + Müllerian structures. Classic **asymmetry**: one palpable testis, impalpable side with flat labioscrotal fold, hemiuterus/tube on streak side (local AMH action). **Second commonest cause** of newborn atypical genitalia. Germ cell neoplasia 15–35% → remove streaks. All 45,X/46,XY need Turner screening (cardiac, renal, growth). ≈ 85% of 45,X/46,XY have male-typical genitalia. Remaining testis: raised male — keep descended testis, bring down intra-abdominal one; raised female — remove or relocate to groin/labioscrotal fold for surveillance. Gender dysphoria in a substantial minority raised female.
- **Ovotesticular DSD**: ovarian + testicular tissue (often ovotestis); mostly 46,XX; urogenital sinus and uterus usual. A vas always accompanies a testis, a tube an ovary. Ovary usually normal, testis dysgenetic; tumour ≈ 3%; polar biopsy for bipolar ovotestis.

7 · EVALUATION OF THE NEWBORN WITH ATYPICAL GENITALIA

> **KEY POINT** **Emergency**: atypical genitalia with bilateral impalpable gonads = salt-wasting CAH until proven otherwise. Check electrolytes and 17-OHP; watch for crisis from the end of week 1. Everything else (designation, surgery) is not an emergency.

- Multidisciplinary review within **24–48 h**; congratulate, say "your baby", never "it"; gender-neutral terms. A urologist without a team should transfer early.
- History: infertility, amenorrhoea, unexplained infant death (CAH), maternal steroids/progestins.
- Examine: stretched phallic length, glans width, orifices, labioscrotal fusion and rugation, anogenital distance, **gonads**.
- **Gonads are the most informative sign**: ovaries do not descend, so a palpable gonad = testicular tissue (excludes typical CAH). 0 palpable → CAH; 1 → MGD, OVT DSD, PAIS, partial dysgenesis; 2 → 46,XY DSD.
- Hypospadias + undescended testis without atypical genitalia (Kaefer 1999): DSD ≈ 30%; 50% if testis impalpable; 65% if proximal meatus.

ALGORITHM

Karyotype (≥45 cells; fluorescence in situ hybridisation for X/Y in hours) + electrolytes + 17-OHP at ≈ 36 h → pelvic ultrasound (US) for uterus (best neonatally) and gonads → at ≈ 1 week: LH, follicle-stimulating hormone (FSH), testosterone, oestrogen, androstenedione, DHT, AMH ± hCG stimulation → genetic panel / microarray / exome → if unclear: magnetic resonance imaging (MRI), genitography, endoscopy, laparoscopy ± gonadal biopsy (**no gonadectomy during diagnosis**)

Test	Interpretation
17-OHP	Raised in CAH (21-OHD)

Test	Interpretation
AMH	Normal → functioning Sertoli (or granulosa) cells
LH, FSH	Raised → gonadal failure
Androstenedione / DHEA sulfate	21-OHD, 17β-HSD3 / 21-OHD, 3β-HSD
Testosterone:DHT	>20:1 → 5α-reductase deficiency
hCG stimulation	No testosterone rise → no testis, Leydig hypoplasia or LH receptor defect

Karyotype	Findings	Likely
46,XX	Impalpable gonads, uterus, ↑17-OHP	CAH
46,XX	Gonads not two ovaries	OVT DSD
46,XY	Uterus ±	Partial dysgenesis, OVT DSD
46,XY	No uterus	Androgen synthesis/action defect
45,X/46,XY	Asymmetric gonads, uterus ±	MGD

Precise diagnosis is nearly always reached in 46,XX DSD but in **<50%** of 46,XY DSD. Late presenters (typical external genitalia): CAIS, Swyer, MRKH, Turner + Y, Klinefelter, 5α-reductase deficiency; basal hormones are informative after puberty.

8 · GENDER ASSIGNMENT AND THE MULTIDISCIPLINARY TEAM

- Team: endocrinologist, urologist/surgeon, gynaecologist, geneticist, psychologist/psychiatrist, social worker, ethicist, nurse, peer support. **Same-day joint clinic**; lifelong care with planned transition.
- Historical **Money (1975)** model — early designation, early surgery, secrecy — failed; replaced by transparency and shared decision-making.
- Designation weighs diagnosis-specific identity data, prenatal androgen exposure, sexual function, fertility, hormonal potential and family/culture. Sex of rearing predicts satisfaction better than androgen exposure; no test predicts identity.
- Female trend: 46,XX CAH, CAIS, Swyer, LH receptor defect. Male trend: PAIS, 5α-reductase and 17β-HSD3 deficiency, partial dysgenesis.
- Gender dysphoria highest in 5α-reductase deficiency raised female (≈ half); lowest in 46,XX CAH raised female.
- Micropenis (stretched length <2.5 SD; <2 cm at 0–5 months) → endocrine work-up; female reassignment no longer recommended.

9 · SURGERY

- **Feminising genitoplasty**: clitoroplasty, vaginoplasty/urogenital sinus mobilisation, labioplasty → *Ch 60*. Clitoral surgery is irreversible and can be catastrophic if identity later differs; benefits of infant surgery unproven. Only urgent indication: urinary retention/obstruction.
- **Masculinising**: mainly proximal hypospadias repair ± penoscrotal transposition, orchidopexy; high complication rates; largely revisable, so infant repair often chosen when raised male.
- **Timing debate**: ideally defer irreversible surgery until the individual can participate; parents often prefer early surgery. CAH surgery is debated separately.
- MRKH: dilation first-line; Vecchietti; buccal or bowel vaginoplasty; uterus transplantation investigational.

10 · GONADAL TUMOUR RISK AND GONADECTOMY

Risk rises with Y material (gonadoblastoma locus, TSPY), dysgenesis and intra-abdominal position; falls with a scrotal gonad. Shift from automatic gonadectomy to individualised risk, function and autonomy (biopsy, relocation, tissue cryopreservation).

Risk	Condition	Germ cell tumour	Management
High	Gonadal dysgenesis + Y, intra-abdominal (incl. Swyer, MGD)	15–35% (Swyer 30–45%)	Gonadectomy at diagnosis
	Frasier	60%	Gonadectomy
	Denys–Drash	40%	Gonadectomy
	PAIS, non-scrotal gonad	50%	Gonadectomy
Intermediate	Turner + Y	12%	Gonadectomy (now debated)
	17β-HSD3 deficiency	28%	Monitor or gonadectomy
	PAIS / dysgenesis, scrotal gonad	Unknown	Pubertal biopsy
Low	CAIS	2%	Keep until after breast development
	Ovotesticular DSD	3%	Remove testicular tissue if discordant
?None	5α-reductase deficiency; Leydig hypoplasia	Unknown	Unresolved

11 · ETHICS

- Principles: bodily autonomy, transparency about known and unknown, avoiding irreversible procedures before consent unless medically necessary, informed parental consent covering limits of predicting identity and fertility.
- Psychosocial morbidity (anxiety, depression, suicidality) is higher → integral psychology input; patient preference for terms (often the diagnosis itself) respected.

12 · MCQ ONE-LINERS

1. "True hermaphrodite" is now ovotesticular DSD.
2. "XY sex reversal" = 46,XY complete gonadal dysgenesis.
3. SRY upregulates SOX9 to drive Sertoli differentiation.
4. Sertoli cells secrete AMH from 7–8 weeks.
5. DHT is the active androgen for external genitalia and prostate.
6. Commonest cause of neonatal atypical genitalia: CAH (21-hydroxylase).
7. Second commonest: mixed gonadal dysgenesis (45,X/46,XY).
8. Measure 17-OHP after 36 hours of age.
9. Salt-wasting crisis usually occurs after day 5–7.
10. 11β-hydroxylase deficiency causes hypertension and hypokalaemia.
11. 17α-hydroxylase deficiency: hypokalaemia, not hyperkalaemia.
12. 3β-HSD deficiency undervirilises 46,XY and virilises 46,XX.
13. Uterus in 46,XY points to gonadal dysgenesis.
14. Testosterone:DHT >20:1 suggests 5α-reductase deficiency.
15. Guevedoces: 5α-reductase deficiency virilising at puberty.
16. CAIS gonads can stay until after breast development (risk 2%).
17. 1–2% of girls with inguinal hernia have CAIS.
18. Frasier syndrome: gonadoblastoma 60%, no Wilms predisposition.
19. Denys–Drash: Wilms 90%, congenital nephrotic syndrome.
20. PMDS: AMH or AMH receptor mutation; hernia uteri inguinale.
21. PMDS underlies 30–50% of transverse testicular ectopia.
22. 46,XX testicular DSD: SRY translocation; azoospermic.
23. Palpable gonad = testicular tissue; ovaries do not descend.
24. Klinefelter: commonest sex chromosome aneuploidy, 1 in 600 males.
25. All 45,X/46,XY patients need Turner screening.
26. Vas always accompanies a testis in ovotesticular DSD.
27. Precise diagnosis reached in <50% of 46,XY DSD.
28. No gonadectomy during the diagnostic phase.
29. Micropenis: stretched length <2.5 SD below mean.
30. Highest dysphoria: 5α-reductase deficiency raised female.

61 Urologic Considerations in Pediatric Renal Transplantation

Reading time ≈ 18 min (slow pace)

1 · CAUSES OF END-STAGE KIDNEY DISEASE (ESKD) IN CHILDREN

- Children needing kidney replacement therapy (KRT) more often have **congenital anomalies of the kidney and urinary tract (CAKUT)**: hypoplasia/aplasia/dysplasia, then obstructive uropathy; acquired: focal segmental glomerulosclerosis. Commonest age at transplant **13–17 years**.
- Younger: dysplasia or obstructive uropathy — mainly **posterior urethral valves (PUV)** in boys. Older: nephrological causes, neuropathic bladder.
- The urologist's task: the condition that destroyed the native kidneys must not destroy the graft. Join the team **before ESKD**, not reactively.

> **KEY POINT** **Five goals**: unimpeded drainage · capacious low-pressure reservoir with social continence · volitional emptying · no infection · fewest operations. Chronic rejection is unavoidable, but every added insult accelerates it.

2 · PRE-TRANSPLANT UROLOGICAL EVALUATION

- **Basic screen (all)**: history + renal ultrasound (US) + uroflow with post-void residual (PVR) if toilet-trained.
- **Voiding cystourethrogram (VCUG)** not routine; do it with neuropathic bladder, obstructive uropathy, vesicoureteral reflux (VUR), prior lower-tract surgery, recurrent febrile urinary tract infection (UTI), worsening hydroureteronephrosis or refractory voiding problems.
- **Urodynamics**: neuropathic bladder, PUV, large ureterocele, refractory dysfunction, high-grade hydronephrosis, recurrent UTI. Measures capacity, compliance, overactivity, emptying, sphincter; **repeat serially**.
- Anuric child with no urological history: no further testing.

ALGORITHM

Positive history → US + VCUG → abnormal dynamics → urodynamics → drugs + clean intermittent catheterisation (CIC) ± bladder cycling → reassess → still unsafe → augmentation → clear for transplant

3 · VALVE BLADDER AND BLADDER MANAGEMENT

- Commonest ESKD pattern: **low capacity, detrusor overactivity, poor compliance** — dangerous because high pressure reaches the kidney even without reflux.
- Valve bladder evolves: overactive small → larger, less contractile → **myogenic failure** (poor emptying).
- **Target**: expected capacity for age at detrusor pressure **< 40 cm H_2O**, socially continent.
- Medical first: oxybutynin (titrate), solifenacin/darifenacin, mirabegron, intradetrusor botulinum toxin; α-blockers aid emptying. Better storage can worsen emptying → CIC. Prove emptying with repeated low PVRs.
- Adherence to pre-transplant CIC tests adherence to the transplant regimen.

Defunctionalised bladder

Does not cycle (anuria or diversion), so response to post-transplant volumes is unpredictable.
Bladder cycling: instil saline in rising volumes via catheter with set dwell → add medication → target expected capacity → confirm by urodynamics → clear, or augment. **A renal allograft must never be used to test whether a bladder works.**

Augmentation

- Indication: cannot reach > **75%** expected capacity at < 40 cm H_2O with CIC no more often than every **3–4 h**, despite maximal medical therapy. No compelling evidence it increases transplant complications.
- PUV bladder → **postpone** (heads to myogenic failure); neuropathic bladder → **earlier** reconstruction (deteriorates under high graft output).

Segment	Point
Dilated ureter (ureterocystoplasty)	Urothelial, no mucus/stones/metabolic problems — always explore first
Ileum, sigmoid	Most used; mucus, stones, infection, malignancy (worse with immunosuppression) → lifelong surveillance
Stomach	Haematuria–dysuria syndrome, especially in anuric patients; rarely used

- **Mitrofanoff/continent catheterisable stoma**: decided by spontaneous emptying; sensate or difficult urethra favours a stoma. Site it **medial to the planned transplant incision**; preserve native ureter for stoma, augment or graft drainage.
- **Timing**: major reconstruction **before** transplant, allowing **6–12 weeks** healing. Exception: transplant before 2 years of age, before continence is expected.
- Peritoneal dialysis (PD): intraperitoneal surgery forces temporary haemodialysis — protect the peritoneum. Avoid concurrent procedures in infants.
- Long-term stoma or transplant ureterostomy only for severe neurodevelopmental impairment or CIC intolerance.
- Circumcision in boys at risk (especially valves) — discuss early; reduces post-transplant UTI.

4 · NATIVE NEPHRECTOMY

Remove (named indication)	Keep (default)
Refractory hypertension; nephrotic syndrome with malnutrition/hypercoagulability; massive polyuria; recurrent upper-tract infection; stones; malignancy risk; very large kidneys; massive VUR (relative)	Water excretion (less fluid restriction); erythropoietin and vitamin D; avoids morbidity; graft usually fits

- On PD: retroperitoneal (bilateral prone) so PD restarts at once. High risk: embolisation.
- Normal ureter: transect high, keep as reserve for salvage ureteroureterostomy. Dilated refluxing/obstructed ureter: remove totally (infection).
- Combined with transplant only for room (small child, huge kidneys) — ideally ipsilateral via transplant incision.

5 · TRANSPLANT INTO THE ABNORMAL BLADDER

- Graft retroperitoneal via Gibson incision; child < **15 kg** may need aorta/vena cava anastomosis and transperitoneal approach. Know augment pedicle position.
- **Extravesical non-refluxing ureteroneocystostomy** (preferred): partially fill bladder → 3–4 cm detrusorotomy → mucosal disc → spatulate 4–5 mm → interrupted mucosa-to-mucosa → anchor stitches → close flaps loosely at entry.
- Alternatives: Barry tunnel; **ureteroureterostomy** to native ureter (small bladder, short donor ureter, hostile pelvis; always the salvage option).
- Stent if difficult reimplant or poor ureter: short double-J out at **4–6 weeks**; early removal at 5–7 days may cut infection. Use the largest urethral catheter tolerated.

6 · UROLOGICAL COMPLICATIONS

Complication	Rate	Timing / management
Ureteric stenosis	**6%**	Most within 100 days; usually distal (ischaemic donor ureter); focal → balloon + stent 4–6 weeks; otherwise open repair with native ureter or bladder flap — do not delay
Urine leak	**3%**	Early; drain-fluid creatinine; check catheter → US → mercaptoacetyltriglycine (MAG3) scan or cystogram → small: observe; large: drain/explore; short necrosis → reimplant; long → native ureter, psoas hitch, flap
Significant VUR	2%	See below
Stones	2%	Treat if > 1 cm and growing, obstructing, infected; one-stage modality

MAG3 localises a leak only if graft function is adequate. Pre-existing obstructive or reflux uropathy raises all complications — lower threshold to intervene.

Rising creatinine with hydronephrosis

ALGORITHM
Indwelling catheter 1–2 weeks → creatinine falls = bladder cause → if not, stent + catheter; signs of rejection → biopsy; failed retrograde stent → nephrostomy + antegrade study

- Graft is unusually sensitive to obstruction; grade of hydronephrosis does not track impairment. Obstruction and rejection can coexist — low threshold for diagnostic stenting.
- Extrinsic causes: **lymphocele**; **post-transplant lymphoproliferative disease (PTLD)** (allograft mass = PTLD until proven otherwise).
- Bladder dysfunction: CIC, medication; augmentation rarely and late.

UTI and graft VUR

- Early UTI: catheter/stent; late UTI reflects bladder function — check emptying, hydronephrosis, selective VCUG. Distinguish colonisation (universal with CIC) from infection.
- Transplant VUR costs more than native VUR (solitary immunosuppressed kidney). Febrile UTI → cystogram.
- Recurrent febrile UTI + optimised bladder + graft VUR → **open/robotic reimplant** (intravesical, transtrigonal or Politano–Leadbetter; stent). Asymptomatic reflux → observe. Endoscopic injection: ≈ one-third success, obstruction risk.

7 · OUTCOMES

- Urological complications are uncommon and rarely lose the graft; thrombosis is a leading cause of early graft loss.
- **≥ 2 UTIs** → lower graft glomerular filtration rate at 1 and 4 years, gap widening with time.
- PUV history → lower obstruction-free survival; link with graft dysfunction inconsistent.

8 · MCQ ONE-LINERS

1. Commonest cause of paediatric ESKD: CAKUT (hypoplasia/dysplasia), then obstructive uropathy.
2. Commonest obstructive cause in boys: posterior urethral valves.
3. Routine pre-transplant VCUG: not required.
4. Safe storage pressure target: < 40 cm H_2O.
5. Augment if < 75% expected capacity despite CIC every 3–4 h and drugs.
6. Valve bladder end-stage: myogenic failure.
7. PUV bladder: postpone augmentation; neuropathic: earlier.
8. Gastrocystoplasty problem: haematuria–dysuria syndrome.
9. Preferred augment material if available: dilated ureter.
10. Never use the allograft to test a bladder.
11. Reconstruction heals 6–12 weeks before transplant.
12. Catheterisable stoma sited medial to transplant incision.
13. Native kidneys: keep by default.
14. Child < 15 kg: graft to aorta/vena cava.

5. Preferred ureteric anastomosis: extravesical non-refluxing ureteroneocystostomy.
6. Salvage for graft ureteric stenosis: ureteroureterostomy to native ureter.
7. Commonest urological complication: ureteric stenosis (≈ 6%).
8. Most obstructions occur within 100 days.
19. Leak fluid identified by high creatinine.
20. First step in creatinine rise + hydronephrosis: indwelling catheter.
21. Allograft mass: PTLD until proven otherwise.
22. Endoscopic injection for graft VUR: ≈ one-third success.
23. Asymptomatic graft VUR: observe.

63 Paediatric Urologic Oncology: Renal and Adrenal

Reading time ≈ 33 min (slow pace)

1 · WILMS TUMOUR (WT): EPIDEMIOLOGY AND GENETICS

- Nephroblastoma: **6–7%** of childhood cancers, **95%** of kidney cancers under 15 y; embryonal tumour from immature renal remnants. Median age **3.5 y**, >80% before 5 y; younger in bilateral and syndromic disease.
- Mostly somatic and multi-hit (Knudson two-hit does not explain most). **10–15%** germline/early epigenetic (11p15); familial 1–2%.
- **Refer for genetics**: bilateral/multifocal, age <2 y, nephrogenic rests, overgrowth/syndromic features, family history.

Gene (locus)	Frequency	Note
IGF2 (11p15) — putative "WT2" locus	70%	Loss of imprinting/LOH (loss of heterozygosity); Beckwith–Wiedemann syndrome
WT1 (11p13)	20%	First WT gene (WAGR deletions); germline or somatic
WTX (Xq11), CTNNB1 (β-catenin)	20%, 15%	Somatic only
TP53 (17p13)	4%	Majority of **anaplastic** tumours

	WAGR	Denys–Drash (DDS)
Features	WT, aniridia, genitourinary anomalies, intellectual disability	46,XY disorder of sex development (DSD) + mesangial sclerosis + WT
WT1 lesion	Deletion (one allele lost)	Zinc-finger point mutation, **dominant-negative** → worse phenotype
Kidney	Renal-failure risk into puberty	Early nephropathy → end-stage renal disease (ESRD)

- **Ideal type I** (WT1 ± CTNNB1): stromal-predominant, intralobar rests, early age, WAGR/DDS. **Type II** (11p15 epigenetic): blastemal/epithelial, perilobar rests, later age, BWS.
- **Beckwith–Wiedemann syndrome (BWS)**/hemihypertrophy: WT risk 4–10%, 21% bilateral; also hepatoblastoma, adrenocortical tumours. Nephromegaly (≥95th centile) = highest risk.
- Prognostic markers (favourable histology): **1p + 16q LOH** (~5%) adverse at all stages → intensified **regimen M**; 1q gain (25–30%) = next stratifier.

Syndrome risk	Examples (WT risk)
High (>20%)	WAGR, DDS (**50%**); Fanconi D1 (BRCA2); Perlman (DIS3L2); mosaic variegated aneuploidy
Moderate (5–10%)	BWS, isolated hemihypertrophy, Frasier (WT1), Simpson–Golabi–Behmel (GPC3)
Low	Li–Fraumeni, neurofibromatosis 1, Sotos, trisomy 18, Bloom

> **KEY POINT** Trap: BWS is the **commonest** predisposing syndrome but only moderate risk; WAGR/DDS are rarer and highest.

Screening protocol

> **ALGORITHM**
> WT risk >1% → renal ultrasound (US) every **3–4 months** until **7 y** → suspicious lesion → computed tomography (CT)/magnetic resonance imaging (MRI)

- Interval does not lengthen with age (growth rate assumed constant).
  BWS/hemihypertrophy/trisomy 18/SGB: abdominal US + α-fetoprotein (AFP) 3-monthly to 4 y (hepatoblastoma).

- Contralateral kidney after nephrectomy: US 3–4-monthly to 5 y (highest risk: infants <12 months with rests).
- Benefit = lower stage → **nephron-sparing surgery (NSS)**, not proven survival. Benign cysts are common in BWS — avoid needless nephrectomy.

2 · PATHOLOGY

- **Favourable histology**: classic **triphasic** — blastema + epithelium (tubules, glomeruloid bodies) + stroma; pseudocapsule; soft, friable → spill risk. 12% multicentric unilateral.
- **Anaplasia** (3 criteria): nuclei ≥**3×** neighbours, hyperchromasia, abnormal mitoses. ~2% under 2 y, ~13% over 5 y. Chemoresistant; **diffuse** worse than focal. Acquired (later age, absent from rests); TP53-driven.
- International Society of Paediatric Oncology (SIOP) post-chemotherapy risk: **low** = completely necrotic; **high** = diffuse anaplasia, **blastemal-predominant**; intermediate = rest. Blastema shrinks but predicts relapse if it survives; differentiated stroma/epithelium barely shrinks yet is cured by excision.

Nephrogenic rests (NRs)

Present in >1/3 of WT kidneys and **>90%** of bilateral WT; most regress. Nephroblastomatosis = multiple rests; diffuse hyperplastic perilobar nephroblastomatosis gives a thick rind, almost always bilateral; ~half develop WT.

	Perilobar	**Intralobar**
Site/timing	Lobar periphery; late embryogenesis	Anywhere in lobe, sinus; early
Syndrome	BWS (11p15)	WAGR, DDS (WT1)

3 · CLINICAL FEATURES AND WORK-UP

- >85% symptomatic: mass, pain; haematuria ~20% (gross → cystoscopy + retrograde pyelogram at surgery for collecting-system extension); hypertension ~25% (renin). Examine for aniridia, hemihypertrophy, genital anomalies.
- Varicocele persisting supine or hepatomegaly → suspect inferior vena cava (IVC) thrombus.
- **Acquired von Willebrand disease** up to 8% — clotting times may be normal; correct with desmopressin (DDAVP).
- Imaging: US → contrast CT or MRI abdomen–pelvis → **chest CT** (lung = commonest metastatic site; do before MRI anaesthesia to avoid atelectasis). No contralateral exploration if imaging normal. Rupture: ascites beyond cul-de-sac is best predictor. Nodes cannot be judged on imaging.
- Imaging cannot separate WT from renal cell carcinoma (RCC) or clear cell sarcoma (CCSK). Clear cell sarcoma and rhabdoid tumour → add bone scan + brain imaging.
- Emergency surgery only for active bleeding/rupture.

4 · STAGING AND COG VS SIOP

Children's Oncology Group (COG) stage	**Definition (surgical–pathological)**
I	Confined, completely resected; capsule intact; no rupture; no renal sinus extension
II	Extracapsular but completely resected; renal sinus or extrarenal vessel involvement
III	Residual abdominal disease: **positive nodes, any spill**, peritoneal implants, positive margin, piecemeal/incomplete removal
IV	Haematogenous metastases (lung, liver, bone, brain)
V	Bilateral at diagnosis

	COG (North America)	SIOP (Europe)
Unilateral	Upfront nephrectomy	Pre-op vincristine (VCR) + dactinomycin (AMD) ~6 weeks, no biopsy
Advantage	Untreated histology and biology (LOH) for risk; very-low-risk cured by surgery alone	Smaller, firmer tumour; fewer complications; spill near zero
Cost	More spill/complications	Loses pre-treatment stage/biology; a minority not WT
Risk basis	Stage + histology + biology	Post-chemotherapy histology
Follow-up	CT-based	US + chest radiograph

KEY POINT Because COG stage is surgical-pathological, the operating surgeon — not the scan — decides chemotherapy intensity and irradiation: one spill converts stage I/II to III.

5 · SURGICAL PRINCIPLES

- **Transperitoneal radical nephrectomy**; inspect liver, peritoneum, nodes. Palpate renal vein/IVC before ligation (intra-op US if equivocal). Spare adrenal if uninvolved.
- **Lymph node sampling mandatory** (formal dissection not needed): hilar, paracaval/para-aortic, interaortocaval (skip metastases). COG requires **≥5 nodes**. Omission is the commonest protocol violation (10–20%) → under-staging.
- **Avoid spill** (~10%; right side, large tumours) — causes local abdominal relapse. Local recurrence risks: spill, unfavourable histology, incomplete removal, no node sampling.
- Complications ~11% (haemorrhage, small-bowel obstruction); higher with >10 cm, caval extension, thoracoabdominal incision, wrong diagnosis.

Chemotherapy

- Backbone VCR + AMD; doxorubicin (DOX, cardiotoxic) for higher stage; cyclophosphamide + etoposide in regimen M and CCSK; carboplatin for anaplastic/relapsed.
- † COG favourable histology: stage I–II → VCR + AMD; stage III–IV → VCR + AMD + DOX with flank/abdominal (and lung) radiotherapy.
- Very-low-risk stage I (AREN0532) → nephrectomy alone.
- CT-only lung nodules: AREN0533 — complete response at week 6 without 1p/16q LOH → lung irradiation omitted safely.

Bilateral WT (stage V) and NSS

ALGORITHM
Synchronous bilateral (5–7%) → **no upfront nephrectomy, no biopsy** → VCR + AMD + DOX 6 weeks → imaging → NSS if feasible → else more chemo but **surgery by week 12** → non-responder: **bilateral open biopsy**

- Why 12 weeks: in National Wilms Tumor Study (NWTS)-4 prolonged chemo delayed surgery to ~7 months and some tumours became anaplastic.
- Open (not needle) biopsy detects anaplasia better; bilateral because anaplasia is discordant between kidneys in ~4/5.
- AREN0534: 4-year event-free survival (EFS) 82%, overall survival (OS) 95%. Commonest cause of renal failure = bilateral nephrectomy for persistent tumour. Transplant after 1–2 y tumour-free.
- Unilateral NSS indications: syndromic/predisposed or solitary kidney, bilateral risk; SIOP UMBRELLA criteria: **<300 mL**, substantial remnant, negative nodes. Use frozen section and a margin of normal kidney.

IVC thrombus

- ~6–10% (renal vein/IVC/atrium); more than half asymptomatic.

- Below hepatic veins → COG upfront nephrectomy with **en bloc** thrombectomy (piecemeal = stage III → radiotherapy). **At/above hepatic veins → pre-op chemotherapy** to avoid cardiopulmonary bypass. Chemo beyond 6–12 weeks adds nothing. Viable cells in thrombus do not affect survival.

6 · RELAPSE AND LATE EFFECTS

Relapse group	Definition	Salvage
Standard	Non-anaplastic, prior VCR/AMD only	70–80%
High	Non-anaplastic, after ≥3 drugs	40–50%
Very high	Anaplastic/blastemal	~10%

- Late effects track **radiotherapy and anthracycline**: heart failure 4.4% after DOX (rare below **150 mg/m²**); second cancers 1% at 10 y (in-field); gonadal failure (alkylators, pelvic/whole-abdominal RT). Renal failure after unilateral nephrectomy only **0.6%** at 20 y — higher in WT1 syndromes.

7 · OTHER RENAL TUMOURS

Tumour	Key facts	Management
Congenital mesoblastic nephroma (CMN)	Commonest **neonatal/antenatal** renal tumour (mean 3½ months); ETV6–NTRK3 fusion (cellular type, shared with congenital fibrosarcoma); classic vs cellular	Radical nephrectomy alone; cellular type can recur — adjuvant only if incompletely resected
Clear cell sarcoma (CCSK)	3%; BCOR duplication; **bone**/brain metastasis; late relapse (up to 10 y); never bilateral	DOX-containing chemo; radiotherapy even at stage I
Rhabdoid tumour (RTK)	2%; **most lethal**; median ~16 months; SMARCB1/INI1 loss (22q), INI1 stain negative; brain tumours/metastases	No standard; EZH2 inhibitor (tazemetostat) under trial
Cystic nephroma / cystic partially differentiated nephroblastoma (CPDN)	Radiologically identical; CPDN has blastema in septa (first 2 y); DICER1	Excision cures (100%); frozen section if partial nephrectomy
RCC	Commonest renal malignancy of 2nd decade (RCC > WT above 13–14 y); **translocation RCC (TFE3/TFEB, Xp11.2)** commonest; renal medullary carcinoma — sickle trait, lethal	Complete resection; nodes less ominous than adults; chemo/radioresistant
Angiomyolipoma	Tuberous sclerosis (50–80%); bleeding risk >3 cm and growing	mTOR inhibitor first-line; embolise for bleeding

> **KEY POINT** Age points to diagnosis: neonate → CMN; toddler → WT/rhabdoid; adolescent → RCC. Metastatic site sets staging tests: lung (WT), bone (CCSK), brain (rhabdoid/CCSK).

8 · NEUROBLASTOMA

- Commonest **extracranial solid tumour** of childhood and commonest malignancy of infancy; neural-crest origin (adrenal medulla, sympathetic chain); median 19 months. 50% adrenal; metastases in ~70% at diagnosis.
- Familial <1% (PHOX2B, ALK). Behaviour ranges from regression to lethal disease.
- Spectrum: neuroblastoma → ganglioneuroblastoma → **ganglioneuroma** (benign; operate only for symptoms). Shimada classification: stroma-poor vs stroma-rich, graded against age.
- Presentation: hard fixed mass, bone pain, periorbital ecchymosis; cord compression; pelvic (organ of Zuckerkandl) → retention. Paraneoplastic: catecholamine excess, VIP (vasoactive intestinal peptide) → watery diarrhoea, **opsoclonus–myoclonus–ataxia**.

Diagnosis

- Urinary **VMA** (vanillylmandelic acid) and **HVA** (homovanillic acid) raised in **90–95%**; used to monitor response/relapse.
- Marrow: two aspirates + two biopsies. CT/MRI (MRI best for intraspinal extension). Calcification and vessel **encasement** favour neuroblastoma over WT.
- **^{123}I-MIBG** (meta-iodobenzylguanidine) scan required for staging (replaces bone scan); non-avid → positron emission tomography (PET). ^{131}I-MIBG is therapeutic.
- Screening (Japan, Quebec, Germany) increased favourable infant tumours without reducing advanced disease or mortality — classic **overdiagnosis**.

Staging and risk

International Neuroblastoma Risk Group (INRG) stage	Definition
L1	Localised, **no** image-defined risk factor (IDRF)
L2	Localised, ≥1 IDRF (mainly vessel/nerve encasement)
M	Distant metastases
MS	Age **<18 months**; metastases to skin, liver, marrow (<10%) only; no bone (old INSS 4S: <12 months)

- INRG is imaging-based, pre-treatment; INSS (International Neuroblastoma Staging System, stages 1–4S) depends on resection and nodes, so surgeon-dependent.
- Risk groups (very low/low/intermediate/high) combine stage, age, histology, **MYCN**, 11q, ploidy. **MYCN amplification** (>10 copies, 2p24; ~20%) = high risk at any stage (except maturing histology); M at ≥18 months always high risk; MS with 11q aberration = high risk.
- Other adverse: 1p36 deletion, unbalanced 11q LOH, 17q gain (strongest on multivariate), **diploidy**. Favourable: **hyperdiploidy**, TrkA expression, age <18 months, thoracic site.

> **KEY POINT** Trap: in neuroblastoma diploid is bad and hyperdiploid good. Stage 4S/MS: OS 80–88% with frequent spontaneous regression; deaths occur in infants <2 months from massive hepatomegaly with respiratory compromise or coagulopathy.

Management by risk

- **Low risk (stage 1, 2, 4S)**: resection alone (stage 1 >90% disease-free); chemo only for symptoms, progression or adverse biology. COG P9641: surgery alone, 5-y OS ~97%. Never remove kidney or vital structures; RT to residual stage 2 does not help.
- **4S**: resection of primary not mandatory (equal survival after biopsy); favourable and asymptomatic → supportive care ± limited chemo.
- **Intermediate**: chemo then resection; pre-op chemo for stage 3/peri-hilar tumours reduces nephrectomy.
- **High risk**: induction (cyclophosphamide, vincristine, doxorubicin alternating with cisplatin–etoposide) → **delayed surgery** after 3–4 courses → myeloablative consolidation with autologous stem-cell rescue (tandem better) → radiotherapy (15–30 Gy) → **13-cis-retinoic acid + anti-GD2 antibody** (with granulocyte-macrophage colony-stimulating factor and interleukin-2) for minimal residual disease.
- Surgical role in stage 4 is debated: no survival benefit from radical resection, but better local control. Delayed surgery gives firmer tumours, fewer ruptures and nephrectomies.
- Surgical complications: diarrhoea (autonomic nerve division at coeliac/superior mesenteric origins), chylous ascites (self-limiting).
- **Cord compression**: chemotherapy first; laminectomy only for progressive deficit (late scoliosis); avoid RT (spinal growth).

KEY POINT Surgical value runs inversely with biology: favourable tumours are cured by little or no resection; unfavourable ones are not rescued by radical resection — so vital structures are never traded for margin.

9 · ADRENOCORTICAL TUMOURS

- BWS predisposes to adrenocortical tumours (and hepatoblastoma).
- † Paediatric adrenocortical tumours (also Li–Fraumeni/TP53) usually present with virilisation or Cushing syndrome; complete open resection without capsular rupture is the mainstay, with mitotane-based chemotherapy for advanced disease.

10 · LANDMARK TRIALS

Trial	Design	Result	Take-home
AREN0532/0533	Combined 1p/16q LOH → regimen M	Stage III–IV EFS ~90% vs 61%	Biology-driven intensification
AREN0533 (lung)	Lung nodule response at week 6	RT omitted, EFS/OS preserved	Response-adapted RT omission
AREN0534	Bilateral WT: 3-drug, surgery by week 12	EFS 82%, OS 95% (vs 56%/81% NWTS-5)	Fixed clock, maximise NSS
COG P9641	Low-risk neuroblastoma, surgery alone	EFS ~89%, OS ~97%	Chemo only if symptomatic
COG anti-GD2	Anti-GD2 + cytokines + retinoid vs retinoid	Antibody superior	Immunotherapy standard in high risk
German NB97	Extent of resection, stage 4 >18 months	No EFS/OS benefit	Radical resection not essential

11 · MCQ ONE-LINERS

1. WT1 is at 11p13; WT2 putative locus at 11p15 (IGF2).
2. Denys–Drash: dominant-negative WT1 point mutation, 46,XY DSD, mesangial sclerosis.
3. WAGR and Denys–Drash carry ~50% WT risk; BWS 5–10%.
4. Screening: renal US every 3–4 months until age 7 if risk >1%.
5. Anaplasia: nuclei ≥3× normal, hyperchromasia, atypical mitoses; TP53 mutated.
6. Nephrogenic rests in >90% of bilateral WT.
7. Any tumour spill upstages WT to COG stage III.
8. Minimum lymph nodes sampled at Wilms nephrectomy (COG): five.
9. Acquired von Willebrand disease in up to 8% of WT.
10. Bilateral WT: definitive surgery by week 12 of chemotherapy.
11. Thrombus above hepatic veins → preoperative chemotherapy.
12. Doxorubicin cardiotoxicity rare below 150 mg/m² cumulative.
13. Commonest neonatal renal tumour: congenital mesoblastic nephroma (ETV6–NTRK3).
14. Clear cell sarcoma metastasises to bone; rhabdoid tumour associated with brain tumours.
15. Rhabdoid tumour: SMARCB1/INI1 loss, most lethal paediatric renal tumour.
16. Commonest paediatric RCC: Xp11.2 translocation (TFE3).
17. Renal medullary carcinoma: sickle cell trait.
18. Neuroblastoma: urinary VMA/HVA raised in 90–95%.
19. 123I-MIBG diagnostic; 131I-MIBG therapeutic.
20. MYCN amplification: >10 copies at 2p24, high risk.
21. Hyperdiploidy favourable in neuroblastoma; diploidy adverse.
22. INRG MS: age <18 months; skin, liver, marrow <10%; no bone.
23. Opsoclonus–myoclonus–ataxia: half have neuroblastoma.
24. Calcification and vessel encasement favour neuroblastoma over Wilms tumour.
25. Cord compression in neuroblastoma: chemotherapy first.

64 Paediatric Urological Oncology: Pelvis, Bladder, Testis and Genitalia

Reading time ≈ 24 min (slow pace)

1 · RHABDOMYOSARCOMA (RMS): BIOLOGY

- Aggressive sarcoma of rhabdomyoblasts; **commonest childhood sarcoma** (~3% of paediatric cancers); 20–30% genitourinary (GU). Bimodal: peak in first 2 y, smaller adolescent peak.
- GU sites: bladder/prostate (BP), paratesticular (PT), vulvovaginal (VV). GU-RMS 5-y survival **>80%**.
- Distant metastases in 16% at diagnosis (lung, bone, marrow).
- Most sporadic; ≥5% syndromic (usually embryonal): **Li–Fraumeni** (TP53), Beckwith–Wiedemann, **DICER1**, neurofibromatosis type 1 (pelvic RMS), Costello (bladder tumours). Diagnosis <3 y → look for germline predisposition.

	Embryonal (ERMS)	**Alveolar (ARMS)**
Site	GU tract (commonest GU histology); **botryoid** variant protrudes from bladder/vagina	Trunk, extremities, perineum
Age	Younger (~70% <10 y)	Older children, adolescents
Fusion	Usually fusion-negative	~80% **PAX3/7–FOXO1** [t(2;13) or t(1;13)]
Prognosis	Better	Fusion-positive worse; fusion-negative ARMS behaves like ERMS

KEY POINT Fusion status predicts outcome better than histology, so current trials assign risk by **fusion status**, not microscopy. MYOD1 mutation (fusion-negative) and circulating tumour DNA at diagnosis are emerging adverse markers.

2 · PRESENTATION AND WORK-UP

- BP: haematuria, stranguria, retention; botryoid mass at meatus. A 3–5-year-old boy with haematuria + new stranguria needs urgent assessment.
- PT: painless scrotal mass, separate from testis. VV: bloody discharge, prolapsing mass.

ALGORITHM
Ultrasound (US) → CT or MRI of primary + nodal basin (BP: pelvic and retroperitoneal to renal vessels; PT: retroperitoneal; VV: inguinal, pelvic, retroperitoneal) → **chest CT** (lung commonest metastatic site) → bone scan + bilateral marrow (omit in fusion-negative ERMS without nodal/lung disease)

CT = computed tomography; MRI = magnetic resonance imaging. Role of positron emission tomography–CT is unsettled.

3 · STAGE, GROUP AND RISK

Three layers: **stage** (pretreatment tumour–node–metastasis, TNM — where the tumour began) → **clinical group** (what the surgeon left behind) → **risk group** (adds fusion status and age).

COG TNM stage	**Site**	**Size / nodes**
1	Favourable: **paratesticular, female GU**, orbit, head–neck	Any, M0
2	Unfavourable: **bladder/prostate**, extremity, parameningeal, trunk	≤5 cm, N0/NX
3	Unfavourable	≤5 cm N1, or >5 cm any N

COG TNM stage	Site	Size / nodes
4	Any	M1

COG = Children's Oncology Group. N1 = node ≥1 cm short axis. BP-RMS can never be below stage 2.

Clinical group (Intergroup Rhabdomyosarcoma Study Group/COG)	Definition
I	Localised, completely resected, nodes negative
IIa / IIb / IIc	Microscopic residual / involved nodes resected, most distal node negative / microscopic residual or most distal node positive
III	Gross residual or **biopsy only**
IV	Distant metastases at onset

Risk	Criteria
Low	Fusion-negative; stage 1 group I–II (III orbit only), or stage 2 group I–II
Intermediate	Fusion-negative stage 1–2 group III (non-orbit), stage 3 group I–II; any fusion-positive non-metastatic; stage 4 fusion-negative <10 y
High	Stage 4 fusion-positive, or ≥10 y

> **KEY POINT** Organ-preserving **biopsy** deliberately places BP-RMS in group III → **intermediate risk**; PT-RMS (resected by orchidectomy) is almost always low risk. Age <1 y or >10 y is independently adverse. Trap: resected nodes with most distal node positive = IIc, not IIb.

4 · TREATMENT PRINCIPLES

- **Chemotherapy is universal**; local control by radiotherapy (RT) and/or surgery; clinical group sets radiotherapy dose. Complete resection (except PT) only if the organ can be preserved.
- Primary exenteration fell from almost all (IRS-I) to **13%** (IRS-IV) with unchanged OS (78–83%). IRS = Intergroup Rhabdomyosarcoma Study; OS = overall survival.

	Pretreatment re-excision (PRE)	Delayed primary excision (DPE)
Timing	Before chemotherapy	After chemotherapy starts
GU survival	**Improves**	Not shown
Group	Can change group	Cannot
Indication	Resectable without organ loss; positive/uncertain margins	Only if R0 (negative-margin) resection likely; may lower RT dose

Bladder/prostate RMS

- Biopsy: **cystoscopic** (preferred; 9–11.5 French resectoscope, undermine with cutting current, remove with forceps, minimal cautery) → percutaneous → open transvesical (last resort; spill).
- Relieve obstruction: **urethral catheter** first (usually removed within weeks as chemo works) → JJ stent → percutaneous nephrostomy if stent impossible. Avoid suprapubic tube (seeding); vesicostomy/ureterostomy delays chemotherapy.
- **End-of-therapy residual mass**: usually stroma or mature rhabdomyoblasts (misread as tumour) → observe if stable and asymptomatic; resection does not improve survival.
- Viable tumour after chemoradiation without metastasis → partial cystectomy (if adequate margin and capacity) or cystoprostatectomy/exenteration can cure.
- Bladder function: only ~40% "normal" after preservation (detrusor overactivity, small capacity). Follow-up: **urodynamics at 6 months**, renal US annually ≥10 y.

Paratesticular RMS

- **Radical inguinal orchidectomy**. Prior scrotal approach → excise remaining cord inguinally as PRE; hemiscrotectomy/scrotal RT only for positive cord margin or scrotal skin invasion. Tumour abutting but not invading the tunica vaginalis is **not** a positive margin.
- **Ipsilateral nerve-sparing retroperitoneal lymph node dissection (RPLND)** before chemotherapy if **age ≥10 y** or radiological N1 at any age; ≥7–13 nodes, up to renal vessels. Open, laparoscopic or robotic — none proven superior.

> **KEY POINT** Why age, not CT? Older boys have far more retroperitoneal disease and CT has a high false-negative rate for nodes.

Female GU, perineal, infants

- Work-up: MRI → examination under anaesthesia → cystoscopy → vaginoscopy + biopsy. Chemotherapy ± brachytherapy controls most; no node dissection; hysterectomy/exenteration only for unresponsive tumours. Cervical RMS needs biopsy after induction even with no visible disease.
- Vulvovaginal (two-thirds of female GU-RMS): MRI and vaginoscopy after 3 and/or 6 cycles; residual positive biopsy → brachytherapy. Uterine: younger, worse OS; hysterectomy only if R0, preserving ovaries.
- Perineal/perianal: mostly alveolar, fusion-positive; 5-y OS ~50%.
- Infants: radiotherapy often withheld (bony pelvis, hips) → more local failure.

Late effects

Radiotherapy causes almost tenfold more sequelae. Bladder dysfunction >60% despite preservation; sexual dysfunction, infertility, ovarian failure, vaginal stenosis; second cancers ~6× (alkylators, anthracyclines). Acute: vincristine neuropathy, doxorubicin cardiotoxicity, cyclophosphamide haemorrhagic cystitis.

5 · OTHER BLADDER TUMOURS

- **Urothelial**: mostly papillary urothelial neoplasm of low malignant potential (PUNLMP)/low-grade; haematuria; transurethral resection of bladder tumour (TURBT) with paediatric resectoscope, fulgurate base; renal–bladder US surveillance. **Recurrence should prompt search for a syndrome** (e.g. Lynch).
- **Spina bifida/exstrophy**: baseline lifetime bladder cancer risk **2–4%**, aggressive, **not increased by intestinal augmentation**; screening asymptomatic patients is neither effective nor cost-effective (cancers arise within months of normal surveillance). Gastrocystoplasty cancers prompted concern.

Benign lesion	Key point	Treatment
Inflammatory myofibroblastic tumour	ALK rearrangement; intermediate malignancy	Complete resection; excellent outcome
Nephrogenic adenoma	After trauma, infection, surgery, transplant; biopsy needed	TURBT/fulguration + treat cause; recurs
Eosinophilic cystitis	Biopsy essential; eosinophilia often absent	Steroids, antihistamines
Haemangioma	Severe haematuria; Klippel–Trenaunay, Sturge–Weber	Propranolol, laser; partial cystectomy
Fibroepithelial polyp	Mimics pelviureteric obstruction (Dietl crisis) or urethral valves	—

6 · TESTICULAR TUMOURS

- ~1/100,000 under 15 y; peak 2–4 y and puberty. Benign far commoner than in adults (38–74%).
- Prepubertal: **teratoma** = commonest tumour (benign); **yolk sac tumour (YST)** = commonest malignancy.

- Painless mass; scrotal Doppler US + markers — lactate dehydrogenase, α-fetoprotein (**AFP**), β-human chorionic gonadotrophin (β-hCG); hormones if precocious puberty.
- US cannot reliably separate benign from malignant: anechoic/cystic with echogenic foci → teratoma; **"onion-skin" rings → epidermoid cyst**.

	AFP	**β-hCG**
Source	Yolk sac, fetal liver, gut — **YST invariably**	Embryonal carcinoma, mixed tumours; rare prepubertal
Half-life	**5 days**	**24 h**
Caveat	Physiologically high in infancy; adult level (<10 ng/mL) by **6–9 months**	—

Surgery

ALGORITHM
Inguinal approach → markers normal + benign-looking US → **testis-sparing partial orchidectomy** with frozen section → benign: close; germ cell malignancy or raised AFP → radical inguinal orchidectomy

- Prepubertal teratoma: partial orchidectomy; **no follow-up** needed. † Postpubertal teratoma is malignant (adult-type) → radical orchidectomy.
- **YST**: radical inguinal orchidectomy; **90%** stage I. CT chest/abdomen/pelvis; follow AFP decline by half-life. Stage I → **surveillance, no RPLND or chemotherapy**. Scrotal violation → stage II (chemo, no hemiscrotectomy). Nodes + raised markers = stage III; stage III–IV → cisplatin, etoposide, bleomycin. Persistent mass → biopsy/RPLND.
- **Stromal tumours**: Leydig (commonest; precocious puberty; Reinke crystals ~40%), Sertoli, juvenile granulosa (benign in infants) → partial orchidectomy if markers normal; radical after puberty (Tanner 5). **Testicular adrenal rests** in poorly controlled congenital adrenal hyperplasia → increase steroid dose first.
- Leukaemia/lymphoma: commonest secondary testicular tumours; testicular relapse now ≤2%.
- Microlithiasis: no childhood surveillance; self-examination from puberty only if risk factors.

Disorders of sex development (DSD)

Risk rises with Y material (SRY/GBY regions) in dysgenetic gonads: complete gonadal dysgenesis ~35%, partial ~24%, Frasier, Denys–Drash, partial androgen insensitivity, Turner with Y = high; complete androgen insensitivity, mixed gonadal dysgenesis = intermediate; no Y = low. Gonadoblastoma is premalignant → dysgerminoma. Gonadectomy is individualised (risk, gender identity, fertility).

> **KEY POINT** Trap: **complete** androgen insensitivity is intermediate (not high) risk; partial is high.

7 · OTHER PELVIC AND GENITAL TUMOURS

- **Sacrococcygeal teratoma**: ~1/40,000, female; neurogenic bladder 36–51%, worse with Altman II–IV (internal). Annual renal–bladder US until toilet training; urodynamics for higher types. **Currarino triad** (anterior sacral mass, sacral defect, anorectal malformation) → neurogenic bladder.
- Paratesticular masses other than RMS (epididymal cyst, adenomatoid tumour, cystadenoma — von Hippel–Lindau): testis-sparing excision, but inguinal approach + frozen section mandatory.
- Ovary: cysts common and regress; malignancy rare (germ cell commonest). Sertoli–Leydig tumour → **screen DICER1**.
- **Urethral prolapse** mimics RMS (doughnut swelling, prepubertal Black girls) → topical oestrogen/steroid; no follow-up.

8 · MCQ ONE-LINERS

1. Commonest paediatric soft-tissue sarcoma: rhabdomyosarcoma.
2. Commonest GU-RMS histology: embryonal; botryoid variant in bladder/vagina.
3. Alveolar RMS fusion: PAX3/7–FOXO1 (chromosome 13).
4. Risk now assigned by fusion status, not histology.
5. Paratesticular and vulvovaginal are favourable sites; bladder/prostate unfavourable.
6. BP-RMS is at least stage 2.
7. Biopsy-only BP-RMS = group III → intermediate risk.
8. Resected nodes, most distal positive = group IIc.
9. Pretreatment re-excision improves GU-RMS survival; delayed excision does not.
0. First drainage in obstructing BP-RMS: urethral catheter.
1. Stable end-of-therapy bladder mass: observe.
2. Paratesticular RMS RPLND: age ≥10 years or N1.
3. Paratesticular RMS: radical inguinal orchidectomy.
4. Exenteration fell to 13% by IRS-IV without survival loss.
5. Recurrent paediatric urothelial tumour: consider Lynch or other syndrome.
6. Neurogenic bladder cancer risk 2–4%, not raised by augmentation.
7. Commonest prepubertal testicular tumour: teratoma; malignant: yolk sac tumour.
8. AFP half-life 5 days; β-hCG 24 hours.
9. AFP reaches adult levels by 6–9 months of age.
20. Onion-skin appearance on US: epidermoid cyst.
21. Stage I prepubertal YST: surveillance after orchidectomy.
22. Reinke crystals: Leydig cell tumour.
23. Testicular adrenal rests: optimise steroid replacement first.
24. Complete androgen insensitivity: intermediate gonadal tumour risk.
25. Sertoli–Leydig ovarian tumour: test for DICER1.
26. Currarino triad: sacral mass, sacral defect, anorectal malformation.

★ Putting it together: common paediatric decisions

Reading time ≈ 8 min (slow pace) · Integrates the paediatric chapters · use these as the skeleton of long-essay answers

A · VESICOURETERAL REFLUX (VUR) AFTER A FEBRILE URINARY TRACT INFECTION (UTI)

ALGORITHM

1. Renal and bladder ultrasound (US) → voiding cystourethrogram (VCUG) if febrile UTIs are multiple, need admission, or US is abnormal.

2. Everyone: treat bladder and bowel dysfunction (BBD), starting with constipation.

3. Low risk (grade I–II, no BBD, boy, older) → observation ± continuous antibiotic prophylaxis (CAP). High risk (grade ≥III, girl, BBD, scars, infant) → CAP.

4. Breakthrough febrile UTI, new scars, or persistent high grade with poor function → correction: endoscopic injection or ureteric reimplantation (→ *Ch 41, 53*).

B · DILATED UPPER TRACT: WHICH DIAGNOSIS?

Condition	Key test	Operate when	Operation
Pelviureteric junction (PUJ) obstruction	Mercaptoacetyltriglycine (MAG3) diuretic renogram	Split function falling or <40%, worsening dilatation, symptoms or infection	Anderson–Hynes dismembered pyeloplasty (open, laparoscopic or robotic)
Primary obstructive megaureter	US + MAG3 (VCUG to exclude reflux)	Falling function, infection, progressive dilatation; most resolve	Tapered or folded reimplant; temporary diversion or balloon dilatation in infants
Duplex with ureterocele / ectopic ureter	US, VCUG, functional scan	Obstruction, infection, incontinence (ectopic ureter in girls)	Endoscopic incision, upper-pole heminephrectomy, or ureteroureterostomy (→ *Ch 52*)
Multicystic dysplastic kidney	US + dimercaptosuccinic acid (DMSA) scan (non-functioning)	Rarely; most involute	Observation; check the contralateral kidney for VUR or PUJ obstruction

C · UNDESCENDED TESTIS (UDT)

ALGORITHM

Palpable → orchidopexy at **6–18 months** (ideally by 12). Imaging is not recommended.

Non-palpable → examination under anaesthesia → still impalpable → diagnostic laparoscopy: blind-ending vessels = vanished testis → stop · vessels and vas entering the ring → groin exploration, excise the nubbin · intra-abdominal testis → laparoscopic orchidopexy (primary or Fowler–Stephens).

Bilateral non-palpable or UDT + hypospadias → work up for a difference of sexual development (DSD) first (→ *Ch 59*).

D · ACUTE SCROTUM

	Testicular torsion	Torsion of an appendage	Epididymitis
Typical	Sudden severe pain, vomiting, high-riding hard testis; neonates and adolescents	Prepubertal; localised upper-pole tenderness, "blue dot"	Gradual onset; urinary symptoms; consider underlying anomaly in young boys

	Testicular torsion	Torsion of an appendage	Epididymitis
Action	**Explore without delay** if suspicion is high (Doppler can miss partial torsion); detorse, fix both sides, remove if non-viable	Analgesia, rest	Urinalysis/culture; antibiotics if infected

E · NEWBORN WITH ATYPICAL GENITALIA

> **KEY POINT** Bilateral impalpable gonads + atypical genitalia = salt-wasting congenital adrenal hyperplasia (CAH) until proven otherwise. Check electrolytes and 17-hydroxyprogesterone (after 36 h) and karyotype; the salt-wasting crisis usually comes at days 5–7. Sex designation and surgery are not emergencies and need a multidisciplinary team.

F · WILMS TUMOUR VS NEUROBLASTOMA

	Wilms tumour	Neuroblastoma
Age	Median 3.5 years	† Younger (median about 18 months)
Imaging	Arises within the kidney († claw sign); may extend into the inferior vena cava (IVC)	Calcification and vessel encasement; † crosses the midline, displaces the kidney
Markers / genetics	WT1, 11p15; syndromes (WAGR — Wilms, aniridia, genitourinary anomalies, intellectual disability; Denys–Drash; Beckwith–Wiedemann)	Urinary catecholamine metabolites — vanillylmandelic acid (VMA), homovanillic acid (HVA) — raised in 90–95%; MYCN amplification = high risk
Strategy	Children's Oncology Group (COG): upfront nephrectomy · International Society of Paediatric Oncology (SIOP): neoadjuvant chemotherapy then surgery	By International Neuroblastoma Risk Group: observe (some infants) → surgery → multimodal therapy