

Paper IV - Recent Advances (MCh Urology, Summer 2015)

Contents

- IV-1. Role of robotics in urology (LAQ, 25)
- IV-2. Marginal kidney donor and newer immunosuppressant drugs in renal transplantation (LAQ, 25)
- IV-3a. Hypospadias fistula: aetiology and management (SAQ, 10)
- IV-3b. Silodosin (SAQ, 10)
- IV-3c. Sacral neuromodulation (SAQ, 10): cross-reference to Paper III
- IV-3d. Urethral stents (SAQ, 10)
- IV-3e. Studer neobladder (SAQ, 10)
- IV-3f. Cryotherapy for carcinoma prostate (SAQ, 10)

IV-1. Role of robotics in urology

Type: LAQ (25 marks) | ~1500 words

Introduction / Definition

- **Robot:** a reprogrammable, multifunctional manipulator. Surgical systems in use today are **master-slave telemanipulators**. The surgeon controls every movement, so the system is not autonomous.
- Urology took up robotics earlier and more widely than any other specialty, largely because of **robot-assisted radical prostatectomy (RARP)**. In many high-income countries RARP is now the commonest approach to radical prostatectomy.
- Campbell: Robotic and laparoscopic surgery chapters (Basics of laparoscopic and robotic surgery).

History (milestones)

Year	Milestone
1985	PUMA 560: CT-guided stereotactic brain biopsy (neurosurgery)
Late 1980s	PROBOT (Imperial College, London): first urological robot, for automated TURP-like prostate resection
1990s	AESOP (Computer Motion): voice-controlled laparoscope holder. ZEUS: master-slave system
2000	da Vinci (Intuitive Surgical) FDA-cleared for general laparoscopic surgery
2000	Binder & Kramer (Frankfurt): first robotic radical prostatectomy
2001	"Lindbergh operation" (Marescaux): transatlantic telesurgical cholecystectomy, New York to Strasbourg, using ZEUS
Early 2000s	Menon (Vattikuti Urology Institute, Detroit): standardised VIP (Vattikuti Institute Prostatectomy) , which made RARP popular

da Vinci system: components

1. **Surgeon console:** 3D high-definition stereoscopic viewer, master controllers, foot pedals (camera, clutch, energy), motion scaling and tremor filtration.
2. **Patient-side cart:** 3-4 robotic arms carrying the camera and **EndoWrist** instruments with 7 degrees of freedom.

3. **Vision cart:** image processing, light source, energy generator, assistant monitor.

Generations: standard (2000), **S** (2006), **Si** (2009; dual console for training, Firefly fluorescence), **Xi** (2014; boom-mounted rotating arms, thinner arms, laser targeting, multiquadrant access), **SP** (single port, single 2.5 cm cannula; FDA approval for urology 2018), **da Vinci 5** (FDA 2024; force-feedback instruments, greater computing power).

Other platforms

- **Hugo RAS** (Medtronic): modular, independent arm carts, open console.
- **Versius** (CMR Surgical, UK): small modular arms, open console.
- **Senhance** (Asensus): haptic feedback, eye-tracking camera control, reusable instruments.
- **SSI Mantra** (SS Innovations, India): Indian low-cost modular system, also used for telesurgery.
- Cheaper competitors are expected to bring costs down, which matters especially for India.

Advantages and disadvantages

Advantages	Disadvantages
3D magnified vision	High capital (~US\$1.5–2.5 million) and disposable costs
7 DOF wristed instruments	Limited haptic feedback (partly addressed in da Vinci 5 and Senhance)
Tremor filtration, motion scaling	Docking time; large footprint
Ergonomic for surgeon	Surgeon remote from patient (emergency undocking)
Shorter learning curve than pure laparoscopy for suturing	Learning curve still exists; credentialing issues
Less blood loss, shorter stay than open	Port-site hernia; specific positioning complications
Ideal for reconstruction (intracorporeal suturing)	Limited availability in public sector

Applications in urology (with evidence)

1. Radical prostatectomy (RARP): the index procedure

- **Yaxley et al., Lancet 2016:** single-surgeon RCT of open retropubic vs robotic prostatectomy. Urinary and sexual function were **similar at 12 weeks**. The robotic arm had less blood loss and a shorter hospital stay.
- **Coughlin et al., Lancet Oncol 2018** (same trial): functional outcomes were still **similar at 24 months**.
- Systematic reviews show lower transfusion rates and shorter stay. Oncological outcomes are comparable to open surgery.
- Technical refinements: Retzius-sparing (Bocciardi) for earlier continence, NeuroSAFE, extended PLND, and single-port/transvesical SP-RARP.

2. Robotic partial nephrectomy (RAPN)

- Compared with laparoscopic partial nephrectomy: **shorter warm ischaemia time, lower conversion to radical nephrectomy**, a shorter learning curve, and the ability to treat complex/hilar tumours.
- Adjuncts include **Firefly (ICG)** for selective clamping, zero ischaemia techniques and 3D virtual models.
- In many centres it has become the preferred minimally invasive approach.

3. Robotic radical cystectomy (RARC)

- **RAZOR trial (Parekh et al., Lancet 2018):** RARC was **non-inferior** to open cystectomy for **2-year progression-free survival**, with less blood loss, fewer transfusions and a slightly shorter stay.

- **iROC trial (Catto et al., JAMA 2022):** RARC with **intracorporeal** diversion increased **days alive and out of hospital at 90 days**, and reduced thromboembolism and wound complications, compared with open surgery.
 - Intracorporeal neobladder: Karolinska (Wiklund) technique.
- 4. Pyeloplasty:** success above 90–95%. Robotic surgery is widely used for adult and paediatric PUJ obstruction and redo pyeloplasty.
- 5. Ureteric reimplantation**
- Adults: Boari flap and psoas hitch for distal strictures.
 - Children: **RALUR** (robot-assisted laparoscopic extravesical ureteral reimplantation, Lich-Gregoir) for VUR, with success rates approaching open surgery in experienced hands.
- 6. Ureteric reconstruction: robotic buccal mucosa graft (BMG) ureteroplasty** for long proximal/mid-ureteric strictures, ureterocalycostomy, ileal ureter, and ureteroureterostomy for retrocaval ureter.
- 7. RPLND (R-RPLND):** primary RPLND for low-volume stage II seminoma or NSGCT, and selected post-chemotherapy cases. Template dissection can preserve ejaculation.
- 8. Adrenalectomy:** equivalent to laparoscopy and helpful in obese patients, partial adrenalectomy and large tumours.
- 9. Robotic simple prostatectomy:** for large glands (>80–100 mL). Less bleeding and shorter stay than open surgery.
- 10. Sacrocolpopexy:** robotic approach for apical prolapse. Also used for VVF/vesicouterine fistula repair, Mitrofanoff and augmentation cystoplasty.
- 11. Robotic kidney transplantation (RAKT):** developed by Menon, Vattikuti and the ERUS-RAKT group (Breda). Lower wound complications and lymphocele rates, useful in obese recipients, with regional hypothermia.
- 12. Other uses:** robotic radical nephrectomy with IVC thrombectomy, nephroureterectomy, ilioinguinal lymph node dissection (VEIL) and diverticulectomy.

Training and credentialing

- Structured pathway: didactics, then **virtual-reality simulators** (dV-Skills Simulator, Mimic dV-Trainer, RobotiX), then dry and wet lab, then modular supervised console time.
- **Fundamentals of Robotic Surgery (FRS):** a validated, multispecialty curriculum similar to FLS.
- **ERUS curriculum** (EAU Robotic Urology Section): modular training with certification.
- **Dual console** (Si and Xi) for mentoring. Credentialing is decided by each institution (minimum case numbers, proctoring).
- **Learning curve for RARP:** commonly quoted as many tens of cases for operative proficiency and more than 150–250 for optimal margins and continence.

Cost

- Costs come from the capital purchase, annual maintenance and limited-use instruments.
- Robotic surgery is cost-effective only at high volume.
- Cheaper alternatives include new platforms, reusable instruments, multispecialty use and indigenous Indian systems (SSI Mantra).

Complications specific to robotics

Problem	Details
Steep Trendelenburg (25–45°)	Raised intraocular pressure; ischaemic optic neuropathy (rare, blindness); corneal abrasion; raised ICP ; facial/laryngeal oedema; reduced FRC and ventilatory difficulty

Problem	Details
Positioning	Brachial plexus injury, lower-limb compartment syndrome / rhabdomyolysis in prolonged lithotomy, peroneal nerve palsy
Instrument-related	Insulation failure causing stray energy and bowel burns; instrument malfunction
Lack of haptics	Tissue tearing, suture breakage
Others	Port-site hernia; CO ₂ embolism; conversion; system failure (rare; most cases are completed or converted)

Prevention: limit Trendelenburg angle and time, padding, shoulder supports avoided or well padded, and lithotomy kept under about 2–3 h where possible.

Recent advances

- **Single-port (SP) surgery:** extraperitoneal and transvesical RARP, with the possibility of same-day discharge.
- **Telesurgery:** since 2023–25, long-distance telesurgery (including RARP and nephrectomy) has been reported from China and India (SSI Mantra) over 5G and fibre networks. Latency remains the main limitation.
- **Haptic / force feedback:** da Vinci 5.
- **Image guidance and augmented reality:** 3D-printed and virtual models overlaid onto the console view; ICG fluorescence; **PSMA-radioguided surgery**.
- **Artificial intelligence:** video-based skill assessment, automated performance metrics (kinematic data predicting continence after RARP), and phase recognition.
- **Autonomy:** graded levels (0–5); current systems are level 0–1. Experimental autonomous suturing (STAR) has been done in animals. Image-guided automated **Aquablation** (AquaBeam) is the first semi-autonomous resection in urology, echoing PROBOT.
- **Robotic endourology:** robotic flexible ureteroscopy (e.g. Avicenna Roboflex) and the Monarch platform for PCNL access.

Diagram to draw: block diagram of da Vinci (surgeon console linked to vision cart and patient-side cart, with EndoWrist 7 DOF inset); port placement for RARP (6-port transperitoneal fan).

Key points to remember: - PUMA 560 (1985) was the first surgical robot. PROBOT (Imperial College) was the first urological robot. da Vinci was FDA-approved in 2000. The first RARP was by Binder & Kramer in 2000. - Yaxley (Lancet 2016) and Coughlin (Lancet Oncol 2018) found no functional difference between open and robotic RP at 12 weeks and 24 months. - RAZOR (Lancet 2018): RARC non-inferior for 2-year PFS. iROC (JAMA 2022): intracorporeal RARC gave more days alive and out of hospital. - Generations: S, Si, Xi (2014), SP (2018), da Vinci 5 (2024). Alternatives: Hugo, Versius, Senhance, SSI Mantra (India). - Steep Trendelenburg causes raised IOP/ICP and ischaemic optic neuropathy; prolonged lithotomy causes compartment syndrome.

IV-2. Discuss marginal kidney donor and describe newer immunosuppressant drugs in renal transplantation

Type: LAQ (25 marks) | ~1600 words

Introduction

- The gap between demand and organ supply has driven wider use of **marginal (extended criteria) donors**. These are donors whose kidneys carry a higher risk of delayed graft function (DGF), primary non-function or shorter graft survival.
- For many wait-listed patients, a marginal kidney still gives better survival than staying on dialysis.
- Campbell: Renal transplantation chapter. Guidelines: KDIGO (candidate evaluation 2020; living donor 2017), EAU Renal Transplantation guideline.

Part A: Marginal kidney donor

Definitions and categories

1. Expanded criteria donor (ECD), UNOS 2002 (deceased, brain-dead):

- Any donor **≥60 years**, or
- A donor aged **50-59 years** with **≥2** of the following:
 - history of **hypertension**
 - **terminal serum creatinine >1.5 mg/dL**
 - **death from cerebrovascular accident**
- ECD kidneys carry about a 70% higher relative risk of graft loss than standard criteria donor (SCD) kidneys (RR >1.7).

2. KDRI / KDPI

- **KDRI** (Kidney Donor Risk Index; Rao 2009) is built from 10 donor factors: age, height, weight, ethnicity, hypertension, diabetes, cause of death, creatinine, HCV status and DCD status.
- **KDPI** is the KDRI expressed as a percentile (0-100%).
- In the **US Kidney Allocation System (December 2014)**, **KDPI >85%** replaced the ECD label.
- Under the same system, the **top 20% EPTS** recipients are matched to kidneys with **KDPI ≤20%** ("longevity matching").

3. Donation after circulatory death (DCD): Maastricht classification

Category	Description	Type
I	Dead on arrival / found dead	Uncontrolled
II	Unsuccessful resuscitation	Uncontrolled
III	Awaiting cardiac arrest after withdrawal of life-sustaining therapy	Controlled (commonest)
IV	Cardiac arrest in a brain-dead donor	Controlled/uncontrolled
V	Euthanasia / medically assisted death (added in some countries)	Controlled

- The **modified Maastricht classification (Paris, 2013)** subdivided the uncontrolled categories.
- Compared with brain-dead donor kidneys, DCD kidneys have **higher DGF rates (roughly 40-50%)** but **similar long-term graft survival**.

4. Other marginal donors

- Donors with **AKI**: graft outcomes are acceptable if renal function was previously normal, and these kidneys should not be discarded on terminal creatinine alone.
- Older, diabetic or hypertensive donors, and donors with proteinuria.
- **HCV-viraemic donor to HCV-negative recipient**, followed by **direct-acting antivirals (DAAs)** with cure rates near 100%. THINKER trial (Goldberg, NEJM 2017).
- **HIV-positive donor to HIV-positive recipient**: permitted in the US by the **HOPE Act (2013)**. South African experience (Muller) came first.
- **Small paediatric donors**: en bloc kidneys (both kidneys with the aorta and IVC) can be transplanted into adults. They carry a higher thrombosis risk but give good long-term function.
- Donors with **treated low-grade malignancy** (e.g. small RCC that has been excised) or those at increased infectious risk.

5. Marginal living donors (KDIGO 2017)

- Older donors; donors with controlled hypertension (on 1-2 drugs, no end-organ damage); obesity (assessed individually, with BMI >30-35 needing caution); isolated microscopic haematuria after a negative work-up; minor anatomical variants; single stone episode.
- KDIGO 2017 recommends a **risk-based approach**: each centre sets an acceptable threshold of projected **lifetime ESKD risk**.

Assessment and optimisation of marginal kidneys

Donor biopsy: Remuzzi score

- Grades glomerulosclerosis, tubular atrophy, interstitial fibrosis and arterial narrowing, each scored 0-3 (total 0-12).
 - **0-3**: single transplant
 - **4-6**: **dual kidney transplant**
 - **≥7**: discard
- **Glomerulosclerosis >20%** has traditionally been linked to poorer outcomes.
- Frozen-section biopsy has limitations, and over-reliance on biopsy leads to unnecessary discards.

Machine perfusion

- **Hypothermic machine perfusion (HMP)**:
 - **Moers et al., NEJM 2009** (Eurotransplant RCT, paired kidneys): HMP **reduced DGF** (about 21% vs 27%) and **improved 1-year graft survival**. The benefit was greatest in ECD kidneys.
 - Resistance indices and flow give an additional viability assessment, which should not be used alone.
- **Normothermic machine perfusion (NMP)** (Nicholson, Hosgood; Cambridge, UK) at 37 °C with oxygenated red cells allows viability assessment and repair (a "quality score"). It is still largely within trials.

Other optimisation measures

- **Short cold ischaemia time**: under 12-18 h is ideal for ECD/DCD kidneys.
- **Dual kidney transplantation (DKT)**: both marginal kidneys go into one recipient to supply enough nephron mass. Placement can be unilateral or bilateral in the iliac fossae.
- **Donor management**: e.g. mild donor hypothermia has been studied to reduce DGF.

Recipient matching

- **"Old-for-old"**: the **Eurotransplant Senior Program (ESP, 1999)** allocates kidneys from donors **≥65 years** to recipients **≥65 years**. It prioritises local allocation and short cold ischaemia time over HLA matching.

- Marginal kidneys suit older, diabetic recipients, who have a long expected wait and high dialysis mortality.
- Patients must give **informed consent** to receive an ECD/high-KDPI kidney.

Outcomes

- **Wolfe et al., NEJM 1999**: transplantation gave a large survival benefit over remaining on the waiting list.
- **Ojo et al. (JASN 2001)**: ECD transplantation also improved survival compared with dialysis, especially in recipients over 40 years, diabetics and at centres with long waiting times.
- Graft survival with ECD/high-KDPI kidneys is lower than with SCD kidneys, but patient survival exceeds that of matched patients staying on dialysis.
- **DGF management**: avoid nephrotoxic early high-dose CNI by delaying CNI under antibody induction.

Part B: Newer immunosuppressant drugs

Overview of the current standard

- **Induction**: basiliximab or rATG.
- **Maintenance (triple therapy)**: **tacrolimus + mycophenolate + steroids**, with or without early steroid withdrawal.
- "Newer" agents aim to avoid calcineurin inhibitor (CNI) nephrotoxicity, improve adherence, desensitise highly sensitised patients and treat antibody-mediated rejection (AMR).

Induction agents

Drug	Mechanism	Dose (usual)	Key points
Basiliximab	Chimeric anti-CD25 (IL-2 receptor α) monoclonal antibody; non-depleting	20 mg IV on day 0 and day 4	Low risk; well tolerated
Rabbit ATG (Thymoglobulin)	Polyclonal T-cell depletion	1-1.5 mg/kg/day; total ~4.5-6 mg/kg	High immunological risk, DGF. Cytokine release, leucopenia, thrombocytopenia, CMV and PTLD
Alemtuzumab	Anti-CD52 (lymphocytes)	30 mg single dose (off-label)	Profound prolonged depletion; allows steroid/CNI minimisation

Maintenance agents (newer and modified)

Drug / class	Mechanism	Points
Extended-release tacrolimus: Advagraf (once daily), Envarsus XR (LCP-tacrolimus, MeltDose)	Calcineurin inhibition reduces IL-2 transcription	Once-daily dosing improves adherence. Envarsus has higher bioavailability (lower dose, flatter peak) and fewer tremors
Mycophenolate (MMF; enteric-coated mycophenolate sodium, EC-MPS)	Inhibits IMPDH, blocking de novo purine synthesis in lymphocytes	GI upset and leucopenia; teratogenic

Drug / class	Mechanism	Points
mTOR inhibitors: sirolimus, everolimus	Bind FKBP12 and block mTOR, arresting the G1 to S cell cycle	Non-nephrotoxic, antiproliferative with fewer skin cancers, and fewer CMV/BK infections . TRANSFORM study (Pascual, JASN 2018) : everolimus + reduced-exposure CNI was non-inferior to MPA + standard CNI, with less CMV/BK viraemia. Side effects: impaired wound healing, lymphocele , proteinuria, hyperlipidaemia, pneumonitis, mouth ulcers. Avoid early after surgery in obese patients (wound problems)
Belatacept	CTLA4-Ig fusion protein that binds CD80/CD86 , blocking the CD28 costimulation "signal 2"	FDA 2011. IV infusion: 10 mg/kg (days 1, 5, then weeks 2, 4, 8, 12), then 5 mg/kg every 4 weeks . BENEFIT (living/SCD) and BENEFIT-EXT (ECD) trials: better GFR than ciclosporin, but higher early acute rejection . 7-year BENEFIT data (Vincenti, NEJM 2016) : improved patient and graft survival. Contraindicated in EBV-seronegative recipients because of the PTLD risk (especially CNS) . Avoids CNI nephrotoxicity, hypertension and diabetes. Monthly infusions help adherence

Desensitisation and antibody-mediated rejection

Drug	Mechanism	Use
IVIG	Immunomodulation, anti-idiotypic effect, FcRn saturation	Desensitisation; AMR (with plasmapheresis)
Rituximab	Anti-CD20 B-cell depletion	ABO-incompatible protocols; AMR adjunct
Bortezomib	Proteasome inhibitor causing plasma-cell apoptosis	AMR. The BORTEJECT RCT (late AMR) was negative. Neuropathy
Eculizumab	Anti-C5, blocking the membrane attack complex	aHUS recurrence prevention and treatment; selected early AMR. Meningococcal vaccination is mandatory
Imlifidase (IdeS)	<i>Streptococcus pyogenes</i> enzyme that cleaves IgG within hours	EMA conditional approval 2020 to desensitise highly sensitised patients with a positive crossmatch to a deceased donor. A single dose of 0.25 mg/kg. Antibody rebound occurs, and ATG/rituximab must be given after imlifidase because it also cleaves therapeutic IgG
Tocilizumab (anti-IL-6 receptor), clazakizumab (anti-IL-6)	Block the IL-6 pathway that drives DSA and inflammation	Chronic active AMR : investigational, not approved
Felzartamab	Anti-CD38 (plasma cells and NK cells)	Phase 2 trial in late AMR (NEJM 2024) showed promising effects; under study
C1-esterase inhibitor	Complement inhibition	AMR trials

Caution about voclosporin

- Voclosporin is a novel CNI that is **approved for lupus nephritis (2021)**.
- It is **not approved for renal transplantation**; earlier transplant studies did not lead to registration.

Tolerance and the future

- **Tolerance induction:** combined kidney and haematopoietic stem-cell transplantation (mixed chimerism) from HLA-matched or haploidentical living donors (Stanford, Northwestern and MGH protocols). Selected patients have been able to stop immunosuppression.
- **Regulatory T-cell (Treg) therapy:** ONE Study and TWO Study.
- **Anti-CD40/CD40L agents** (e.g. tegoprubart): investigational, and also used in xenotransplantation.
- **Xenotransplantation:** gene-edited pig kidney into a living human recipient (MGH, March 2024), followed by further compassionate-use cases and trials.

Side-effect summary

Drug	Signature toxicity
Tacrolimus	Nephrotoxicity, NODAT, tremor, hypertension, hyperkalaemia, hypomagnesaemia
MMF	GI upset, cytopenia, teratogenicity
mTOR inhibitors	Wound healing problems, lymphocele, proteinuria, dyslipidaemia, pneumonitis
Belatacept	PTLD (EBV-), early rejection
Depleting antibodies	Infection (CMV), PTLD, cytokine release

Recent advances

- Normothermic perfusion with ex vivo repair.
- Using HCV-positive kidneys with DAAs.
- Using **imlifidase** for highly sensitised patients.
- **Donor-derived cell-free DNA** and molecular microscope (MMDx) diagnostics for rejection.
- Pig-kidney xenotransplantation.

Diagram to draw: three-signal model of T-cell activation (signal 1 TCR-MHC, blocked by ATG; signal 2 CD28-CD80/86, blocked by belatacept; signal 3 IL-2R-mTOR, blocked by basiliximab and sirolimus/everolimus), with the CNI and MMF sites marked.

Key points to remember: - ECD (UNOS 2002): age ≥ 60 , or 50–59 with ≥ 2 of HTN, creatinine >1.5 mg/dL, CVA death. Replaced by KDPI $>85\%$ in December 2014. - Remuzzi score: 0–3 single, 4–6 dual, ≥ 7 discard. HMP reduces DGF (Moers, NEJM 2009). Eurotransplant Senior Program: ≥ 65 to ≥ 65 . - Belatacept (CTLA4-Ig): BENEFIT/BENEFIT-EXT show better GFR. Contraindicated in EBV-seronegative recipients (PTLD). - TRANSFORM: everolimus + low-dose CNI is non-inferior, with fewer CMV/BK infections. - Imlifidase (IgG-cleaving enzyme) has EMA conditional approval (2020) for highly sensitised recipients. Voclosporin is for lupus nephritis, not transplantation.

IV-3a. Etiology and management of hypospadias fistula

Type: SAQ (10 marks) | ~550 words

Definition and incidence

- A **urethrocutaneous fistula (UCF)** is an abnormal communication between the neourethra and the penile skin. It is the **commonest complication of hypospadias repair**.
- Incidence:
 - **Distal TIP (Snodgrass)**: about **<5-10%**.
 - **Proximal and staged repairs**: higher, often **10-25%** or more.
 - Highest in **redo** repairs.
- Most fistulas appear in the early postoperative period. Some present late, so long-term follow-up is needed.
- Campbell: Hypospadias chapter (paediatric).

Aetiology (risk factors)

Local or technical factors

- **Distal obstruction**: meatal stenosis, neourethral stricture, or a narrow neourethra causing high voiding pressure. This is the **most important correctable cause**.
- **Ischaemia** from poor flap vascularity, excess diathermy or tight dressing.
- **Infection** and **haematoma**.
- **Tension** on the suture line, and **overlapping suture lines** (urethral and skin lines on top of each other).
- **No barrier (interposition) layer** between the neourethra and the skin.
- Poor-quality tissue: a small glans, a shallow urethral plate, previous surgery, or a deficient spongiosum.
- Suture material: subcuticular, inverting, absorbable fine suture (6-0/7-0 PGA) is preferred.

Patient factors

- Proximal hypospadias, redo surgery, and small glans size (under about 14 mm). The role of age and preoperative androgens is debated.
- **Catheter or stent problems**: early removal, blockage or **bladder spasm**.

Prevention

- **Barrier layer**:
 - **dorsal dartos (preputial) pedicled flap**
 - **tunica vaginalis flap** (especially for proximal or redo repairs)
 - spongioplasty
- These layers reduce fistula rates significantly.
- Other measures:
 - non-overlapping suture lines, with a tension-free glanuloplasty
 - meticulous haemostasis and magnification
 - adequate urinary drainage

Evaluation

- Examine during voiding: site, number and size. A coronal fistula may mean glans dehiscence.
- **Rule out distal obstruction**:
 - calibrate the meatus and neourethra

- urethroscopy/cystoscopy (often at the time of repair)
- uroflowmetry
- Saline or methylene-blue injection through the meatus at surgery (with the urethra compressed proximally) identifies small or multiple fistulas.
- Skin cover: is there enough local dartos?

Management

- **Timing: wait at least 6 months** after the primary repair for inflammation to settle and tissue to mature. A few tiny early fistulas close on their own.
- **Always correct distal obstruction** (meatotomy, meatoplasty, stricture management) at the same sitting. Otherwise the fistula recurs.

Fistula type	Procedure
Small (<2-3 mm), single, away from glans	Excise the tract, close in layers (inverting subcuticular 6-0/7-0 absorbable), add a dartos/subcutaneous barrier flap , rotate skin so suture lines do not overlap
Large, multiple, or with a thin urethra	Redo urethroplasty (TIP redo, onlay flap, or staged BMG)
Coronal / subcoronal (glans dehiscence)	Re-do glanuloplasty or meatoplasty (redo TIP), rather than simple closure
Barrier options	Local dartos, tunica vaginalis flap , de-epithelialised skin flap; Y-V skin advancement for skin cover

- **Diversion:** small simple closures can be left without a catheter or with a short-term stent. Larger or redo repairs need a stent or catheter for 5–10 days.
- **Success:** simple layered closure succeeds in about **80-90%**. Results are worse for large, multiple or recurrent fistulas.
- **Recurrence:** look for missed distal obstruction.

Recent advances

- Routine use of **barrier layers** and **tunica vaginalis** flaps in proximal repairs.
- **Tissue sealants** (fibrin glue) and buccal grafts in staged redo surgery.
- Validated outcome scores (**HOSE, HOPE**) for standardised reporting of complications.

UCF → wait ≥6 mo → calibrate/cystoscopy
 └ distal obstruction? → correct same sitting
 └ small single → excise + layered closure + dartos/TV flap
 └ large/multiple/glans dehiscence → redo urethroplasty

Diagram to draw: cross-section of the repair showing the neourethra, dartos barrier flap and skin, with suture lines offset.

Key points to remember: - Fistula is the commonest complication. TIP distal rate is under 5–10%; proximal repairs carry a higher rate. - Distal obstruction (meatal stenosis) must always be excluded and corrected. - Wait at least 6 months, then do a layered closure with a barrier (dartos or tunica vaginalis) and non-overlapping suture lines. - Coronal fistula equals glans dehiscence and needs redo glanuloplasty.

IV-3b. Silodosin

Type: SAQ (10 marks) | ~550 words

Definition

- Silodosin is a **highly uroselective α 1A-adrenoceptor antagonist** (indoline derivative), used mainly for **LUTS due to BPH**.
- Campbell: Medical management of BPH chapter. EAU Non-neurogenic male LUTS guideline. AUA BPH guideline.

Pharmacology

- Receptor selectivity:**
 - α 1A : α 1B about **162 : 1**
 - α 1A : α 1D about **55 : 1**
- Practical consequence: very little effect on vascular α 1B receptors, so **orthostatic hypotension is minimal**. The α 1A blockade in the prostate, bladder neck, urethra and seminal tract causes **ejaculatory dysfunction**.

Pharmacokinetics

- Oral bioavailability about **32%**. It is taken **with a meal**.
- Metabolised by **CYP3A4**, **UGT2B7** (forming the active glucuronide KMD-3213G, which has a long half-life) and alcohol/aldehyde dehydrogenase.
- It is a **P-glycoprotein** substrate.
- Excreted in urine and faeces.

Dose and contraindications

Item	Detail
Dose	8 mg once daily with a meal (4 mg capsules also available)
Renal impairment	Moderate (CrCl 30-50 mL/min): 4 mg daily
Contraindicated	Severe renal impairment (CrCl <30), severe hepatic impairment (Child-Pugh C), strong CYP3A4 inhibitors (ketoconazole, clarithromycin, ritonavir), strong P-gp inhibitors (ciclosporin)
Caution	Other α -blockers, PDE5 inhibitors (minor additive BP effect), planned cataract surgery

Efficacy

- US phase III RCTs (Marks et al., 2009):** compared with placebo, silodosin improved **IPSS and Qmax**, with onset within **2-4 days** (Qmax improved within hours of the first dose).
- Chapple et al., Eur Urol 2011** (European RCT, silodosin 8 mg vs tamsulosin 0.4 mg vs placebo, 12 weeks):
 - silodosin was **non-inferior to tamsulosin** for IPSS
 - only silodosin significantly reduced **nocturia** compared with placebo
 - abnormal ejaculation was more frequent with silodosin
- Long-term open-label extensions show sustained efficacy.
- Silodosin does not reduce prostate volume or the risk of acute urinary retention or surgery. Combine it with a 5-ARI for large glands.

Adverse effects

- **Retrograde ejaculation / anejaculation:** about **28%** in US trials. This is the **most common** side effect and is reversible.
 - Mechanism: reduced seminal emission (vas, seminal vesicles) rather than true retrograde flow.
- Dizziness, orthostatic hypotension (<3%), nasal congestion, diarrhoea, headache.
- **Intraoperative floppy iris syndrome (IFIS):** warn the patient and inform the ophthalmologist before cataract surgery.

Uses

1. **BPH/LUTS:** first-line α -blocker. Useful in elderly and cardiac patients because of low hypotension risk.
2. **Medical expulsive therapy (MET), off-label:** for **distal ureteric stones >5 mm**.
 - Meta-analyses and small RCTs suggest silodosin may give expulsion rates similar to or higher than tamsulosin.
 - Large tamsulosin RCTs (**SUSPEND, Lancet 2015**) showed no benefit overall.
 - **EAU** supports α -blockers for distal stones >5 mm.
3. **Ureteric stent-related symptoms:** improves pain and urinary scores.
4. **Chronic prostatitis / chronic pelvic pain syndrome:** as part of multimodal UPOINT therapy, especially in α -blocker-naïve men.
5. Also used for premature ejaculation (exploiting the anejaculation effect) and to facilitate ureteral access sheath placement (studied).

Comparison of α -blockers

Drug	Selectivity	Dose	Titration	Orthostatic hypotension	Ejaculatory dysfunction
Silodosin	α 1A (very high)	8 mg OD	No	Least	Highest (~28%)
Tamsulosin	α 1A = α 1D > α 1B	0.4 mg OD	No	Low	Moderate (~10%)
Alfuzosin	Non-selective (clinically uroselective)	10 mg OD (XL)	No	Low-moderate	Least
Doxazosin	Non-selective	1-8 mg OD (GITS 4-8 mg)	Yes (standard form)	Higher	Low
Terazosin	Non-selective	1-10 mg OD at night	Yes	Highest ("first-dose" effect)	Low

Recent advances

- Fixed-dose combinations with dutasteride or tadalafil (in some markets).
- Use of silodosin in MET and stent symptoms.
- Interest in silodosin for premature ejaculation.

Diagram to draw: α 1 receptor subtype distribution (α 1A in prostate/bladder neck, α 1B in vessels, α 1D in bladder/spinal cord) with the action of silodosin marked.

Key points to remember: - Selectivity α 1A : α 1B is about 162 : 1. Dose 8 mg OD with a meal; 4 mg if CrCl 30-50; contraindicated if CrCl <30, Child-Pugh C, or with strong CYP3A4 inhibitors. - Commonest adverse effect is anejaculation/retrograde ejaculation (~28%). Lowest hypotension risk of the α -blockers. - Chapple 2011: non-inferior to tamsulosin and also reduced nocturia. - Warn about IFIS before cataract surgery.

IV-3c. Sacral neuromodulation

Type: SAQ (10 marks)

See Paper III, Q3a (Sacral neuromodulation): identical question.

Recent advances angle (emphasise in a Recent Advances paper): - **Rechargeable, miniaturised IPGs:** - **Axonics r-SNM** (FDA 2019; about 15-year life) - **Medtronic InterStim Micro** (rechargeable) - recharge-free long-life devices (**InterStim X, Axonics F15**) - **Full-body MRI-conditional** systems (1.5 T and 3 T with SureScan/compatible leads), which solve the main historical limitation of SNM. - **Standardised tined-lead placement** with curved stylet and fluoroscopic landmarks (Matzel), improving test-phase success and reducing reoperation. - **Closed-loop / sensing SNM** using evoked compound action potential (ECAP) recordings to auto-adjust stimulation is at the research stage. - **Implantable tibial nerve stimulators:** **eCoin** (FDA 2022), **Revi** (BlueWind, FDA 2023) give less-invasive alternatives for refractory OAB/UUI.

IV-3d. Urethral stents

Type: SAQ (10 marks) | ~550 words

Definition

- A urethral stent is a tubular endoprosthesis placed endoscopically in the **prostatic or bulbar urethra** to keep the lumen open. It is a **minimally invasive alternative** to surgery in men unfit for definitive treatment.
- Campbell: Urethral stricture / reconstruction and BPH minimally invasive therapy chapters.

Classification

Type	Examples	Features
Temporary, non-epithelialising	Memokath (thermo-expandable nickel-titanium shape-memory spiral; e.g. Memokath 028 prostatic, 044 bulbar), Allium (covered, self-expanding nitinol; BUS bulbar and prostatic stents), polyurethane / Spanner temporary prostatic stent	Do not become epithelialised; removable; need replacement or removal over months to years
Permanent, epithelialising	UroLume (woven superalloy mesh)	Covered by urothelium in 3–6 months; difficult removal. UroLume has been withdrawn from the market
Biodegradable	PLGA spiral stents (experimental)	Dissolve; avoid removal
By site	Prostatic (BPH, retention) vs bulbar urethral (stricture) vs sphincteric (DSD)	

Indications

- BPH with retention or LUTS in high-risk patients** unfit for TURP or anaesthesia, as an alternative to a long-term catheter.
- Recurrent bulbar urethral stricture** in men **unfit for or unwilling to undergo urethroplasty**. Short bulbar strictures only; avoid in the penile urethra (pain, erection problems).
- Detrusor-sphincter dyssynergia (DSD) in spinal cord injury**: sphincteric stent (UroLume, Memokath) as a reversible alternative to **external sphincterotomy**.
- Others: palliation of obstruction from prostate cancer, post-brachytherapy and post-radiation stenosis (limited).

Technique

- Done under local or regional anaesthesia: urethroscopy, measurement of stricture length, placement under endoscopic or fluoroscopic guidance.
- Memokath expands at about 55 °C and softens when cooled for removal.
- The stent must not cross the external sphincter in a stricture patient, to avoid incontinence. Bulbar stents start distal to the sphincter.

Complications

Complication	Comment
Migration	Commonest with temporary stents
Encrustation, stones	Especially in prostatic stents

Complication	Comment
Hyperplastic tissue ingrowth / restenosis	Major problem with UroLume (end ingrowth); this needs resection
UTI, haematuria	
Incontinence, post-void dribbling	Especially if the stent crosses the sphincter
Perineal pain, painful erections	Bulbar stents
Difficult removal	Epithelialised stents; may need open urethroplasty, which becomes more complex

Results

- **Prostatic stents** in unfit men: most become catheter-free in the short term, but a significant proportion need removal for migration, encrustation or symptoms.
- **UroLume for strictures:**
 - early results were good
 - long-term follow-up showed high rates of **recurrent obstruction from tissue hyperplasia**
 - the need for repeat procedures, and the difficult salvage urethroplasty, led to its abandonment
- **DSD sphincteric stents:** comparable to sphincterotomy in the short term. Later migration and ingrowth are common.

Place in current guidelines

- **AUA Urethral Stricture guideline (2016, amended 2023): urethral stents should not be placed in patients who are considering urethroplasty.** A stent makes later reconstruction harder.
- **EAU Urethral Stricture guideline:** stents are **not recommended as routine** therapy. They are reserved for selected unfit patients.
- Urethroplasty remains the standard for recurrent strictures.

Recent advances

- **Optilume drug-coated balloon** (paclitaxel-coated urethral balloon; FDA 2021) for recurrent anterior strictures of 3 cm or less.
 - **ROBUST III RCT (Elliott, J Urol 2022):** much higher stricture-free rates than DVIU or dilation at 6 months (about 75% vs 27%).
 - It is a balloon, not a stent.
- **iTIND** (temporary implantable nitinol device; FDA 2020): placed in the prostatic urethra for 5–7 days, remodelling the bladder neck and prostatic urethra by ischaemic incision, then removed. It is a MIST option for BPH.
- **Zenflow Spring:** an investigational prostatic implant for BPH.
- **Allium** covered stents and biodegradable stents are under evaluation.

Diagram to draw: sagittal urethra showing a prostatic stent positioned between the bladder neck and verumontanum, and a bulbar stent distal to the external sphincter.

Key points to remember: - Two types: temporary (Memokath nitinol thermo-expandable, Allium) and permanent epithelialising (UroLume, now withdrawn). - Indications: unfit BPH patients, unfit bulbar stricture patients, DSD as an alternative to sphincterotomy. - Main problems: migration, encrustation, hyperplastic ingrowth. - AUA: do not place a stent in a man considering urethroplasty. - Optilume (ROBUST III) is the recent advance for recurrent short strictures.

IV-3e. Studer Neobladder

Type: SAQ (10 marks) | ~550 words

Introduction

- The Studer neobladder is an **orthotopic ileal low-pressure reservoir** anastomosed to the urethra. It was developed by **Urs E. Studer (Bern, Switzerland)** in the mid-1980s and first published in the late 1980s.
- Campbell: Orthotopic urinary diversion chapter.

Principles

- **Detubularisation:** interrupts coordinated peristaltic contractions, giving **low pressure**.
- **Spherical reconfiguration** (Laplace's law, $P = 2T/r$): the largest volume for the surface area gives **high capacity at low pressure**.
- An **isoperistaltic afferent tubular limb** works as an antireflux mechanism and allows a tension-free ureteric anastomosis with ureters of different lengths.
- The pouch lies orthotopically and the patient **voids by abdominal straining (Valsalva) with pelvic floor relaxation**, maintaining continence through the intact **rhabdosphincter**.

Technique (Campbell)

1. Isolate **54-56 cm of terminal ileum**, ending about **25 cm proximal to the ileocaecal valve** (to preserve bile-salt and B12 absorption).
2. Restore bowel continuity.
3. **Distal ~40-44 cm:** opened along the **antimesenteric border**, arranged in a **U**. The medial borders are sutured to form the back wall.
4. **Proximal ~12-14 cm:** left **tubular** as the **isoperistaltic afferent (chimney) limb**.
5. **Ureteroileal anastomosis:** end-to-side, refluxing (Nesbit technique) of each ureter onto the afferent limb, over stents (Wallace anastomosis is an alternative).
6. Fold the U-shaped plate (the bottom of the U folded up between the arms) to make a **spherical reservoir**. The **most dependent point** gets a small opening for the **urethral anastomosis** (6 sutures), tension-free.
7. Stents and a suprapubic/urethral catheter go in. The initial capacity is about 120-150 mL, increasing to **400-500 mL** over months.

Indications and contraindications

- **Indications:** radical cystectomy (MIBC, high-risk NMIBC) in a motivated patient with a **negative urethral margin** (frozen section at the apical urethra), and in selected women (bladder neck free of tumour).
- **Contraindications:** **impaired renal function** (commonly quoted thresholds: eGFR below ~35-50 mL/min or serum creatinine >2 mg/dL), **hepatic dysfunction**, **positive urethral margin**, IBD/short bowel/radiation enteritis, inability to perform **CISC** or poor compliance, incompetent external sphincter.

Postoperative care

- Daily catheter **irrigation** for mucus; **sodium bicarbonate** for acidosis; catheter out after a cystogram (~10-14 days).
- **Voiding training:** timed voiding every 2 h, extended to 3-4 h; sit, relax the pelvic floor and use Valsalva.

- **Nocturnal alarm** (void once or twice at night) to prevent enuresis and overdistension. Pelvic floor exercises.

Outcomes

- **Daytime continence: about 90%** at 1 year.
- **Night-time continence: about 70-80%.** This improves over time but declines with age.
- **CISC** needed in a minority of men (under 10-20%). The need is **higher in women** (hypercontinence, which can be up to 20-40% in some series).

Complications

Early	Late
Urine leak, ileus, UTI, pyelonephritis	Hyperchloraemic metabolic acidosis (ileal absorption of NH_4Cl)
Mucus retention	Vitamin B12 deficiency (start screening at 3-5 years)
	Ureteroileal anastomotic stricture (~3-10%)
	Stones (in pouch: mucus, staples, infection)
	Incontinence (day/night) or hypercontinence/retention
	Pouch-vaginal fistula (women); urethral recurrence; bowel dysfunction (diarrhoea)

Comparison with other orthotopic neobladders

Neobladder	Key features
Studer	U-shaped detubularised pouch with an isoperistaltic afferent limb , refluxing ureteric anastomosis
Hautmann (W)	Longer ileal segment (~60-70 cm) folded into a W ; larger initial capacity; ureters implanted directly (or with chimneys)
Robotic intracorporeal (Karolinska, Wiklund)	Studer-type pouch built intracorporeally

Recent advances

- Fully **intracorporeal robotic Studer neobladder**; **ERAS** pathways.
- Nerve-sparing and pelvic-organ-sparing cystectomy in women to reduce hypercontinence.

Diagram to draw: (1) 54-56 cm ileum marked into a 12-14 cm afferent limb and 40-44 cm detubularised segment, 25 cm from the ileocaecal valve. (2) The final spherical pouch with the afferent limb receiving both ureters and the urethral anastomosis at the most dependent point.

Key points to remember: - Segment: 54-56 cm of ileum ending 25 cm proximal to the ICV. Distal 40-44 cm detubularised into a U; proximal 12-14 cm isoperistaltic afferent limb. - Principles: detubularisation plus spherical shape (Laplace) gives a low-pressure, high-capacity reservoir. - Continence about 90% by day and 70-80% at night. Women have more hypercontinence and need CISC more often. - Main metabolic issues: hyperchloraemic acidosis and B12 deficiency. Avoid with significantly impaired renal function (eGFR below ~35-50 mL/min).

IV-3f. Cryotherapy for carcinoma prostate

Type: SAQ (10 marks) | ~600 words

Definition

- Cryotherapy (cryoablation, cryosurgical ablation of the prostate, CSAP) is **image-guided transperineal freezing** of prostate tissue to cause **cell death**.
- Campbell: Ablative therapies for prostate cancer chapter (Focal therapy and cryotherapy).

Physical principle

- **Joule-Thomson effect:** a high-pressure gas expanding through a narrow orifice changes temperature.
 - **Argon** produces freezing to about **-185 °C** at the needle tip.
 - **Helium** produces warming and is used for **active thawing**.
- **Thermocouples** (apex, rectal wall, external sphincter, neurovascular bundles) monitor temperatures in real time.
- A **urethral warming catheter** protects the urethra from sloughing.

Generations

Generation	Features
1st (1960s-70s)	Liquid nitrogen , large probes, no imaging; high fistula/incontinence
2nd (1990s)	TRUS guidance, urethral warmer , thermocouples (Onik)
3rd	Argon/helium, thin 17-gauge needles , template, multiple thermocouples
4th	Computer planning, precise ice-ball shaping; enables focal / hemi-gland therapy with MRI fusion

Mechanisms of cell death

1. **Direct cellular injury: intracellular ice** (rapid freezing) ruptures membranes; extracellular ice causes osmotic dehydration.
 2. **Vascular injury:** endothelial damage, platelet aggregation and microthrombosis lead to **ischaemic necrosis**. This is the main effect at the margins.
 3. **Apoptosis** at the periphery of the ice ball.
 4. **Immunological effect:** release of tumour antigens may produce a "cryo-immunologic" response.
- **Critical parameters:** target **-40 °C**; **rapid freeze, slow thaw; two freeze-thaw cycles** (the second enlarges the lethal zone).
 - On TRUS the ice-ball edge (~0 °C) is a hyperechoic rim; the lethal isotherm lies inside it.

Indications

1. **Salvage cryoablation after radiation failure:** the **most important current role**.
 - Requires biopsy-proven local recurrence, no metastases on PSMA PET/CT or MRI, and preferably PSA <10 ng/mL.
2. **Primary whole-gland therapy:** low- to intermediate-risk disease in men unfit or unwilling for RP/RT, or with a small gland (<40-50 mL; larger glands need ADT downsizing).
 - **AUA** lists whole-gland cryosurgery as an option. **EAU** does not recommend it as standard.

3. **Focal / hemi-gland therapy:** MRI-visible unilateral intermediate-risk lesion, within a registry or trial.

Technique

- TRUS-guided transperineal template placement of cryoneedles, thermocouples and urethral warmer; freeze anterior to posterior to -40°C , helium thaw, repeat cycle.

Outcomes

- Low PSA nadir predicts success; failure is usually judged by the **Phoenix definition (nadir + 2 ng/mL)** (not validated specifically for cryotherapy).
- **Donnelly et al., Cancer 2010** (Canadian RCT, cryoablation vs external beam RT): cryotherapy was **non-inferior** for biochemical progression, with fewer positive 36-month biopsies but worse sexual function.
- **Salvage cryotherapy:** 5-year biochemical disease-free survival is reported at roughly 50-60% in selected patients.

Complications

Complication	Primary whole gland	Salvage
Erectile dysfunction	~ 80-90% (focal: much less)	Very high
Urinary incontinence	~1-5%	~5-20%
Rectourethral fistula	<0.5-1%	~1-3%
Urethral sloughing	<5% with a warmer	Higher
Urinary retention	~2-5%	
Perineal pain, penile numbness, scrotal oedema	Common, transient	

Comparison with other ablative energies

	Cryo	HIFU	IRE	Focal laser (FLA)
Energy	Freezing (Joule-Thomson)	Focused ultrasound heat ($>60^{\circ}\text{C}$)	Electric pulses $\rightarrow$ nanopores (non-thermal)	Thermal, MRI-guided
Access	Transperineal	Transrectal	Transperineal	Transperineal / transrectal
Notes	Anterior tumours treatable	AP height limit (~4 cm)	Spares vessels/nerves	Small lesions

Guideline positions

- **EAU 2024/2025:** primary whole-gland or focal cryotherapy/HIFU **only within trials or registries**. Salvage cryotherapy is an option for selected local recurrence after RT in experienced centres.
- **AUA/ASTRO 2022:** whole-gland cryosurgery may be considered for low/intermediate-risk disease. Focal therapy carries the caveat of limited long-term data.
- **NCCN:** cryotherapy is listed among **local salvage options** after RT failure.

Recent advances

- MRI-TRUS fusion **focal/hemi-gland cryotherapy**; **PSMA PET** to select salvage patients; cryo plus immunotherapy under study.

Diagram to draw: sagittal prostate with transperineal cryoneedles, urethral warmer, thermocouples at the apex and rectal wall, and the ice ball with the -40°C isotherm inside the 0°C edge.

Key points to remember: - Argon freezes and helium thaws (Joule-Thomson). Target -40°C , with two freeze-thaw cycles. - Cell death comes from intracellular ice, vascular ischaemia (main), apoptosis and an immunological effect. - The main role is salvage after radiation failure. ED is about 80-90% after whole-gland treatment. Rectourethral fistula is under 1% in primary treatment and higher in salvage. - EAU: primary or focal cryotherapy only within trials or registries. Phoenix definition = nadir + 2.

Rapid Revision Sheet - Paper IV

All the key points from this paper in one place. Read this in the last 30 minutes before the exam.

IV-1. Role of robotics in urology (LAQ)

- PUMA 560 (1985) was the first surgical robot. PROBOT (Imperial College) was the first urological robot. da Vinci was FDA-approved in 2000. The first RARP was by Binder & Kramer in 2000.
- Yaxley (Lancet 2016) and Coughlin (Lancet Oncol 2018) found no functional difference between open and robotic RP at 12 weeks and 24 months.
- RAZOR (Lancet 2018): RARC non-inferior for 2-year PFS. iROC (JAMA 2022): intracorporeal RARC gave more days alive and out of hospital.
- Generations: S, Si, Xi (2014), SP (2018), da Vinci 5 (2024). Alternatives: Hugo, Versius, Senhance, SSI Mantra (India).
- Steep Trendelenburg causes raised IOP/ICP and ischaemic optic neuropathy; prolonged lithotomy causes compartment syndrome.

IV-2. Discuss marginal kidney donor and describe newer immunosuppressant drugs in renal transplantation (LAQ)

- ECD (UNOS 2002): age ≥ 60 , or 50–59 with ≥ 2 of HTN, creatinine >1.5 mg/dL, CVA death. Replaced by KDPI $>85\%$ in December 2014.
- Remuzzi score: 0–3 single, 4–6 dual, ≥ 7 discard. HMP reduces DGF (Moers, NEJM 2009). Eurotransplant Senior Program: ≥ 65 to ≥ 65 .
- Belatacept (CTLA4-Ig): BENEFIT/BENEFIT-EXT show better GFR. Contraindicated in EBV-seronegative recipients (PTLD).
- TRANSFORM: everolimus + low-dose CNI is non-inferior, with fewer CMV/BK infections.
- Imlifidase (IgG-cleaving enzyme) has EMA conditional approval (2020) for highly sensitised recipients. Voclosporin is for lupus nephritis, not transplantation.

IV-3a. Etiology and management of hypospadias fistula (SAQ)

- Fistula is the commonest complication. TIP distal rate is under 5–10%; proximal repairs carry a higher rate.
- Distal obstruction (meatal stenosis) must always be excluded and corrected.
- Wait at least 6 months, then do a layered closure with a barrier (dartos or tunica vaginalis) and non-overlapping suture lines.
- Coronal fistula equals glans dehiscence and needs redo glanuloplasty.

IV-3b. Silodosin (SAQ)

- Selectivity $\alpha 1A : \alpha 1B$ is about 162 : 1. Dose 8 mg OD with a meal; 4 mg if CrCl 30–50; contraindicated if CrCl <30 , Child-Pugh C, or with strong CYP3A4 inhibitors.
- Commonest adverse effect is anejaculation/retrograde ejaculation ($\sim 28\%$). Lowest hypotension risk of the α -blockers.
- Chapple 2011: non-inferior to tamsulosin and also reduced nocturia.
- Warn about IFIS before cataract surgery.

IV-3c. Sacral neuromodulation (SAQ)

- **Rechargeable, miniaturised IPGs:**
 - **Axonics r-SNM** (FDA 2019; about 15-year life)
 - **Medtronic InterStim Micro** (rechargeable)
 - recharge-free long-life devices (**InterStim X, Axonics F15**)

- **Full-body MRI-conditional** systems (1.5 T and 3 T with SureScan/compatible leads), which solve the main historical limitation of SNM.
- **Standardised tined-lead placement** with curved stylet and fluoroscopic landmarks (Matzel), improving test-phase success and reducing reoperation.
- **Closed-loop / sensing SNM** using evoked compound action potential (ECAP) recordings to auto-adjust stimulation is at the research stage.
- **Implantable tibial nerve stimulators: eCoin** (FDA 2022), **Revi** (BlueWind, FDA 2023) give less-invasive alternatives for refractory OAB/UUI.

IV-3d. Urethral stents (SAQ)

- Two types: temporary (Memokath nitinol thermo-expandable, Allium) and permanent epithelialising (UroLume, now withdrawn).
- Indications: unfit BPH patients, unfit bulbar stricture patients, DSD as an alternative to sphincterotomy.
- Main problems: migration, encrustation, hyperplastic ingrowth.
- AUA: do not place a stent in a man considering urethroplasty.
- Optilume (ROBUST III) is the recent advance for recurrent short strictures.

IV-3e. Studer Neobladder (SAQ)

- Segment: 54–56 cm of ileum ending 25 cm proximal to the ICV. Distal 40–44 cm detubularised into a U; proximal 12–14 cm isoperistaltic afferent limb.
- Principles: detubularisation plus spherical shape (Laplace) gives a low-pressure, high-capacity reservoir.
- Continence about 90% by day and 70–80% at night. Women have more hypercontinence and need CISC more often.
- Main metabolic issues: hyperchloraemic acidosis and B12 deficiency. Avoid with significantly impaired renal function (eGFR below ~35–50 mL/min).

IV-3f. Cryotherapy for carcinoma prostate (SAQ)

- Argon freezes and helium thaws (Joule–Thomson). Target -40°C , with two freeze–thaw cycles.
- Cell death comes from intracellular ice, vascular ischaemia (main), apoptosis and an immunological effect.
- The main role is salvage after radiation failure. ED is about 80–90% after whole-gland treatment. Rectourethral fistula is under 1% in primary treatment and higher in salvage.
- EAU: primary or focal cryotherapy only within trials or registries. Phoenix definition = nadir + 2.