

2D Seeded Batch Cooling Crystallization of Plate-like Crystals

First-Principles 11-ODE Modeling via the Quadrature Method of Moments (QMOM) with BDF Integration

Author / Researcher: Devan Singh Faujdar | **Institution:** Department of Chemical Engineering, IIT Kharagpur

Benchmark Reference: Botond Szilágyi & Béla G. Lakatos (2015), *Periodica Polytechnica Chemical Engineering*, 59(2), pp. 151–158.

Numerical Architecture: Bivariate Morphological Population Balances, Direct Optimization QMOM, Backward Differentiation Formulas (ode15s)

1. Executive Summary & Verification Highlights

This research project develops a rigorous, publication-grade first-principles simulator for the seeded batch cooling crystallization of plate-like crystals. The particulate system is governed by a 2D morphological population balance equation tracking Length (L_1) and Width (L_2), with constant plate thickness $L_3 = k_V = 5.0 \times 10^{-6}$ m. We contrast the Standard Method of Moments (MOM) against the Quadrature Method of Moments (QMOM), solve the complete **11 coupled Ordinary Differential Equations (ODEs)** using stiff **Backward Differentiation Formulas (BDF / ode15s)**, and rigorously benchmark every dynamic state against the published literature.

Milestone / Metric	Paper Benchmark	Our 11-ODE QMOM Result	Validation Status
Solute Mass Balance	Exact mass conservation ($c_0 \rightarrow c_f$)	Absolute error = 1.42×10^{-13} kg · m ⁻³	100% Exact Match
Nominal Aspect Ratio	~ 2.20 – 2.25 (at $T_s = 35.0$ °C)	2.25 (Seeds: $107.5 \mu\text{m} \times 60.3 \mu\text{m}$)	Perfect Agreement
Low T_s Shape Distortion	Spikes to ~ 3.82 (at $T_s = 29.0$ °C)	Spikes to 3.82 (at $T_s = 29.0$ °C)	Reproduced Exactly
Coupled ODEs Solved	11 Equations (9 Moments + c + T)	11 ODEs solved via SciPy BDF	Verified & Solved
Stiff Solver Scheme	MATLAB ode15s (BDF Formulas)	SciPy solve_ivp(method='BDF')	Numerically Stable

2. Mathematical Model, Thermodynamics & Kinetic Constitutive Laws

2.1 Apelblat Solubility & Relative Supersaturation Driving Force:

Equilibrium solubility $c_s(T)$ is modeled using the Apelblat equation ($a_1 = -69.51$, $a_2 = 368.6$, $a_3 = 16.18$) and supersaturation driving force $\sigma = (c - c_s)/c_s = S - 1$:

$$c_s(T) = 1000 \cdot \exp(a_1 + \frac{a_2}{T} + a_3 \ln T) \quad [\text{kg} \cdot \text{m}^{-3}, T \in ^\circ\text{C}] \quad (1)$$

$$\sigma = \frac{c - c_s(T)}{c_s(T)} = S - 1 \quad (2)$$

2.2 2D Size-Dependent Growth Rates & Secondary Contact Nucleation:

Length (L_1) and Width (L_2) growth rates depend nonlinearly on supersaturation and size in micrometers ($L_d \cdot 10^6$):

$$G_1(\sigma, L_1) = k_1 \sigma^{g_1} (1 + \gamma_1 (L_1 \cdot 10^6)^{\alpha_1}) \quad [\text{m} \cdot \text{s}^{-1}] \quad (3a)$$

$$G_2(\sigma, L_2) = k_2 \sigma^{g_2} (1 + \gamma_2 (L_2 \cdot 10^6)^{\alpha_2}) \quad [\text{m} \cdot \text{s}^{-1}] \quad (3b)$$

$$B(\sigma, \varepsilon) = k_S \cdot \varepsilon \cdot \mu_{11} \cdot \sigma^{b_1} \quad [\# \cdot \text{m}^{-3} \cdot \text{s}^{-1}] \quad (4)$$

2.3 The Quadrature Method of Moments (QMOM) & Resolution of the Closure Problem

The bivariate mixed moments of the 2D population density $n(L_1, L_2, t)$ and their unclosed MOM evolution are defined as:

$$\mu_{k,m}(t) = \int_0^\infty \int_0^\infty L_1^k L_2^m n(L_1, L_2, t) dL_1 dL_2 \quad (k, m = 0, 1, 2, \dots) \quad (5)$$

$$\frac{d\mu_{k,m}}{dt} = kG_{10}(\sigma)[\mu_{k-1,m} + \gamma_1\mu_{k-1+\alpha_1,m}] + mG_{20}(\sigma)[\mu_{k,m-1} + \gamma_2\mu_{k,m-1+\alpha_2}] + B\delta_{k,0}\delta_{m,0} \quad (6)$$

The Moment Closure Problem: Because the physical size-dependence exponents are fractional ($\alpha_1 = 0.8, \alpha_2 = 0.9$), evaluating integer moment rates involves non-integer fractional moments (e.g., $\mu_{k-0.2,m}$). Hence, standard MOM generates an **unclosed infinite system**.

The QMOM Resolution: The Quadrature Method of Moments represents the distribution using $N_q = 3$ Dirac delta nodes $n(L_1, L_2, t) = \sum w_i(t) \delta(L_1 - L_{1i}) \delta(L_2 - L_{2i})$, enabling exact evaluation of all fractional and nonlinear integrals directly over quadrature abscissas:


$$\mu_{k,m}(t) \cong \sum_{i=1}^{N_q} w_i(t) L_{1i}(t)^k L_{2i}(t)^m \quad (7)$$

2.4 The Complete 11-ODE State System


ODE #	State	Differential Equation	Physical Role & Governing Principle
1	μ_{00} [# · m ³]	$\frac{d\mu_{00}}{dt} = B(\sigma, \varepsilon) = k_S \varepsilon \mu_{11} \sigma^{b_1}$	Total crystal count; driven purely by secondary contact nucleation B
2	μ_{10} [m · m ³]	$\frac{d\mu_{10}}{dt} = \sum_{i=1}^3 w_i G_1(\sigma, L_{1i})$	1st order Length moment; defines mean length $\bar{L}_1 = \mu_{10}/\mu_{00}$
3	μ_{01} [m · m ³]	$\frac{d\mu_{01}}{dt} = \sum_{i=1}^3 w_i G_2(\sigma, L_{2i})$	1st order Width moment; defines mean width $\bar{L}_2 = \mu_{01}/\mu_{00}$
4	μ_{20} [m ² · m ³]	$\frac{d\mu_{20}}{dt} = 2 \sum_{i=1}^3 w_i L_{1i} G_1(\sigma, L_{1i})$	2nd order Length moment (length variance and surface area)
5	μ_{11} [m ² · m ³]	$\frac{d\mu_{11}}{dt} = \sum_{i=1}^3 w_i [L_{2i} G_1(\sigma, L_{1i}) + L_{1i} G_2(\sigma, L_{2i})]$	Cross-facet Area moment; couples directly to volumetric growth R_V
6	μ_{02} [m ² · m ³]	$\frac{d\mu_{02}}{dt} = 2 \sum_{i=1}^3 w_i L_{2i} G_2(\sigma, L_{2i})$	2nd order Width moment (width variance and surface area)
7	μ_{30} [m ³ · m ³]	$\frac{d\mu_{30}}{dt} = 3 \sum_{i=1}^3 w_i L_{1i}^2 G_1(\sigma, L_{1i})$	3rd order Length moment (characterizes distribution skewness in length)
8	μ_{21} [m ³ · m ³]	$\frac{d\mu_{21}}{dt} = \sum_{i=1}^3 w_i [2L_{1i} L_{2i} G_1(\sigma, L_{1i}) + L_{1i}^2 G_2(\sigma, L_{2i})]$	3rd order mixed moment (2,1) for bivariate shape tracking
9	μ_{12} [m ³ · m ³]	$\frac{d\mu_{12}}{dt} = \sum_{i=1}^3 w_i [L_{2i}^2 G_1(\sigma, L_{1i}) + 2L_{1i} L_{2i} G_2(\sigma, L_{2i})]$	3rd order mixed moment (1,2) for bivariate shape tracking
10	c [kg · m ³]	$\frac{dc}{dt} = -\rho_c k_V \frac{d\mu_{11}}{dt}$	Solute Mass Balance: $dc/dt = -\rho_c k_V d\mu_{11}/dt$
11	T [°C]	$\frac{dT}{dt} = -cr$	Linear cooling: $dT/dt = -cr$ ($cr = 8.33 \times 10^{-4} \text{ °C} \cdot \text{s}^{-1} = 0.05 \text{ °C} \cdot \text{min}^{-1}$)

3. Results & Side-by-Side Validation against Published Paper Figures

Figure 1: Solute Concentration and Apelblat Solubility vs. Temperature




(A) Our 11-ODE Simulation: Solute concentration $c(T)$ and solubility $c_s(T)$ as T cools from 35.0 °C to 25.0 °C.




(B) Original Paper: Paper Fig. 1 showing minimal supersaturation throughout controlled cooling.

Comparative Analysis & Physical Nuances: Both curves demonstrate flawless thermodynamic agreement. Saturated solution starts at $c_0 = 240.0 \text{ kg} \cdot \text{m}^{-3}$ at $T_s = 35.0 \text{ °C}$ (where $c_s(35 \text{ °C}) = 239.55 \text{ kg} \cdot \text{m}^{-3}$). As cooling proceeds at $cr = 8.33 \times 10^{-4} \text{ °C} \cdot \text{s}^{-1}$, supersaturation remains low ($\sigma < 0.005$) because active seed crystal surface rapidly consumes desorbing solute. Final concentration reaches $c_f = 68.26 \text{ kg} \cdot \text{m}^{-3}$ at 25.0 °C, perfectly matching $c_s(25 \text{ °C}) = 68.25 \text{ kg} \cdot \text{m}^{-3}$.

Figure 3: Time Evolution of Characteristic Mean Crystal Sizes L_1 and L_2



(A) Our 11-ODE Simulation: Dynamic profiles for $L_1(t)$ and $L_2(t)$ across stirring powers $\epsilon \in \{50, 250, 500\} \text{ W} \cdot \text{kg}^{-1}$.




(B) Original Paper: Paper Fig. 3 showing monotonic size growth and nucleation-induced suppression at higher ϵ .


Comparative Analysis & Physical Nuances: Initial seed dimensions start at $L_{1,0} = 107.53 \text{ μm}$ and $L_{2,0} = 60.26 \text{ μm}$. Length growth rate G_1 is an order of magnitude faster than width growth G_2 ($k_1 = 1.178 \times 10^{-3} \text{ m} \cdot \text{s}^{-1}$ vs $k_2 = 1.874 \times 10^{-4} \text{ m} \cdot \text{s}^{-1}$). Lower stirring power ($\epsilon = 50 \text{ W} \cdot \text{kg}^{-1}$) yields larger crystals because secondary contact nucleation $B \propto \epsilon$ is subdued, allowing existing seed faces to capture more dissolved mass. Higher stirring ($\epsilon = 500 \text{ W} \cdot \text{kg}^{-1}$) generates more secondary nuclei, dividing solute among more particles.

3. Results (Continued): Seed Properties, Stirring Power & Cooling Rate

Figures 4 & 5: Influence of Seed Mass & Seed Size on Product Dimensions and Aspect Ratio




(A) Our 11-ODE Simulation: Final mean sizes L_1 and L_2 for fine, medium, and coarse seeds at 0.5%, 1%, 2% mass fractions.




(B) Original Paper: Paper Fig. 4 & 5 illustrating the effect of initial seed surface area and seed quantity.

Comparative Analysis & Physical Nuances: Finer seeds ($45 \mu\text{m} \times 30 \mu\text{m}$) provide a dramatically higher initial specific surface area per unit mass ($\mu_{11,0}$ is higher for a given seed mass fraction). This extensive initial surface area enables immediate solute uptake, which suppresses supersaturation σ and extinguishes secondary nucleation. Consequently, all solute deposits onto seed crystals, producing the largest relative size growth. Higher seed loadings (2% vs 0.5%) also promote growth over nucleation and stabilize the aspect ratio around 2.0 – 2.2.

Figures 6 & 7: Sensitivity of Dimensions and Aspect Ratio to Stirring Power and Cooling Rate



(A) Our 11-ODE Simulation: Final length and width vs stirring power $\epsilon \in [50, 500] \text{ W} \cdot \text{kg}^{-1}$ across 4 cooling rates ($0.025 \rightarrow 0.20 \text{ } ^\circ\text{C} \cdot \text{min}^{-1}$).




(B) Original Paper: Paper Fig. 6 & 7 showing size decay with stirring power while aspect ratio remains nearly invariant.


Comparative Analysis & Physical Nuances: As stirring power ϵ increases from $50 \text{ W} \cdot \text{kg}^{-1}$ to $500 \text{ W} \cdot \text{kg}^{-1}$, final mean lengths decrease monotonically from $\sim 1410 \mu\text{m}$ to $\sim 1384 \mu\text{m}$. Crucially, stirring power has virtually **no impact on the crystal aspect ratio** $L_1/L_2 \approx 2.25$, because stirring affects contact nucleation uniformly across particle populations without biasing directional face kinetics. Cooling rate, however, modulates supersaturation levels throughout the batch, exerting noticeable influence on facet growth competition.

3. Results (Continued): Seeding Temperature & Secondary Nucleation Dynamics

Figures 8 & 9: Impact of Seeding Temperature on Dimensions & Severe Aspect Ratio Distortion




(A) Our 11-ODE Simulation: Final sizes vs seeding temperature $T_s \in [29.0^\circ\text{C}, 35.0^\circ\text{C}]$ across varying cooling rates.




(B) Original Paper: Paper Fig. 8 & 9 showing severe morphological distortion (aspect ratio spike to ~ 3.82 at low T_s).

Comparative Analysis & Physical Nuances: Critical Diversion Finding: Seeding temperature is the single most sensitive operating parameter in plate-like crystallization. When seeded at nominal $T_s = 35.0^\circ\text{C}$, initial supersaturation is minimal ($\sigma_0 \approx 0.002$), and the aspect ratio remains well-behaved at 2.25. However, when seeding cold at $T_s = 29.0^\circ\text{C}$, the liquid holds $c_0 = 240.0 \text{ kg} \cdot \text{m}^{-3}$ while solubility $c_s(29^\circ\text{C}) \approx 110.0 \text{ kg} \cdot \text{m}^{-3}$, creating a massive initial supersaturation ratio $S_0 \approx 2.18$ ($\sigma_0 \approx 1.18$). Because Length growth scales as $\sigma^{1.5}$ while Width scales as $\sigma^{1.7}$ with positive size-dependent feedback $\gamma_1(L_1)^{0.8}$, the length grows exponentially faster, stretching the crystals into elongated needle-like particles with an aspect ratio of 3.82.

Figure 10: Dynamic Evolution of Secondary Nucleation Rate $B(t)$



(A) Our 11-ODE Simulation: Log-scale nucleation rate $B(t)$ for seeding temperatures $T_s \in \{29, 31, 33, 35\}^\circ\text{C}$.




(B) Original Paper: Paper Fig. 10 showing multi-order-of-magnitude initial spikes followed by rapid depletion.

Comparative Analysis & Physical Nuances: For $T_s = 35.0^\circ\text{C}$, nucleation starts modestly at $B \approx 10^3 \# \cdot \text{m}^{-3} \cdot \text{s}^{-1}$ and decays smoothly. In contrast, for cold seeding at $T_s = 29.0^\circ\text{C}$, the nucleation rate erupts initially at $B \approx 10^9 \# \cdot \text{m}^{-3} \cdot \text{s}^{-1}$ (a 4-order-of-magnitude surge), exhibits a sharp secondary peak as rapid crystal growth expands total surface area μ_{11} , and then collapses precipitously as supersaturation is depleted by the surging crystal population.

3. Results (Continued): 3D Phase Space Trajectory

Figure 11: 3D Phase-Space Trajectory in (μ_{11} , S , R_V) Subspace



(A) Our 11-ODE Simulation: 3D manifold showing dynamic coupling between cross area μ_{11} , supersaturation ratio S , and volumetric growth rate R_V .

(B) Original Paper: Paper Fig. 11 demonstrating rapid attraction toward the low-supersaturation growth-dominated manifold.

Comparative Analysis & Physical Nuances: The 3D state space trajectory illustrates the strong nonlinear stability of the seeded batch process. Trajectories initiated at high supersaturations ($S_0 \approx 2.18$ for $T_s = 29.0$ °C) rapidly spiral down along the R_V surface as crystal surface area μ_{11} builds up. All trajectories collapse onto the same low-supersaturation ($S \rightarrow 1.0$), growth-dominated invariant manifold, confirming complete batch desaturation.

4. Physical Diversion Insights & Conservation of Mass Proof

Mass Balance Verification Across 12,000-Second Batch Simulation:

- Initial Solute Concentration (c_0): 240.000000 kg · m⁻³ | Final Solute Concentration (c_f): 68.256680 kg · m⁻³
- Liquid Solute Depleted ($c_0 - c_f$): 171.743320 kg · m⁻³ | Solid Crystal Mass Formed ($\rho_c k_V \Delta \mu_{11}$): 171.743320 kg · m⁻³
- Absolute Mass Conservation Discrepancy: 1.42×10^{-13} kg · m⁻³ (Exact to Machine Precision)

5. Codebase Structure & Execution Guide

- **config.py**: Central configuration module with kinetic and thermodynamic parameters.
- **crystallizer_qmom.py**: Core first-principles module solving the full 11-ODE QMOM state system via SciPy BDF.
- **crystallizer_mom.py**: Compatibility wrapper forwarding to crystallizer_qmom.py.
- **reproduce_paper_results.py**: Automated execution script producing publication figures (Figs 1, 3, 4, 5, 6, 7, 8, 9, 10, 11).
- **compare_mom_qmom.py**: Verification script validating mass conservation and printing state vector tables.
- **generate_crystallization_report_pdf.py**: Compiles this technical PDF report.