

# Market Segmentation

In marketing, the purpose is to align the needs and desires of consumers with the offerings of suppliers. This matching process is crucial for both consumers and suppliers and forms the basis of marketing planning. Marketing planning involves a sequence of activities that lead to the establishment of marketing objectives and the formulation of plans to achieve them.

A marketing plan consists of two main components: a strategic marketing plan and a tactical marketing plan. The strategic plan provides the long-term direction for the organization and sets out the overall goals and objectives. It focuses on determining where the organization wants to go and why. On the other hand, the tactical plan translates the strategic plan into specific instructions for short-term marketing actions. It outlines the detailed steps and decisions needed to reach the desired goals.

The strategic marketing plan involves identifying consumer needs and desires, assessing the organization's strengths and weaknesses, and examining external opportunities and threats. This analysis often includes a SWOT analysis (strengths, weaknesses, opportunities, and threats) to understand the organization's position in the market.

- market research plays a crucial role in understanding consumer needs and desires. It utilizes various sources of information, such as surveys, focus groups, interviews, observations, and experiments, to gain insights into consumer preferences.
- Once the strategic marketing plan is established, the organization focuses on two key decisions: segmentation and targeting of specific consumer groups and creating a favorable market positioning.
- In summary, strategic and tactical marketing are both essential for organizational success. The strategic plan provides the overall direction, while the tactical plan translates the strategy into actionable steps.

Market segmentation is a crucial decision-making tool for marketing managers that involves dividing a heterogeneous market into smaller, homogeneous segments based on consumer characteristics. It is an essential aspect of strategic marketing and plays a significant role in driving marketing success. Market segmentation helps in selecting a target market for a product and designing an appropriate marketing mix. By understanding the differences between consumers and aligning organizational strengths with consumer needs, market segmentation can lead to improved competitive advantage and long-term success. It offers benefits such as a better understanding of consumers, improved match of organizational strengths and consumer needs, and the potential for market dominance. Effective market segmentation allows for a more efficient allocation of resources, higher return on investment, and targeted sales efforts. Additionally, it can contribute to team building and improved communication within an organization.

## Market Segmentation Analysis

Market segmentation analysis involves grouping consumers into segments based on their shared product preferences or characteristics. It is a statistical and exploratory process that requires the involvement of a

competent data analyst and a user who understands the organization's mission. The analysis consists of three layers: the core layer focuses on extracting market segments, the second layer involves tasks such as collecting good data, exploring data, and profiling segments, and the third layer includes non-technical tasks related to organizational implementation.

Approaches to market segmentation analysis can be systematized in different ways. One approach, proposed by Dibb and Simkin, is based on organizational constraints. It distinguishes three approaches: quantitative survey-based segmentation, segment evolution, and segment revolution. The quantitative survey-based approach assumes a willingness to start from scratch and develop a new marketing plan based on the segmentation findings. Segment evolution involves refining and sharpening existing segment focus, while the emergence of segments from qualitative research or exploratory research represents a more random discovery.

In summary, market segmentation analysis is a multi-layered process that requires statistical analysis, data exploration, and user-driven decision-making. The choice of approach depends on organizational constraints and the desired level of change in marketing strategies.

In market segmentation analysis, the choice of segmentation variables plays a crucial role. Segmentation variables are consumer characteristics that are used to extract market segments. There are two types of segmentation variables: unidimensional and multidimensional.

- Unidimensional segmentation variables involve using a single piece of information about consumers to create segments. For example, age can be used as a unidimensional segmentation variable, resulting in age groups as market segments. Older consumers can then be selected as a target segment. Other examples of unidimensional segmentation variables include gender, country of origin, and prior purchase behavior.
- Multidimensional segmentation variables involve using multiple pieces of information about consumers. This creates a multidimensional statistical problem. For instance, expenditure patterns can be used as multidimensional segmentation variables. For a tourist destination, analyzing tourists' expenditure patterns on various vacation activities like theme parks, dining out, and shopping can help identify target segments.

In reality, naturally existing and well-separated segments are rare. This has led to the recognition of different conceptual approaches to data-driven segmentation: natural segmentation, reproducible segmentation, and constructive segmentation.

Natural segmentation assumes that distinct market segments exist in the data and the aim is to find them. Reproducible segmentation recognizes that natural segments may not exist, but the data contain some structure that allows for the generation of consistent segmentation solutions. Constructive segmentation deals with data where neither cluster structure nor any other structure exists. In this case, segmentation solutions are created even if they are random.

It is important to conduct data structure analysis before extracting segments to understand the nature of the data and make informed segmentation decisions. Data structure analysis helps identify whether natural segments, reproducible segments, or constructive segments can be extracted from the data.

Market segmentation analysis is a systematic process that involves dividing a market into distinct groups of consumers with similar characteristics and needs. This analysis helps organizations identify target segments and develop customized marketing strategies to effectively reach and serve those segments. The ten-step approach to market segmentation analysis includes the following:

1. Step 1: Deciding (not) to segment: Assessing the suitability of the market for segmentation and determining if the organization can commit to a long-term segmentation strategy.
2. Step 2: Specifying the ideal target segment: Defining the characteristics and attributes of the ideal market segment that the organization wants to target.
3. Step 3: Collecting data: Gathering relevant data on segmentation variables (e.g., demographics, behavior) and descriptor variables (e.g., preferences, needs) either through primary research or existing sources.
4. Step 4: Exploring data: Analyzing and preprocessing the collected data to gain insights and identify patterns or trends that can inform the segmentation process.
5. Step 5: Extracting segments: Using segmentation variables to split consumers into distinct segments. This can be achieved through various algorithms such as distance-based, model-based, or hybrid methods.
6. Step 6: Profiling segments: Identifying and understanding the key features and characteristics of each extracted market segment.
7. Step 7: Describing segments: Providing a detailed description of each segment, including their demographics, behavior, preferences, and other relevant attributes.
8. Step 8: Selecting the target segment(s): Evaluating the segments and choosing one or a few segments to focus on based on their attractiveness and compatibility with the organization's goals and resources.
9. Step 9: Customizing the marketing mix: Developing a tailored marketing mix (product, price, promotion, place) that addresses the specific needs and preferences of the selected target segment(s).
10. Step 10: Evaluation and monitoring: Assessing the success of implementing the segmentation strategy and continuously monitoring the segments for any changes in size or characteristics that may require adjustments to the strategy.

## Market Segmentation Analysis steps

Before pursuing a market segmentation strategy, organizations must understand the implications and commit to a long-term approach. Market segmentation requires substantial changes and investments, including research, data collection, product development, pricing and distribution adjustments, and tailored communication with target segments. The decision to pursue market segmentation should be made at the highest executive level and communicated across the organization.

Implementing market segmentation may face various barriers. Lack of leadership and involvement from senior management, inadequate allocation of resources, and resistance to change can hinder successful implementation. Organizational culture, such as a lack of market orientation, resistance to new ideas, poor communication, and short-term thinking, can also impede progress. Lack of understanding and training on market segmentation foundations, as well as the absence of a qualified marketing expert or data manager in the organization, pose additional obstacles. Objective restrictions like financial limitations or structural constraints can be challenging. Process-related barriers include undefined objectives, poor planning, lack of structured processes, unclear responsibilities, and time pressure. To overcome these barriers, it is crucial to remove them proactively or consider abandoning the segmentation strategy if they cannot be resolved.

To move forward with market segmentation, organizations need a resolute sense of purpose, dedication, patience, and a willingness to address encountered problems. Clear communication, understanding of the segmentation analysis, and graphical visualizations can facilitate the interpretation and acceptance of the results. By addressing and overcoming barriers, organizations can increase the likelihood of successful implementation and reap the benefits of market segmentation.

## Exploratory Data Analysis

We utilize a McDonalds dataset to demonstrate data exploration with actual data. There are 15 columns and more than 1000 rows in this data collection. The owner discovers whether the consumer enjoys the cuisine in the dataset. Likewise, the age group they prefer. Exploratory data analysis cleans, identifies, and corrects errors in the data after it has been collected. The handling of missing values, outliers, removing duplicate values, transforming categorical data to numerical data (Label Encoder, One-hot Encoding), etc. are all examples of errors. The most appropriate algorithm for extracting valuable market segments is also provided guidance during this exploratory stage. It aids in determining the variables' measurement levels, investigating each variable's univariate distribution, and evaluating the dependency patterns between variables. We examine the data using visualization after cleaning and transforming it.

Using graphs, maps, and other visual aids to illustrate the facts and data is helpful. Data professionals may easily comprehend any patterns, trends, or anomalies in a data set thanks to these visualizations. Histograms, Boxplots, Scatterplots, Pie Charts, Bar Charts, Line Charts, Area Charts, Heat Maps, and Tree Maps are some examples of visualization. These techniques help us visualize and comprehend the data.

## Collecting Data

This can be done using a variety of methods, but the most common are distance-based methods, model-based methods, and algorithms integrating variable selection with the task of extracting market segments. Distance-based methods use a distance measure to group consumers who are like each other. The most common distance measure is Euclidean distance, which measures the distance between two

points in Euclidean space. Other distance measures include Manhattan distance, Hamming distance, and Jaccard distance. Model-based methods use statistical models to group consumers. The most common model-based method is cluster analysis. Cluster analysis uses a variety of algorithms to group consumers into clusters. The most common cluster analysis algorithms are k-means clustering, hierarchical clustering, and density based clustering. Algorithms integrating variable selection with the task of extracting market segments use a combination of distance-based methods and model-based methods to extract market segments. These algorithms are typically used when the number of variables is large. The choice of which method to use depends on the data and the desired characteristics of the segments. Once the segments have been extracted, they need to be evaluated. This can be done by examining the characteristics of the segments and by comparing the segments to the ideal target segment. If the segments are not satisfactory, then the segmentation process may need to be repeated.

The passage discusses the nature of data-driven market segmentation analysis, which is exploratory and typically involves unstructured consumer data. The results of a segmentation analysis are determined by both the data and the segmentation algorithm chosen, and different algorithms can impose different structures on the extracted segments. The passage illustrates this point with an example of k-means cluster analysis and single linkage hierarchical clustering on a data set containing two spiraling segments. The passage also emphasizes the importance of selecting a suitable clustering method that matches the data analytic features of the resulting clustering with the context-dependent requirements desired by the researcher.

## Exploring Data

In this section, the importance of exploratory data analysis (EDA) in market segmentation is highlighted. EDA involves cleaning and preprocessing the data and identifying the measurement levels of the variables. It also involves investigating the distribution of each variable and assessing the dependency structures between variables. The results from EDA provide insights into the most suitable segmentation methods for extracting market segments. To illustrate EDA using real data, the authors use a travel motives dataset that contains 20 travel motives reported by 1000 Australian residents in relation to their last vacation. The data is available in a CSV file, which can be accessed through the R package MSA or downloaded from the book's website. The CSV file can be explored using a spreadsheet program before commencing analyses in R. In addition, missing values need to be handled appropriately. Missing values

can occur for a variety of reasons, including non-response, respondents not providing a valid answer, or technical problems during data collection. The most common approach to handling missing values is to impute them, which means replacing the missing values with plausible estimates based on the available data. There are several techniques for imputing missing values, including mean imputation, hot deck imputation, and multiple imputation. Overall, data cleaning is a crucial step in preparing data for analysis, as it ensures that the data is accurate, complete, and consistent. Without proper data cleaning, analysis results may be unreliable and misleading, which can have serious consequences for decision making.

Distance measures are used to measure the distance between two vectors. A distance measure is a function with two arguments,  $x$  and  $y$ , which are the two vectors being compared, and it returns a non-negative value representing the distance between them. The distance measures have to comply with certain criteria, such as symmetry, that is,  $d(x, y) = d(y, x)$ , and the distance of a vector to itself is 0, that is,  $d(x, y) = 0$  if and only if  $x = y$ . Most distance measures also fulfil the triangle inequality, which states that the distance between two points via an intermediate point is at least as long as the direct distance between them.

The most common distance measures used in market segmentation analysis are the Euclidean distance, the Manhattan or absolute distance, and the asymmetric binary distance. Euclidean distance is the straight-line distance between two points in a two-dimensional space, while Manhattan distance is the distance between two points on a grid where streets have to be used to get from one point to another. The asymmetric binary distance is used only for binary vectors, where all elements are either 0 or 1, and it only uses dimensions where at least one vector contains a value of 1.

Hierarchical clustering methods are a way of grouping data that mimics how a human would approach dividing a set of observations into groups. The process starts by defining a distance measure between pairs of observations and a linkage method to obtain distances between groups of observations. Divisive clustering starts with the complete data set and splits it into two segments in each step until each observation has its own segment. Agglomerative clustering starts with each observation representing its own segment and merges the two segments closest to each other until the complete data set forms one large segment. Both methods result in a sequence of nested partitions ranging from one segment to  $n$  segments. The linkage method generalizes how distances between groups of observations are obtained, and different combinations of distance and linkage methods can reveal different features of the data. Single linkage uses a next-nearest neighbor approach to join sets and can reveal non-convex, non-linear structures, while average and complete linkage extract more compact clusters. Ward clustering is another popular method based on squared Euclidean distances that joins the two sets of observations with the minimal weighted squared Euclidean distance between cluster centers. The result of hierarchical clustering is typically presented as a dendrogram, which is a tree diagram that shows how the segments are nested

The k-means clustering algorithm can be improved by using smart starting values instead of randomly selecting consumers from the dataset. Randomly selected consumers may not be representative of the data space, increasing the likelihood of the algorithm getting stuck in a local optimum. To avoid this problem, the algorithm should be initialized using starting points that are evenly spread across the entire data space.

- Model-based methods for market segmentation rely on a statistical model to identify segments. The model assumes that there are underlying unobserved (latent) variables that determine consumer behavior.
- One of the most commonly used model-based methods for market segmentation is latent class analysis (LCA). LCA assumes that there are unobserved (latent) classes of consumers with different preferences for products or services.
- Finite mixture models (FMM) are another model-based method for market segmentation. FMM is a generalization of LCA that allows for more flexible distributional assumptions for the latent variables.
- Bayesian methods are another class of model-based methods for market segmentation. Bayesian methods provide a framework for estimating the posterior distribution of the model parameters, given the observed data and prior information.
- Model-based clustering is a more general approach to clustering that encompasses both distance-based and model-based methods. Model-based clustering assumes that the data are generated by a mixture of several underlying distributions, each corresponding to a different cluster.
- The auto-encoding neural network is a type of cluster analysis that uses a single hidden layer perceptron. The network has three layers - input, hidden, and output. The hidden layer takes the input data and produces a weighted linear combination of the inputs using non-linear functions. The output layer gives the response of the network, which in the case of clustering, is the same as the input. The network is trained to predict the inputs as accurately as possible by minimizing the squared Euclidean distance between inputs and outputs.
- Bagged clustering is a segmentation method that combines partitioning and hierarchical clustering algorithms with bootstrapping. The method starts by creating multiple bootstrapped samples from the original data set using random drawing with replacement.
- Model-based methods are viewed as an additional extraction method that offers a different approach to segment extraction. They are based on the assumption that market segments have certain sizes and specific characteristics that are unknown, and use empirical data to find the best values for these segment sizes and characteristics. Model-based methods are seen as selecting a general structure and fine-tuning it based on consumer data.

## Model Based Methods

In market segmentation, it is not always the case that each segmentation variable contributes to determining the segmentation solution. Redundant or noisy variables might exist, and pre-processing methods can be used to identify and remove them.

There are different methods of market segmentation, and in your previous message, you provided an example of variable selection in market segmentation using the VSBD algorithm. Another method of market segmentation that I can tell you about is factor-cluster analysis. This approach involves factor analyzing the segmentation variables in the first step and then discarding the original segmentation variables. In the second step, the factor scores resulting from the factor analysis are used to extract market segments

External cluster indices require two market segmentation solutions as input, and measure the similarity of these solutions. The similarity is defined as the extent to which the two solutions classify observations in the same way. The closer the two solutions are in terms of segment assignments, the higher the similarity. External cluster indices are useful when the correct market segmentation solution is known, or when it can be generated artificially. In the latter case, the known segmentation solution can be used as a benchmark against which to compare any other solutions obtained from the same data set. External cluster indices require two market segmentation solutions as input, and measure the similarity of these solutions.

## Selecting Target Segments

The eighth step in market segmentation analysis is to select the target segments. Companies should choose the segments that are most attractive and that align with their marketing objectives. The selected segments should be large enough to be profitable but not so large that they require a generic marketing approach.

In conclusion, market segmentation analysis is a critical process for developing successful marketing strategies. By following these steps, companies can identify the most profitable segments and tailor their marketing efforts to meet the specific needs of those customers. Through market segmentation analysis, companies can increase their sales, profits, and market share while building long-term relationships with their customers.

## Metric Descriptor Variables

Visualizations for metric descriptor variables (such as age, number of nights at the tourist destinations, money spent on accommodation) are commonly based on density plots, box plots or histograms. The density plot visualizes the distribution of a variable by estimating a smoothed density function. The box plot summarizes the distribution by showing the median, the quartiles, and outliers. The histogram is similar to the density plot, but it discretizes the data and shows the distribution by counting the number of observations in each interval.

In market research, a mosaic plot is a visualization tool that displays a cross-tabulation between two or more categorical variables. The plot is used to illustrate the relationship between the variables, with the size of each rectangle in the plot being proportional to the number of cases in that cell. The plot is divided into columns, with each column representing a level of one of the variables being plotted. Each rectangle within a column is divided into a number of smaller rectangles, with each smaller rectangle representing a level of the other variable being plotted. The shading or color of each rectangle is used to indicate the proportion of cases falling into that category.



In the context of market segmentation, a mosaic plot can be used to display the relationship between segment membership and a particular descriptor variable. This allows for the identification of patterns in the data that may not be immediately apparent from a simple table of frequencies. For example, a mosaic plot might show that members of one segment are more likely to be male, while members of another segment are more likely to be female. By identifying these patterns, marketers can gain a better understanding of the characteristics of each segment and develop more effective marketing strategies to target them.

This is an excerpt from a statistical analysis of consumer behavior in the Australian travel industry. The analysis uses a logistic regression model to predict membership in a specific segment based on various demographic and behavioral variables.

The first part of the excerpt describes the process of including all available variables in the model and then using stepwise selection to select the most relevant variables. The selected model includes Education, NEP, and Vacation.Behavior variables.

The second part of the excerpt compares the predictive performance of the full model with the stepwise-selected model using boxplots of predicted probabilities of segment membership for consumers in and not in the segment. The comparison shows that neither model performs optimally in predicting segment membership, as the median predicted probabilities are not well-differentiated between consumers in and not in the segment.

Overall, this excerpt provides a brief overview of the statistical analysis process used to understand consumer behavior in the Australian travel industry, highlighting the importance of model selection and evaluating model performance.

Classification and regression trees (CARTs) are a machine learning approach used for predicting a categorical or binary dependent variable based on a set of independent variables. CARTs are advantageous because they can perform variable selection, are easy to interpret, and can incorporate interaction effects. Additionally, they work well with a large number of independent variables. However, they have the disadvantage of being unstable, as small changes in the data can lead to completely different trees.

CARTs use a stepwise procedure to fit the model. At each step, consumers are split into groups based on one independent variable, with the aim of making the resulting groups as pure as possible with respect to the dependent variable. The resulting tree shows the nodes that emerge from each splitting step, with the root node containing all consumers and the terminal nodes being those that are not split further. The segment membership can be predicted by moving down the tree based on the independent variables of the consumers, and the final prediction is made at the terminal node.

Tree constructing algorithms differ in several ways, including the splits into two or more groups at each node, the selection criterion for the independent variable for the next split, the selection criterion for the split point of the independent variable, the stopping criterion for the stepwise procedure, and the final prediction at the terminal node.

R packages such as `rpart` and `partykit` implement different tree constructing algorithms. The `partykit` package also allows for the visualization of fitted tree models. A conditional inference tree can be fitted using the `ctree()` function from the `partykit` package, as shown in the example provided in the text.

The passage describes the use of decision trees for predicting segment membership based on descriptor variables in the Australian travel motives data set. The `partykit` package is used to fit classification trees, with the `ctree()` and `ctree_control()` functions used to specify the tree construction parameters.

The passage provides an example of fitting a tree with segment 6 membership as the dependent variable, with the minimum terminal node size set to 100 and the minimum criterion value set to 0.99. The resulting tree is visualized using the `plot()` function.

The passage also describes fitting a tree for categorical dependent variables with more than two categories using the `ctree()` function, with segment membership as a factor with six levels as the dependent variable. The resulting tree is shown using a text-based output.

The output of the tree models shows the splitting variables and the resulting terminal nodes, with the proportion of segment members in each node indicated by stacked bar charts. The accuracy of predicting segment membership for each terminal node is also reported.

This is an excerpt from a statistical analysis of consumer behavior in the Australian travel industry. The analysis uses a logistic regression model to predict membership in a specific segment based on various demographic and behavioral variables.

The first part of the excerpt describes the process of including all available variables in the model and then using stepwise selection to select the most relevant variables. The selected model includes Education, NEP, and Vacation.Behavior variables.

The second part of the excerpt compares the predictive performance of the full model with the stepwise-selected model using boxplots of predicted probabilities of segment membership for consumers in and not in the segment. The comparison shows that neither model performs optimally in predicting segment membership, as the median predicted probabilities are not well-differentiated between consumers in and not in the segment.

Overall, this excerpt provides a brief overview of the statistical analysis process used to understand consumer behavior in the Australian travel industry, highlighting the importance of model selection and evaluating model performance.

Classification and regression trees (CARTs) are a machine learning approach used for predicting a categorical or binary dependent variable based on a set of independent variables. CARTs are advantageous because they can perform variable selection, are easy to interpret, and can incorporate interaction effects. Additionally, they work well with a large number of independent variables. However, they have the disadvantage of being unstable, as small changes in the data can lead to completely different trees.

CARTs use a stepwise procedure to fit the model. At each step, consumers are split into groups based on one independent variable, with the aim of making the resulting groups as pure as possible with respect to the dependent variable. The resulting tree shows the nodes that emerge from each splitting step, with the root node containing all consumers and the terminal nodes being those that are not split further. The segment membership can be predicted by moving down the tree based on the independent variables of the consumers, and the final prediction is made at the terminal node.

Tree constructing algorithms differ in several ways, including the splits into two or more groups at each node, the selection criterion for the independent variable for the next split, the selection criterion for the split point of the independent variable, the stopping criterion for the stepwise procedure, and the final prediction at the terminal node.

R packages such as `rpart` and `partykit` implement different tree constructing algorithms. The `partykit` package also allows for the visualization of fitted tree models. A conditional inference tree can be fitted using the `ctree()` function from the `partykit` package, as shown in the example provided in the text.

The passage describes the use of decision trees for predicting segment membership based on descriptor variables in the Australian travel motives data set. The `partykit` package is used to fit classification trees, with the `ctree()` and `ctree_control()` functions used to specify the tree construction parameters.

The passage provides an example of fitting a tree with segment 6 membership as the dependent variable, with the minimum terminal node size set to 100 and the minimum criterion value set to 0.99. The resulting tree is visualized using the `plot()` function.

The passage also describes fitting a tree for categorical dependent variables with more than two categories using the `ctree()` function, with segment membership as a factor with six levels as the dependent variable. The resulting tree is shown using a text-based output.

The output of the tree models shows the splitting variables and the resulting terminal nodes, with the proportion of segment members in each node indicated by stacked bar charts. The accuracy of predicting segment membership for each terminal node is also reported.

## Team Members

- Manikandan C - TeamLeader
- Deepika Pratap
- Devansh
- Bhumeni Sai Purushottham

- Aryan

## Github

- <https://github.com/Mani512996/internship.git>
- <https://github.com/DevanshG7/DevanshG7.git>
- <https://github.com/Purush0704/Internship>
- [https://github.com/deepikapbarve/Project\\_2](https://github.com/deepikapbarve/Project_2)
- <https://github.com/Aryann29>