

**Table 4. Results - Main Outcomes**

	Any Overtime	Any overtime	Conditional overtime	RVUs (trimmed)	Door-to- decision	Patients
<b>Panel a: Baseline regression adjusted</b>						
Scribe present	-0.187*** [0.052]	-0.078** [0.033]	-0.426*** [0.125]	1.515* [0.781]	-0.185 [0.156]	0.530 [0.324]
<b>Panel b: Excluding overnight shifts</b>						
Scribe present	-0.176*** [0.059]	-0.063* [0.036]	-0.411*** [0.142]	1.585* [0.894]	-0.181 [0.167]	0.386 [0.382]
<b>Panel c: 19 or fewer patients</b>						
Scribe present	-0.095 [0.066]	-0.048 [0.042]	-0.261* [0.150]	2.070** [1.043]	-0.145 [0.223]	0.485 [0.330]
<b>Panel d: Greater than 19 patients</b>						
Scribe present	-0.313*** [0.087]	-0.108* [0.056]	-0.561*** [0.156]	0.573 [1.182]	-0.141 [0.216]	0.082 [0.324]
<b>Panel e: Physicians in the top half of pre-study outcome</b>						
Scribe present	-0.165** [0.071]	-0.043 [0.040]	-0.326*** [0.102]	1.088 [1.124]	-0.080 [0.286]	0.288 [0.471]
<b>Panel f: Physicians in the bottom half of pre-study outcome</b>						
Scribe present	-0.141** [0.067]	-0.107** [0.042]	-0.604** [0.238]	1.928* [1.073]	-0.153 [0.183]	0.768* [0.436]

Notes: All regressions include physician, month, type of shift, and location of shift fixed effects. Regressions also include controls for whether or not the shift is on a weekend and, for all estimates except patients, patients. Standard errors are shown in brackets, and for all except the tobit regressions are White's robust standard errors. The tobit standard errors are bootstrapped with 100 repetitions. \* Statistically significant at the 10% level, \*\* at the 5% level, and \*\*\* at the 1% level