

Latent 10/12/2018



GRADE 12 EXAMINATION  
NOVEMBER 2018

**ADVANCED PROGRAMME MATHEMATICS: PAPER I  
MODULE 1: CALCULUS AND ALGEBRA**


**MARKING GUIDELINES**

Time: 2 hours

200 marks

These marking guidelines are prepared for use by examiners and sub-examiners, all of whom are required to attend a standardisation meeting to ensure that the guidelines are consistently interpreted and applied in the marking of candidates' scripts.

The IEB will not enter into any discussions or correspondence about any marking guidelines. It is acknowledged that there may be different views about some matters of emphasis or detail in the guidelines. It is also recognised that, without the benefit of attendance at a standardisation meeting, there may be different interpretations of the application of the marking guidelines.

  
PLEASE TURN OVER  
7/12/2018

## QUESTION 1

## 1.1 (a) METHOD 1

$$|x^2 + x| = -2x - 2$$

$$\therefore (x^2 + x)^2 = (-2x - 2)^2 \quad \checkmark \text{ m - squaring both sides}$$

$$\therefore x^4 + 2x^3 + x^2 = 4x^2 + 8x + 4$$

$$\therefore x^4 + 2x^3 - 3x^2 - 8x - 4 = 0 \quad \checkmark \text{ a}$$

$$\text{by inspection } x = -1 \text{ is a root } \checkmark \text{ m}$$

$$\therefore (x+1)(x^3 + x^2 - 4x - 4) = 0 \quad \checkmark \text{ a}$$

$$\text{by inspection } x = -1 \text{ is a root } \checkmark \text{ m}$$

$$\therefore (x+1)(x+1)(x^2 - 4) = 0 \quad \checkmark \text{ a}$$

$$\therefore (x+1)(x+1)(x+2)(x-2) = 0 \quad \checkmark \text{ a } \text{OR}$$

$$\therefore x = -1 \text{ or } \pm 2$$

$$\text{a check reveals } x = -1 \text{ or } -2 \text{ only } \checkmark \text{ m - checking - can be implied } \checkmark \text{ a}$$

$$\begin{aligned} (-2)^2 - x &= -2x - 2 \\ x^2 + 2 + 2 &= 0 \\ \text{N/S} \\ \text{with other side} \\ \text{correct } 4/8 \end{aligned}$$

## ALTERNATIVE 1

$$|x^2 + x| = -2x - 2$$

$$\therefore x^2 + x = -2x - 2 \text{ or } -x^2 - x = -2x - 2 \quad \checkmark \text{ m - 2 cases } \checkmark \text{ a}$$

$$\therefore x^2 + 3x + 2 = 0 \text{ or } x^2 - x - 2 = 0 \quad \checkmark \text{ a } \checkmark \text{ a}$$

$$\therefore x = -1 \text{ or } -2 \text{ or } x = -1 \text{ or } 2 \quad \checkmark \text{ ca } \checkmark \text{ ca}$$

$$\text{but a check reveals } x = -1 \text{ or } -2 \quad \checkmark \text{ m - checking - can be implied } \checkmark \text{ a}$$

if only 1 case  
max 4

## ALTERNATIVE 2

$$|x^2 + x| = -2x - 2 \quad \checkmark \text{ a}$$

$$\text{if } x^2 + x < 0 \text{ then } -1 < x < 0 \text{ and we have } \checkmark \text{ m - 2 cases}$$

$$-(x^2 + x) = -2x - 2 \quad \checkmark \text{ a}$$

$$\therefore -x^2 - x = -2x - 2$$

$$\therefore x^2 - x - 2 = 0$$

$$\therefore x = 2 \text{ or } -1 \text{ (a contradiction)} \quad \checkmark \text{ m checking } \checkmark \text{ a discarding}$$

$$\text{if } x^2 + x \geq 0 \text{ then } x \leq -1 \text{ or } x \geq 0 \text{ and we have}$$

$$x^2 + x = -2x - 2 \quad \checkmark \text{ a}$$

$$\therefore x^2 + 3x + 2 = 0$$

$$\therefore (x+2)(x+1) = 0$$

$$\therefore x = -2 \text{ or } -1 \quad \checkmark \text{ a}$$

$$\text{Ans only } 2/8 \quad x = -2 \text{ or } x = -1$$

(8)

(8)

10

answer only (b)  
→ 1  
if mess up line 1  
B/D max 4

$$\begin{aligned} \ln x^3 + 2 \ln x^2 &= 7 \\ \therefore \ln x^3 + \ln x^4 &= 7 \quad \checkmark \text{m log laws, } \checkmark \text{a} \\ \therefore \ln x^7 &= 7 \quad (\text{can go straight to the answer from here}) \quad \checkmark \text{a} \\ \therefore 7 \ln x &= 7 \quad \checkmark \text{a} \\ \therefore \ln x &= 1 \quad \checkmark \text{ca} \\ \therefore x &= e \quad \checkmark \text{ca} \end{aligned}$$

(overlook  $x = 2,718$ )

$$\begin{aligned} \checkmark \text{a} \quad \checkmark \text{m log law} \\ 3 \ln x + 4 \ln x &= 7 \\ \therefore 7 \ln x &= 7 \quad \checkmark \text{ca} \quad \checkmark \text{a} \\ \therefore \ln x &= 1 \quad \checkmark \text{ca} \\ \therefore x &= e \quad \checkmark \text{ca} \\ (6) \end{aligned}$$

1.2 (a)  
 $\log_e x$  is okay

$$\begin{aligned} y &= y_0 e^{-kt} \\ \therefore e^{-kt} &= \frac{y}{y_0} \quad \checkmark \text{a} \\ \therefore -kt &= \ln \frac{y}{y_0} \quad \checkmark \text{m logs } \checkmark \text{a} \end{aligned}$$

1 further  
alternative  
forms for k

$$\therefore k = \frac{\ln \frac{y}{y_0}}{-t} \quad \checkmark \text{ca} \quad \text{OR} \quad \frac{\ln y - \ln y_0}{-t} \quad \text{OR} \quad \frac{\ln y_0 - \ln y}{t} \quad (4)$$

$$(b) \quad k = \frac{\ln \frac{0,5 y_0}{y_0}}{-5700} \approx 1,216 \times 10^{-4} \quad \checkmark \text{ca} \quad 0,000122 \quad (\text{to 6 dp}) \quad (2)$$

not necessary

$$\begin{aligned} (c) \quad 0,9 y_0 &= y_0 e^{-kt} \quad \checkmark \text{m} \quad \text{indicating 90\% ideal ratio} \\ \therefore -kt &= \ln 0,9 \quad \checkmark \text{m logs} \\ \therefore t &= \frac{\ln 0,9}{-k} \quad \checkmark \text{a} \\ \therefore t &\approx 866 \text{ years} \quad \checkmark \text{ca} \end{aligned}$$

accept "exact" value for k (Will get 864 if use rounded)  
answers [24]



QUESTION 2

2.1 If  $3+2i$  is a root then so is  $3-2i$  ✓ m – can be implied  
so our equation is:

B/g Max 4  $(x+3)(x-(3+2i))(x-(3-2i))=0$  ✓ m ✓ a

if x out directly 2-process

$\therefore (x+3)((x-3)-2i)((x-3)+2i)=0$  ✓ m  
 $\therefore (x+3)((x-3)^2-4i^2)=0$  ✓ a  
 $\therefore (x+3)(x^2-6x+13)=0$  ✓ ca  
 $\therefore x^3-3x^2-5x+39=0$  ✓ ca ✓ ca

2 pairs  
✓ 1 complex/im.  
✓ 1 hence real

2.2 A cubic equation will have three roots. ✓ a ✓ a Complex roots of polynomials with real coefficients occur in (conjugate) pairs so there must be at least one real root. ✓ a ✓ a

2.3  $\frac{a+bi}{-b+ai} \times \frac{-b-ai}{-b-ai}$  ✓ m ✓ a  
 $= \frac{-ab-a^2i-b^2i-abi^2}{b^2-a^2i^2}$  ✓ ca  
 $= \frac{ab-ab-i(a^2+b^2)}{b^2+a^2}$  ✓ a  
 $= \frac{-i(a^2+b^2)}{b^2+a^2}$  ✓ a  
 $= -i$  ✓ a

Alternative 1:

$x = 3+2i$   
 $x-3 = 2i$   
 $x^2-6x+9 = -4$   
 $x^2-6x+13 = 0$


5 directly

Alternative 2

(8)

OR NB

OR Graph



(4)  
or explanation

$\frac{a}{ai} \times \frac{i}{i} = \frac{ai}{-a} = -i$   
 $\frac{a+bi}{a-b} = \frac{a+bi}{a-b} \times \frac{a+b}{a+b} = \frac{a^2-b^2+2abi}{a^2-b^2}$   
 $\frac{a^2-b^2+2abi}{a^2-b^2} = -i$   
 $\frac{8}{8}$

(8)  
[20]

Sum/Product

$x^2 - \text{sum } x + \text{Prod} = 0$

Sum =  $\sqrt{a}$  Prod =  $(3-2i)(3+2i) = 9-4i^2 = 13$  ✓ m

$\therefore x^2 - 6x + 13 = 0$  ✓ ca ✓ ca 5 to here

2.1)  $x = 3+2i$  ✓  
 $x = 3-2i$  ✓  
 $(x-3-2i)(x-3+2i) = 0$   
 $(x-3)^2 - (2i)^2 = 0$   
 $(x-3)^2 + 4 = 0$   
 $x^2 - 6x + 9 + 4 = 0$   
 $x^2 - 6x + 13 = 0$   
if 0 is left by any of these three steps  
2.2 Add wording: crossing x-axis at least once (real root - something extra)

2.2 A cubic function intersects with the x-axis at least once as it has a range that extends to  $+\infty$  and  $-\infty$  making its range  $\mathbb{R}$  with no restrictions. It is therefore impossible for a cubic equation to not have at least one real root (an x-intercept).

PLEASE TURN OVER

Q

Alternate:

③ Let  $n=1$   $2^{3(1)} - 3^1 = 5$  ✓ v.a.  
 $\therefore$  Divisible by 5 ✓ or

Let's assume that (the statement is true for  $n=k$ )  
 $\therefore 2^{3k} - 3^k$  is divisible by 5.

Let  $n=k+1$ : ✓

$$2^{3(k+1)} - 3^{k+1} \quad \checkmark$$

$$= 2^{3k} \cdot 2^3 - 3^k \cdot 3 \quad \checkmark$$

$$= 8 \cdot 2^{3k} - 3 \cdot 3^k$$

$$= (8 \cdot 2^{3k}) - (8-5) \cdot 3^k \quad \checkmark \text{ a.}$$

$$= 8 \cdot 2^{3k} - 8 \cdot 3^k + 5 \cdot 3^k \quad \checkmark \text{ a.}$$

$$= 8(2^{3k} - 3^k) + 5 \cdot 3^k \quad \checkmark \text{ factorise.}$$

$$\therefore 2^{3k} - 3^k \text{ is divisible by } 5 \quad \checkmark$$

$$\text{and } 5 \cdot 3^k$$

Conclusion (2.3)

Q

## QUESTION 3

Strict - proof

if  $n=1$  we have  $2^3 - 3 = 5$  which is clearly divisible by 5.  $\checkmark_m \checkmark_a$

Assume true for  $n=k$  viz. that

$$2^{3k} - 3^k = 5p \text{ where } p \in \mathbb{N} \quad (*) \quad \checkmark_m$$

for  $n=k$  $\checkmark$  for  $p \in \mathbb{N}$  or  $\mathbb{Z}$ .

Now if  $n=k+1$  we have:  $\checkmark_m$  idea of  $(k+1)$

$$2^{3(k+1)} - 3^{k+1} \quad \checkmark_a$$

$$= 2^{3k+3} - 3^{k+1}$$

$$= 2^{3k} \times 2^3 - 3 \times 3^k$$

$$= 8 \times 2^{3k} - 3 \times 3^k \quad \checkmark_m \text{ splitting using } *$$

from  $(*)$  we have  $2^{3k} = 5p + 3^k$  so if  $n=k+1$  we have

$$= 8(5p) + 8(3^k) - 3 \times 3^k \quad \checkmark_a$$

$$= 5(8p) + 5(3^k) \quad \checkmark_a$$

$$= 5(8p + 3^k) \quad \checkmark_a \text{ OR}$$



which is clearly divisible by 5

so, by the Principle of Mathematical Induction we have proved the result for  $n \in \mathbb{N} \quad \checkmark_a \quad \checkmark_a$

OR PMI (or a summary of their argument) [14]

## QUESTION 4

4.1 (a)

 $\checkmark \checkmark$  shape $\checkmark \checkmark$  symmetry about y-axis $\checkmark \checkmark$  range $\checkmark$  y-intif a full  $e^x$   
max 3 $\checkmark$  another pt. if only one  
branch max 4

(b)

 $x=0 \quad \checkmark \checkmark$ 

(0; 1) is fine

(8)

(2)

his/her  
consistent with graph if it has a pt. of non-differentiability

2

4.2 if  $f$  is differentiable at  $x = 2$

it must be continuous at  $x=2$  ✓m

so  $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x)$  **OK**

$$\text{so } 2a - b - 1 = 4b - 2a + 5 \quad \text{or} \quad 4a - 5b = 6 \quad (1) \quad \checkmark \text{ca}$$

but also,  $\lim_{x \rightarrow 2^-} f'(x) = \lim_{x \rightarrow 2^+} f'(x) \quad \checkmark_m \checkmark_a$

$$a = 4b - a \quad \text{or} \quad a = 2b \quad (2) \checkmark_a$$

(solving (1) and (2) simultaneously)  $\checkmark_m$  - solving simultaneously  
 $8b - 5b = 6 \checkmark_a$  so  $b = 2 \checkmark_{ca}$  and  $a = 4 \checkmark_{ca}$

$$8b - 5b = 6 \sqrt{a} \quad \text{so} \quad b = 2 \sqrt{ca} \quad \text{and} \quad a = 4 \sqrt{ca}$$

(10)

[20]

Max 8/10  
for notation  
abuse

### QUESTION 5

5.1  $segment = sector - \Delta \sqrt{m'}$

$$\therefore 308 = \frac{1}{2} (18^2) \theta - \frac{1}{2} (18^2) \sin \theta$$

✓m – equating to 308

$$\therefore 162\theta - 162\sin\theta - 308 = 0$$

(6)

5.2  $f(\theta) = 162\theta - 162\sin\theta - 308 \text{ } \checkmark \text{ m}$

$$\therefore \theta_{n+1} = \theta_n - \frac{\sqrt{n} \cdot 162\theta - 162\sin\theta - 308\sqrt{a}}{162 - 162\cos\theta \sqrt{a}}$$

$\theta = 2.49984 \checkmark \checkmark$   $\sqrt{m}$  subst. 2 - optional  
to 5 or more d.p.

(8)

114

### QUESTION 6

6.1  $f(0) = \frac{1}{2}$ , so  $y$ -int  $\left(0, \frac{1}{2}\right)$  or  $y = \frac{1}{2}$

$$\frac{2x^2 - 3x - 2}{x - 4} = 0 \quad \checkmark_m \text{ letting}$$

$$\therefore 2x^2 - 3x - 2 = 0 \quad \checkmark m \text{ for numerator} = 1$$

$$\therefore (2x+1)(x-2)=0 \quad \text{---a}$$

$$\therefore x\text{-ints } \left(-\frac{1}{2}; 0\right) \in a \text{ and } (2; 0) \in a$$

Accept answers only (6)

vert. asymptote

$$y = 4 \cdot 6.2$$


vertical asymptote:  $x = 4 \sqrt{a} \sqrt{a}$

$$2x^2 - 3x - 2 = (x - 4)(2x + 5) + R \quad \checkmark m \checkmark a \checkmark a \checkmark a$$

so, oblique asymptote is  $y = 2x + 5 \sqrt{a} \sqrt{a}$

Accept answers only (8)

(be on the lookout for method involving limit)



\* If they mess up long division / method )  
May  $\frac{3}{b}$ .

1



6.3  $f(x) = \frac{2x^2 - 3x - 2}{x - 4}$

$\therefore f'(x) = \frac{(4x-3)(x-4) - 1(2x^2-3x-2)}{(x-4)^2} = 0 \checkmark m = 0$

$\therefore 4x^2 - 19x + 12 - 2x^2 + 3x - 2 = 0 \checkmark a$

$\therefore 2x^2 - 16x + 14 = 0$

$\therefore x^2 - 8x + 7 = 0$

$\therefore (x-1)(x-7) = 0$

$\therefore x = 1 \text{ or } 7 \checkmark ca$

$\therefore (1;1) \text{ and } (7;25) \text{ are stationary points}$   
 $\checkmark ca \quad \checkmark ca$

Product rule

$f(x) = (2x^2 - 3x - 2)(x-4)^{-1}$   
 $\therefore f'(x) = (4x-3)(x-4)^{-1} - (x-4)^{-2}(2x^2-3x-2)$

$\therefore \frac{4x-3}{x-4} - \frac{2x^2-3x-2}{(x-4)^2} = 0$

$f''(x) = \frac{(x-4)^2(4x-3) - (2x^2-3x-2)(2x-4)}{(x-4)^4}$   
 $= \frac{(x-4)^2(4x-3) - (2x^2-3x-2)(2x-4)}{(x-4)^4}$   
 $= \frac{(x-4)^2(4x-3) - (2x^2-3x-2)(2x-4)}{(x-4)^4}$   
 $= \frac{(x-4)^2(4x-3) - (2x^2-3x-2)(2x-4)}{(x-4)^4}$

(8)

6.4  $f''(1) < 0$  so  $(1;1)$  is a local maximum  $\checkmark m \checkmark a$   
 $-\frac{4}{3}$   $f''(7) > 0$  so  $(7;25)$  is a local minimum  $\checkmark m \checkmark a$   
 $\frac{4}{3}$

can't use other methods

(4)

[26]

### QUESTION 7

7.1  $x^2 + xy + y^2 = 1$   
 $\therefore 2x + y + x \frac{dy}{dx} + 2y \frac{dy}{dx} = 0 \checkmark a$

Be on lookout

$\therefore \frac{dy}{dx}(x+2y) = -2x-y \checkmark m$  factorising

$\therefore \frac{dy}{dx} = \frac{-2x-y}{x+2y}$

(5)

7.2 At A,  $y = 0 \therefore x = 1 \checkmark m \checkmark a$  for  $y=0$   $(2;0) \quad x = \pm 1$

so, at A,  $\frac{dy}{dx} = \frac{-2}{1} = -2 \checkmark a$  (for wrong) max 3 for normal

$\therefore y = -2(x-1) \checkmark a$  for str. line

$\therefore y = -2x + 2$  (not necessary)

(5)

[10]

2



latest 10/12/2018



GRADE 12 EXAMINATION  
NOVEMBER 2018

**ADVANCED PROGRAMME MATHEMATICS: PAPER I  
MODULE 1: CALCULUS AND ALGEBRA**


**MARKING GUIDELINES**

Time: 2 hours

200 marks

These marking guidelines are prepared for use by examiners and sub-examiners, all of whom are required to attend a standardisation meeting to ensure that the guidelines are consistently interpreted and applied in the marking of candidates' scripts.

The IEB will not enter into any discussions or correspondence about any marking guidelines. It is acknowledged that there may be different views about some matters of emphasis or detail in the guidelines. It is also recognised that, without the benefit of attendance at a standardisation meeting, there may be different interpretations of the application of the marking guidelines.

  
7/12/2018

**QUESTION 1**

**1.1 (a) METHOD 1**

$$|x^2 + x| = -2x - 2$$

$$\therefore (x^2 + x)^2 = (-2x - 2)^2 \quad \checkmark \text{ m – squaring both sides}$$

$$\therefore x^4 + 2x^3 + x^2 = 4x^2 + 8x + 4$$

$$\therefore x^4 + 2x^3 - 3x^2 - 8x - 4 = 0 \quad \checkmark \text{ a}$$

$$\text{by inspection } x = -1 \text{ is a root } \checkmark \text{ m}$$

$$\therefore (x+1)(x^3 + x^2 - 4x - 4) = 0 \quad \checkmark \text{ a}$$

$$\text{by inspection } x = -1 \text{ is a root } \checkmark \text{ m}$$

$$\therefore (x+1)(x+1)(x^2 - 4) = 0 \quad \checkmark \text{ a}$$

$$\therefore (x+1)(x+1)(x+2)(x-2) = 0 \quad \checkmark \text{ a } \text{or}$$

$$\therefore x = -1 \text{ or } \pm 2$$

$$\text{a check reveals } x = -1 \text{ or } -2 \text{ only } \checkmark \text{ m – checking – can be implied } \checkmark \text{ a}$$

$$\begin{aligned} (-2)^2 - x &= -2x - 2 \\ x^2 + x + 2 &= 0 \\ \text{N/S} \\ \text{with other side} \\ \text{correct } 4/8 \end{aligned}$$

**ALTERNATIVE 1**

$$|x^2 + x| = -2x - 2$$

$$\therefore x^2 + x = -2x - 2 \text{ or } -x^2 - x = -2x - 2 \quad \checkmark \text{ m – 2 cases } \checkmark \text{ a}$$

$$\therefore x^2 + 3x + 2 = 0 \text{ or } x^2 - x - 2 = 0 \quad \checkmark \text{ a } \checkmark \text{ a}$$

$$\therefore x = -1 \text{ or } -2 \text{ or } x = -1 \text{ or } 2 \quad \checkmark \text{ ca } \checkmark \text{ ca}$$

$$\text{but a check reveals } x = -1 \text{ or } -2 \quad \checkmark \text{ m – checking – can be implied } \checkmark \text{ a}$$

if only 1 case  
max 4

**ALTERNATIVE 2**

$$|x^2 + x| = -2x - 2$$

$$\text{if } x^2 + x < 0 \text{ then } -1 < x < 0 \text{ and we have } \checkmark \text{ m – 2 cases } \checkmark \text{ a}$$

$$-(x^2 + x) = -2x - 2 \quad \checkmark \text{ a}$$

$$\therefore -x^2 - x = -2x - 2$$

$$\therefore x^2 - x - 2 = 0$$

$$\therefore x = 2 \text{ or } -1 \text{ (a contradiction) } \checkmark \text{ m checking } \checkmark \text{ a discarding}$$

$$\text{if } x^2 + x \geq 0 \text{ then } x \leq -1 \text{ or } x \geq 0 \text{ and we have } \checkmark \text{ a}$$

$$x^2 + x = -2x - 2 \quad \checkmark \text{ a}$$

$$\therefore x^2 + 3x + 2 = 0$$

$$\therefore (x+2)(x+1) = 0$$

$$\therefore x = -2 \text{ or } -1 \quad \checkmark \text{ a}$$

$$\text{Ans only } 2/8 \quad x = -2 \text{ or } x = -1$$

(8)

(8)

Q

answer only  
→  
if mess up line 1  
B/D max 4

(b)  $\ln x^3 + 2 \ln x^2 = 7$   
 $\therefore \ln x^3 + \ln x^4 = 7$   $\checkmark$  m log laws,  $\checkmark$  a  
 $\therefore \ln x^7 = 7$  (can go straight to the answer from here)  $\checkmark$  a  
 $\therefore 7 \ln x = 7$   $\checkmark$  a  
 $\therefore \ln x = 1$   $\checkmark$  ca  
 $\therefore x = e$   $\checkmark$  ca  
 (overlook  $x = 2,718$ )

$\checkmark$  a  
 $\checkmark$  m log lav  
 $3 \ln x + 4 \ln x = 7$   
 $\therefore 7 \ln x = 7$   $\checkmark$  ca  $\checkmark$  a  
 $\therefore \ln x = 1$   $\checkmark$  ca  
 $\therefore x = e$   $\checkmark$  ca  
 (6)

1.2 (a)  
 $\log_e x$  is okay

$y = y_0 e^{-kt}$   
 $\therefore e^{-kt} = \frac{y}{y_0}$   $\checkmark$  a  
 $\therefore -kt = \ln \frac{y}{y_0}$   $\checkmark$  m  $\checkmark$  a

1 further  
alternative  
forms for k

$\therefore k = \frac{\ln \frac{y}{y_0}}{-t}$   $\checkmark$  ca OR  $\frac{\ln y - \ln y_0}{-t}$  OR  $\frac{\ln y_0 - \ln y}{t}$  (4)

(b)  $k = \frac{\ln \frac{0,5y_0}{y_0}}{-5700}$   $\checkmark$  m indicating  $\frac{1}{2}$   
 $\approx 1,216 \times 10^{-4}$   $\checkmark$  ca 0,000122 (to 6 dp) (2)  
 not necessary

(c)  $0,9y_0 = y_0 e^{-kt}$   $\checkmark$  m indicating 90% idea / ratio  
 $\therefore -kt = \ln 0,9$   $\checkmark$  m logs  
 $\therefore t = \frac{\ln 0,9}{-k}$   $\checkmark$  a  
 $\therefore t \approx 866$  years  $\checkmark$  ca (Will get 864 if use rounded value for k) (4)  
 accept "exact" answers [24]

1

QUESTION 2

2.1 If  $3+2i$  is a root then so is  $3-2i$  ✓ m - can be implied  
so our equation is:

B/d Max 4

$$(x+3)(x-(3+2i))(x-(3-2i))=0 \quad \text{Factors} \quad \checkmark m \checkmark a$$

if  $x$  out  
directly  
2-process

$$\therefore (x+3)((x-3)-2i)((x-3)+2i)=0 \quad \checkmark m$$

$$\therefore (x+3)((x-3)^2-4i^2)=0 \quad \checkmark a$$

$$\therefore (x+3)(x^2-6x+13)=0 \quad \checkmark ca$$

$$\therefore x^3-3x^2-5x+39=0 \quad \checkmark ca \checkmark ca$$

2 pairs  
✓ 1 complex/im.  
✓ 1 hence real

2.2

A cubic equation will have three roots. ✓ a ✓ a Complex roots of polynomials with real coefficients occur in (conjugate) pairs so there must be at least one real root. ✓ a ✓ a not necessary


2.3

$$\begin{aligned} \frac{a+bi}{-b+ai} \times \frac{-b-ai}{-b-ai} & \quad \checkmark m \checkmark a \\ & = \frac{-ab-a^2i-b^2i-abi^2}{b^2-a^2i^2} \quad \checkmark ca \\ & = \frac{ab-ab-i(a^2+b^2)}{b^2+a^2} \quad \checkmark a \\ & = \frac{-i(a^2+b^2)}{b^2+a^2} \quad \checkmark a \\ & = -i \checkmark a \end{aligned}$$

NB

OR

Graph



explanation

$$\begin{aligned} \frac{a}{ai} \times \frac{i}{i} & = \frac{ai}{-a} = -i \\ ai - b & \quad \frac{a+bi}{a+bi} \cdot \frac{-i}{-i} = \frac{-i(a+bi)}{-i^2(a^2+b^2)} = \frac{-i(a+bi)}{a^2+b^2} = \frac{-ia-bi}{a^2+b^2} = \frac{-i(a+b)}{a^2+b^2} \end{aligned}$$

(8)  
[20]

Sum/Product

$$x^2 - \text{sum } x + \text{Prod} = 0$$

$$\text{Sum} = 6$$

$$\text{Prod} = (3-2i)(3+2i) = 9-4i^2 = 13$$

$$\therefore x^2 - 6x + 13 = 0$$

5 to here

2.1  $x=3+2i$   $x=3-2i$   $x=3-2i$  ✓  
 $(x-3+2i)(x-3-2i)(x-3-2i)=0$   
 $(x-3)(x^2-6x+13-4i^2)=0$   
 $(x-3)(x^2-6x+17)=0$   
 $x^3-3x^2-5x+39=0$   
 $x^3-3x^2-5x+39=0$  ✓ ca  
if 0 is left of any of these three signs  
2.2 All working crossing x-axis at least once  
1 real root ✓  
(Need to say something extra)

2.2 A cubic function intersects with the x-axis at least once as it has 'arms' that extend to + and - infinity making its range  $\mathbb{R}$  with no restrictions. It is therefore impossible for a cubic equation to not have at least one real root (an x-intercept).

PLEASE TURN OVER

Q

Alternate:

③ Let  $n=1$   $2^{2(1)} - 3^1 = 5$  ✓ v.a.  
 $\therefore$  Divisible by 5 ✓ v.a.

Let's assume that (the statement is true for  $n=k$ )  
 $\therefore 2^{2k} - 3^k$  is divisible by 5.

Let  $n=k+1$  ✓

$$2^{2(k+1)} - 3^{k+1} \quad \checkmark$$

$$= 2^{2k} \cdot 2^2 - 3^k \cdot 3 \quad \checkmark$$

$$= 4 \cdot 2^{2k} - 3 \cdot 3^k$$

$$= (8 \cdot 2^{2k}) - (8-5) \cdot 3^k \quad \checkmark$$

$$= 8 \cdot 2^{2k} - 8 \cdot 3^k + 5 \cdot 3^k \quad \checkmark$$

$$= 8(2^{2k} - 3^k) + 5 \cdot 3^k \quad \checkmark$$

$\therefore 2^{2k} - 3^k$  is divisible by 5.  
and  $5 \cdot 3^k$

Conclusion (2.7)

**QUESTION 3**

strict - proof

if  $n=1$  we have  $2^3 - 3 = 5$  which is clearly divisible by 5.  $\checkmark_m \checkmark_a$

Assume true for  $n=k$  viz. that

$$2^{3k} - 3^k = 5p \text{ where } p \in \mathbb{N} \quad (*) \quad \checkmark_m$$

for  $n=k$

$\checkmark$  for  $p \in \mathbb{N}$  or  $\mathbb{Z}$

Now if  $n=k+1$  we have:  $\checkmark_m$  idea of  $(k+1)$

$$2^{3(k+1)} - 3^{k+1} \quad \checkmark_a$$

$$= 2^{3k+3} - 3^{k+1}$$

$$= 2^{3k} \times 2^3 - 3 \times 3^k$$

$$= 8 \times 2^{3k} - 3 \times 3^k \quad \checkmark_m \text{ splitting using } *$$

from  $(*)$  we have  $2^{3k} = 5p + 3^k$  so if  $n=k+1$  we have

$$= 8(5p) + 8(3^k) - 3 \times 3^k \quad \checkmark_a$$

$$= 5(8p) + 5(3^k) \quad \checkmark_a$$

$$= 5(8p + 3^k) \quad \checkmark_a \text{ OR}$$


which is clearly divisible by 5

so, by the Principle of Mathematical Induction we have proved the result for  $n \in \mathbb{N} \checkmark_a \checkmark_a$

OR PMI (or a summary of their argument) [14]

**QUESTION 4**

4.1 (a)



$\checkmark \checkmark$  shape


$\checkmark \checkmark$  symmetry about y-axis

$\checkmark \checkmark$  range

$\checkmark$  y-int

if a full  $e^x$   
max 3

$\checkmark$  another pt. if only one  
branch max 4



(b)

$x=0 \checkmark \checkmark$

$(0;1)$  is fine

his/her

consistent with graph if it has a pt. of non-differentiability

(8)

(2)

12

4.2 if  $f$  is differentiable at  $x = 2$

it must be continuous at  $x = 2$  ✓m

so  $\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x)$  **OK**

$$\text{so } 2a - b - 1 = 4b - 2a + 5 \quad \text{or} \quad 4a - 5b = 6 \quad (1) \quad \checkmark \text{ca}$$

but also,  $\lim_{x \rightarrow 2^-} f'(x) = \lim_{x \rightarrow 2^+} f'(x) \quad \checkmark m \quad \checkmark a$

$$a = 4b - a \quad \text{or} \quad a = 2b \quad (2) \checkmark a$$

(solving (1) and (2) simultaneously)  $\checkmark$  m - solving simultaneously  
 $8h - 5h = 6\sqrt{ca}$  so  $h = 2\sqrt{ca}$  and  $a = 4\sqrt{ca}$

$$8b - 5b = 6 \sqrt{a} \quad \text{so} \quad b = 2 \sqrt{ca} \quad \text{and} \quad a = 4 \sqrt{ca}$$

Max 8/10  
for notation  
abuse

(10)

[20]

### QUESTION 5

5.1  $\text{segment} = \text{sector} - \Delta \sqrt{m'}$

$$\therefore 308 = \frac{1}{2} (18^2) \theta - \frac{1}{2} (18^2) \sin \theta$$

$$\therefore 162\theta - 162\sin\theta - 308 = 0$$

✓m – equating to 308

(6)

5.2  $f(\theta) = 162\theta - 162\sin\theta - 308 \text{ } \checkmark \text{ m}$

$$\therefore \theta_{n+1} = \theta_n - \frac{\sqrt{n} \cdot 162\theta - 162\sin\theta - 308\sqrt{a}}{162 - 162\cos\theta \sqrt{a}}$$

$\theta = 2.49984 \checkmark \checkmark$  in subst. 2 - optional  
to 5 or more d.p.

(8)

[14]

### QUESTION 6

6.1  $f(0) = \frac{1}{2}$ , so  $y$ -int  $\left(0; \frac{1}{2}\right)$  or  $y = \frac{1}{2}$

$$\frac{2x^2 - 3x - 2}{x - 4} = 0 \quad \checkmark \text{m letting } y = 0$$

$$\therefore 2x^2 - 3x - 2 = 0 \quad \checkmark m \text{ for numerator} = 1$$

$$\therefore (2x+1)(x-2)=0 \quad \text{---a}$$

$\therefore x\text{-ints } \left(-\frac{1}{2}; 0\right) \in a \text{ and } (2; 0) \in a$

Accept answers only (6)

vert. asymptote

$$y = 4 \cdot 6.2^x$$


vertical asymptote:  $x = 4 \sqrt{a} \sqrt{a}$

$$2x^2 - 3x - 2 = (x - 4)(2x + 5) + R \quad \checkmark m \checkmark a \checkmark a \checkmark a$$

so, oblique asymptote is  $y = 2x + 5 \sqrt{a} \sqrt{a}$

Accept answers only (8)

(be on the lookout for method involving limit)



\* If they mess up long division / method )  
Max  $\frac{3}{8}$ .

19



6.3  $f(x) = \frac{2x^2 - 3x - 2}{x - 4}$  Product rule

$$\therefore f'(x) = \frac{(4x-3)(x-4) - 1(2x^2-3x-2)}{(x-4)^2} = 0 \checkmark m = 0$$

$$\therefore 4x^2 - 19x + 12 - 2x^2 + 3x + 2 = 0 \checkmark a$$

$$\therefore 2x^2 - 16x + 14 = 0$$

$$\therefore x^2 - 8x + 7 = 0$$

$$\therefore (x-1)(x-7) = 0$$

$$\therefore x = 1 \text{ or } 7 \checkmark ca$$

$$\therefore (1;1) \text{ and } (7;25) \text{ are stationary points}$$

$$\checkmark ca \quad \checkmark ca$$

(8)

6.4  $f''(1) < 0$  so  $(1;1)$  is a local maximum  $\checkmark m \checkmark a$

$f''(7) > 0$  so  $(7;25)$  is a local minimum  $\checkmark m \checkmark a$

(4)

can't use other methods

(4)

[26]

### QUESTION 7

7.1  $x^2 + xy + y^2 = 1$

$$\therefore 2x + y + x \frac{dy}{dx} + 2y \frac{dy}{dx} = 0 \checkmark a$$

$$\therefore \frac{dy}{dx}(x+2y) = -2x-y \checkmark m \text{ factorising}$$

$$\therefore \frac{dy}{dx} = \frac{-2x-y}{x+2y}$$

(5)

7.2 At A,  $y=0 \therefore x=1 \checkmark m \checkmark a$  for  $y=0$   $(2;0) x = \pm 1$

so, at A,  $\frac{dy}{dx} = \frac{-2}{1} = -2 \checkmark a$  (for wrong)

$\therefore y = -2(x-1) \checkmark a$  max 3 for normal

$\therefore y = -2x + 2$  (not necessary)

(5)

[10]

P

QUESTION 8

8.1  $\cos \theta = \frac{FC}{CD} \checkmark m$  — using trig.

$\therefore FC = 0,4 \cos \theta \checkmark a$

$\therefore A = \Delta CDF + \Delta BEG + BCFG \checkmark m$  — resolving into areas/trap.

$\therefore A = 2 \left( \frac{1}{2} \times 0,4 \times 0,4 \cos \theta \sin \theta \right) + 0,4 \times 0,4 \cos \theta \quad (\Delta CDF = \Delta BEG)$

$\therefore A = 0,16 \sin \theta \cos \theta + 0,16 \cos \theta$

$\therefore A = 0,08 \sin 2\theta + 0,16 \cos \theta \checkmark ca$

$\therefore V = 20(0,08 \sin 2\theta + 0,16 \cos \theta) \checkmark m$  — (for  $\times$  by 20)

$\therefore V = 1,6 \sin 2\theta + 3,2 \cos \theta$

$FC = 0,4 \cos \theta \checkmark$

$GE = 0,4 \sin \theta \checkmark$

putting 2 triangles together

$A = 2 \times \frac{1}{2} (0,4)^2 \sin 2\theta + 0,4 \times 0,4 \cos \theta \checkmark$   
 $= 0,08 \sin 2\theta + 0,16 \cos \theta$   
 (8)

N.B.  
Answer  
Given

8.2  $V = 1,6 \sin 2\theta + 3,2 \cos \theta$

a.  $\frac{dV}{d\theta} = 3,2 \cos 2\theta - 3,2 \sin \theta = 0 \checkmark m$  — equating to zero

$\therefore \cos 2\theta = \sin \theta \checkmark a$

$\therefore \cos 2\theta = \cos \left( \frac{\pi}{2} - \theta \right) \checkmark m \checkmark a$

$\therefore 2\theta = \frac{\pi}{2} - \theta$

$\therefore 3\theta = \frac{\pi}{2}$

$\therefore \theta = \frac{\pi}{6} \checkmark a$

Alternative:

$1 - 2\sin^2 \theta = \sin \theta \checkmark m$

$2\sin^2 \theta + \sin \theta - 1 = 0 \checkmark a$

$(2\sin \theta - 1)(\sin \theta + 1) = 0$

$\therefore \sin \theta = \frac{1}{2} \text{ or } -1$

$\therefore \theta = \frac{\pi}{6} \quad (7/8 \text{ if keep } -\frac{\pi}{2}) \quad (8)$

if solve for "turning point"

$f(\theta) = 162 - 162 \cos \theta$   
 $f'(\theta) = 162 \sin \theta$   
 $0 = 162 \sin \theta$   
 $\sin \theta = 0$   
 $\theta = 0$   
 $\theta = \pi$   
 $\theta = 2\pi$   
 Max 5

QUESTION 9

9.1 (a)  $\sin^3 \theta = \sin \theta \times \sin^2 \theta \checkmark a$

$= \sin \theta (1 - \cos^2 \theta) \checkmark a \checkmark a$

$= \sin \theta - \sin \theta \cos^2 \theta \text{ as required } \checkmark a$

(b)  $\int \sin^3 \theta d\theta = \int \sin \theta d\theta - \int \sin \theta \cos^2 \theta d\theta$

$= -\cos \theta + \frac{\cos^3 \theta}{3} + c \checkmark a$

May go  
"backwards"

(8.2)  $\frac{dV}{d\theta} = 3,2 \cos 2\theta - 3,2 \sin \theta$   
 $0,2(1 - 2\sin^2 \theta) - 3,2 \sin \theta = 0$   
 $1 - 2\sin^2 \theta - 16 \sin \theta = 0$   
 $2\sin^2 \theta + 16 \sin \theta - 1 = 0$   
 $(2\sin \theta - 1)(\sin \theta + 1) = 0$   
 $\sin \theta = \frac{1}{2} \checkmark a$   
 $\theta = \frac{\pi}{6} \checkmark a$   
 (8)

(4)

(8)

*(Handwritten mark)*

$$(8.1) \quad \frac{FC}{\sin(\frac{\pi}{2} - \theta)} = \frac{0,4}{\sin \frac{\pi}{2}} \quad \checkmark$$

$$\frac{FC}{\cos \theta} = 0,4$$

$$FC = 0,4 \cos \theta \quad \checkmark$$

$$DF^2 = 0,4^2 - (0,4 \cos \theta)^2$$

$$DF = \sqrt{0,16 - 0,16 \cos^2 \theta} \quad \checkmark$$

$$A = \frac{1}{2} (\text{Sum of } \parallel \text{ sides}) \times \text{Height}$$

$$= \frac{1}{2} (2\sqrt{0,16 - 0,16 \cos^2 \theta} + 2(0,4)) \times 0,4 \cos \theta$$

$$= 0,4 \cos \theta \sqrt{0,16 - 0,16 \cos^2 \theta} + 0,16 \cos \theta$$

$$= 0,4 \cos \theta \times 0,4 \sqrt{1 - \cos^2 \theta} + 0,16 \cos \theta$$

$$= \cancel{0,16} \cos \theta \sqrt{\sin^2 \theta} + \cancel{0,16} \cos \theta$$

$$= \cancel{0,16} \cos \theta \cdot \sin \theta + \cancel{0,16} \cos \theta$$

$$= 0,08 \sin 2\theta + \cancel{0,16} \cos \theta$$

$$V = 20 (0,08 \sin 2\theta + \cancel{0,16} \cos \theta)$$

$$= 1,6 \sin 2\theta + 3,2 \cos \theta$$

10

$$(9.1) \quad (b) \quad \int \sin^3 \theta d\theta = \int \sin \theta d\theta - \int \sin \theta \cos^2 \theta d\theta$$

$$\left. \begin{aligned} \text{Set } u &= \cos \theta \\ \frac{du}{d\theta} &= -\sin \theta \\ -\sin \theta d\theta &= du \end{aligned} \right\} \checkmark M$$

$$= -\cos \theta + \int u^2 du \quad \checkmark$$

$$= -\cos \theta + \frac{u^3}{3} + C$$

$$= -\cos \theta + \frac{\cos^3 \theta}{3} + C \quad \checkmark$$

(8)

*[Handwritten signature]*

$$\begin{aligned}
9.1 \quad b) \quad & \int (\sin \theta - \sin \theta \cos^2 \theta) d\theta \\
&= \int \sin \theta d\theta - \frac{1}{2} \int \sin \theta (1 + \cos 2\theta) d\theta \\
&= -\cos \theta - \frac{1}{2} \int (\sin \theta + \sin \theta \cos 2\theta) d\theta \\
&= -\cos \theta + \frac{1}{2} \cos \theta - \frac{1}{4} \int (\sin 3\theta + \sin(-\theta)) d\theta \\
&= -\cos \theta + \frac{1}{2} \cos \theta + \frac{1}{4} \left( \frac{\cos 3\theta}{3} \right) - \frac{1}{4} (\cos \theta) + C \\
&= -\frac{3}{4} \cos \theta + \frac{\cos 3\theta}{12} + C \checkmark
\end{aligned}$$

19

## 9.2 METHOD 1

$$\int \frac{x}{\sqrt{2+x}} dx \quad \text{substituting}$$

let  $u = 2+x$  then  $x = u-2$  and  $du = dx$

$$\therefore \int \frac{u-2}{u^{\frac{1}{2}}} du \quad \checkmark a$$

$$= \int u^{\frac{1}{2}} - 2u^{-\frac{1}{2}} du \quad \checkmark m \text{ splitting}$$

$$= \frac{2}{3} u^{\frac{3}{2}} - 4u^{\frac{1}{2}} + c$$

$$= \frac{2}{3} (2+x)^{\frac{3}{2}} - 4(2+x)^{\frac{1}{2}} + c \quad \checkmark m \text{ subst back}$$

## ALTERNATIVE 1

$$\int \frac{x}{\sqrt{2+x}} dx$$

$$= \int x(2+x)^{-\frac{1}{2}} dx$$

$$\text{by parts } f = x \quad \checkmark a \text{ and } g' = (2+x)^{-\frac{1}{2}} \quad \checkmark a$$

$$\text{so } f' = 1 \quad \checkmark a \text{ and } g = 2(2+x)^{\frac{1}{2}} \quad \checkmark a$$

$$= 2x(2+x)^{\frac{1}{2}} - \int 2(2+x)^{\frac{1}{2}} dx \quad \checkmark ca$$

$$= 2x(2+x)^{\frac{1}{2}} - \frac{4(2+x)^{\frac{3}{2}}}{\frac{3}{2}} + c \quad \checkmark ca$$

(don't penalise c)

(8)

[20]



**QUESTION 10**

10.1  $Area = \frac{10}{3} + \frac{3}{2(4)} + \frac{1}{6(4^2)}$  ✓m substituting 4 for n ✓a

$= \frac{119}{32}$  ✓ca or 3,719 (3)

10.2 It will be an over-approximation. ✓a As  $n$  gets larger the answer decreases  
✓ca ✓ca under-only  $\rightarrow 0$

10.3  $Area = \lim_{n \rightarrow \infty} \left( \frac{10}{3} + \frac{3}{2n} + \frac{1}{6n^2} \right)$  ✓m for  $n$  tending to infinity

$= \frac{10}{3}$  units<sup>2</sup> ✓a ✓a (3) — — —

10.4  $\frac{10}{3}$  units<sup>2</sup> ✓a since this is simply a reflection of the shaded area in the y-axis. ✓a ✓a (3)

[12]





## QUESTION 11

$$\begin{aligned}
 \text{Area} &= \int_1^7 g(x) - f(x) dx \quad \checkmark \text{ limits} \quad \checkmark \text{m sub. areas} \\
 &= \int_1^7 f(x) + kx + 1 - f(x) dx \quad \checkmark a \\
 &= \int_1^7 kx + 1 dx \quad \checkmark a \\
 &= \left[ \frac{kx^2}{2} + x \right]_1^7 \quad \checkmark a \text{-notation} \\
 \text{so, } \frac{49k}{2} + 7 - \frac{k}{2} - 1 &= 54 \quad \checkmark \\
 \text{so, } 49k + 14 - k - 2 &= 108 \quad \checkmark ca \\
 \text{so, } 48k &= 96 \\
 \text{so } k &= 2 \quad \checkmark ca
 \end{aligned}$$

if he/she does:  
 $\frac{[f(x)]^2}{2}$  Penalise 2

[12]

## QUESTION 12

$$\begin{aligned}
 \text{vol} &= \pi \int_a^b [f(x)]^2 dx \\
 \text{for volume formula} \quad \therefore 175 &= \pi \int_0^h (-x^2 + 6x + 4) dx \quad \checkmark a \\
 \therefore 175 &= \pi \left[ -\frac{x^3}{3} + 3x^2 + 4x \right]_0^h \quad \checkmark a \\
 \therefore 175 &= \pi \left( -\frac{h^3}{3} + 3h^2 + 4h \right) \quad \checkmark a \quad \checkmark \text{m substitution} \\
 \therefore -h^3 + 9h^2 + 12h - \frac{525}{\pi} &= 0 \quad \checkmark ca \\
 \therefore h &= 5,28 \text{ cm} \quad \checkmark ca
 \end{aligned}$$

If double up Max 8  
B/D  
 Max of 9  
 for use

[12]

Total: 200 marks

alternative for Q11

$$\begin{aligned}
 & f = g(x) = \ln(x) + 1 \quad y = \ln(x) \\
 & u = \frac{2x-1}{x} = \frac{2}{x} - 1 \quad w = \frac{2x-1}{x} = \frac{2}{x} - 1 \\
 & h_1 = f\left(1 + \frac{2}{n}\right) \quad h_2 = f\left(1 + \frac{2}{n}\right) \\
 & = f\left(1 + \frac{2}{n}\right) + f\left(1 + \frac{2}{n}\right) + 1 = f\left(1 + \frac{2}{n}\right) \\
 & S_4 = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left( f\left(1 + \frac{2}{n}\right) + f\left(1 + \frac{2}{n}\right) + 1 \right) = f\left(1 + \frac{2}{n}\right) \\
 & S_4 = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left[ \frac{6k}{n} + \frac{36k}{n^2} + \frac{6}{n} \right] \quad (*) \\
 & S_4 = \lim_{n \rightarrow \infty} \left[ \frac{6k}{n} \cdot n + \frac{6}{n} \cdot n + \frac{36k}{n^2} \left( \frac{n^2}{2} + n \right) \right] \quad \checkmark \\
 & S_4 = \lim_{n \rightarrow \infty} \left[ 6k + 6 + 18k + \frac{36k}{n} \right] \quad \checkmark \\
 & S_4 = 6k + 6 + 18k \\
 & 6k = 24k \\
 & 2 = k \quad \checkmark
 \end{aligned}$$

12

Q